

SMITH | LANGSFORD WILLING | BARNES | UTBER

JACARANDA MATHS QUEST 9

AUSTRALIAN CURRICULUM | FIFTH EDITION

AUSTRALIAN
CURRICULUM
v9.0



learn on

www.jacplus.com.au

jacaranda
A Wiley Brand

JACARANDA
MATHS QUEST 9

AUSTRALIAN CURRICULUM | FIFTH EDITION

JACARANDA MATHS QUEST 9

AUSTRALIAN CURRICULUM | FIFTH EDITION

CATHERINE SMITH

BEVERLY LANGSFORD WILLING

MARK BARNES

CHRISTINE UTBER

CONTRIBUTING AUTHOR:

James Smart | Geetha James | Caitlin Mahony | Rebecca Charlesworth
Michael Sheedy | Kahni Burrows | Paul Menta



jacaranda
A Wiley Brand

Fifth edition published 2023 by
John Wiley & Sons Australia, Ltd
Level 4, 600 Bourke Street, Melbourne, Vic 3000

First edition published 2011
Second edition published 2014
Third edition published 2018
Fourth edition published 2021

Typeset in 10.5/13 pt TimesLT Std

© John Wiley & Sons Australia, Ltd 2011, 2014, 2018,
2021, 2023

The moral rights of the authors have been asserted.

ISBN: 978-1-394-19408-7

Reproduction and communication for educational purposes

The Australian *Copyright Act 1968* (the Act) allows a maximum of one chapter or 10% of the pages of this work, whichever is the greater, to be reproduced and/or communicated by any educational institution for its educational purposes provided that the educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL).

Reproduction and communication for other purposes

Except as permitted under the Act (for example, a fair dealing for the purposes of study, research, criticism or review), no part of this book may be reproduced, stored in a retrieval system, communicated or transmitted in any form or by any means without prior written permission. All inquiries should be made to the publisher.

Trademarks

Jacaranda, the JacPLUS logo, the learnON, assessON and studyON logos, Wiley and the Wiley logo, and any related trade dress are trademarks or registered trademarks of John Wiley & Sons Inc. and/or its affiliates in the United States, Australia and in other countries, and may not be used without written permission. All other trademarks are the property of their respective owners.

Front cover images: © AlexanderTrou/Shutterstock; © Marish/Shutterstock; © alexdndz/Shutterstock; © Vector Juice/Shutterstock

Illustrated by various artists, diacriTech and Wiley Composition Services

Typeset in India by diacriTech



A catalogue record for this
book is available from the
National Library of Australia

The Publishers of this series acknowledge and pay their respects to Aboriginal Peoples and Torres Strait Islander Peoples as the traditional custodians of the land on which this resource was produced.

This suite of resources may include references to (including names, images, footage or voices of) people of Aboriginal and/or Torres Strait Islander heritage who are deceased. These images and references have been included to help Australian students from all cultural backgrounds develop a better understanding of Aboriginal and Torres Strait Islander Peoples' history, culture and lived experience.

It is strongly recommended that teachers examine resources on topics related to Aboriginal and/or Torres Strait Islander Cultures and Peoples to assess their suitability for their own specific class and school context. It is also recommended that teachers know and follow the guidelines laid down by the relevant educational authorities and local Elders or community advisors regarding content about all First Nations Peoples.

All activities in this resource have been written with the safety of both teacher and student in mind. Some, however, involve physical activity or the use of equipment or tools. **All due care should be taken when performing such activities.** To the maximum extent permitted by law, the author and publisher disclaim all responsibility and liability for any injury or loss that may be sustained when completing activities described in this resource.

The Publisher acknowledges ongoing discussions related to gender-based population data. At the time of publishing, there was insufficient data available to allow for the meaningful analysis of trends and patterns to broaden our discussion of demographics beyond male and female gender identification.

Contents

About this resource.....	vii
Acknowledgements.....	xiv

NAPLAN practice

online only

- Set A** Calculator allowed
- Set B** Non-calculator
- Set C** Calculator allowed
- Set D** Non-calculator
- Set E** Calculator allowed
- Set F** Non-calculator

1	Number skills and index laws	1
1.1	Overview	2
1.2	Real numbers	4
1.3	Rounding numbers and significant figures	11
1.4	Review of index laws	16
1.5	Raising powers	24
1.6	Negative indices	28
1.7	Scientific notation	34
1.8	Square roots and cube roots	41
1.9	Surds (extending)	44
1.10	Review	56
	Answers	64
2	Algebra	71
2.1	Overview	72
2.2	Using pronumerals	74
2.3	Algebra in worded problems	83
2.4	Simplification of algebraic expressions	88
2.5	Expanding brackets	95
2.6	Difference of two squares and perfect squares	104
2.7	Further expansions	109
2.8	The highest common factor	112
2.9	The highest common binomial factor	118
2.10	Solving worded problems	122
2.11	Review	131
	Answers	138

3	Linear equations	147
3.1	Overview	148
3.2	Solving linear equations and inequations	150
3.3	Solving linear equations with brackets	161
3.4	Solving linear equations with pronumerals on both sides	165
3.5	Solving problems with linear equations	169
3.6	Rearranging formulas	175
3.7	Review	181
	Answers	187
4	Congruence and similarity	191
4.1	Overview	192
4.2	Ratio and scale	195
4.3	Congruent figures	204
4.4	Nets, polyhedra and Euler's formula	212
4.5	Similar figures	217
4.6	Area and volume of similar figures	229
4.7	Review	238
	Answers	247
5	Pythagoras and trigonometry	253
5.1	Overview	254
5.2	Pythagoras' theorem	258
5.3	Applications of Pythagoras' theorem	267
5.4	Trigonometric ratios	274
5.5	Calculating unknown side lengths	282
5.6	Calculating unknown angles	291
5.7	Angles of elevation and depression	299
5.8	Review	305
	Answers	315
6	Linear and non-linear graphs	321
6.1	Overview	322
6.2	Plotting linear graphs	325
6.3	Features of linear graphs	331
6.4	The equation of a straight line	339
6.5	Sketching linear graphs	348
6.6	Technology and linear graphs	356
6.7	Practical applications of linear graphs	360
6.8	Midpoint of a line segment and distance between two points	365
6.9	Non-linear relations	371
6.10	Review	385
	Answers	393

Semester review 1	online only	410
--------------------------	--------------------	------------

7 Proportion and rates 413

7.1 Overview	414
7.2 Direct linear proportion	416
7.3 Proportionality	421
7.4 Introduction to rates	426
7.5 Constant and variable rates	429
7.6 Review	436
Answers	444

8 Financial mathematics (extending) 447

8.1 Overview	448
8.2 Salaries and wages	451
8.3 Penalty rates	455
8.4 Simple interest	461
8.5 Compound interest	469
8.6 Review	478
Answers	483

9 Measurement 487

9.1 Overview	488
9.2 Measurement	492
9.3 Area	501
9.4 Area and perimeter of a sector	512
9.5 Surface area of rectangular and triangular prisms	519
9.6 Surface area of a cylinder	525
9.7 Volume of prisms and cylinders	530
9.8 Review	543
Answers	552

10 Probability 557

10.1 Overview	558
10.2 Theoretical probability	564
10.3 Experimental probability	571
10.4 Venn diagrams and two-way tables	580
10.5 Compound events	597
10.6 Review	610
Answers	617

11 Statistics 625

11.1 Overview	626
11.2 Collecting data	630
11.3 Organising data	639
11.4 Displaying data	649
11.5 Measures of centre	664
11.6 Measures of spread	680
11.7 Review	689
Answers	695

12 Quadratic equations and graphs 707

12.1 Overview	708
12.2 Quadratic equations (extending)	710
12.3 The Null Factor Law (extending)	715
12.4 Solving quadratic equations with two terms (extending)	719
12.5 Factorising and solving monic quadratic equations (extending)	723
12.6 Graphs of quadratic functions	729
12.7 Sketching parabolas of the form $y = ax^2$	738
12.8 Sketching parabolas of the form $y = (x - h)^2 + k$	744
12.9 Sketching parabolas of the form $y = (x + a)(x + b)$ (extending)	756
12.10 Applications	760
12.11 Review	766
Answers	775


Semester review 2 **online only** 800

13 Algorithmic thinking **online only**

13.1 Overview
13.2 Programs
13.3 Arrays
13.4 Loops
13.5 Set structures
13.6 Sorting algorithms
13.7 The Monte Carlo method
13.8 Review
Answers

Glossary	802
Index	808

About this resource



NEW FOR

AUSTRALIAN CURRICULUM V9.0



JACARANDA MATHS QUEST 9

AUSTRALIAN CURRICULUM
FIFTH EDITION

Developed by teachers for students

Tried, tested and trusted. The fifth edition of the *Jacaranda Maths Quest* series, revised fourth edition, continues to focus on helping teachers achieve learning success for every student – ensuring no student is left behind, and no student is held back.

Because both what and how students learn matter



Learning is personal

Whether students need a challenge or a helping hand, you'll find what you need to create engaging lessons.

Whether in class or at home, students can get unstuck and progress! Scaffolded lessons, with detailed worked examples, are all supported by teacher-led video eLessons. Automatically marked, differentiated question sets are all supported by detailed worked solutions. And Brand-new Quick Quizzes support in-depth skill acquisition.



Learning is effortful

Learning happens when students push themselves. With learnON, Australia's most powerful online learning platform, students can challenge themselves, build confidence and ultimately achieve success.



Learning is rewarding

Through real-time results data, students can track and monitor their own progress and easily identify areas of strength and weakness.

And for teachers, Learning Analytics provide valuable insights to support student growth and drive informed intervention strategies.

Learn online with Australia's most

Everything you need
for each of your lessons
in one simple view

- Trusted, curriculum-aligned content
- Engaging, rich multimedia
- All the teacher support resources you need
- Deep insights into progress
- Immediate feedback for students
- Create custom assignments in just a few clicks.

Practical teaching advice and ideas for each lesson provided in teachON

Teaching videos for all lessons

Reading content and rich media including embedded videos and interactivities

The screenshot displays the learnON interface for Jacaranda Maths Quest 7 AC 5e. The main content area is titled "3.2 The real number system" and includes a "LEARNING INTENTION" section with the following points:

- At the end of this subtopic you should be able to:
- classify real numbers as irrational or rational
- classify rational numbers as integers or non-integers
- place real numbers on a number line.

Below this is section "3.2.1 Introduction to real numbers" with a bullet point: "The real number system contains the set of rational and irrational numbers. The real number system is denoted by the symbol R ." An "Interactivity" section titled "Real numbers" follows, with instructions: "Drag and drop each number into its appropriate real number category. Select the Submit button to check your work." The numbers to be categorized are $\sqrt{5}$, -5 , 0 , 2 , and 0.375 . The categories are: Irrational numbers, Non-integer rationals, Zero, Negative integers, and Positive integers. Each category has a "Drop here" button. At the bottom of the interactivity are "Submit" and "Try again" buttons.

On the right side, there is a sidebar with "3.2 Ex 1" and "SELECT YOUR" options. Below that, there are two questions: "Q1 Place the following numbers on a number line. a. -4 " and "b. $\frac{1}{4}$ ". Each question has a number line and a "SUBMIT ANSWER" button.

At the bottom left of the interface, there is a "3.1 OVERVIEW" button.

powerful learning tool, learnON

The image shows a screenshot of the learnON software interface. The interface is divided into several sections: a top navigation bar with 'Teacher' and 'Student' tabs, a main content area with '3.2 Ex 2' and '3.2 TBQ' tabs, and a 'RESOURCES' sidebar. The 'RESOURCES' sidebar lists various content types with counts and dropdown menus: Topic PDF (1), eWorkbook (29), Solutions (1), Digital documents (1), Video eLessons (8), Interactivities (17), Weblinks (6), TEACHER eWorkbook (26), TEACHER Digital documents (9), and TEACHER Weblinks (2). The main content area shows a 'PRACTISE' section with 'CONSOLIDATE' and 'MASTER' sub-sections, each containing a list of question numbers. Below this is a number line with a point marked at 1 and a '1 mark(s)' indicator. The bottom of the screen shows '3.3 ADDING AND SUBTRACTING FRACTIONS' with a right-pointing arrow.

Teacher **Student** Year7A-Jacaranda-VC-... Help Samwell Tarty

3.2 Ex 2 3.2 TBQ 1

RESOURCES

- Topic PDF 1
- eWorkbook 29
- Solutions 1
- Digital documents 1
- Video eLessons 8
- Interactivities 17
- Weblinks 6
- TEACHER eWorkbook 26
- TEACHER Digital documents 9
- TEACHER Weblinks 2

PRACTISE

CONSOLIDATE

Questions: 1, 6, 9, 12, 14, 17, 18, 21, 24, 26

MASTER

Questions: 2, 7, 10, 15, 22, 23, 25, 28

1 mark(s)

3.3 ADDING AND SUBTRACTING FRACTIONS

New! Quick Quiz questions for skill acquisition

Differentiated question sets

Teacher and student views

Textbook questions

Fully worked solutions

eWorkbook

Digital documents

Video eLessons

Interactivities

Extra teaching support resources


Interactive questions with immediate feedback

Get the most from your online resources

Online, these new editions are the complete package


Trusted Jacaranda theory, plus tools to support teaching and make learning more engaging, personalised and visible.

Embedded interactivities and videos enable students to explore concepts and learn deeply by 'doing'.






New teaching videos for every lesson are designed to help students learn concepts by having a 'teacher at home', and are flexible enough to be used for pre- and post-learning, flipped classrooms, class discussions, remediation and more.

Brand new! Quick Quiz questions for skill acquisition in every lesson.




Three differentiated question sets, with immediate feedback in every lesson, enable students to challenge themselves at their own level. Instant reports give students visibility into progress and performance.



Every question has immediate, corrective feedback to help students overcome misconceptions as they occur and get unstuck as they study independently – in class and at home.


Maths Quest 7 includes Powering up for Year 7

A six-week 'Powering up for Year 7' online program is designed to plug any gaps from earlier years.




NAPLAN Online Practice

Go online to complete practice NAPLAN tests. There are 6 NAPLAN-style question sets available to help you prepare for this important event. They are also useful for practising your Mathematics skills in general.




Also available online is a video that provides strategies and tips to help with your preparation.

eWorkbook



The eWorkbook enables teachers and students to download additional activities to support deeper learning.

A wealth of teacher resources




Enhanced teacher support resources for every lesson, including:

- work programs and curriculum grids
- practical teaching advice
- three levels of differentiated teaching programs
- quarantined topic tests (with solutions)

Customise and assign


An inbuilt testmaker enables you to create custom assignments and tests from the complete bank of thousands of questions for immediate, spaced and mixed practice.



Reports and results

Data analytics and instant reports provide data-driven insights into progress and performance within each lesson and across the entire course.

Show students (and their parents or carers) their own assessment data in fine detail. You can filter their results to identify areas of strength and weakness.



Acknowledgements

The authors and publisher would like to thank the following copyright holders, organisations and individuals for their assistance and for permission to reproduce copyright material in this book.

© Australian Curriculum, Assessment and Reporting Authority (ACARA) 2010 to present, unless otherwise indicated. This material was downloaded from the ACARA website (www.acara.edu.au) (**Website**) (accessed January, 2023) and was not modified. The material is licensed under CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>). Version updates are tracked in the 'Curriculum version history' section on the 'About the Australian Curriculum' page (<https://www.australiancurriculum.edu.au/about-the-australian-curriculum/>) of the Australian Curriculum website.

ACARA does not endorse any product that uses the Australian Curriculum or make any representations as to the quality of such products. Any product that uses material published on this website should not be taken to be affiliated with ACARA or have the sponsorship or approval of ACARA. It is up to each person to make their own assessment of the product, taking into account matters including, but not limited to, the version number and the degree to which the materials align with the content descriptions and achievement standards (where relevant). Where there is a claim of alignment, it is important to check that the materials align with the content descriptions and achievement standards (endorsed by all education Ministers), not the elaborations (examples provided by ACARA).

Images

• © Aila Images/Shutterstock: **10** • © Albachiaraa/Shutterstock: **512** • © Alex Mit/Shutterstock: **179** • © ALPA PROD/Shutterstock: **460** • © BlueBarronPhoto/Shutterstock: **64** • © Brian A Jackson/Shutterstock: **4** • © Business stock/Shutterstock: **461** • © CGN089/Shutterstock: **55** • © ClaudioValdes/Shutterstock: **160** • © Creativa Images/Shutterstock: **450** • © Daxiao Productions/Shutterstock: **174** • © Dean Drobot/Shutterstock: **90** • © DenisProduction.com/Shutterstock: **180** • © Dima Moroz/Shutterstock: **261** • © Diyana Dimitrova/Shutterstock: **64** • © djgis/Shutterstock: **370** • © Dragon Images/Shutterstock: **454** • © Drazen Zigic/Shutterstock: **458** • © Fablok/Shutterstock: **40** • © fizkes/Shutterstock: **477** • © Galina Barskaya/Shutterstock: **81** • © Ground Picture/Shutterstock: **175, 456** • © Harvepino/Shutterstock: **462** • © hidesy/Shutterstock: **448** • © ITTIGallery/Shutterstock: **469** • © Jiripravda/Shutterstock: **569** • © jocic/Shutterstock: **15** • © Kuki Ladron de Guevara/Shutterstock: **15** • © Lightspring/Shutterstock: **311** • © Marques/Shutterstock: **53** • © Merydolla/Shutterstock: **192** • © michaeljung/Shutterstock: **460** • © Mila Supinskaya Glashchenko/Shutterstock: **168** • © Monkey Business Images/Shutterstock: **56, 85, 454, 468** • © New Africa/Shutterstock: **464** • © NikoNomad/Shutterstock: **40, 63** • © Nilobon Sweeney/Shutterstock: **15** • © Norman Pogson/Shutterstock: **254** • © pikselstock/Shutterstock: **449** • © Pius Lee/Shutterstock.com: **307** • © PlusONE/Shutterstock: **236** • © Potstock/Shutterstock: **164** • © puhhha/Shutterstock: **2** • © rachel ko/Shutterstock: **169** • © Rawpixel.com/Shutterstock: **137, 185** • © REDPIXEL.PL/Shutterstock: **468** • © Robert Kneschke/Shutterstock: **21** • © Sergey Nivens/Shutterstock: **36** • © Serhiy Stakhnyk/Shutterstock: **148** • © Shutterstock: **203** • © Shutterstock/Bankoo: **72** • © Skylines/Shutterstock: **125** • © Sofiaworld/Shutterstock: **233** • © Stuart Monk/Shutterstock: **164** • © szefei/Shutterstock: **476** • © ThomasLENNE/Shutterstock: **122** • © UfaBizPhoto/Shutterstock: **469** • © Walter Cicchetti/Shutterstock: **137** • © wavebreakmedia/Shutterstock: **42, 186** • © XiXinXing/Shutterstock: **459** • © John Wiley & Sons Australia: **569, 570** • © Adisa/Shutterstock: **637** • © Africa Studio/Shutterstock: **577** • © Alan Poulson Photography/Shutterstock: **612** • © Alex James Bramwell/Shutterstock: **613** • © Alexander Rath/Shutterstock: **774** • © AlexanderTrou/Shutterstock: Marish/Shutterstock, vectorstudi/Shutterstock, **1, 71, 147, 191, 253, 321, 413, 447, 487, 557, 625** • © AlexanderTrou/Shutterstock:Marish/Shutterstock, vectorstudi/Shutterstock, **707** • © alexsl/Getty Images: **281** • © Analia Aguilar Camacho/Shutterstock: **525** • © andrea crisante/Shutterstock: **523** • © Andres/Shutterstock: **359, 592** • © Andrii Yalanskyi/Shutterstock: **426** • © anek.soowannaphoom/Shutterstock: **631** • © Anel Alijagic/Shutterstock: **171** • © antoniodiaz/Shutterstock: **453** • © Aquarius Studio/Shutterstock: **664** • © ARTvektor/Shutterstock: **609** • © Ashley Whitworth/Shutterstock: **509** • © Barney Magrath/NASA: **41** • © bikeriderlondon/Shutterstock: **568** • © Binikins/Shutterstock: **480** • © Bo Valentino/Shutterstock: **586** • © cassiede alain/Shutterstock: **743** • © Chatchai Somwat/Shutterstock: **641** • © CHRISTOPHE ROLLAND/Shutterstock: **678** • © chrupka/Shutterstock: **606** • © Ciprian Vladut/Shutterstock: **322** • © Corbis: **763** • © Dario Lo Presti/Shutterstock: **392** • © David H.Seymour/Shutterstock: **540** • © defpicture/Shutterstock: **511** • © Desmos: Inc., **750, 751** • © Dimedrol68/Shutterstock: **424** • © Dmitry Natashin/Shutterstock: **55** • © dotshock/Shutterstock: **764** • © Edw/Shutterstock: **183** • © Eric Gevaert/Shutterstock: **438**

• © ESB Professional/Shutterstock: **419, 467** • © freesoulproduction/Shutterstock: **29** • © Galina Barskaya/Shutterstock: **77**
 • © Hallgerd/Shutterstock: **239** • © hddigital/Shutterstock: **479** • © Henri et George/Shutterstock: **183** • © ILYA
 GENKIN/Shutterstock: **364** • © Image Addict: **549** • © inspiredbyart/Shutterstock: **423** • © J.D.S/Shutterstock: **125** • ©
 Janaka Dharmasena/Shutterstock: **414** • © Jennifer Wright/John Wiley & Sons Australia: **528** • ©
 jessicakirsh/Shutterstock: **612** • © Jesus Cervantes/Shutterstock: **680** • © jiris/Shutterstock: **23** • © John Wiley & Sons
 Australia: **87, 103** • © Juergen Faelchle/Shutterstock: **500** • © Kahan Shan/Shutterstock: **417** • © Kaspars
 Grinvalds/Shutterstock: **418** • © Khakimullin Aleksandr/Shutterstock: **182** • © Kitch Bain/Shutterstock: **598** • ©
 ksy9/Shutterstock: **765** • © kurhan/Shutterstock: **179** • © Laitr Keiows/Shutterstock: **499** • © Larry-Rains/Shutterstock: **129**
 • © Leah-Anne Thompson/Shutterstock: **392** • © leolintang/Shutterstock: **434** • © Lerner Vadim/Shutterstock: **372** • ©
 lucadp/Shutterstock: **524** • © LuckyImages/Shutterstock: **658** • © Luna Vandoorne/Shutterstock: **174** • © Lyn Elms/John
 Wiley & Sons Australia: **307** • © Magdalena Kucova/Shutterstock: **488** • © Maksim Kabakou/Shutterstock: **129** • ©
 Martin Charles Hatch/Shutterstock: **688** • © Maxisport/Shutterstock: **228** • © metamorworks/Shutterstock: **558** • © Michael
 William/Shutterstock: **171** • © michaelangelus/Getty Images: **773** • © MilanB/Shutterstock: **217** • © Mmaxer/Shutterstock:
82 • © Monkey Business Images/Shutterstock: racorn/Shutterstock, Kari-Ann Tapp/John Wiley & Sons Australia, **728** •
 © monticello/Shutterstock: **596** • © Nata-Lia/Shutterstock: **540** • © Naypong Studio/Shutterstock: **203** • © Neale
 Cousland/Shutterstock: **86** • © Nikodash/Shutterstock: **498** • © Olaf Speier/Shutterstock: **566** • © OSTILL is Franck
 Camhi/Shutterstock: **774** • © OZBEACHES/Shutterstock: **708** • © PavleMarjanovic/Shutterstock: **773** • ©
 PeopleImages/E+/Getty ImagesDean Drobot/Shutterstock: **451** • © Petr Toman/Shutterstock: **428** • © Philip
 Yuan/Shutterstock: **202** • © photka/Shutterstock: **416** • © PhotoDisc: **216, 285, 289, 539, 541** • © Phovoir/Shutterstock: **302**
 • © pics4sale/Shutterstock: **512** • © PinkBlue/Shutterstock: **524** • © Pressmaster/Shutterstock: **768** • ©
 przyzmat/Shutterstock: **363** • © quka/Shutterstock: **197** • © Radu Bercan/Shutterstock: **759** • © Ramon grosso
 dolarea/Shutterstock: **86** • © Rawpixel.com/Shutterstock: **626** • © Redshinestudio/Shutterstock: **169** • © Renee Bryon/John
 Wiley & Sons: **298** • © Renee Bryon/John Wiley & Sons Australia: **272, 600** • © Renee Bryon/John Wiley &
 Sons Australia: **94, 245** • © Robyn Mackenzie/Shutterstock: **422, 598** • © Roma Koschel/Shutterstock: **565** • ©
 s74/Shutterstock: **768** • © Samuel Borges Photography/Shutterstock: **433** • © SensorSpot/Getty Images: **660** • © SERGEI
 PRIMAKOV/Shutterstock: **425** • © Shenjun Zhang/Shutterstock: **755** • © Sinisa Botas/Shutterstock: **578** • ©
 smallblackcat/Shutterstock: **227** • © Sonja Foos/Shutterstock: **662** • © Source: Wiley Art: **297** • © Stefan
 Schurr/Shutterstock: **437** • © StepStock/Shutterstock: **719** • © StockPhotosLV/Shutterstock: **330** • ©
 sunsetman/Shutterstock: **428** • © Suzanne Tucker/Shutterstock: **179, 608** • © Taras Vyshnya/Shutterstock: **425** • © Texas
 Instruments: **356, 357** • © Tom Hirtreiter/Shutterstock: **442** • © Torderiul/Shutterstock: **518** • © Tyler Olson/Shutterstock:
338 • © Val Thoermer/Shutterstock: **578** • © Vitalii Vodolazskiy/Shutterstock Rtimages/Shutterstock: **456** • ©
 Vitalinka/Shutterstock: **606** • © Vladitto/Shutterstock: **363** • © Volt Collection/Shutterstock: **262** • ©
 wavebreakmedia/Shutterstock: **571** • © weknow/Shutterstock: **133** • © Wiley art: **123** • © WilleeCole
 Photography/Shutterstock: **240** • © Worytko Pawel/Shutterstock: **612** • © Yellow duck/Shutterstock: pattang/Shutterstock,
 Denis Tabler/Shutterstock, Nattika/Shutterstock, rangizzz/Shutterstock, **630** • © Yermek/Shutterstock: **691** • ©
 Avesun/Shutterstock: **271** • © CASIO: **221–223, 283, 284, 292, 326, 327, 341, 366, 368, 373, 377, 378, 380, 381, 403, 404,**
494, 495, 574, 712, 717, 725, 733, 750, 762 • © ChiccoDodiFC/Shutterstock: **290** • © Christina Richards/Shutterstock: **298**
 • © Haali/Shutterstock: **58, 182, 239, 306, 386, 479, 544** • © John Wiley & Sons Australia: **103** • ©
 LightSecond/Shutterstock: **338** • © PhotoDisc: **298** • © Polka Dot Images: **298** • © Renee Bryon/John Wiley & Sons
 Australia: **271, 290** • © Texas Instruments: **221–223, 326, 327, 341, 368, 373, 377, 378, 380, 381, 403, 404, 494–496, 574,**
575, 712, 717, 725, 733, 750, 762 • © Texas Instruments Incorporated: **283, 284, 292**


Every effort has been made to trace the ownership of copyright material. Information that will enable the publisher to rectify any error or omission in subsequent reprints will be welcome. In such cases, please contact the Permissions Section of John Wiley & Sons Australia, Ltd.

NAPLAN practice

online only

Go online to complete practice NAPLAN tests. There are 6 NAPLAN-style question sets available to help you prepare for this important event. They are also useful for practising your Mathematics skills in general.

Also available online is a video that provides strategies and tips to help with your preparation.



SET A
Calculator allowed

SET B
Non-calculator

SET C
Calculator allowed

SET D
Non-calculator

SET E
Calculator allowed

SET F
Non-calculator

NAPLAN practice


Overview

Why learn this?

Our lives are interwoven with mathematics. Counting, measuring and pattern-making are all part of everyday life. We use numbers to mark significant events (such as birthdays) and for identification (such as passports and credit cards). We use numbers to describe ourselves (for example our height and weight). Shopping involves understanding numbers, and tallying scores in sports requires a comparison of numbers. You may not realise just how much you rely on numbers.



on Resources


 **Video eLesson** NAPLAN: Strategies and tips (eles-1688)

Set A

Calculator allowed

- Which of the following is the longest time?
A. 95 000 seconds B. 22 hours C. 1800 minutes D. 1 day
- Anne bought 20 pens and Jo bought 30 pens. Each pen costs the same amount, and the pens cost \$75 altogether. How much did Jo pay for her pens?
- When a ball is dropped it bounces on the ground. The maximum height reached after each bounce is half the distance the ball drops. If a ball is dropped from a ladder 6 m above the ground, what is the total distance travelled by the time the ball hits the ground the third time?
A. 9 m B. 18 m C. 10.5 m D. 15 m
- Last year at Fern Hill High School, there were 985 students. The student population has increased by 20% this year. How many students are at Fern Hill High School this year?
A. 1000 B. 1182 C. 1218 D. 1320
- The length of a rectangle is 9 cm and its perimeter is 30 cm. What is the area of the rectangle?

- The graph shows the number of juices sold each day of a particular week at a school. Calculate the mean daily number of juices sold at the school that week.



- $(10^2 + 12 \div 3) \div (-8) = ?$
A. 13 B. 14
C. -13 D. -14
- $\frac{3}{8}$ expressed as a percentage is:
A. 30.8% B. 13%
C. 37.5% D. 0.375%

- Lena wants to make 2 chocolate cakes. The recipe for a chocolate cake is:

$1\frac{1}{2}$ cups self-raising flour

$\frac{1}{2}$ cup sugar

2 tablespoons cocoa

1 egg


125 grams butter.

How much of each ingredient will she need to use?

- cups self-raising flour
- cup sugar
- tablespoons cocoa
- eggs
- grams butter



- In the diagram shown, what is the value of x ?
A. 110° B. 105° C. 120° D. 100°
- Express in binary form.
 113_{ten}



12. Jen bought a car on sale for 15% off the original price, which was \$12 500. How much did she pay for the car?

- A. \$1875 B. \$8505 C. \$10 625 D. \$10 510




13. Connor climbs to the summit of a mountain 4250 m above sea level. Inga is scuba diving at the bottom of the ocean. The ocean floor is 27 metres below sea level. What is the vertical distance between Connor and Inga?



14. The ratio of cordial to water in Shinji's favourite drink is 2 : 5. How much water does he need to add to 80 mL of cordial to make his favourite drink?

- A. 100 mL B. 200 mL C. 300 mL D. 400 mL


15. Which of the following represents the right side view of the 3-dimensional figure shown?



16. A rectangular paddock is surrounded by a fence. The perimeter of the paddock is 870 m. If the length of the paddock is 65 m longer than its width, calculate the dimensions of the paddock.

17. Calculate the area of material required to make the tent shown, including the floor.

- A. 12 m² B. 12.5 m²
C. 12.45 m² D. 24.45 m²



18. Write the rule for the following pattern made from sticks.

Let n = number of sticks.
Let s = number of squares.




- A. $n = 2s + 2$ B. $n = 3s + 1$ C. $n = 4s$ D. $n = s + 3$

19. If 1 teaspoon = 5 mL and 1 tablespoon = 20 mL, how many teaspoons are there in a tablespoon?

- A. 3 B. 4 C. 5 D. 2

20. The total surface area (TSA) of a cylinder can be found using the formula $TSA = 2\pi r(r + h)$ where r is the radius and h is the height of the cylinder. Find h when $r = 4$ cm and the TSA is 377 cm².



21. A car travelled at 60 kilometres per hour for 2 hours and 15 minutes. How far did it travel?

- A. 165 km B. 120 km C. 75 km D. 135 km

22. The table below shows the times for 4 students to run 100 m.

Student	Mitch	Josh	Ben	Liam
Time (s)	14.2	15.4	14.8	14.4

What is the average (mean) time the 4 students ran 100 m?


23. Which of the following is the smallest number?

- A. $\sqrt[3]{6}$ B. 0.18 C. 0.5^2 D. $\frac{3}{8}$





24. Complete the following to make this equation true. $13 \times (15 - 1) = 180 + ?$

25. The diagram shows a bag of tokens. Aneko chooses a token from the bag without looking. What is the chance of Aneko choosing a star token?


- A. $\frac{3}{10}$ B. $\frac{1}{3}$ C. $\frac{1}{4}$ D. $\frac{1}{2}$



26. Which of the following diagrams shows an equilateral triangle?



27. What is the volume of this rectangular prism?




28. What is the longest pencil that can fit inside the box shown? (Round to 1 decimal place.)

- A. 26.9 mm B. 25 mm C. 22.4 mm D. 24.6 mm

29. Which coordinates will the graph of $y = -2x + 1$ pass through?

- A. (5, 11) B. (5, -9) C. (1, 3) D. (1, -3)



30. If $a = -3$, then $3a^3$ equals:

- A. 27 B. -27 C. 81 D. -81

Set B

Non-calculator

1. How many faces does a rectangular prism have?

- A. 4 B. 6
C. 8 D. 12

2. Oranges sell for \$2.50 per kilogram. What is the total cost of 3 kg of oranges?

- A. \$2.50 B. \$5.00
C. \$4.50 D. \$7.50



3. A fair, six-sided die is rolled. What is the chance of it landing on a 2?

- A. $\frac{1}{6}$ B. $\frac{1}{3}$ C. $\frac{1}{2}$ D. 2


4. $7 + (2 \times 5) - 9 = ?$

- A. 36 B. 17 C. 8 D. 7

5. Evaluate 1.3×0.6 .

6. What is the equation of the graph shown?

- A. $y = x + 3$
B. $y = x - 3$
C. $y = 3x + 3$
D. $y = 3x - 3$



7. Draw a linear graph using the table of values below.

x	-3	-2	-1	0	1	2	3
y	2	1	0	-1	-2	-3	-4


8. Sivan recorded the colour of cars travelling along her street between 9 am and 10 am.

Colour	Blue	White	Red	Silver	Black	Gold
Number	1	6	2	5	3	3

What was the percentage of white cars?


9. What type of angle is this?

- A. Acute
- B. Obtuse
- C. Reflex
- D. Straight



10. What is the inequation that represents the interval shown on the number line?

- A. $1 \leq x \leq 6$
- B. $1 \leq x < 6$
- C. $1 \geq x \geq 6$
- D. $1 < x \leq 6$



11. A store is having a 25% off sale. What would be the reduced price of an item that was originally \$120?

- A. \$40
- B. \$30
- C. \$100
- D. \$90



12. Two friends share 24 lollies. One friend gets twice as many as the other. How many lollies do they each receive?

- A. 10 and 14
- B. 8 and 16
- C. 6 and 18
- D. 4 and 20

13. Calculate the product of $\frac{3}{4}$ and $\frac{5}{6}$.

- A. $1\frac{7}{12}$
- B. $\frac{9}{10}$
- C. $\frac{19}{12}$
- D. $\frac{5}{8}$

14. The sum of 3 consecutive odd numbers is 21. Which equation best represents this situation?


- A. $x + 3 = 21$
- B. $3x = 21$
- C. $3x + 6 = 21$
- D. $6x + 3 = 21$

15. Find the mean and the median of the data below. 2, 1, 5, 5, 4, 0, 2, 5, 2, 6

- A. Mean = 3, median = 2
- B. Mean = 3.2, median = 4
- C. Mean = 3, median = 3
- D. Mean = 3.2, median = 3

16. The Venn diagram shows the number of students in Year 9 who play basketball, tennis and hockey. Find $n(\text{tennis} \cap \text{hockey})$.

- A. 61
- B. 78
- C. 70
- D. 64



17. Evaluate $3^2 + 2^3 + 3^3$.

- A. 40
- B. 44
- C. 36
- D. 38

18. What is the missing number?


$$5000 + 700 + ? + 6 = 5746$$

19. A piece of string 30 cm long forms a rectangle. If the length of the rectangle formed is 10 cm, what is its width?

- A. 5 cm
- B. 10 cm
- C. 15 cm
- D. 20 cm


20. Which of the following is the top view of the 3-dimensional figure shown?

- A.
- B.
- C.
- D.




21. This ruler is used to measure the height of a teacher. How tall is the teacher?

- A. 1.6 m
- B. 1.75 m
- C. 1.075 m
- D. 175 m



22. A roof makes an angle of 40° with the ceiling. What is the value of x ?

- A. 50°
- B. 40°
- C. 45°
- D. 90°



23. The stem-and-leaf plot shows the number of goals scored in each netball match over a season. The range is:

- A. 48
- B. 50
- C. 14
- D. 42

Stem	Leaf
1	49
2	579
3	1248
4	04
5	47
6	2

24. A circular spinner with 5 equal-sized sectors is labelled from 1 to 5. The spinner is spun. What is the probability of it landing on an odd number?

- A. 3
- B. $\frac{1}{2}$
- C. $\frac{3}{5}$
- D. $\frac{1}{3}$

25. Write an expression to represent the following:
 P is added to the square of q and then the sum is divided by y .






26. If one hour of parking costs \$4.50, how much will it cost to park from 9 am until 2 pm?

- A. \$45
- B. \$22.50
- C. \$20
- D. \$16.50

27. A map has a scale showing that 1 cm on the map represents an actual distance of 2 km. Write the appropriate scale ratio for this map.


- A. 1:2
- B. 1:200
- C. 1:2000
- D. 1:200 000

28. The shape shown is rotated 90° in a clockwise direction around the black dot. Which of the following choices represents the result?






29. ABC is an isosceles triangle. What is the value of y ?

- A. 5
 B. 7
 C. 9
 D. 11



30. Which of the following is the net of a cube?



Set C

Calculator allowed

1. The cube root of 64 is between:

- A. 1^2 and 1^3
 B. 1^2 and 3^2
 C. 2^3 and 4^2
 D. 3^2 and 4^2

2. A javelin is thrown 3 times. The mean distance of the 3 throws is 55 m. The fourth throw reaches 63 m. What is the mean distance after 4 throws?

- A. 52 m
 B. 57 m
 C. 55 m
 D. 60 m

3. The monthly cost (C) of renting a mobile phone is given by the equation $C = 20 + 0.2x$, where x is the call time in minutes. If February's bill was \$120, the call time was:

- A. 300 minutes
 B. 100 minutes
 C. 500 minutes
 D. 450 minutes

4. A music store has two promotions running at the same time:

- Buy 2 CDs for \$26.95 each and get the third free.
- 30% off all CDs which normally retail at \$24.95 each.

If Emma wants to buy three CDs, what is the best price she can get?

- A. \$53.90
 B. \$52.40
 C. \$51.90
 D. \$54.50

5. A number is multiplied by 3, then 12 is subtracted. The final answer is the square of 6. What is the original number?

6. A painter quotes \$448 to paint the walls of a bedroom with the following dimensions:

length = 4 m

width = 3 m

height = 3.2m.

How much extra will he charge to paint the ceiling as well (at the same original cost per square metre)?


- A. \$100
 B. \$110
 C. \$120
 D. \$150

7. Solve the equation for q . $\frac{3-4q}{2} = \frac{1}{4}$



8. What is the equation of the graph shown?

- A. $y = -2x - 2$
- B. $y = 2x - 2$
- C. $y = -2x + 2$
- D. $y = 2x + 2$




9. For the triangle shown, x equals:

- A. 16
- B. 17
- C. 18
- D. 19

10. The scale $200\text{cm} \Leftrightarrow 4\text{m}$ simplifies to the ratio:

- A. 1 : 2
- B. 1 : 4
- C. 2 : 4
- D. 50 : 1




11. A house is bought for \$450 000 and sold for \$525 000. Express the profit as a percentage of the cost price. (Round to 2 decimal places.)

- A. 14.29%
- B. 16.67%
- C. 85.71%
- D. 16.7%


12. A 300-g block of chocolate is to be broken into its individual squares. What is the mass of 4 squares of chocolate?

- A. 10 g
- B. 16 g
- C. 5 g
- D. 20 g



13. The area of the composite shape shown is:

- A. 30cm^2
- B. 40cm^2
- C. 50cm^2
- D. 45cm^2



14. What is the next term in the following pattern? 1, 4, 9, 16, ...

- A. 20
- B. 25
- C. 29
- D. 36

15. A cyclist travels 15km in 30 minutes. The speed (in km/h) she is travelling is:

- A. 40 km/h
- B. 60 km/h
- C. 80 km/h
- D. 30 km/h

16. The intersection point for $y = 2x + 4$ and $x + y = 1$ is:

- A. $(-2, 1)$
- B. $(-1, -2)$
- C. $(-1, 2)$
- D. $(2, 1)$


17. The factorised form of $2x^2y^3 - 6xy^2$ is:

- A. $2xy^2(xy - 3)$
- B. $2xy(x^2y - 3)$
- C. $2xy(xy - 6)$
- D. $2x^2y(xy - 3)$



18. The graph shows the frequency of the number of bread rolls bought at a bakery. What is the total number of rolls bought?

- A. 200
- B. 188
- C. 150
- D. 168



19. What are the mean and median of the data shown in the table?

Score (x)	Frequency
1	1
2	2
3	3
4	4




- A. The mean is 3 and median is 4.
- B. The mean is 3 and median is 3.
- C. The mean is 4 and median is 3.
- D. The mean is 4 and median is 4.

20. Which number is exactly halfway between 3.25 and 4.75?

- A. 3.95
- B. 4.15
- C. 4.25
- D. 4

21. Write a fraction equivalent to 0.75.

22. Which diagram is the net of a square-based pyramid?




23. The table summarises how much time Lian spent on her Maths project. What was the average amount of time each day Lian worked on her project?

- A. 25 minutes
- B. 30 minutes
- C. 43 minutes
- D. 1 hour

Time spent on project	
Day	Time
Monday	30 minutes
Tuesday	15 minutes
Wednesday	$1\frac{1}{4}$ hours
Thursday	50 minutes
Friday	45 minutes


24. Draw a line of symmetry through the diagram.



25. If $x = -2$, what is the value of $\frac{2x}{3x-2}$?

- A. $\frac{1}{2}$
- B. $\frac{1}{4}$
- C. $\frac{2}{3}$
- D. $\frac{1}{3}$

26. A map of Fern Island is shown below.




What direction is the treasure from Point Shipwreck?

27. The shape at right is to be enlarged by a factor of 3.

The new area will be:

- A. 3 times the original area
- B. 6 times the original area
- C. 9 times the original area
- D. 12 times the original area



28. Complete the table of values for the rule $y = 3x^2$.

x	-3	-2	-1	0	1	2	3
y							

29. The population of a city is expected to increase by 2.5% each year for the next 5 years. If this city's current population is 300000, what is it expected to be in 5 years' time?

- A. 339422
- B. 755000
- C. 3750345
- D. 355406

30. Find the area of material (m^2) needed to make a circular cushion with a diameter of 110cm.

Note: Two pieces of material are cut out and sewn together to make the cushion.

- A. $2.5 m^2$
- B. $2.0 m^2$
- C. $1.9 m^2$
- D. $1.5 m^2$

Set D

Non-calculator

1. What is the equivalent fraction to $3\frac{2}{5}$?


- A. $\frac{17}{3}$
- B. $\frac{11}{5}$
- C. $\frac{17}{5}$
- D. $\frac{11}{2}$

2. The value of $\left[\left(\frac{1}{2}\right)^2 + 2\frac{1}{4}\right] \div \frac{5}{7}$ is:

- A. $\frac{25}{14}$
- B. $3\frac{1}{2}$
- C. $4\frac{1}{2}$
- D. $2\frac{1}{3}$


3. A large cube, shown at right, is painted blue. How many cubes, inside this large cube, have no painted faces?

- A. 8
- B. 16
- C. 36
- D. 64




4. A glass of water (250 mL) is poured from the jug. How much water is left in the jug?

- A. 1.75L
- B. 1.5L
- C. 1.25L
- D. 1L



5. Measure the size of the unknown angle, x , using the protractor.



- A. 125°
- B. 66°
- C. 54°
- D. 126°

6. $14.076 - 9.25$ equals:

- A. 4.826
- B. 4.51
- C. 4.926
- D. 0.482

7. Jake recorded his classmates' preferred ice-cream flavours in the table below.

Ice-cream flavour	Number of students
Chocolate	14
Vanilla	3
Strawberry	7
Choc-mint	6






What percentage of students prefers choc-mint flavoured ice-cream?

- A. 60%
- B. 20%
- C. 6%
- D. 25%


8. A car travels at 75 kilometres per hour. How far will it travel in 4 hours?

- A. 300 km
- B. 150 km
- C. 100 km
- D. 250 km

9. Which of the following diagrams shows the arrow below multiplied by a scale factor of $\frac{1}{2}$?




10. \overline{XZ} is 32 cm. \overline{XY} is a quarter of the distance of \overline{XZ} . Find \overline{YZ} .



- A. 8 cm B. 16 cm C. 20 cm D. 24 cm
11. How many sticks would be needed to make the 5th shape in this pattern?

- A. 10
B. 16
C. 28
D. 34



12. To convert from degrees Fahrenheit ($^{\circ}\text{F}$) to degrees Celsius ($^{\circ}\text{C}$), the following formula is used:


$$C = \frac{5}{9}(F - 32).$$

What is the Fahrenheit temperature equivalent to 35°C ?

- A. 86°F B. 95°F C. 70°F D. 100°F

13. The two points A and B are on which line?

- A. $y = 1 - x$ B. $y = x + 1$
C. $y = x - 1$ D. $y = 3 - x$




14. Solve the equation $\frac{4m}{5} + 7 = 11$.

- A. $m = 5$ B. $m = 4$
C. $m = -5$ D. $m = 2$


15. A football team played 22 games during a season. The team won 4 more games than it lost. How many games did the team win?

- A. 9 B. 11 C. 13 D. 14

16. $\angle BAD$ and $\angle BCD$ are supplementary angles. $\angle ABC$ is 130° . Find the size of $\angle ADC$.



17. Draw the reflection of the shape shown below.



18. If $p = 3$ and $q = -5$, evaluate $q^2 - 2p$.


- A. -31 B. -19 C. 21 D. 19

19. The area of a circle is $A = \pi r^2$. Which value best estimates the area of a circle with a radius of 5 cm?

- A. 75 cm^2 B. 15 cm^2 C. 50 cm^2 D. 7.5 cm^2

20. This dot plot shows the ages of people in a choir. What is the most common age of the choir members?

- A. 50 B. 58
C. 59 D. 60



21. What is the probability of getting Tails when a fair coin is tossed?

- A. 0 B. $\frac{1}{2}$
C. 1 D. $\frac{1}{3}$

22. Insert ' $<$ ' or ' $>$ ' to make this statement true. $4^3 _ 2^5$

23. The following scores were recorded in a test: $\frac{18}{20}$ and 88%. Which is the higher score?

24. The cost of two different books is in the ratio 4 : 5. If the more expensive book costs \$25, what does the cheaper book cost?

- A. \$5 B. \$10
C. \$15 D. \$20

25. Calculate the value of 3 m 75 cm + 5 m 36 cm.

- A. 8 m 95 cm B. 9 m 11 cm
C. 9 m 20 cm D. 8 m 11 cm



26. There are 60 squares in a chocolate block. If Stef eats $\frac{2}{3}$ of the block, how many squares are left?


- A. 30 B. 25 C. 20 D. 15

27. The length of a rectangular sporting field is four times its width. If the perimeter of the field is 200 m, find the length and width.


28. A bag of marbles contains 22 red, 20 yellow, 15 green and 18 blue marbles. Darcy chooses one marble without looking. What is the chance it will be green?

- A. $\frac{1}{3}$ B. $\frac{1}{4}$ C. $\frac{1}{5}$ D. $\frac{1}{6}$

29. Which pair of shapes fits together to make a cube?



30. What shape is revealed when a triangular prism is cut vertically?



- A. Triangle B. Square C. Rectangle D. Trapezium


Set E

Calculator allowed

1. The volume of a cylindrical glass with diameter 7.2 cm and height 15.4 cm is:


- A. 2 L B. 627 cm³
 C. 550 cm³ D. 750 mL

2. Use the diagram below to determine the values of x and y .



3. A square window is to be inserted into a square wall. The side length of the window is 70 cm. What is the area of the remaining wall?

- A. $39\,200\text{ m}^2$
 B. 3.92 m^2
 C. 4.41 m^2
 D. 6.8 m^2




4. The formula $y = \frac{3x-4}{5}$ rearranged to make x the subject is:

- A. $x = \frac{5y-3}{4}$ B. $x = \frac{4y-5}{3}$ C. $x = \frac{3y+4}{5}$ D. $x = \frac{5y+4}{3}$

5. How many hours are there in 4 weeks and 3 days?

6. The value of x in the diagram is equal to:

- A. 107°
 B. 110°
 C. 77°
 D. 90°




7. An unknown number is multiplied by 6 and then added to 4. The result is 46. Which equation best describes this?

- A. $6 + 4x = 46$ B. $46 + 6x = 4$ C. $6x + 4 = 46$ D. $6(x + 4) = 46$

8. The number line shown can be described by the equation:

- A. $x < -1$ B. $x \geq -1$ C. $x > -1$ D. $x < 1$



9. Arrange the following in descending order.

$2.202, 0.12, \left(\frac{1}{2}\right)^2, \sqrt{\frac{1}{4}}, 0.01$

10. A car uses 10 L of fuel to travel 80 km. How much fuel is needed to travel 400 km?

- A. 30 L B. 40 L C. 50 L D. 60 L

11. A length of rope is 7.5 m. It is to be cut into 2 pieces in the ratio of 2:3. What will be the two new lengths?

- A. 2.5 m and 5 m
 B. 3 m and 4.5 m
 C. 3.5 m and 4 m
 D. 1 m and 6.5 m



12. A 5-kg bag of potatoes cost \$7.99. The cost of loose potatoes is \$1.55 per kg. Is it cheaper to buy:


- A. 5 kg of loose potatoes?
 B. a 5-kg bag of potatoes?

13. Which diagram of the net of a cube has opposite faces that sum to 7?

- A. B. C. D.

14. Which of the following shapes is represented by the net shown at right?


- A. A triangular prism
- B. A triangular pyramid
- C. A square pyramid
- D. A cube



15. If $A = \{\text{odd numbers between 0 and 10}\}$ and $B = \{3, 6, 9\}$ then $A \cup B$ is equal to:

- A. $\{1, 3, 5, 7, 9\}$
- B. $\{3, 9\}$
- C. $\{2, 4, 8, 10\}$
- D. $\{1, 3, 5, 6, 7, 9\}$



16. The following map shows campsites in a campground.



Jon is camped at grid reference C2. If he is facing north and looks due east, what is he looking at?

- A. Toilets
- B. Campsite C
- C. Campsite A
- D. Campsite E


17. Using the similar triangles shown below, find the value of x .



- A. 1.3
- B. 2.6
- C. 6.3
- D. 9.3

18. The perimeter of the race track shown to the right is:

- A. 390.23 m
- B. 395.2 m
- C. 405 m
- D. 309.35 m



19. There are 1500 people in a fun run; 1125 are over 18 years of age. What percentage of people are aged 18 years and under?

- A. 25%
- B. 55%
- C. 45%
- D. 35%

20. The vertices of a triangle are A (1, 1), B (3, 4) and C (4, 2). Give the vertices of the triangle if it is reflected in the y -axis.


21. The data in the table below shows the number of books read by 6 students in Year 9 during one month.

Number of books read	3, 6, 8, 4, 4, 2
-----------------------------	------------------

Which of the following is true?

- A. Mean = median
 - B. Median = mode
 - C. Mean = mode
 - D. Mean < mode
22. A bike worth \$420 has been reduced by \$ x . The new value of the bike (in dollars) is:
- A. $420 - x$
 - B. $420x$
 - C. $\frac{420}{x}$
 - D. $420 + x$


23. X and Y are 12.5 cm apart on a map. Calculate the actual distance between X and Y using the given key.



Key: 1 cm \Leftrightarrow 6 km


24. Pat bought a packet of 100 marbles for \$5.00. The cost of one marble is:
- A. \$0.50
 - B. \$0.20
 - C. \$0.02
 - D. \$0.05
25. A bag contains 6 red, 8 green and 4 blue counters. What is the ratio of red to blue counters?
- A. 3:2
 - B. 2:3
 - C. 3:4
 - D. 2:1

26. The graph at right shows the distance travelled by a hiker. The average speed of the hiker is closest to:







- A. 15 km/h
- B. 1 km/h
- C. 2 km/h
- D. 8 km/h

27. What is the value of x on the number line shown?




- A. -1.5
- B. -0.5
- C. 0.5
- D. 1.5

28. Which window has the largest area?



29. What is the size of $\angle XYZ$?



30. $24 \times ? = 16$

The missing value is:

- A. $\frac{1}{3}$ B. $\frac{2}{3}$ C. $\frac{1}{2}$ D. $\frac{3}{4}$

Set F

Non-calculator

1. The sum of 11 and a number, r , is 37. What is r ?

- A. 27 B. 14 C. 26 D. 8

2. $\sqrt{150}$ is between:

- A. 8 and 10 B. 10 and 12 C. 12 and 14 D. 14 and 16


3. The arrow on the spinner is spun 50 times. Which table is most likely to represent the results?

A.

Colour	Number of spins
Red	25
Blue	17
Yellow	8

B.

Colour	Number of spins
Red	16
Blue	25
Yellow	9



C.

Colour	Number of spins
Red	8
Blue	15
Yellow	27

D.

Colour	Number of spins
Red	25
Blue	12
Yellow	13

4. You are going to share $3\frac{1}{4}$ pizzas with your friends. How many $\frac{1}{6}$ sized slices can you cut from the pizzas?

- A.** 20 **B.** $19\frac{1}{2}$ **C.** 18 **D.** $16\frac{1}{2}$

5. $2 \times 2 \times 2 \times 2 \times 2$ is equal to:

- A.** $2^2 \times 2^2$ **B.** 8×2 **C.** 2×5 **D.** $4^2 \times 2$

6. There are 12 boys and 14 girls in a class. What fraction of the class is boys?

- A.** $\frac{6}{13}$ **B.** $\frac{6}{7}$ **C.** $\frac{1}{2}$ **D.** $\frac{3}{4}$

7. Of 100 000 newspapers printed, 5% are delivered to homes, with the remaining papers going to the newsagent. How many papers are delivered to the newsagent?

- A.** 5000 **B.** 50 000 **C.** 95 000 **D.** 75 000

8. A cricketer played 10 innings and scored the following number of runs.

45, 0, 33, 7, 11, 25, 57, 80, 33, 47

What will change if the score of 80 is removed from the set?


- A.** Mean and range
B. Mean and mode
C. Mean and median
D. Mean only



9. The side view of an A-frame swing looks like this.


$\angle BCD = 68^\circ$

Find $\angle BDC$ and $\angle DEF$.




10. How many faces does this pyramid have?






- A. 4
- B. 5
- C. 6
- D. 7



11. For the following graph, state the domain (possible x -values) and range (possible y -values).



12. Which of the following represents the reflection of this triangle in the mirror shown below?



13. Which of the following is the rearranged version of the formula $m = \frac{n - qr}{p}$?

- A. $p = mn - qr$ B. $n = mp + qr$ C. $n = \frac{m}{p} + \frac{q}{r}$ D. $n = \frac{m + qr}{p}$

14. If a number is halved and then increased by 4, it will be equal to 2 less than the original number. What is the original number?

- A. 8 B. 10 C. 12 D. 20

15. When $a = 1$ and $b = -1$, the value of $(a - b)^2$ is:

- A. 0 B. 1 C. 2 D. 4

16. Which linear graph joins the points $(5, -9)$ and $(-3, 7)$?





- A. $y = x + 4$ B. $y = 2x + 1$ C. $y = -x - 4$ D. $y = -2x + 1$

17. A T-shirt worth \$35 is discounted by 15%. Which of the following calculations gives the new price?

- A. 35×0.85 B. 35×0.15 C. $35 + 0.15$ D. $35 - 15$

18. A fair coin is tossed twice. Write down the four possible outcomes.

19. Which of these patterns does not possess a line of symmetry?



20. Arrange the following in ascending order.

$$\frac{1}{2}, 0.2^2, \sqrt{4}, \left(\frac{1}{2}\right)^2$$

21. The solution to $\frac{4-x}{3} - 5 > 3$ is:

- A. $x > 20$ B. $x > -20$ C. $x < -20$ D. $x < 20$

22. Fill in the missing value to make the following equation true.


$$2(2h - 1) = 3h - 2 + ?$$

23. Evaluate $\sqrt{(-2)^2 + (-4)^2} + 5$.

- A. 5 B. 25 C. 10 D. 15


24. What is the probability of choosing a 5 in a regular deck of cards (no jokers)?

- A. $\frac{5}{52}$
 B. $\frac{1}{4}$
 C. $\frac{1}{13}$
 D. $\frac{1}{2}$



25. What percentage of this shape is shaded?

- A. 8%
- B. 20%
- C. 50%
- D. 40%



26. The probability of hiring a certain coloured bicycle is shown in the table below.

Colour	White	Red	Silver	Blue
Probability	0.55	0.25	?	0.05

What is the probability of hiring a silver bicycle if there are no colours other than the ones shown?

- A. 0.15
 - B. 0.25
 - C. 0.2
 - D. 0.5
27. Two people buy a \$5 ticket in a \$200 000 lottery. They each contribute \$2 and \$3. How much would each collect if they won?
28. Andrew rides x km to school. Isabelle rides 3 km more. Together they ride 13 km. Which equation represents this situation?
- A. $x - 3 = 13$
 - B. $2x + 3 = 13$
 - C. $x + 3 = 13$
 - D. $2x - 3 = 13$
29. The maximum temperature recorded for a June day was 14°C . The minimum temperature recorded that day was -2°C . What is the difference between the maximum and minimum temperatures recorded that day?
30. The mean, median and mode for the following set of data is:
0, 1, 1, 2, 3, 3, 3, 3
- A. Mean = 2, median = 3, mode = 1
 - B. Mean = 2, median = 3, mode = 3
 - C. Mean = 2, median = 2.5, mode = 3
 - D. Mean = 3, median = 2.5, mode = 3

Answers


NAPLAN practice

Set A Calculator allowed

1. C
2. \$45
3. D
4. B
5. 54 cm^2
6. 21
7. C
8. C
9. 3, 1, 4, 2, 250
10. A
11. $1110\ 001_2$
12. C
13. 4277 m
14. B
15. C
16. $l = 250 \text{ m}$ $w = 185 \text{ m}$
17. C
18. B
19. B
20. $h \approx 11 \text{ cm}$
21. D
22. 14.7 s
23. B
24. 2
25. C
26. D
27. 360 mm^3
28. A
29. B
30. D

Set B Non-calculator

1. B
2. D
3. A
4. C
5. 0.78
6. C




8. 30%
9. B
10. D
11. D
12. B
13. D
14. C
15. D
16. D
17. B
18. 40
19. A
20. C
21. B
22. A
23. A
24. C
25. $\frac{q^2 + p}{y}$
26. B
27. D
28. D
29. B
30. D

Set C Calculator allowed

1. B
2. B
3. C
4. B
5. 16
6. C
7. $q = \frac{5}{8}$
8. A
9. C
10. A
11. B
12. D
13. B
14. B
15. D


- 16. C
- 17. A
- 18. B
- 19. B
- 20. D
- 21. $\frac{3}{4}$
- 22. D
- 23. C



- 25. A
- 26. North-east
- 27. C
- 28. 27, 12, 3, 0, 3, 12, 27
- 29. A
- 30. C

Set D Non-calculator

- 1. C
- 2. B
- 3. A
- 4. B
- 5. D
- 6. A
- 7. B
- 8. A
- 9. D
- 10. D
- 11. C
- 12. B
- 13. A
- 14. A
- 15. C
- 16. 50°
- 17.



- 18. D
- 19. A
- 20. D
- 21. B

- 22. >
- 23. $\frac{18}{20}$
- 24. D
- 25. B
- 26. C
- 27. $l = 80 \text{ m}, w = 20 \text{ m}$
- 28. C
- 29. A
- 30. C

Set E Calculator allowed

- 1. B
- 2. $x = 20^\circ, y = 120^\circ$
- 3. B
- 4. D
- 5. 744 hours
- 6. A
- 7. C
- 8. C
- 9. 2.202, $\sqrt{\frac{1}{4}}, \left(\frac{1}{2}\right)^2, 0.12, 0.01$
- 10. C
- 11. B
- 12. A
- 13. C
- 14. C
- 15. D
- 16. D
- 17. C
- 18. A
- 19. A
- 20. $A'(-1, 1), C'(-4, 2), B'(-3, 4)$
- 21. B
- 22. A
- 23. C
- 24. D
- 25. A
- 26. C
- 27. B
- 28. C
- 29. 132°
- 30. B

Set F Non-calculator

- 1. C
- 2. C
- 3. A
- 4. B
- 5. D
- 6. A
- 7. C
- 8. A

9. $\angle BDC = 56^\circ$
 $\angle DEF = 124^\circ$
10. D
11. Domain: $-4 \leq x < 8$
Range: $3 < y \leq 9$
12. D
13. B
14. C
15. D
16. D
17. A
18. HH, HT, TH, TT
19. B
20. 0.2^2 , $\left(\frac{1}{2}\right)^2$, $\frac{1}{2}$, $\sqrt{4}$
21. C
22. h
23. A
24. C
25. D
26. A
27. \$80 000 and \$120 000
28. B
29. 16°C
30. C

1 Number skills and index laws

LESSON SEQUENCE

1.1 Overview	2
1.2 Real numbers	4
1.3 Rounding numbers and significant figures	11
1.4 Review of index laws	16
1.5 Raising powers	24
1.6 Negative indices	28
1.7 Scientific notation	34
1.8 Square roots and cube roots	41
1.9 Surds (extending)	44
1.10 Review	56



LESSON

1.1 Overview

1.1.1 Why learn this?

The earliest mathematics that humans used involved simple counting. Knowing that two goats could be swapped for six chickens was an important part of life twelve thousand years ago, so being able to count was a big deal.

Of course, being able to count is still a big deal today, but over thousands of years we have moved on from counting sheep. Ancient Egypt was the first recorded civilisation to think about fractions. Later on, ancient Greek mathematicians became the first people we know of to think about numbers that can't be expressed as fractions — these thoughts gave rise to the idea of irrational numbers. The value of π (pi) — which is the ratio of the circumference of a circle to its diameter — is one such irrational number. The value of π can only be approximated — if you were to try to say all of the digits in the value of π (that we presently know of) at a rate of 10 digits a second it would take you around 200 000 years to say them!



As new kinds of numbers have been discovered, new ways of writing them have been developed. These new ways of writing numbers can make it easier to work with numbers like the massively large ones that describe the distances between stars, or the incredibly small ones that describe the size of subatomic particles.

In this topic we will continue our mathematical journey by exploring a range of different types of numbers and the ways they can be written.

Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 1.1 Pre-test

- State whether the following statement is True or False.
Decimals that do not terminate and also do not recur are irrational numbers.
- State the kind of number that $7.\overline{324}$ is an example of.
- State whether the following statements are True or False.
 - π is an irrational number.
 - Every surd is an irrational number.
- Calculate $\sqrt[3]{27}$.
- MC** Select the simplest form of $\sqrt{50}$ from the following.
 - $5\sqrt{10}$
 - $2\sqrt{5}$
 - $5\sqrt{2}$
 - $10\sqrt{5}$
 - $25\sqrt{2}$
- MC** Select the correct answer for the expression $\sqrt{5} - 2\sqrt{45} + 3\sqrt{20}$.
 - $7\sqrt{5}$
 - $\sqrt{5}$
 - $2\sqrt{70}$
 - $19\sqrt{5}$
 - $18\sqrt{5}$
- The length of the hypotenuse of a right-angled triangle uses the formula $c^2 = a^2 + b^2$.
Calculate the length of the hypotenuse for a triangle with side lengths of $\sqrt{2}$ and $\sqrt{3}$. Leave your answer as a surd.
- MC** Select how many significant figures there are in 0.0305.
 - 1
 - 2
 - 3
 - 4
 - 5
- Round 7.43861 to 3 decimal places.
- Round 7.43861 to 3 significant figures.
- MC** Select the correct scientific notation to 3 significant figures for 0.000 000 021 37.
 - 2.137×10^8
 - 2.14×10^8
 - 2.14×10^{-8}
 - 2.137×10^{-8}
 - 21.4×10^{-9}
- MC** Select the series of numbers that is correctly ordered from smallest to largest.
 - $-\sqrt[3]{8}, -\sqrt[3]{27}, \sqrt[4]{81}$
 - $\sqrt{3}, e, 5 \times 10^2$
 - $2.1 \times 10^{-5}, 2.13 \times 10^2, 2.75 \times 10^{-1}$
 - $\left(\frac{2}{5}\right)^2, \sqrt{0.16}, 0.3$
 - $\sqrt{3}, e, 5 \times 10^2, 3.\overline{14}, \pi, \frac{22}{7}$
- An electron has a mass of $9.1093819 \times 10^{-31}$ kg, correct to 8 significant figures. A neutron is 1836 times the size of an electron. Evaluate the mass of a neutron correct to 4 significant figures.
- Rationalise $\frac{4}{\sqrt{3}}$.
- Simplify $\frac{5}{2}\sqrt{\frac{10}{5}}$.

LESSON

1.2 Real numbers

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- distinguish between natural, integer, rational, irrational and real numbers
- remember and use the symbols used to represent natural, integer, rational, irrational and real numbers
- write integers and decimals as rational number fractions
- write recurring decimals using dot notation
- distinguish between recurring and terminating decimals.

1.2.1 Natural numbers


eles-4374

- The earliest use of numbers by humans was for counting things. The numbers we used were the positive whole numbers: 1, 2, 3, 4 ...
- These counting numbers are now known as the **natural numbers**.
- We use the symbol N to represent the set of natural numbers.



Integers

- **Integers** include the entire set of whole numbers. This means that the integer group contains:
 - the positive whole numbers (i.e. the natural numbers) 1, 2, 3, 4 ...
 - the negative whole numbers $-1, -2, -3, -4 \dots$
 - 0 (which is neither positive nor negative).
- We use the symbol Z to represent the full set of integers. You may also see Z^- used to represent only the group of negative integers and Z^+ used for only the group of positive integers. This means that Z^+ and N represent the same group of numbers.



Rational numbers

- Integers, along with recurring decimals, are part of a larger group called the **rational numbers** represented by the symbol, Q .
- Rational numbers are numbers that can be expressed as a fraction with whole numbers both above and below the dividing sign.

Examples of rational numbers include $\frac{3}{7}$, -5 (which can be written as $\frac{-5}{1}$),
 3.2 (which can be written as $\frac{16}{5}$) and -0.036 (which can be written as $\frac{-36}{1000}$).

The denominator in a fraction

The denominator (bottom number) in a fraction cannot be 0. The quotient $9 \div 0$ has no answer. Because of this, numbers such as $\frac{9}{0}$ are said to be **undefined**. Because they do not exist, they are not rational numbers!

WORKED EXAMPLE 1 Writing rational numbers as fractions

By writing each of the following in fraction form, show that these numbers are rational.

- a. 7 b. -11 c. 0 d. $-4\frac{2}{5}$ e. 1.2

THINK

Each number must be written as a fraction using integers.

- a. The number 7 has to be written in fraction form. To write a number as a fraction, it must be written with a numerator and denominator. For a whole number like 7, the denominator is always 1.
- b. The number -11 has to be written in fraction form. To write a number as a fraction, it must be written with a numerator and denominator. Because -11 is a whole number, the denominator is 1.
- c. The number 0 has to be written in fraction form. To write a number as a fraction, it must be written with a numerator and a denominator. Because 0 is a whole number, the denominator is 1.
- d. Write $-4\frac{2}{5}$ as an improper fraction.
- e. The number 1.2 can be expressed as $1 + 0.2$. This can then be expressed as $1 + \frac{2}{10}$. Write this as an improper fraction.

WRITE

- a. $7 = \frac{7}{1}$. This number is rational.
- b. $-11 = \frac{-11}{1}$. This number is rational.
- c. $0 = \frac{0}{1}$. This number is rational.
- d. $-4\frac{2}{5} = \frac{-22}{5}$. This number is rational.
- e. $1.2 = \frac{12}{10}$. This number is rational.



eles-4377

1.2.2 Recurring decimals

- When a rational number is written in decimal form, there are two possibilities:
 - The decimal will come to an end (or **terminate**).
For example, $\frac{5}{4} = 1.25$, which is a decimal that terminates (ends) after two decimal places.
 - The decimal will repeat itself endlessly. For example, $\frac{7}{6} = 1.16666\dots$
The 6 in this number is a repeating digit, which means the decimal does not terminate – it has no end.
- Decimals that endlessly repeat digits (or blocks of digits) are called **recurring decimals**.
- Recurring decimals are written by placing dots or a line above the repeating digits.

Writing recurring decimals

Number of recurring digits	Example	How to write the decimal
1 recurring digit	1.615 555 555 ... (Only the digit 5 repeats.)	1.615̇
2 recurring digits	1.615 151 515 ... (The digits 1 and 5 repeat.)	1.615̇ or 1.615̄
3 or more recurring digits	1.615 615 615 ... (The digits 6, 1 and 5 repeat.)	1.615̇ or 1.615̄

WORKED EXAMPLE 2 Writing recurring decimals in extended form

Write the first 8 digits of each of the following recurring decimals.

a. $3.0\dot{2}$

b. $47.\dot{1}$

c. $11.54\dot{9}$

THINK

a. In $3.0\dot{2}$, the digits 0 and 2 repeat.

b. In $47.\dot{1}$, the digit 1 repeats.

c. In $11.54\dot{9}$, the digits 4 and 9 repeat.

WRITE

a. 3.020 202 0

b. 47.111 111

c. 11.549 494

WORKED EXAMPLE 3 Writing fractions as recurring decimals

Write each fraction as a recurring decimal using dot notation. Use a calculator for the division.

a. $\frac{5}{6}$

b. $\frac{57}{99}$

c. $\frac{25}{11}$

d. $\frac{4}{7}$

THINK

a. $5 \div 6 = 0.833\ 333\ 3$
3 repeats, so a dot goes above the 3.

b. $57 \div 99 = 0.575\ 757\ 575\ 7$
5 and 7 repeat, so dots go above the 5 and the 7.

c. $25 \div 11 = 2.272\ 727\ 27$
2 and 7 repeat, so dots go above the 2 and the 7.

d. $4 \div 7 = 0.571\ 428\ 571$
It looks as though 571 428 will recur, so dots go above the 5 and the 8.

WRITE

a. $\frac{5}{6} = 0.8\dot{3}$

b. $\frac{57}{99} = 0.\dot{5}\dot{7}$

c. $\frac{25}{11} = 2.\dot{2}\dot{7}$

d. $\frac{4}{7} = 0.\dot{5}71\ 42\dot{8}$

1.2.3 Irrational numbers

- A number is **irrational** if it cannot be written either as a fraction or as a terminating or recurring decimal.
- Irrational numbers may be expressed as decimals. In irrational numbers the digits do not repeat themselves in any particular order.

For example:

$$\sqrt{18} = 4.242\ 640\ 687\ 12\ \dots$$

$$\sqrt{0.03} = 0.173\ 205\ 080\ 757\ \dots$$

- We use the symbol I to represent the set of irrational numbers.
- Surds are irrational numbers that cannot be expressed as the ratio of two integers. They are typically written in the form \sqrt{n} , where n is a positive integer that is not a perfect square.


Identifying irrational numbers

To classify a number as either rational or irrational:

1. Determine whether it can be expressed as a whole number, a fraction or a terminating or recurring decimal.
2. If the answer to step 1 is yes, then the number is rational. If the answer is no, then the number is irrational.

- Exact representations of irrational numbers refer to representations that are precise numbers not in a decimal form.
 - They are expressed as roots, logarithms, or trigonometric functions.
 - Commonly used in theoretical or mathematical contexts where precision is crucial.
- Approximate decimal representations of irrational numbers are decimal numbers that are close to the actual value of the irrational number, but not exact.
 - These representations are obtained by truncating or rounding off the decimal expansion of the number to a certain number of digits.



For example, π is an irrational number that can be approximated to different degrees of accuracy, such as 3.14, 3.1416, or 3.14159265359.



- The main difference between exact representations and approximate decimal representations of irrational numbers is the level of precision.

Real numbers

- The set of **real numbers** is made up of the group of rational numbers and the group of irrational numbers. These groups can, in turn, be broken up into subsets of other number groups.
- A real number is any number that lies on the number line.
- We use the symbol **R** to represent the set of real numbers.



DISCUSSION



Determine how many recurring digits there will be for $\frac{1}{13}$.

π (pi) — a special number

The symbol π (**pi**) is used for a particular number. That number is the circumference of a circle whose diameter length is 1 unit. Pi can be approximated as a decimal that is non-terminating and non-recurring. Therefore, π is classified as an irrational number. It is also called a transcendental number.

In decimal form, $\pi = 3.141\ 592\ 653\ 589\ 793\ 23 \dots$ It has been calculated to 62.8 trillion decimal places with the aid of a computer.

Resources

-  **eWorkbook** Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)
-  **Interactivity** Individual pathway interactivity: Real numbers (int-4476)
The number system (int-6027)

Individual pathways

PRACTISE

1, 4, 9, 11, 14

CONSOLIDATE

2, 5, 6, 8, 12, 15

MASTER

3, 7, 10, 13, 16, 17

Fluency

- WE1** Show that the following numbers are rational by writing each of them in fraction form.
 - 15
 - 8
 - $2\frac{2}{3}$
 - $-5\frac{1}{8}$
- Show that the following numbers are rational by writing each of them in fraction form.
 - $\sqrt{16}$
 - $7\frac{3}{10}$
 - 0.002
 - 87.2
- Show that the following numbers are rational by writing each of them in fraction form.
 - 0
 - 1.56
 - 3.612
 - 0.08
- WE2** Write the first 8 digits of each of the following recurring decimals.
 - $0.\dot{5}$
 - $0.\dot{5}\dot{1}$
 - $0.5\dot{1}$
 - $6.0\dot{3}\dot{1}$
 - $5.\dot{1}8\dot{3}$
- Write the first 8 digits of each of the following recurring decimals.
 - $-7.02\dot{4}$
 - $8.\dot{9}12\dot{4}$
 - $5.12\dot{3}\dot{4}$
 - $5.\dot{1}23\dot{4}$
 - $3.0\dot{0}\dot{2}$
- WE3** Write the following fractions as recurring decimals using dot notation. (Use a calculator.)
 - $\frac{5}{9}$
 - $\frac{3}{11}$
 - $2\frac{2}{11}$
 - $\frac{3}{7}$
 - $\frac{-173}{99}$
- Write the following fractions as recurring decimals using dot notation. (Use a calculator.)
 - $\frac{73}{990}$
 - $-3\frac{5}{6}$
 - $\frac{7}{15}$
 - $\frac{46}{99}$
 - $\frac{46}{990}$

Understanding

- From the following list of numbers:

$$-3, \frac{-3}{7}, 0, 2.\dot{3}, 2\frac{3}{5}, 15$$
 - write down the natural numbers
 - write down the integers
 - write down the rational numbers
 - write down the recurring decimals.
- Write these numbers in order from smallest to largest.

$$2.\dot{1}, 2.\dot{1}\dot{2}, 2.\dot{1}2\dot{1}, 2.1\dot{2}\dot{1}, 2.1\dot{2}$$
- State which of the following statements are true.
 - All natural numbers are integers.
 - All integers are rational numbers.
 - All rational numbers have recurring decimals.
 - Natural numbers can be negative integers.
 - All numbers with decimals are rational numbers.
 - Irrational numbers are not real numbers.

Reasoning

11. Explain why all integers are rational numbers.
12. **a.** If positive integers are represented by Z^+ , explain what symbol we would use to represent the set of negative rational numbers.
b. Explain why we could not write N^- .
13. Are all fractions in which both the numerator and denominator are integers rational? Explain why or why not.

Problem solving

14. **a.** Using the fraction $\frac{a}{b}$, where a and b are natural numbers, write 3 recurring decimals in fractional form using the smallest natural numbers possible.
b. Determine the largest natural number you used.
15. Write 2 fractions that have the following number of repeating digits in their decimal forms.
 - a.** 1 repeating digit
 - b.** 2 repeating digits
 - c.** 3 repeating digits
 - d.** 4 repeating digits
16. If a \$2 coin weighs 6 g, a \$1 coin weighs 9 g, a 50 c coin weighs 15 g, a 20 c coin weighs 12g, a 10 c coin weighs 5 g and a 5 c coin weighs 3 g, determine what the maximum value of the coins would be if a bundle of them weighed exactly 10 kg.
17. A Year 9 student consumes two bottles of water every day, except on every fifth day when they only have one. Evaluate their annual bottled-water consumption.



LESSON

1.3 Rounding numbers and significant figures

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- round decimal numbers to a given number of decimal places
- identify the significant figures in a decimal number
- round decimal numbers to a given number of significant figures.



eles-4380

1.3.1 Rounding decimals

- If a number can't be written as an exact decimal, or if it has a lot of digits after the decimal point, we can use a shorter, approximate value.
- A number's approximate value can be found by using a calculator and then rounding it off to the desired level of accuracy.
- When we round numbers, we write them to either a certain number of decimal places or a certain number of significant figures.

Rounding to a given number of decimal places

1. Decide how many decimal places you need to round the number to.
2. Identify the digit that is located at the position you need to round to. This digit is called the rounding digit.
3. Look at the digit that comes after the rounding digit.
 - If the digit's value is 0, 1, 2, 3 or 4, leave the rounding digit as it is.
 - If the digit's value is 5, 6, 7, 8 or 9, then increase the rounding digit by adding 1 to its value.
4. Remove all of the numbers after the rounding digit.

WORKED EXAMPLE 4 Rounding a decimal number to three decimal places

Round 3.456 734 correct to three decimal places (3 d.p.).

THINK

1. Identify the rounding digit located at the third decimal place. In this case it is 6.
2. Look at the digit that comes after the rounding digit.
3. The number after the rounding digit is a 7, so increase the value of the rounding digit by adding 1. Next, remove the digits after it.

WRITE

3.456734

↓
3.456734
3.457

WORKED EXAMPLE 5 Writing a real number to three decimal places

Write these numbers correct to three decimal places (3 d.p.).

a. $\sqrt{3}$

b. π

c. $5.\dot{1}9$

d. $\frac{2}{3}$

e. 7.123 456

THINK

- a. $\sqrt{3} = 1.7320 \dots$ The rounding digit is 2. The next digit is 0, so leave 2 as it is.
- b. $\pi = 3.1415 \dots$ The rounding digit is 1. The next digit is 5, so add 1 to the rounding digit.
- c. $5.\dot{1}9 = 5.1999 \dots$ The rounding digit is 9. The next digit is 9, so add 1 to the rounding digit.
- d. $\frac{2}{3} = 0.6666 \dots$ The rounding digit is 6. The next digit is 6, so add 1 to the rounding digit.
- e. 7.123 456. The rounding digit is 3. The next digit is 4, so leave 3 as is.

WRITE

- a. $\sqrt{3} \approx 1.732$
- b. $\pi \approx 3.142$
- c. $5.\dot{1}9 \approx 5.200$
- d. $\frac{2}{3} \approx 0.667$
- e. $7.1234 \approx 7.123$



eles-4381

1.3.2 Rounding to significant figures

- Another method of rounding decimals is to write them so that they are correct to a certain number of significant figures.
- When decimals are rounded, they don't always have the same number of decimal places as significant figures.

For example, the number 1.425 has three decimal places but four significant figures.

Counting significant figures in a decimal number

1. Identify the first non-zero digit in the number. This digit may be located either before or after the decimal point.

This digit is counted as the first significant figure.

2. Continue to count all digits in the number to the end of the number as it is written.

This will give you the total number of significant figures.

WORKED EXAMPLE 6 Counting significant figures

State the number of significant figures in each of the following numbers.

a. 25

b. 0.04

c. 3.02

d. 0.100

THINK

- a. In the number 25, the first significant figure is 2. There is 1 more significant figure after the first one.
- b. In the number 0.04, the first significant figure is 4. There are no more significant figures after the first one.
- c. In the number 3.02, the first significant figure is 3. There are 2 more significant figures after the first one.
- d. In the number 0.100, the first significant figure is 1. There are 2 more significant figures after the first one.

WRITE

- a. 2 significant figures
- b. 1 significant figure
- c. 3 significant figures
- d. 3 significant figures

WORKED EXAMPLE 7 Rounding real numbers to five significant figures

Round these numbers so that they are correct to five significant figures.

- a. π b. $\sqrt{200}$ c. $0.\dot{0}\dot{3}$ d. 2530.166

THINK

- a. $\pi \approx 3.14159 \dots$ The first significant figure is 3. Starting with 3, write down the next 4 digits. The 6th digit is 9, so add 1 to the value of 5, the 5th digit.
- b. $\sqrt{200} \approx 14.1421 \dots$ The first significant figure is 1. Starting with 1, write down the next 4 digits. The 6th digit is 1, so leave 2, the 5th digit, as it is.
- c. $0.\dot{0}\dot{3} = 0.0303030 \dots$ The first significant figure is 3. Starting with 3, write down the next 4 digits. The 6th digit is 0, so leave 3, the 5th digit, as it is.
- d. 2530.16... The first significant figure is 2. Starting with 2, write the next 4 digits. The 6th digit is 6, so add 1 to the value of 1, the 5th digit.

WRITE

- a. 3.1416
- b. 14.142
- c. 0.030303
- d. 2530.2

DISCUSSION

Is there an equal number of rational numbers and irrational numbers? Explain.

on Resources



eWorkbook Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)



Interactivity Individual pathway interactivity: Rounding numbers and significant figures (int-4478)
Rounding to significant figures (int-6030)

Exercise 1.3 Rounding numbers and significant figures

learn **on**

1.3 Quick quiz **on**

1.3 Exercise

Individual pathways

■ PRACTISE

1, 4, 7, 10, 12, 15, 18

■ CONSOLIDATE

2, 5, 8, 11, 13, 16, 19

■ MASTER

3, 6, 9, 14, 17, 20

Fluency

1. **WE4&5** Write each of the following correct to 3 decimal places.

a. $\frac{\pi}{2}$

b. $\sqrt{5}$

c. $\sqrt{15}$

d. 5.12×3.21

e. $5.\dot{1}$

f. $5.1\dot{5}$

2. Write each of the following correct to 3 decimal places.

a. $5.\dot{1}\dot{5}$

b. 11.72^2

c. $\frac{3}{7}$

d. $\frac{1}{13}$

e. $2\frac{3}{7}$

f. 0.999999

3. Write each of the following correct to three decimal places.
 a. 4.000 01 b. $2.79 \div 11$ c. 0.0254 d. 0.000 913 6 e. 5.000 01 f. 2342.156
4. **WE6** State how many significant figures there are in each of the following numbers.
 a. 36 b. 207 c. 1631 d. 5.04
5. State how many significant figures there are in each of the following numbers.
 a. 176.2 b. 95.00 c. 0.21 d. 0.01
6. State how many significant figures there are in each of the following numbers.
 a. 0.000 316 b. 0.1007 c. 0.010 d. 0.0512
7. **WE7** Round each number so that it is correct to five significant figures.
 a. $\frac{\pi}{2}$ b. $\sqrt{5}$ c. $\sqrt{15}$ d. 5.12×3.21 e. $5.\dot{1}$ f. $5.1\dot{5}$
8. Round each number so that it is correct to five significant figures.
 a. $5.\dot{1}\dot{5}$ b. 11.72^2 c. $\frac{3}{7}$
 d. $\frac{1}{13}$ e. $2\frac{3}{7}$ f. 0.999 999
9. Write each number so that it is correct to 5 significant figures.
 a. 6.581 29 b. 4.000 01 c. $2.79 \div 11$ d. 0.0254 e. 0.000 913 6 f. 5.000 01

Understanding

10. Write the value of π correct to 4, 5, 6 and 7 decimal places.
11. State whether each of the following numbers has more significant figures than decimal places.
 a. 17.26 b. 0.0032 c. 1.06 d. 0.010 005
12. State whether each statement is True or False.
 a. Every surd is a rational number.
 b. Every surd is an irrational number.
 c. Every irrational number is a surd.
 d. Every surd is a real number.
13. State whether each statement is True or False.
 a. π is a rational number.
 b. π is an irrational number.
 c. π is a surd.
 d. π is a real number.
14. State whether each statement is True or False.
 a. $1.\dot{3}\dot{1}$ is a rational number.
 b. $1.\dot{3}\dot{1}$ is an irrational number.
 c. $1.\dot{3}\dot{1}$ is a surd.
 d. $1.\dot{3}\dot{1}$ is a real number.

Reasoning

15. Write a decimal number that has 3 decimal places, but 4 significant figures. Show your working.
16. Explain why the number 12.995 412 3 becomes 13.00 when rounded to 2 decimal places.

17. The rational numbers 3.1416 and $\frac{22}{7}$ are both used as approximations to π .
- Determine the largest number of decimal places these numbers can be rounded to in order to give the same value.
 - Explain which of these two numbers gives the best approximation to π .



Problem solving

18. The area of a circle is calculated using the formula $A = \pi \times r^2$, where r is the radius of the circle. Pi (π) is sometimes rounded to 2 decimal places to become 3.14. A particular circle has a radius of 7 cm.
- Use $\pi = 3.14$ to calculate the area of the circle to 2 decimal places.
 - Use the π key on your calculator to calculate the area of the circle to 4 decimal places.
 - Round your answer for part **b** to 2 decimal places.
 - Are your answers for parts **a** and **c** different? Discuss why or why not.
19. The volume of a sphere (a ball shape) is calculated using the formula $v = \frac{4}{3} \times \pi \times r^3$, where r is the radius of the sphere. A beach ball with a radius of 25 cm is bouncing around the crowd at the MCG during the Boxing Day Test.
- Calculate the volume of the beach ball to 4 decimal places.
 - Calculate the volume to 4 decimal places and determine how many significant figures the result has.
 - Explain whether the calculated volume is a rational number.
20. In a large sample of written English there are about 7 vowels for every 11 consonants. The letter e accounts for about one-third of the occurrence of vowels. Explain how many times you would expect the letter e to occur in a passage of 100 000 letters. Round your answer to the nearest 100.



LESSON

1.4 Review of index laws

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify the base and the exponent of a number written in index form
- recall the first three index laws
- use the first three index laws to simplify expressions involving indices.




1.4.1 Index notation

eles-4382

- The product of factors can be written in a shorter form called **index notation** (also known as exponent notation).

$$\begin{array}{c} \text{Index (exponent)} \\ \swarrow \\ \text{Base} \rightarrow 6^4 = 6 \times 6 \times 6 \times 6 \leftarrow \text{Factor form} \\ = 1296 \end{array}$$

- Any composite number can be written as a product of powers of prime factors using a factor tree, or by other methods, such as repeated division.



$$\begin{aligned} 100 &= 2 \times 2 \times 5 \times 5 \\ &= 2^2 \times 5^2 \end{aligned}$$

WORKED EXAMPLE 8 Using index notation to express a product of prime factors

Express 360 as a product of powers of prime factors using index notation.

THINK

1. Express 360 as a product of a factor pair.
2. Further factorise 6 and 60.
3. Further factorise 4 and 15. They are both composite numbers.
4. There are no more composite numbers.
5. Write the answer using index notation.

Note: The factors are generally expressed with their bases written in ascending order.

WRITE

$$\begin{aligned} 360 &= 6 \times 60 \\ &= 2 \times 3 \times 4 \times 15 \\ &= 2 \times 3 \times 2 \times 2 \times 3 \times 5 \\ &= 2 \times 2 \times 2 \times 3 \times 3 \times 5 \\ 360 &= 2^3 \times 3^2 \times 5 \end{aligned}$$

1.4.2 Multiplication using indices

- When multiplying **terms** with the same bases, add the indices. This is the **First Index Law**.
- When more than one base is involved, apply the First Index Law to each base separately.

The First Index Law

$$a^m \times a^n = a^{m+n}$$

For example,
 $2x^2 \times x^3 = 2x^{2+3} = 2x^5$

WORKED EXAMPLE 9 Simplifying an expression using the First Index Law

Simplify $5e^{10} \times 2e^3$.

THINK

1. The order is not important when multiplying, so place the coefficients first.
2. Simplify by multiplying the coefficients and then applying the First Index Law (add the indices). Write the answer.

WRITE

$$\begin{aligned} 5e^{10} \times 2e^3 \\ = 5 \times 2 \times e^{10} \times e^3 \\ = 10e^{13} \end{aligned}$$

WORKED EXAMPLE 10 Simplifying using the First Index Law on multiple bases

Simplify $7m^3 \times 3n^5 \times 2m^8n^4$.

THINK

1. The order is not important when multiplying, so place the coefficients first and group all similar pronumerals together.
2. Simplify by multiplying the coefficients and applying the First Index Law (add the indices). Write the answer.

WRITE

$$\begin{aligned} 7m^3 \times 3n^5 \times 2m^8n^4 \\ = 7 \times 3 \times 2 \times m^3 \times m^8 \times n^5 \times n^4 \\ = 42m^{11}n^9 \end{aligned}$$

1.4.3 Division using indices

- When dividing terms with the same bases, subtract the indices. This is the **Second Index Law**.
- When the coefficients do not divide evenly, simplify by cancelling.

The Second Index Law

$$a^m \div a^n = a^{m-n} \text{ or } \frac{a^m}{a^n} = a^{m-n}$$

For example,
 $2x^5 \div x^3 = \frac{2x^5}{x^3} = 2x^{5-3} = 2x^2$

WORKED EXAMPLE 11 Simplifying using the Second Index Law

Simplify $\frac{25v^6 \times 8w^9}{10v^4 \times 4w^5}$.

THINK

1. Simplify the numerator and the denominator by multiplying the coefficients.
2. Simplify further by dividing the coefficients and applying the Second Index Law (subtract the indices). Write the answer.

WRITE

$$\begin{aligned} & \frac{25v^6 \times 8w^9}{10v^4 \times 4w^5} \\ &= \frac{200v^6w^9}{40v^4w^5} \\ &= \frac{\overset{5}{\cancel{200}}}{\underset{1}{\cancel{40}}} \times \frac{v^6}{v^4} \times \frac{w^9}{w^5} \\ &= 5v^2w^4 \end{aligned}$$

WORKED EXAMPLE 12 Simplifying using the Second Index Law with non-cancelling coefficients

Simplify $\frac{7t^3 \times 4t^8}{12t^4}$.

THINK

1. Simplify the numerator by multiplying the coefficients.
2. Simplify further by dividing the coefficients by the highest common factor. Then apply the Second Index Law (subtract the indices). Write the answer.

WRITE

$$\begin{aligned} & \frac{7t^3 \times 4t^8}{12t^4} \\ &= \frac{28t^{11}}{12t^4} \\ &= \frac{28}{12} \times \frac{t^{11}}{t^4} \\ &= \frac{7t^7}{3} \end{aligned}$$



1.4.4 Zero index

eles-4385

- Any number divided by itself is 1.
- This means that $\frac{a^m}{a^m} = 1$. So, $a^{m-m} = a^0 = 1$.
- In general, any number (except zero) to the power of 0 is equal to 1. This is the **Third Index Law**.

The Third Index Law

$$a^0 = 1, \text{ where } a \neq 0$$

For example,
 $2^0 = 1$

WORKED EXAMPLE 13 Evaluating using the Third Index Law

Evaluate the following.

a. t^0

b. $(xy)^0$

c. 17^0

d. $5x^0$

e. $(5x)^0 + 2$

f. $5^0 + 3^0$

THINK

- Apply the Third Index Law.
- Apply the Third Index Law.
- Apply the Third Index Law.
- Apply the Third Index Law.
- Apply the Third Index Law.
- Apply the Third Index Law.

WRITE

- $t^0 = 1$
- $(xy)^0 = 1$
- $17^0 = 1$
- $5x^0 = 5 \times x^0$
 $= 5 \times 1$
 $= 5$
- $(5x)^0 + 2 = 1 + 2$
 $= 3$
- $5^0 + 3^0 = 1 + 1$
 $= 2$

WORKED EXAMPLE 14 Simplifying using the First and Third index laws

Simplify $\frac{9g^7 \times 4g^4}{6g^3 \times 2g^8}$.

THINK

- Simplify the numerator and the denominator by applying the First Index Law.
- Simplify the fraction further by applying the Second Index Law.
- Simplify by applying the Third Index Law.
- Write the answer.

WRITE

$$\begin{aligned} & \frac{9g^7 \times 4g^4}{6g^3 \times 2g^8} \\ &= \frac{36g^{11}}{12g^{11}} \\ &= \frac{\cancel{3}3\cancel{6}g^{11}}{\cancel{1}1\cancel{2}g^{11}} \\ &= 3g^0 \\ &= 3 \times 1 \\ &= 3 \end{aligned}$$

1.4.5 Cancelling fractions

eles-4386

- Consider the fraction $\frac{x^3}{x^7}$. This fraction can be cancelled by dividing the denominator and the numerator by the highest common factor (HCF), which is x^3 , so $\frac{x^3}{x^7} = \frac{1}{x^4}$.
Note: $\frac{x^3}{x^7} = x^{-4}$ when we apply the Second Index Law.

We will study negative indices in a later section.

DISCUSSION

How do the First, Second and Third index laws help with calculations?

WORKED EXAMPLE 15 Using the first three index laws to simplify expressions

Simplify these fractions by cancelling.

a. $\frac{x^5}{x^7}$

b. $\frac{6x}{12x^8}$

c. $\frac{30x^5y^6}{10x^7y^3}$

THINK

- a. Divide the numerator and denominator by the highest common factor (HCF), which is x^5 .
- b. 1. Divide the numerator and denominator by the HCF, which is $6x$.
2. Simplify and write the answer.
- c. 1. Divide the numerator and denominator by the HCF, which is $10x^5y^3$.
2. Simplify and write the answer.

WRITE

a. $\frac{x^5}{x^7} = \frac{1}{x^2}$

b. $\frac{6x}{12x^8} = \frac{6}{12} \times \frac{x}{x^8}$
 $= \frac{1}{2} \times \frac{1}{x^7}$

$$= \frac{1}{2x^7}$$

c. $\frac{30x^5y^6}{10x^7y^3} = \frac{30}{10} \times \frac{x^5}{x^7} \times \frac{y^6}{y^3}$
 $= \frac{3}{1} \times \frac{1}{x^2} \times \frac{y^3}{1}$
 $= \frac{3y^3}{x^2}$

on Resources



eWorkbook

Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)



Video eLesson

Index notation (eles-1903)



Interactivities

Individual pathway interactivity: Review of index laws (int-4516)

Review of index form (int-3708)

First Index Law (int-3709)

Second Index Law (int-3711)

Third Index Law (int-3713)

Individual pathways

PRACTISE

1, 3, 6, 10, 13, 15, 19, 20, 21, 26

CONSOLIDATE

2, 4, 7, 8, 11, 18, 22, 23, 27, 28

MASTER

5, 9, 12, 14, 16, 17, 24, 25, 29, 30

Fluency

1. **WE8** Use index notation to express each of the following as a product of powers of prime factors.

- a. 12 b. 72 c. 75 d. 240 e. 640 f. 9800

2. **WE9** Simplify each of the following.

- a. $4p^7 \times 5p^4$ b. $2x^2 \times 3x^6$ c. $8y^6 \times 7y^4$ d. $3p \times 7p^7$ e. $12t^3 \times t^2 \times 7t$ f. $6q^2 \times q^5 \times 5q^8$

3. **WE10** Simplify each of the following.

- a. $2a^2 \times 3a^4 \times e^3 \times e^4$ b. $4p^3 \times 2h^7 \times h^5 \times p^3$ c. $3k^2 \times 2k^{11} \times k$ d. $3^4 \times 3^4 \times 3^4$

4. Simplify each of the following.

- a. $2m^3 \times 5m^2 \times 8m^4$ b. $2gh \times 3g^2 h^5$
 c. $5p^4 q^2 \times 6p^2 q^7$ d. $8u^3 w \times 3uw^2 \times 2u^5 w^4$

5. Simplify each of the following.

- a. $9y^8 d \times y^5 d^3 \times 3y^4 d^7$
 b. $7b^3 c^2 \times 2b^6 c^4 \times 3b^5 c^3$
 c. $4r^2 s^2 \times 3r^6 s^{12} \times 2r^8 s^4$
 d. $10h^{10} v^2 \times 2h^8 v^6 \times 3h^{20} v^{12}$



6. **WE11** Simplify each of the following.

- a. $\frac{15p^{12}}{5p^8}$ b. $\frac{18r^6}{3r^2}$ c. $\frac{45a^5}{5a^2}$ d. $\frac{60b^7}{20b}$

7. Simplify each of the following.

- a. $\frac{100r^{10}}{5r^6}$ b. $\frac{9q^2}{q}$ c. $\frac{130d^3}{d^5}$ d. $\frac{21t^{11}}{3t^7}$

8. **WE12** Simplify each of the following.

- a. $\frac{8p^6 \times 3p^4}{16p^5}$ b. $\frac{12b^5 \times 4b^2}{18b^2}$ c. $\frac{25m^{12} \times 4n^7}{15m^2 \times 8n}$ d. $\frac{27x^9 y^3}{12xy^2}$

9. Simplify each of the following.

- a. $\frac{12j^8 \times 6f^5}{8j^3 \times 3f^2}$ b. $\frac{8p^3 \times 7r^2 \times 2s}{6p \times 14r}$ c. $\frac{27a^9 \times 18b^5 \times 4c^2}{18a^4 \times 12b^2 \times 2c}$ d. $\frac{81f^{15} \times 25g^{12} \times 16h^{34}}{27f^9 \times 15g^{10} \times 12h^{30}}$

10. **WE13** Evaluate the following.

- a. m^0 b. $6m^0$ c. $(6m)^0$ d. $(ab)^0$ e. $5(ab)^0$

11. Evaluate the following.

- a. $w^0 x^0$ b. 85^0 c. $85^0 + 15^0$ d. $x^0 + 1$ e. $5x^0 - 2$

12. Evaluate the following.

a. $\frac{x^0}{y^0}$ b. $x^0 - y^0$ c. $3x^0 + 11$ d. $3a^0 + 3b^0$ e. $3(a^0 + b^0)$

13. **WE14** Simplify each of the following.

a. $\frac{2a^3 \times 6a^2}{12a^5}$ b. $\frac{3c^6 \times 6c^3}{9c^9}$ c. $\frac{5b^7 \times 10b^5}{25b^{12}}$ d. $\frac{8f^3 \times 3f^7}{4f^5 \times 3f^5}$ e. $\frac{9k^{12} \times 4k^{10}}{18k^4 \times k^{18}}$

14. Simplify each of the following.

a. $\frac{2h^4 \times 5k^2}{20h^2 \times k^2}$ b. $\frac{p^3 \times q^4}{5p^3}$ c. $\frac{m^7 \times n^3}{5m^3 \times m^4}$ d. $\frac{8u^9 \times v^2}{2u^5 \times 4u^4}$ e. $\frac{9x^6 \times 2y^{12}}{3y^{10} \times 3y^2}$

Understanding

15. **WE15** Simplify each of the following by cancelling.

a. $\frac{x^7}{x^{10}}$ b. $\frac{m}{m^9}$ c. $\frac{m^3}{4m^9}$ d. $\frac{12x^6}{6x^8}$

16. Simplify each of the following by cancelling.

a. $\frac{12x^8}{6x^6}$ b. $\frac{24t^{10}}{t^4}$ c. $\frac{5y^5}{10y^{10}}$ d. $\frac{35x^2y^{10}}{20x^7y^7}$

17. Simplify each of the following by cancelling.

a. $\frac{12m^2n^4}{30m^5n^8}$ b. $\frac{16m^5n^{10}}{8m^5n^{12}}$ c. $\frac{20x^4y^5}{10x^5y^4}$ d. $\frac{a^2b^4c^6}{a^6b^4c^2}$

18. Calculate the value of each of the following expressions if $a = 3$.

a. $2a$ b. a^2 c. $2a^2$ d. $a^2 + 2$ e. $a^2 + 2a$ f. $a^2 - a$

Reasoning

19. Explain why x^2 and $2x$ are not the same number. Use an example to illustrate your reasoning.

20. **MC**

a. $12a^8b^2c^4(de)^0f$, when simplified, is equal to:

A. $12a^8b^2c^4$ B. $12a^8b^2c^4f$ C. $12a^8b^2f$ D. $12a^8b^2$ E. $12a^8b^2c^4f^2$

b. $\left(\frac{6}{11}a^2b^7\right)^0 \times - (3a^2b^{11})^0 + 7a^0b$, when simplified, is equal to:

A. $7b$ B. $1 + 7b$ C. $-1 + 7ab$ D. $-1 + 7b$ E. $-1 - 7b$

c. You are told that there is an error in the statement $3p^7q^3r^5s^6 = 3p^7s^6$.

Explain what the left-hand side should be for this statement to be correct.

A. $(3p^7q^3r^5s^6)^0$ B. $(3p^7)^0q^3r^5s^6$ C. $3p^7(q^3r^5s^6)^0$ D. $3p^7(q^3r^5)^0s^6$ E. $6p^7(q^3r^5)^0s^6$

21. **MC**

a. You are told that there is an error in the statement $\frac{8f^6g^7h^3}{6f^4g^2h} = \frac{8f^2}{g^2}$.

Explain what the left-hand side should be for this statement to be correct.

A. $\frac{8f^6(g^7h^3)^0}{(6)^0f^4g^2(h)^0}$ B. $\frac{8(f^6g^7h^3)^0}{(6f^4g^2h)^0}$ C. $\frac{8(f^6g^7)^0h^3}{(6f^4)^0g^2h}$ D. $\frac{8f^6g^7h^3}{(6f^4g^2h)^0}$ E. $\frac{8f^8(g^7h^3)^0}{(6)^0f^4g(h)^0}$

b. Select what $\frac{6k^7m^2n^8}{4k^7(m^6n)^0}$ is equal to.

A. $\frac{6}{4}$

B. $\frac{3}{2}$

C. $\frac{3n^8}{2}$

D. $\frac{3m^2n^8}{2}$

E. $\frac{3m^2n^6}{2}$

22. Explain why $5x^5 \times 3x^3$ is not equal to $15x^{15}$.

23. A multiple-choice question requires a student to multiply 5^6 by 5^3 . The student is having trouble deciding which of these four answers is correct: 5^{18} , 5^9 , 25^{18} or 25^9 .

a. State the correct answer.

b. Explain your answer by using another example to explain the First Index Law.

24. A multiple-choice question requires a student to divide 5^{24} by 5^8 . The student is having trouble deciding which of these four answers is correct: 5^{16} , 5^3 , 1^{16} or 1^3 .

a. Determine the correct answer.

b. Explain your answer by using another example to explain the Second Index Law.

25. a. Calculate the value of $\frac{5^7}{5^7}$.

b. Determine the value of any number divided by itself.

c. According to the Second Index Law, which deals with exponents and division, $\frac{5^7}{5^7}$ should equal 5 raised to what index?

d. Use an example to explain the Third Index Law.

Problem solving

26. Answer the following questions.

a. For $x^2x^\Delta = x^{16}$ to be true, determine what number must replace the triangle.

b. For $x^\Delta x^0 x^\circ = x^{12}$ to be true, there are 55 ways of assigning positive whole numbers to the triangle, circle, and diamond. Give at least four of these.

27. Answer the following questions.

a. Determine a pattern in the units digit for powers of 3.

b. The units digit of 3^6 is 9. Determine the units digit of 3^{2001} .

28. Answer the following questions.

a. Determine a pattern in the units digit for powers of 4.

b. Determine the units digit of 4^{105} .

29. Answer the following questions.

a. Investigate the patterns in the units digit for powers of 2 to 9.

b. Predict the units digit for the following.

i. 2^{35}

ii. 3^{16}

iii. 8^{51}

30. Write $4^{n+1} + 4^{n+1}$ as a single power of 2.



LESSON

1.5 Raising powers

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- apply the Fourth, Fifth and Sixth index laws to raise a power to another power
- apply the first six index laws to simplify expressions involving indices.

1.5.1 More index laws

eles-4387

- When a power is raised to another power the indices are multiplied. This is the **Fourth Index Law**.
- The Fourth Index Law can then be applied to products and quotients of mixed bases to develop the Fifth and Sixth index laws.

The Fourth Index Law

$$(a^m)^n = a^{m \times n}$$

For example, $(x^2)^3 = x^{2 \times 3} = x^6$

The Fifth Index Law

$$(a \times b)^m = a^m \times b^m$$

For example, $(2x)^3 = 2^3 \times x^3 = 8x^3$

The Sixth Index Law

$$\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$$

For example, $\left(\frac{4x}{2}\right)^2 = \frac{4^2 x^2}{2^2} = 4x^2$

WORKED EXAMPLE 16 Simplifying using the Fourth Index Law

Simplify the following.

a. $(7^4)^8$

THINK

- Simplify by applying the Fourth Index Law (multiply the indices).
1. Write the expression, including all indices.
2. Simplify by applying the Fourth Index Law (multiply the indices) for each term inside the brackets.
3. Write the answer.

b. $(3a^2b^5)^3$

WRITE

a. $(7^4)^8$
 $= 7^{4 \times 8}$
 $= 7^{32}$

b. $(3^1 a^2 b^5)^3$
 $= 3^{1 \times 3} a^{2 \times 3} b^{5 \times 3}$
 $= 3^3 a^6 b^{15}$
 $= 27a^6b^{15}$

WORKED EXAMPLE 17 Simplifying using the Fifth Index Law

Simplify $(2b^5)^2 \times (5b)^3$.

THINK

1. Write the expression, including all indices.
2. Simplify by applying the Fifth Index Law.
3. Calculate the number values and bring them to the front. Then use the First Index Law to simplify the powers of b .
4. Write the answer.

WRITE

$$\begin{aligned} & (2^1 b^5)^2 \times (5^1 b^1)^3 \\ &= 2^{1 \times 2} b^{5 \times 2} \times 5^{1 \times 3} b^{1 \times 3} \\ &= 2^2 \times b^{10} \times 5^3 \times b^3 \\ &= 4 \times 125 \times b^{10+3} \\ &= 500b^{13} \end{aligned}$$

WORKED EXAMPLE 18 Simplifying using the Sixth Index Law

Simplify $\left(\frac{2a^5}{d^2}\right)^3$.

THINK

1. Write the expression, including all indices.
2. Simplify by applying the Sixth Index Law for each term inside the brackets.
3. Simplify and then write the answer.


WRITE

$$\begin{aligned} & \left(\frac{2^1 a^5}{d^2}\right)^3 \\ &= \frac{2^{1 \times 3} a^{5 \times 3}}{d^{2 \times 3}} \\ &= \frac{2^3 a^{15}}{d^6} \\ &= \frac{8 a^{15}}{d^6} \end{aligned}$$

TI | THINK

In a new problem, on a Calculator page, complete the entry line as: $\left(\frac{2a^5}{d^2}\right)^3$
Press ENTER.

WRITE




$$\left(\frac{2a^5}{d^2}\right)^3 = \frac{8a^{15}}{d^6}$$

CASIO | THINK

On the Main screen complete the entry line as: $\left(\frac{2a^5}{d^2}\right)^3$
Press EXE.

WRITE



$$\left(\frac{2a^5}{d^2}\right)^3 = \frac{8a^{15}}{d^6}$$

DISCUSSION

What difference, if any, is there between the operation of the index laws on numeric terms compared with similar operations on algebraic terms?

Resources



eWorkbook Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)



Interactivity Individual pathway interactivity: Raising powers (int-4517)

Fourth Index Law (int-3716)

Fifth and sixth index laws (int-6063)

Exercise 1.5 Raising powers

learn **on**

1.5 Quick quiz



1.5 Exercise

Individual pathways

PRACTISE

1, 3, 7, 9, 14, 15, 20

CONSOLIDATE

2, 5, 8, 10, 12, 16, 17, 21

MASTER

4, 6, 11, 13, 18, 19, 22, 23

Fluency

1. **WE16** Simplify each of the following.

a. $(e^2)^3$

b. $(f^8)^{10}$

c. $(p^{25})^4$

d. $(r^{12})^{12}$

e. $(2a^2)^3$

2. Simplify each of the following.

a. $(a^2b^3)^4$

b. $(pq^3)^5$

c. $(g^3h^2)^{10}$

d. $(3w^9q^2)^4$

e. $(7e^5r^2q^4)^2$

3. **WE17** Simplify each of the following.

a. $(p^4)^2 \times (q^3)^2$

b. $(r^5)^3 \times (w^3)^3$

c. $(b^5)^2 \times (n^3)^6$

d. $(j^6)^3 \times (g^4)^3$

e. $(2a)^3 \times (b^2)^2$

4. Simplify each of the following.

a. $(q^2)^2 \times (r^4)^5$

b. $(h^3)^8 \times (j^2)^8$

c. $(f^4)^4 \times (a^7)^3$

d. $(r^5)^2 \times (u^4)^2$

e. $(i^3)^5 \times (j^2)^6$

5. **WE18** Simplify each of the following.

a. $\left(\frac{3b^4}{d^3}\right)^2$

b. $\left(\frac{5h^{10}}{2j^2}\right)^2$

c. $\left(\frac{2k^5}{3t^8}\right)^3$

d. $\left(\frac{7p^9}{8q^{22}}\right)^2$

6. Simplify each of the following.

a. $\left(\frac{5y^7}{3z^{13}}\right)^3$

b. $\left(\frac{4a^3}{7c^5}\right)^4$

c. $\left(\frac{-4k^2}{7m^6}\right)^3$

d. $\left(\frac{-2g^7}{3h^{11}}\right)^4$

Understanding

7. Simplify each of the following.

a. $(2^3)^4 \times (2^4)^2$

b. $(t^7)^3 \times (t^3)^4$

c. $(a^4)^0 \times (a^3)^7$

d. $(e^7)^8 \times (e^5)^2$

8. Simplify each of the following.

a. $(g^7)^3 \times (g^9)^2$

b. $(3a^2) \times (2a^6)^2$

c. $(2d^7)^3 \times (3d^2)^3$

d. $(10r^2)^4 \times (2r^3)^2$

9. **MC** $(p^7)^2 \div p^2$ is equal to:

A. p^7

B. p^{12}

C. p^{16}

D. $p^{4.5}$

E. p^{13}

10. **MC** $\frac{(w^5)^2 \times (p^7)^3}{(w^2)^2 \times (p^3)^5}$ is equal to:

A. w^2p^6

B. $(wp)^6$

C. $w^{14}p^{36}$

D. w^2p^2

E. $(wp)^7$

11. **MC** $(r^6)^3 \div (r^4)^2$ is equal to:

A. r^3

B. r^4

C. r^8

D. r^{10}

E. r^{12}

12. Simplify each of the following.

a. $(a^3)^4 \div (a^2)^3$

b. $(m^8)^2 \div (m^3)^4$

c. $(n^5)^3 \div (n^6)^2$

d. $(b^4)^5 \div (b^6)^2$

e. $(f^7)^3 \div (f^2)^2$

f. $(g^8)^2 \div (g^5)^2$

13. Simplify each of the following.

a. $(p^9)^3 \div (p^6)^3$

b. $(y^4)^4 \div (y^7)^2$

c. $\frac{(c^6)^5}{(c^5)^2}$

d. $\frac{(f^5)^3}{(f^2)^4}$

e. $\frac{(k^3)^{10}}{(k^2)^8}$

f. $\frac{(p^{12})^3}{(p^{10})^2}$

Reasoning

14. a. Replace the triangle with the correct index for the equation $4^7 \times 4^7 \times 4^7 \times 4^7 \times 4^7 = (4^7)^\Delta$.

b. The expression $(p^5)^6$ means to write p^5 as a factor how many times?

c. If you rewrote the expression from part b without any exponents, in the format $p \times p \times p \dots$, determine how many factors you would need.

d. Explain the Fourth Index Law.

15. a. Simplify each of the following.

i. $(-1)^{10}$

ii. $(-1)^7$

iii. $(-1)^{15}$

iv. $(-1)^6$

b. Write a general rule for the result obtained when -1 is raised to a positive power. Explain your answer.

16. Jo and Danni are having an algebra argument. Jo is sure that $-x^2$ is equivalent to $(-x)^2$, but Danni thinks otherwise.

Explain who is correct and justify your answer.

17. A multiple-choice question requires a student to calculate $(5^4)^3$. The student is having trouble deciding which of these three answers is correct: 5^{64} , 5^{12} or 5^7 .

a. Determine the correct answer.

b. Explain your answer by using another example to illustrate the Fourth Index Law.

18. a. Without using your calculator, simplify each side of the following equations to the same base and then solve each of them.
- $8^x = 32$
 - $27^x = 243$
 - $1000^x = 100\,000$
- b. Explain why all 3 equations have the same solution.
19. Consider the expression 4^{3^2} . Identify which is the correct answer, 4096 or 262 144. Justify your choice.

Problem solving

20. The diameter of a typical atom is so small that it would take about 10^8 atoms arranged in a line to reach a length of just 1 centimetre. Estimate how many atoms are contained in a cubic centimetre. Write this number as a power of 10.
21. Writing a base as a power itself can be used as a way to simplify an expression. Copy and complete the following calculations.

a. $16^{\frac{1}{2}} = (4^2)^{\frac{1}{2}} = \underline{\hspace{2cm}}$

b. $343^{\frac{2}{3}} = (7^3)^{\frac{2}{3}} = \underline{\hspace{2cm}}$

22. Simplify the following using index laws.

a. $8^{\frac{1}{3}}$

b. $27^{\frac{4}{3}}$

c. $125^{-\frac{2}{3}}$

d. $512^{\frac{2}{9}}$

e. $16^{-\frac{1}{2}}$

f. $4^{-\frac{1}{2}}$

g. $32^{-\frac{1}{5}}$

h. $49^{-\frac{1}{2}}$

23. a. Use the index laws to simplify the following.

i. $(3^2)^{\frac{1}{2}}$

ii. $(4^2)^{\frac{1}{2}}$

iii. $(8^2)^{\frac{1}{2}}$

iv. $(11^2)^{\frac{1}{2}}$

- b. Use your answers from part a to calculate the value of the following.


i. $9^{\frac{1}{2}}$

ii. $16^{\frac{1}{2}}$

iii. $64^{\frac{1}{2}}$

iv. $121^{\frac{1}{2}}$

- c. Use your answers to parts a and b to write a sentence describing what happens when you raise a number to a power of one-half.



LESSON

1.6 Negative indices

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- apply the Seventh Index Law to evaluate expressions using negative indices
- apply the first seven index laws to simplify expressions involving indices.

1.6.1 The Seventh Index Law

eles-4388

- Negative indices occur when the power (or exponent) is a negative number.
For example, 3^{-2} .
- To explain the meaning or value of negative indices it is useful to consider patterns of numbers written in index form.
For example, in the sequence $3^4 = 81$, $3^3 = 27$, $3^2 = 9$, $3^1 = 3$, $3^0 = 1$ each number is $\frac{1}{3}$ of the number before it.

Powers decrease by 1

3^4	3^3	3^2	3^1	3^0
81	27	9	3	1

$\times \frac{1}{3}$ $\times \frac{1}{3}$ $\times \frac{1}{3}$ $\times \frac{1}{3}$

- It is logical, then, that the next numbers in this sequence are:

$$3^{-1} = \frac{1}{3}, 3^{-2} = \frac{1}{9}, 3^{-3} = \frac{1}{27}, 3^{-4} = \frac{1}{81}$$

Powers decrease by 1

3^0	3^{-1}	3^{-2}	3^{-3}	3^{-4}
1	$\frac{1}{3}$	$\frac{1}{9}$	$\frac{1}{27}$	$\frac{1}{81}$

$\times \frac{1}{3}$ $\times \frac{1}{3}$ $\times \frac{1}{3}$ $\times \frac{1}{3}$

The Seventh Index Law

$$a^{-n} = \frac{1}{a^n}, \text{ where } a \neq 0$$

For example, $2^{-3} = \frac{1}{2^3} = \frac{1}{8}$

WORKED EXAMPLE 19 Evaluating using the Seventh Index Law

Evaluate the following.

a. 5^{-2}

b. 7^{-1}

c. $\left(\frac{3}{5}\right)^{-1}$

THINK

a. 1. Apply the rule $a^{-n} = \frac{1}{a^n}$.

2. Simplify and write the answer.

b. Apply the rule $a^{-n} = \frac{1}{a^n}$.

WRITE

a. $5^{-2} = \frac{1}{5^2}$
 $= \frac{1}{25}$

b. $7^{-1} = \frac{1}{7^1}$
 $= \frac{1}{7}$

- c. 1. Apply the Sixth Index Law, $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$.
2. Apply the Seventh Index Law, $a^{-n} = \frac{1}{a^n}$, to the numerator and denominator.
3. Simplify and write the answer.

$$\begin{aligned} \text{c. } \left(\frac{3^1}{5^1}\right) &= \frac{3^{-1}}{5^{-1}} \\ &= \frac{1}{3} \div \frac{1}{5} \\ &= \frac{1}{3} \times \frac{5}{1} \\ &= \frac{5}{3} \end{aligned}$$

WORKED EXAMPLE 20 Writing with positive indices

Write the following with positive indices.

a. x^{-3}

b. $5x^{-6}$

c. $\frac{x^{-3}}{y^{-2}}$

THINK

- a. Apply the Seventh Index Law, $a^{-n} = \frac{1}{a^n}$.
- b. 1. Write in expanded form and then apply the Seventh Index Law, $a^{-n} = \frac{1}{a^n}$.
2. Simplify and write the answer.
- c. 1. Write the fraction using division.
2. Apply the Seventh Index Law, $a^{-n} = \frac{1}{a^n}$.
3. Simplify and write the answer.

WRITE

a. $x^{-3} = \frac{1}{x^3}$

b. $5x^{-6} = 5 \times x^{-6}$
 $= 5 \times \frac{1}{x^6}$
 $= \frac{5}{x^6}$

c. $\frac{x^{-3}}{y^{-2}} = x^{-3} \div y^{-2}$
 $= \frac{1}{x^3} \div \frac{1}{y^2}$
 $= \frac{1}{x^3} \times \frac{y^2}{1}$
 $= \frac{y^2}{x^3}$

WORKED EXAMPLE 21 Simplifying products of powers with positive and negative indices

Simplify the following expressions, writing your answers with positive indices.

a. $x^3 \times x^{-8}$

b. $3x^{-2}y^{-3} \times 5xy^{-4}$

THINK

- a. 1. Apply the First Index Law $a^n \times a^m = a^{m+n}$.
2. Write the answer with a positive index.

WRITE

a. $x^3 \times x^{-8} = x^{3+(-8)}$
 $= x^{-5}$
 $= \frac{1}{x^5}$

b. 1. Write in expanded form and apply the First Index Law.

2. Apply the rule $a^{-n} = \frac{1}{a^n}$.

3. Simplify and write the answer.

$$\begin{aligned} \text{b. } 3x^{-2}y^{-3} \times 5xy^{-4} &= 3 \times 5 \times x^{-2} \times x^1 \times y^{-3} \times y^{-4} \\ &= 15x^{-1}y^{-7} \\ &= \frac{15}{1} \times \frac{1}{x} \times \frac{1}{y^7} \\ &= \frac{15}{xy^7} \end{aligned}$$

WORKED EXAMPLE 22 Simplifying quotients of powers with positive and negative indices

Simplify the following expressions, writing your answers with positive indices.

a. $\frac{t^2}{t^{-5}}$

b. $\frac{15m^{-5}}{10m^{-2}}$

THINK

a. Apply the Second Index Law, $\frac{a^n}{a^m} = a^{n-m}$.

b. 1. Apply the Second Index Law and simplify.

2. Write the answer with positive indices.

WRITE

$$\begin{aligned} \text{a. } \frac{t^2}{t^{-5}} &= t^{2-(-5)} \\ &= t^{2+5} \\ &= t^7 \end{aligned}$$

$$\begin{aligned} \text{b. } \frac{15m^{-5}}{10m^{-2}} &= \frac{15}{10} \times \frac{m^{-5}}{m^{-2}} \\ &= \frac{3}{2} \times m^{-5-(-2)} \\ &= \frac{3}{2} \times m^{-3} \\ &= \frac{3}{2} \times \frac{1}{m^3} \\ &= \frac{3}{2m^3} \end{aligned}$$

DISCUSSION

What strategy will you use to remember the index laws?

Resources



eWorkbook

Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)



Video eLesson

Negative indices (eles-1910)



Interactivity

Individual pathway interactivity: Negative indices (int-4518)
Negative indices (int-6064)

Individual pathways

PRACTISE

1, 2, 5, 8, 11, 14, 17, 21

CONSOLIDATE

3, 6, 9, 12, 15, 18, 19, 22

MASTER

4, 7, 10, 13, 16, 20, 23, 24

Fluency

1. Copy and complete the following patterns.

a. $3^5 = 243$

$3^4 = 81$

$3^3 = 27$

$3^2 =$

$3^1 =$

$3^0 =$

$3^{-1} = \frac{1}{3}$

$3^{-2} = \frac{1}{9}$

$3^{-3} =$

$3^{-4} =$

$3^{-5} =$

b. $5^4 = 625$

$5^3 =$

$5^2 =$

$5^1 =$

$5^0 =$

$5^{-1} =$

$5^{-2} =$

$5^{-3} =$

$5^{-4} =$

c. $10^4 = 10\ 000$

$10^3 =$

$10^2 =$

$10^1 =$

$10^0 =$

$10^{-1} =$

$10^{-2} =$

$10^{-3} =$

$10^{-4} =$

2. **WE19** Evaluate each of the following expressions.

a. 2^{-5}

b. 3^{-3}

c. 4^{-1}

d. 10^{-2}

3. Evaluate each of the following expressions.

a. 5^{-3}

b. $\left(\frac{1}{7}\right)^{-1}$

c. $\left(\frac{3}{4}\right)^{-1}$

d. $\left(\frac{3}{4}\right)^{-2}$

4. Evaluate each of the following expressions.

a. $\left(\frac{1}{3}\right)^{-3}$

b. $\left(\frac{3}{2}\right)^{-1}$

c. $\left(2\frac{1}{4}\right)^{-2}$

d. $\left(\frac{2}{7}\right)^{-2}$

5. **WE20** Write each expression with positive indices.

a. x^{-4}

b. y^{-5}

c. z^{-1}

d. a^2b^{-3}

e. $m^{-2}n^{-3}$

6. Write each expression with positive indices.

a. $(m^2n^3)^{-1}$

b. $\frac{x^2}{y^{-2}}$

c. $\frac{5}{x^{-3}}$

d. $\frac{x^{-2}}{w^{-5}}$

e. $\frac{1}{x^{-2}y^{-2}}$

7. Write each expression with positive indices.

a. $a^2b^{-3}cd^{-4}$

b. $\frac{a^2b^{-2}}{c^2d^{-3}}$

c. $10x^{-2}y$

d. $3^{-1}x$

e. $\frac{m^{-3}}{x^2}$

Understanding

8. **WE21** Simplify the following expressions, writing your answers with positive indices.

a. $a^3 \times a^{-8}$ b. $m^7 \times m^{-2}$ c. $m^{-3} \times m^{-4}$ d. $2x^{-2} \times 7x$ e. $2b^{-2} \times 3b^2$ f. $ab^{-5} \times 5b^2$

9. Simplify the following expressions, writing your answers with positive indices.

a. $x^5 \times x^{-8}$ b. $3x^2y^{-4} \times 2x^{-7}y$ c. $10x^5 \times 5x^{-2}$
d. $x^5 \times x^{-5}$ e. $10a^2 \times 5a^{-7}$ f. $10a^{10} \times a^{-6}$

10. Simplify the following expressions, writing your answers with positive indices.

a. $16w^2 \times -2w^{-5}$ b. $4m^{-2} \times 4m^{-2}$ c. $(3m^2n^{-4})^3$
d. $(a^2b^5)^{-3}$ e. $(a^{-1}b^{-3})^{-2}$ f. $(5a^{-1})^2$

11. **WE22** Simplify the following expressions, writing your answers with positive indices.

a. $\frac{x^3}{x^8}$ b. $\frac{x^{-3}}{x^8}$ c. $\frac{x^3}{x^{-8}}$ d. $\frac{x^{-3}}{x^{-8}}$ e. $\frac{6a^2c^5}{a^4c}$

12. Simplify the following expressions, writing your answers with positive indices.

a. $10a^2 \div 5a^8$ b. $5m^7 \div m^8$ c. $\frac{a^5b^6}{a^5b^7}$ d. $\frac{a^2b^8}{a^5b^{10}}$ e. $\frac{a^{-3}bc^3}{abc}$

13. Simplify the following expressions, writing your answers with positive indices.

a. $\frac{4^{-2}ab}{a^2b}$ b. $\frac{m^{-3} \times m^{-5}}{m^{-5}}$ c. $\frac{2t^2 \times 3t^{-5}}{4t^6}$ d. $\frac{t^3 \times t^{-5}}{t^{-2} \times t^{-3}}$ e. $\frac{(m^2n^{-3})^{-1}}{(m^{-2}n^3)^2}$

14. Write the following numbers as powers of 2.

a. 1 b. 8 c. 32 d. 64 e. $\frac{1}{8}$ f. $\frac{1}{32}$

15. Write the following numbers as powers of 4.

a. 1 b. 4 c. 64 d. $\frac{1}{4}$ e. $\frac{1}{16}$ f. $\frac{1}{64}$

16. Write the following numbers as powers of 10.

a. 1 b. 10 c. 10 000 d. 0.1 e. 0.01 f. 0.000 01

Reasoning

17. Answer the following questions.

- The result of dividing 3^7 by 3^3 is 3^4 . Determine the result of dividing 3^3 by 3^7 .
- Explain what it means to have a negative index.
- Explain how you write a negative index as a positive index.

18. Indices are encountered in science, where they help to deal with very small and large numbers. The diameter of a proton is 0.000 000 000 000 3 cm.

Explain why it is logical to express this number in scientific notation as 3×10^{-13} .

19. Answer the following questions.

- When asked to write an expression with positive indices that is equivalent to $x^3 + x^{-3}$, a student gave the answer x^0 . Is this answer correct?
Explain why or why not.
- When asked to write an expression with positive indices that is equivalent to $(x^{-1} + y^{-1})^{-2}$, a student gave the answer $x^2 + y^2$. Is this answer correct?
Explain why or why not.

20. Answer the following questions.

- a. When asked to write an expression with positive indices that is equivalent to $x^8 - x^{-5}$, a student gave the answer x^3 . Is this answer correct? Explain why or why not.
- b. Another student said that $\frac{x^2}{x^8 - x^5}$ is equivalent to $\frac{1}{x^6} - \frac{1}{x^3}$. Is this answer correct? Explain why or why not.

Problem solving

21. Write the following numbers as basic numerals.

- a. 4.8×10^{-2} b. 7.6×10^3 c. 2.9×10^{-4} d. 8.1×10^0

22. Determine the value of n in the following expressions.

- a. $4793 = 4.793 \times 10^n$ b. $0.631 = 6.31 \times 10^n$ c. $134 = 1.34 \times 10^n$ d. $0.000\ 56 = 5.6 \times 10^n$

23. Evaluate the following.

- a. Half of 2^{20} b. One-third of 3^{21}

24. Simplify the following expressions.

- a. $(2^{-1} + 3^{-1})^{-1}$ b. $\frac{3^{400}}{6^{200}}$

LESSON

1.7 Scientific notation

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify numbers that are written in scientific notation
- write numbers in scientific notation as decimal numbers
- write decimal numbers using scientific notation.

1.7.1 Using scientific notation


eles-4389

- Scientists often work with extremely large and extremely small numbers. These can range from numbers as large as 1 000 000 000 000 000 km (the approximate diameter of our galaxy) down to numbers as small as 0.000 000 000 06 mm (the diameter of an atom).
- Doing calculations with numbers written this way can be challenging — it can be easy to lose track of all of those zeros.
- It is more useful in these situations to use a notation system based on powers of 10. This system is called **scientific notation** (or **standard form**).
- A number written in scientific notation looks like this:

$$\begin{array}{ccc} & 5.316 \times 10^2 & \\ & \swarrow \quad \searrow & \\ \text{A number between} & \times & \text{A power of 10} \\ \text{1 and 10} & & \end{array}$$

Using a calculator for scientific notation

- Numbers written in scientific notation can be entered into a calculator using its scientific notation buttons.
- Some calculators have different ways of displaying numbers written in scientific notation.
For example, some calculators display 5.71×10^4 as 5.71E4.
- When you are writing down scientific notation yourself, you should always show this number as 5.71×10^4 .



Writing any number in scientific notation

1. Move the decimal point so that it is between the first and second significant figures.
2. Consider how many places the decimal point has moved — this number of places corresponds to the appropriate power of 10.
3. If the decimal point has moved to the left, the power of 10 will have a positive index (this happens with very large numbers).
4. If the decimal point has moved to the right, the power of 10 will have a negative index (this happens with very small numbers).

WORKED EXAMPLE 23 Writing decimal numbers in scientific notation

Write each of these numbers in scientific notation.

- 827.2
- 51 920 000 000
- 0.0051
- 0.000 000 007 648

THINK

1. The numerical part must be written so it is a number between 1 and 10. This means we must move the decimal point 2 steps to the left.
 2. Moving the decimal point 2 steps left corresponds to the power 10^2 .
Note that the number of steps moved is equal to the index.
1. Even though the decimal point is not written, we know that it lies after the final zero.
 2. To get a number between 1 and 10 we move the decimal 10 steps to the left. The corresponding power will be 10^{10} .
1. To get a number between 1 and 10 we must move the decimal point 3 steps to the right. This corresponds to the power 10^{-3} .
Note that moving to the right gives a negative index.
1. To get a number between 1 and 10 we must move the decimal point 9 steps to the right. This corresponds to the power 10^{-9} .

WRITE

a. 8.272
←
Decimal moves 2 steps left
 8.272×10^2

b. 5.192×10^{10}

c. 5.1×10^{-3}

d. 7.648×10^{-9}

1.7.2 Converting scientific notation to decimal notation

- Converting scientific notation to decimal notation is simply the reverse of the process outlined in section 1.7.1.

WORKED EXAMPLE 24 Converting scientific notation with positive indices to decimals

The following numbers are written in scientific notation. Write them in decimal notation.

a. 7.136×10^2

b. 5.017×10^5

c. 8×10^6

THINK

- The index on the power of 10 is positive 2, therefore we move the decimal point 2 steps to the right.
- The index on the power of 10 is positive 5, therefore we move the decimal point 5 steps to the right. To do this we will need to add extra zeros.
- The index on the power of 10 is positive 6, therefore we move the decimal point 6 steps to the right. While the decimal point has not been written as part of the scientific notation, we know it lies after the 8. We will need to add extra zeros.

WRITE

a. $7.136 \times 10^2 = 713.6$

b. $5.017 \times 10^5 = 501\,700$

c. $8 \times 10^6 = 8\,000\,000$

WORKED EXAMPLE 25 Converting scientific notation with negative indices to decimals

Write these numbers in decimal notation.

a. 9.12×10^{-1}

b. 7.385×10^{-2}

c. 6.32×10^{-7}

THINK

- The index on the power of 10 is negative 1, therefore we move the decimal point 1 step to the left.
- The index on the power of 10 is negative 2, therefore we move the decimal point 2 steps to the left. To do this we will need to add extra zeros.
- The index on the power of 10 is negative 7, therefore we move the decimal point 7 steps to the left. To do this we will need to add extra zeros.

WRITE

a. $9.12 \times 10^{-1} = 0.912$

b. $7.3857 \times 10^{-2} = 0.073\,857$

c. $6.32 \times 10^{-7} = 0.000\,000\,632$

WORKED EXAMPLE 26 Representing decimals in exponential form

Write these numbers in exponential form.

a. 0.475

b. 0.03

c. 0.00023

THINK

- Write 0.475 in expanded fraction form.
- Express the expanded form in the exponential form.

WRITE

a. $0.475 = \frac{4}{10} + \frac{7}{100} + \frac{5}{1000}$
 $= 4 \times 10^{-1} + 7 \times 10^{-2} + 5 \times 10^{-3}$

- b. 1. Write 0.03 in expanded fraction form.
2. Express the expanded fraction from in the exponential form.

$$\begin{aligned} \text{b. } 0.03 &= \frac{0}{10} + \frac{3}{100} \\ &= 3 \times 10^{-2} \end{aligned}$$

- c. 1. Write 0.00023 in expanded form.
2. Express the expanded form in the exponential form.

$$\begin{aligned} \text{c. } 0.00023 &= \frac{0}{10} + \frac{0}{100} + \frac{0}{1000} + \\ &\quad \frac{2}{10\,000} + \frac{3}{100\,000} \\ &= 2 \times 10^{-4} + 3 \times 10^{-5} \end{aligned}$$



Metric prefixes for SI units

The following prefixes allow us to describe tiny numbers as well as incredibly large numbers of units succinctly.

Symbol	Prefix	Multiplication Factor	
T	tera	10^{12}	1 000 000 000 000
G	giga	10^9	1 000 000 000
M	mega	10^6	1 000 000
k	kilo	10^3	1 000
h	hecto	10^2	100
da	deka	10^1	10
d	deci	10^{-1}	0.1
c	centi	10^{-2}	0.01
m	milli	10^{-3}	0.001
μ	micro	10^{-6}	0.000 001
n	nano	10^{-9}	0.000 000 001
p	pico	10^{-12}	0.000 000 000 001

For example, 1 pico-gram = 10^{-12} gram and 1 terabyte = 10^{12} bytes

Resources

-  **eWorkbook** Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)
-  **Interactivities** Individual pathway interactivity: Scientific notation (int-4479)
Scientific notation (int-6031)

1.7 Quick quiz

1.7 Exercise

Individual pathways

PRACTISE

1, 4, 6, 7, 10, 13, 14, 19, 23

CONSOLIDATE

2, 5, 8, 11, 15, 16, 20, 24

MASTER

3, 9, 12, 17, 18, 21, 22, 25, 26

Fluency

1. **WE23** Write these numbers in scientific notation.

- a. 5000 b. 431 c. 38 d. 350 000

2. Write these numbers in decimal notation.

- a. 1×10^9 b. $3.926\ 73 \times 10^2$ c. 5.911×10^2 d. 5.1×10^3

3. Write these numbers in decimal notation.

- a. 7.34×10^5 b. 7.1414×10^6 c. 3.51×10 d. 8.05×10^4

4. **WE24** Write these numbers in decimal notation.

- a. 6.14×10^2 b. 6.14×10^3 c. 6.14×10^4 d. 3.518×10^2

5. Write these numbers in scientific notation.

- a. 72.5 b. 725 c. 7250 d. 725 000 000

6. Write these numbers in scientific notation, correct to 4 significant figures.

- a. 43.792 b. 5317 c. 258.95 d. 110.11 e. 1 632 000 f. 1 million

7. **WE25** Write these numbers in decimal notation.

- a. 2×10^{-1} b. 4×10^{-3} c. 7×10^{-4} d. 3×10^{-2}

8. Write these numbers in decimal notation.

- a. 8.273×10^{-2} b. 7.295×10^{-2} c. 2.9142×10^{-3} d. 3.753×10^{-5}

9. Write these numbers in decimal notation.

- a. 5.29×10^{-4} b. 3.3333×10^{-5} c. 2.625×10^{-9} d. 1.273×10^{-15}

10. Write these numbers in scientific notation.

- a. 0.7 b. 0.005 c. 0.000 000 3 d. 0.000 000 000 01

11. **WE26** Write these numbers in exponential form.

- a. 0.231 b. 0.003 62 c. 0.000 7 d. 0.063

12. Write these numbers in scientific notation, correct to 3 significant figures.

- a. 0.006 731 b. 0.142 57 c. 0.000 068 3
d. 0.000 000 005 12 e. 0.0509 f. 0.012 46

Understanding


13. Write each of the following sets of numbers in ascending order.

- a. 8.31×10^2 , 3.27×10^3 , 9.718×10^2 , 5.27×10^2
b. 7.95×10^2 , 4.09×10^2 , 7.943×10^2 , 4.37×10^2
c. 5.31×10^{-2} , 9.29×10^{-3} , 5.251×10^{-2} , 2.7×10^{-3}
d. 8.31×10^2 , 3.27×10^3 , 7.13×10^{-2} , 2.7×10^{-3}

14. One carbon atom weighs 1.994×10^{-23} g.
- Write this weight as a decimal.
 - Calculate how much one million carbon atoms will weigh.
 - Calculate how many carbon atoms there are in 10 g of carbon. Give your answer correct to 4 significant figures.



15. The distance from Earth to the Moon is approximately 3.844×10^5 km. If you could drive there at a constant speed of 100 km/h, calculate how long it would take. Determine how long that is in days, correct to 2 decimal places.
16. Earth weighs 5.97×10^{24} kg and the Sun weighs 1.99×10^{30} kg. Calculate how many Earths it would take to balance the Sun's weight. Give your answer correct to 2 decimal places.
17. Inside the nucleus of an atom, a proton weighs 1.6703×10^{-28} kg and a neutron weighs 1.6726×10^{-27} kg. Determine which one is heavier and by how much.
18. Earth's orbit has a radius of 7.5×10^7 km and the orbit of Venus has a radius of 5.4×10^7 km.



Calculate how far apart the planets are when:

- they are closest to each other
- they are furthest apart from each other.

Reasoning

19. A USB stick has 8 MB (megabytes) of storage.
1 MB = 1 048 576 bytes.
- Determine the number of bytes in 8 MB of storage correct to the nearest 1000.
 - Write the answer to part **a** in scientific notation.
20. The basic unit of electric current is the ampere. It is defined as the constant current flowing in 2 parallel conductors 1 metre apart in a vacuum, which produces a force between the conductors of 2×10^{-7} newtons (N) per metre.
Complete the following statement. Write the answer as a decimal.
1 ampere = 2×10^{-7} N/m = _____ N/m
21. **a.** Without performing the calculation, state the power(s) of 10 that you believe the following equations will have when solved.
- $5.36 \times 10^7 + 2.95 \times 10^3$
 - $5.36 \times 10^7 - 2.95 \times 10^3$
- Evaluate equations **i** and **ii**, correct to 3 significant figures.
 - Were your answers to part **a** correct? Why or why not?
22. Explain why $2.39 \times 10^{-3} + 8.75 \times 10^{-7} = 2.39 \times 10^{-3}$, correct to 3 significant figures.


Problem solving

- 23.** Distance is equal to speed multiplied by time. If we travelled at 100 km/h it would take us approximately 0.44 years to reach the Moon, 89.6 years to reach Mars, 1460 years to reach Saturn and 6590 years to reach Pluto.
- Assuming that there are 365 days in a year, calculate the distance (as a basic numeral) between Earth and:
 - the Moon
 - Mars
 - Saturn
 - Pluto.
 - Write your answers to part **a**, correct to 3 significant figures.
 - Write your answers to part **a** using scientific notation, correct to 3 significant figures.
- 24.** A light-year is the distance that light travels in one year. Light travels at approximately 300 000 km/s.
- Calculate the number of seconds in a year (assuming 1 year = 365 days).
 - Write your answer to part **i** using scientific notation.
 - Calculate the distance travelled by light in one year. Express your answer:
 - as a basic numeral
 - using scientific notation.
 - The closest star to Earth (other than the Sun) is in the star system Alpha Centauri, which is 4.3 light-years away.
 - Determine how far this is in kilometres, correct to 4 significant figures.
 - If you were travelling at 100 km/h, determine how many years it would take to reach Alpha Centauri.



- 25.** Scientists used Earth's gravitational pull on nearby celestial bodies (for example, the Moon) to calculate the mass of Earth. Their answer was that Earth weighs approximately 5.972 sextillion metric tonnes.
- Write 5.972 sextillion using scientific notation.
 - Determine how many significant figures this number has.
- 26.** Atoms are made up of smaller particles called protons, neutrons and electrons. Electrons have a mass of $9.109\,381\,88 \times 10^{-31}$ kilograms, correct to 9 significant figures.

- Write the mass of an electron correct to 5 significant figures.
- Protons and neutrons are the same size. They are both 1836 times the size of an electron. Use the mass of an electron (correct to 9 significant figures) and your calculator to evaluate the mass of a proton correct to 5 significant figures.
- Use the mass of an electron (correct to 3 significant figures) to calculate the mass of a proton, correct to 5 significant figures.
- Explain why it is important to work with the original amounts and then round to the specified number of significant figures at the end of a calculation.



LESSON

1.8 Square roots and cube roots

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- write square and cube roots using the radical symbol
- use fractional indices to represent square roots and cube roots
- simplify expressions that use square roots and cube roots.



1.8.1 Square roots

eles-4391

- The symbol $\sqrt{\quad}$ (the radical symbol) means ‘**square root**’ — a number that multiplies by itself to give the original number.
- Every number actually has a positive and negative square root. For example, $(2)^2 = 4$ and $(-2)^2 = 4$.
- This means that the square root of 4 is +2 or -2. For this topic, however, assume that $\sqrt{\quad}$ is positive unless otherwise indicated.
- A square root is the inverse of squaring (raising to the power 2).
- Because of this, a square root of a number is equivalent to raising that number to an index of $\frac{1}{2}$.
- In general, $\sqrt{a} = a^{\frac{1}{2}}$.



WORKED EXAMPLE 27 Evaluating square roots

Evaluate $\sqrt{16p^2}$.

THINK

1. We need to work out the square root of both 16 and p^2 .
2. Remember that 4 is multiplied by itself to give 16, so the square root of 16 is 4. For the square root of the pronumeral, replace the square root sign with a power of $\frac{1}{2}$.
3. Use the Fourth Index Law.
4. Simplify and write the answer.

WRITE

$$\begin{aligned}\sqrt{16p^2} &= \sqrt{16} \times \sqrt{p^2} \\ &= 4 \times (p^2)^{\frac{1}{2}} \\ &= 4 \times p^{2 \times \frac{1}{2}} \\ &= 4 \times p^1 \\ &= 4p\end{aligned}$$



1.8.2 Cube roots

eles-4392

- The symbol $\sqrt[3]{\quad}$ means ‘**cube root**’ — a number that multiplies by itself 3 times to give the original number.
- The cube root is the inverse of cubing (raising to the power 3).
- Because of this, a cube root is equivalent to raising a number to an index of $\frac{1}{3}$.
- In general, $\sqrt[3]{a} = a^{\frac{1}{3}}$.
- In more general terms, $a^{\frac{n}{m}} = \sqrt[m]{a^n}$.

WORKED EXAMPLE 28 Evaluating cube roots

Evaluate $\sqrt[3]{8j^6}$

THINK

1. We need to work out the cube root of both 8 and j^6 .
2. Remember that 2 multiplied by itself 3 times gives 8, so the cube root of 8 is 2. For the cube root of the pronumeral, replace the cube root sign with a power of $\frac{1}{3}$.
3. Use the Fourth Index Law.
4. Simplify and write the answer.

WRITE

$$\begin{aligned}\sqrt[3]{8j^6} &= \sqrt[3]{8} \times \sqrt[3]{j^6} \\ &= 2 \times (j^6)^{\frac{1}{3}} \\ &= 2 \times j^{6 \times \frac{1}{3}} \\ &= 2 \times j^2 \\ &= 2j^2\end{aligned}$$

DISCUSSION

Why does the cube root of a number always have the same sign (positive or negative) as the number itself?

on Resources



eWorkbook Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)



Interactivities Individual pathway interactivity: Square roots and cube roots (int-4519)

Cube roots (int-6065)

Square roots (int-6066)

Exercise 1.8 Square roots and cube roots

learn **on**

1.8 Quick quiz **on**

1.8 Exercise

Individual pathways

PRACTISE

1, 4, 7, 10, 14

CONSOLIDATE

2, 5, 8, 11, 15, 16

MASTER

3, 6, 9, 12, 13, 17

Fluency

1. Write the following in index form.

a. $\sqrt{15}$

b. \sqrt{m}

c. $\sqrt[3]{t}$

d. $\sqrt[3]{w^2}$

2. Evaluate the following.

a. $49^{\frac{1}{2}}$

b. $4^{\frac{1}{2}}$

c. $27^{\frac{1}{3}}$

d. $125^{\frac{1}{3}}$

3. Evaluate the following.

a. $1000^{\frac{1}{3}}$

b. $64^{\frac{1}{2}}$

c. $64^{\frac{1}{3}}$

d. $1\,000\,000^{\frac{1}{2}}$

e. $1\,000\,000^{\frac{1}{3}}$

f. $\left(27^{\frac{1}{3}}\right)^2$

Understanding

4. **WE27** Simplify the following expressions.

a. $\sqrt{m^2}$

b. $\sqrt[3]{b^3}$

c. $\sqrt{36t^4}$

d. $\sqrt[3]{m^3n^6}$

5. Simplify the following expressions.

a. $\sqrt[3]{125t^6}$

b. $\sqrt[5]{x^5y^{10}}$

c. $\sqrt[4]{a^8m^{40}}$

d. $\sqrt[3]{216y^6}$

6. Simplify the following expressions.

a. $\sqrt[3]{64x^6y^6}$

b. $\sqrt{25a^2b^4c^6}$

c. $\sqrt[7]{b^{49}}$

d. $\sqrt[3]{b^3} \times \sqrt{b^4}$

7. **WE28 MC** $\sqrt[3]{8000m^6n^3p^3q^6}$ is equal to:

A. $2666.6m^2npq^2$

B. $20m^2npq^2$

C. $20m^3n^0p^0q^3$

D. $7997m^2npq^2$

E. $2m^2npq^2$

8. **MC** $\sqrt[3]{3375a^9b^6c^3}$ is equal to:

A. $1125a^3b^2c$

B. $1125a^6b^3c^0$

C. $1123a^6b^3$

D. $15a^3b^2c$

E. $5a^3b^2c$

9. **MC** $\sqrt[3]{15\,625f^3g^6h^9}$ is equal to:

A. $25fg^2h^3$

B. $25f^0g^3h^6$

C. $25g^3h^6$

D. $5208.3fg^2h^3$

E. $250fg^2h^3$

Reasoning

10. Answer the following questions.

a. Using the First Index Law, explain how $3^{\frac{1}{2}} \times 3^{\frac{1}{2}} = 3$.

b. Express $3^{\frac{1}{2}}$ in another way.

c. Write $\sqrt[n]{a}$ in index form.

d. Without a calculator, evaluate $8^{\frac{1}{3}}$.

11. Answer the following questions.

a. Explain why calculating $z^{2.5}$ is a square root problem.

b. Explain whether $z^{0.3}$ is a cube root problem.

12. Prithvi and Yasmin are having an algebra argument. Prithvi is sure that $\sqrt{x^2}$ is equivalent to x , but Yasmin thinks otherwise. State who is correct.

Explain how you would resolve this disagreement.

13. Verify that $(-8)^{\frac{1}{3}}$ can be evaluated and explain why $(-8)^{\frac{1}{4}}$ cannot be evaluated.

Problem solving

14. The British mathematician Augustus de Morgan enjoyed telling his friends that he was x years old in the year x^2 . Determine the year of Augustus de Morgan's birth, given that he died in 1871.

15. Kepler's Third Law describes the relationship between the distance of planets from the Sun and their orbital periods. It is represented by the equation $d^{\frac{1}{2}} = t^{\frac{1}{3}}$. Solve for:
- d in terms of t
 - t in terms of d .
16. If $n^{\frac{3}{4}} = \frac{8}{27}$, evaluate the value of n .
17. An unknown number is multiplied by 4 and then has 5 subtracted from it. It is now equal to the square root of the original unknown number, squared.
- Determine how many solutions to this problem are possible. Explain why.
 - Determine all possible values for the unknown number.

LESSON

1.9 Surds (extending)

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify when a square root forms a surd
- place surds in their approximate position on a number line
- simplify expressions containing surds.




1.9.1 Identifying surds

eles-4393

- In many cases the square root or cube root of a number results in a rational number. For example, $\sqrt{25} = \pm 5$ and $\sqrt[3]{1.728} = 1.2$.
- When the square root of a number is an irrational number, it is called a **surd**.
- For example, $\sqrt{10} \approx 3.162\ 277\ 660\ 17\ \dots$. Since $\sqrt{10}$ cannot be written as a fraction, a recurring decimal or a **terminating decimal**, it is irrational and therefore it is a surd.
- The value of a surd can be approximated using a number line.

For example, we know that $\sqrt{21}$ lies between 4 and 5, because it lies between $\sqrt{16}$ (which equals 4) and $\sqrt{25}$ (which equals 5).

We can show its approximate position on the number line like this:



- Geometric constructions can be used to locate rational numbers and square root on a number line. For example, $\sqrt{2}$ is located at the intersection of an arc and the number line, where the radius of the arc is the length of the diagonal of a one-unit square.





WORKED EXAMPLE 29 Locating the square root on a number line

Represent $\sqrt{2}$ on the number line.

THINK

1. Start with a horizontal number line from 0 to 2.
2. Extend a vertical number line up from the origin.
3. Draw in a unit square with its lower left corner at the origin. All sides of the unit square are 1 unit.
4. Draw a diagonal of the unit square from the origin, O to the opposite corner, B.

WRITE/ DRAW



5. Apply the Pythagoras theorem to determine the length of the side OB:

$$OB^2 = OA^2 + AB^2$$


$$OB^2 = OA^2 + AB^2$$

$$= 1^2 + 1^2$$


$$= 2$$

$$OB = \sqrt{2}$$


The length of the diagonal is $\sqrt{2}$.



6. Now to locate $\sqrt{2}$ accurately on the number line, draw an arc with its centre at the origin, O and its radius (diagonal) of $\sqrt{2}$ using a compass.



7. Label $\sqrt{2}$ at the intersections of the arc on both the horizontal and vertical number lines.



$\sqrt{2}$ is located at the intersection of an arc and the number line, where the radius of the arc is the length of the diagonal of a one-unit square.

WORKED EXAMPLE 30 Identifying surds

Identify which of the following are surds.

- | | |
|----------------|------------------|
| a. $\sqrt{0}$ | b. $\sqrt{20}$ |
| c. $-\sqrt{9}$ | d. $\sqrt[3]{6}$ |

THINK

- a. $\sqrt{0} = 0$. This is a rational number and therefore not a surd.
- b. $\sqrt{20} \approx 4.472\ 135\ 955\dots$ This is an irrational number and therefore a surd.
- c. $-\sqrt{9} = -3$. This is a rational number and therefore not a surd.
- d. $\sqrt[3]{6} \approx 1.817\ 120\ 592\ 83\dots$ This is an irrational number and therefore a surd.

WRITE

- a. $\sqrt{0} = 0$. This is not a surd.
- b. $\sqrt{20}$ is a surd.
- c. $-\sqrt{9} = -3$. This is not a surd.
- d. $\sqrt[3]{6}$ is a surd.

1.9.2 Multiplying and dividing surds

eles-4394

- Multiplication of surds is done by multiplying the numerical parts of each surd under a single radical sign.
- Division of surds is done by placing the quotient of the numerical parts of each surd under a single radical sign.

Multiplying surds

$$\sqrt{a} \times \sqrt{b} = \sqrt{ab}$$

Dividing surds

$$\frac{\sqrt{a}}{\sqrt{b}} = \sqrt{\frac{a}{b}}$$

WORKED EXAMPLE 31 Multiplying surds

Evaluate the following, leaving your answer in surd form.

a. $\sqrt{7} \times \sqrt{2}$

b. $5 \times \sqrt{3}$

c. $\sqrt{5} \times \sqrt{5}$

d. $-2\sqrt{3} \times 4\sqrt{5}$

THINK

a. Apply the rule $\sqrt{a} \times \sqrt{b} = \sqrt{ab}$.

b. Only $\sqrt{3}$ is a surd. It is multiplied by 5, which is not a surd.

c. Apply the rule $\sqrt{a} \times \sqrt{b} = \sqrt{ab}$.

d. Multiply the whole numbers. Then multiply the surds.

WRITE

a. $\sqrt{7} \times \sqrt{2} = \sqrt{14}$

b. $5 \times \sqrt{3} = 5\sqrt{3}$

c. $\sqrt{5 \times 5} = \sqrt{25}$
 $= 5$

d. $-2\sqrt{3} \times 4\sqrt{5} = -2 \times 4 \times \sqrt{3} \times \sqrt{5}$
 $= -8 \times \sqrt{15}$
 $= -8\sqrt{15}$

WORKED EXAMPLE 32 Dividing surds

Evaluate the following, leaving your answer in surd form.

a. $\frac{\sqrt{10}}{\sqrt{5}}$

b. $\sqrt{\frac{10}{5}}$

c. $\frac{-6\sqrt{8}}{4\sqrt{4}}$

d. $\frac{\sqrt{20}}{\sqrt{5}}$

e. $\frac{5}{\sqrt{5}}$

THINK

a. Apply the rule $\sqrt{a} \div \sqrt{b} = \sqrt{\frac{a}{b}}$.

b. Simplify the fraction.

c. Simplify the whole numbers. Then apply the rule

$$\sqrt{a} \div \sqrt{b} = \sqrt{\frac{a}{b}}$$

d. Apply the rule $\sqrt{a} \div \sqrt{b} = \sqrt{\frac{a}{b}}$.

e. Rewrite the numerator as the product of two surds and then simplify.

WRITE

a. $\frac{\sqrt{10}}{\sqrt{5}} = \sqrt{\frac{10}{5}}$
 $= \sqrt{2}$

b. $\sqrt{\frac{10}{5}} = \sqrt{2}$

c. $\frac{-6\sqrt{8}}{4\sqrt{4}} = \frac{-3\sqrt{2}}{2}$

d. $\frac{\sqrt{20}}{\sqrt{5}} = \sqrt{\frac{20}{5}} = \sqrt{4}$
 $= 2$

e. $\frac{5}{\sqrt{5}} = \frac{\sqrt{5} \times \sqrt{5}}{\sqrt{5}}$
 $= \sqrt{5}$

1.9.3 Simplifying surds

eles-4395

- Just as a rational number can be written many different ways (e.g. $\frac{1}{2} = \frac{5}{10} = \frac{7}{14}$), so can a surd. It is expected that surds should normally be written in their simplest form.
- A surd is in its simplest form when the number inside the radical sign has the smallest possible value.
- Note that $\sqrt{24}$ can be factorised several ways.

For example:

$$\sqrt{24} = \sqrt{2} \times \sqrt{12}$$

$$\sqrt{24} = \sqrt{3} \times \sqrt{8}$$

$$\sqrt{24} = \sqrt{4} \times \sqrt{6}$$

In the last example, $\sqrt{4} = 2$, which means:

$$\begin{aligned}\sqrt{24} &= 2 \times \sqrt{6} \\ &= 2\sqrt{6}\end{aligned}$$

$2\sqrt{6}$ is equal to $\sqrt{24}$, and $2\sqrt{6}$ is $\sqrt{24}$ written in its simplest form.

- To simplify a surd you must determine a factor that is also a perfect square. For example, 4, 9, 16, 25, 36 or 49.
- A surd like $\sqrt{22}$, for example, cannot be simplified because 22 has no perfect square factors.
- Surds can be simplified in more than one step.

$$\begin{aligned}\sqrt{72} &= \sqrt{4} \times \sqrt{18} \\ &= 2\sqrt{18} \\ &= 2 \times \sqrt{9} \times \sqrt{2} \\ &= 2 \times 3\sqrt{2} \\ &= 6\sqrt{2}\end{aligned}$$

WORKED EXAMPLE 33 Simplifying surds

Simplify the following surds.

a. $\sqrt{18}$

b. $6\sqrt{20}$

THINK

a. 1. Rewrite 18 as the product of two numbers, one of which is a perfect square (9).

2. Simplify.

b. 1. Rewrite 20 as the product of two numbers, one of which is square (4).

2. Simplify.

WRITE

a. $\sqrt{18} = \sqrt{9} \times \sqrt{2}$

$$\begin{aligned}&= 3 \times \sqrt{2} \\ &= 3\sqrt{2}\end{aligned}$$

b. $6\sqrt{20} = 6 \times \sqrt{4} \times \sqrt{5}$

$$\begin{aligned}&= 6 \times 2 \times \sqrt{5} \\ &= 12\sqrt{5}\end{aligned}$$

1.9.4 Entire surds

- The surd $\sqrt{45}$, when simplified, is written as $3\sqrt{5}$.
The surd $\sqrt{45}$ is called an entire surd because it is written entirely inside the radical sign. The surd $3\sqrt{5}$, however, is not an entire surd.
- Writing a surd as an entire surd reverses the process of simplification.

WORKED EXAMPLE 34 Writing entire surds

Write $3\sqrt{7}$ as an entire surd.

THINK

- In order to place the 3 inside the radical sign it has to be written as $\sqrt{9}$.
- Apply the rule $\sqrt{a} \times \sqrt{b} = \sqrt{ab}$.

WRITE

$$\begin{aligned} 3\sqrt{7} &= \sqrt{9} \times \sqrt{7} \\ &= \sqrt{63} \end{aligned}$$

WORKED EXAMPLE 35 Comparing surds

Determine which number is larger, $3\sqrt{5}$ or $5\sqrt{3}$.

THINK

- Write $3\sqrt{5}$ as an entire surd.
- Write $5\sqrt{3}$ as an entire surd.
- Compare the values of each surd.
- Write your answer.

WRITE

$$\begin{aligned} 3\sqrt{5} &= \sqrt{9} \times \sqrt{5} \\ &= \sqrt{45} \\ 5\sqrt{3} &= \sqrt{25} \times \sqrt{3} \\ &= \sqrt{75} \\ \sqrt{75} &> \sqrt{45} \\ 5\sqrt{3} &\text{ is the larger number.} \end{aligned}$$

1.9.5 Addition and subtraction of surds

- Surds can be added or subtracted if they have like terms.
- Surds should be simplified before adding or subtracting like terms.

WORKED EXAMPLE 36 Adding and subtracting surds

Simplify each of the following.

a. $6\sqrt{3} + 2\sqrt{3} + 4\sqrt{5} - 5\sqrt{5}$

b. $3\sqrt{2} - 5 + 4\sqrt{2} + 9$

THINK

- Collect the like terms ($\sqrt{3}$ and $\sqrt{5}$).
- Collect the like terms and simplify.

WRITE

$$\begin{aligned} \text{a. } 6\sqrt{3} + 2\sqrt{3} + 4\sqrt{5} - 5\sqrt{5} &= 8\sqrt{3} - \sqrt{5} \\ \text{b. } 3\sqrt{2} - 5 + 4\sqrt{2} + 9 &= 3\sqrt{2} + 4\sqrt{2} - 5 + 9 \\ &= 7\sqrt{2} + 4 \end{aligned}$$

WORKED EXAMPLE 37 Simplifying surd expressions

Simplify $5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3}$.

THINK

1. Simplify $5\sqrt{75}$.
2. Next, simplify $6\sqrt{12}$.
3. Lastly, simplify $\sqrt{8}$.
4. Rewrite the original expression and simplify by adding like terms.

WRITE

$$\begin{aligned} 5\sqrt{75} &= 5 \times \sqrt{25} \times \sqrt{3} \\ &= 5 \times 5 \times \sqrt{3} \\ &= 25\sqrt{3} \end{aligned}$$


$$\begin{aligned} 6\sqrt{12} &= 6 \times \sqrt{4} \times \sqrt{3} \\ &= 6 \times 2 \times \sqrt{3} \\ &= 12\sqrt{3} \end{aligned}$$

$$\begin{aligned} \sqrt{8} &= \sqrt{4} \times \sqrt{2} \\ &= 2\sqrt{2} \end{aligned}$$

$$\begin{aligned} 5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3} \\ &= 25\sqrt{3} - 12\sqrt{3} + 2\sqrt{2} - 4\sqrt{3} \\ &= 9\sqrt{3} + 2\sqrt{2} \end{aligned}$$

TI | THINK

In a new problem, on a Calculator page, complete the entry line as: $5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3}$
Press ENTER.




$$\begin{aligned} 5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3} \\ &= 9\sqrt{3} + 2\sqrt{2} \end{aligned}$$

WRITE

CASIO | THINK

Make sure the calculator is set to Standard mode. On the Main screen, complete the entry line as: $5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3}$
Press EXE.




WRITE



$$\begin{aligned} 5\sqrt{75} - 6\sqrt{12} + \sqrt{8} - 4\sqrt{3} \\ &= 9\sqrt{3} + 2\sqrt{2} \end{aligned}$$

DISCUSSION

Are all square root numbers surds?

-  **eWorkbook** Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)
-  **Video eLesson** Surds (eles-1906)
-  **Interactivities** Individual pathway interactivity: Surds (int-4477)
Simplifying surds (int-6028)
Surds on the number line (int-6029)

Exercise 1.9 Surds (extending)

1.9 Quick quiz **on**

1.9 Exercise

Individual pathways

PRACTISE

1, 5, 7, 10, 13, 16, 19, 21, 23, 24,
27, 30, 35, 39

CONSOLIDATE

2, 4, 9, 11, 15, 17, 22, 25, 29, 31,
32, 36, 40

MASTER

3, 6, 8, 12, 14, 18, 20, 26, 28, 33,
34, 37, 38, 41, 42

Fluency

1. Write down the square roots of each of the following.

a. 1	b. 4	c. 0	d. $\frac{1}{9}$	e. $1\frac{9}{16}$
------	------	------	------------------	--------------------
2. Write down the square roots of each of the following.

a. 0.16	b. 400	c. 10 000	d. $\frac{4}{25}$	e. 1.44
---------	--------	-----------	-------------------	---------
3. Write down the square roots of each of the following.

a. 20.25	b. 1 000 000	c. 0.0009	d. 256	e. $2\frac{23}{49}$
----------	--------------	-----------	--------	---------------------
4. Write down the value of each of the following.

a. $\sqrt{81}$	b. $-\sqrt{81}$	c. $\sqrt{121}$	d. $-\sqrt{441}$
----------------	-----------------	-----------------	------------------
5. Write down the value of each of the following.

a. $\sqrt[3]{8}$	b. $\sqrt[3]{64}$	c. $\sqrt[3]{343}$	d. $\sqrt[4]{81}$
------------------	-------------------	--------------------	-------------------
6. Write down the value of each of the following.

a. $\sqrt[5]{1024}$	b. $\sqrt[3]{125}$	c. $-\sqrt{49}$	d. $\sqrt[3]{-27}$
---------------------	--------------------	-----------------	--------------------
7. **WE29** Place the following on a number line.

a. $\sqrt{3}$	b. $\sqrt{7}$	c. $\sqrt{12}$	d. $\sqrt{34}$
---------------	---------------	----------------	----------------
8. **WE30** Identify which of the following are surds.

a. $-\sqrt[3]{216}$	b. $\sqrt[3]{-216}$	c. $-\sqrt{2}$	d. $1 + \sqrt{2}$
---------------------	---------------------	----------------	-------------------
9. Identify which of the following are surds.

a. 1.32	b. $1.\dot{3}2$	c. $\sqrt[4]{64}$	d. 1.752 16
---------	-----------------	-------------------	-------------

10. **WE31** Simplify each of the following.

a. $\sqrt{3} \times \sqrt{7}$ b. $-\sqrt{3} \times \sqrt{7}$ c. $2 \times \sqrt{6}$
 d. $2 \times 3\sqrt{7}$ e. $2\sqrt{7} \times 5\sqrt{2}$

11. Simplify each of the following.

a. $3\sqrt{7} \times 4$ b. $\sqrt{7} \times 9$ c. $-2\sqrt{5} \times -11\sqrt{2}$
 d. $2\sqrt{3} \times 11$ e. $\sqrt{3} \times \sqrt{3}$

12. Simplify each of the following.

a. $-3\sqrt{2} \times -5\sqrt{5}$ b. $2\sqrt{3} \times 4\sqrt{3}$ c. $\sqrt{6} \times \sqrt{6}$
 d. $\sqrt{11} \times \sqrt{11}$ e. $\sqrt{15} \times 2\sqrt{15}$

13. **WE32** Simplify each of the following.

a. $\sqrt{\frac{12}{4}}$ b. $-\sqrt{\frac{10}{5}}$
 c. $\frac{\sqrt{18}}{\sqrt{3}}$ d. $\frac{-\sqrt{15}}{-\sqrt{3}}$

14. Simplify each of the following.

a. $\frac{15\sqrt{6}}{5\sqrt{2}}$ b. $\frac{15\sqrt{6}}{10}$
 c. $\frac{15\sqrt{6}}{\sqrt{3}}$ d. $\frac{5}{3}\sqrt{\frac{15}{3}}$



15. Simplify each of the following.

a. $\frac{-10\sqrt{10}}{5\sqrt{2}}$ b. $\frac{\sqrt{9}}{\sqrt{3}}$ c. $\frac{3}{\sqrt{3}}$ d. $\frac{7}{\sqrt{7}}$

16. **WE33** Simplify each of the following.

a. $\sqrt{20}$ b. $\sqrt{8}$ c. $\sqrt{18}$ d. $\sqrt{49}$

17. Simplify each of the following.

a. $\sqrt{30}$ b. $\sqrt{50}$ c. $\sqrt{28}$ d. $\sqrt{108}$

18. Simplify each of the following

a. $\sqrt{288}$ b. $\sqrt{48}$ c. $\sqrt{500}$ d. $\sqrt{162}$

19. Simplify each of the following.

a. $2\sqrt{8}$ b. $5\sqrt{27}$ c. $6\sqrt{64}$ d. $7\sqrt{50}$ e. $10\sqrt{24}$

20. Simplify each of the following.

a. $5\sqrt{12}$ b. $4\sqrt{42}$ c. $12\sqrt{72}$ d. $9\sqrt{45}$ e. $12\sqrt{242}$

21. **WE34** Write each of the following in the form \sqrt{a} (that is, as an entire surd).

a. $2\sqrt{3}$ b. $5\sqrt{7}$ c. $6\sqrt{3}$ d. $4\sqrt{5}$ e. $8\sqrt{6}$

22. Write each of the following in the form \sqrt{a} (that is, as an entire surd).

a. $3\sqrt{10}$ b. $4\sqrt{2}$ c. $12\sqrt{5}$ d. $10\sqrt{6}$ e. $13\sqrt{2}$

23. **WE35** Determine which number in each of the following pairs of numbers is larger.
- a. $\sqrt{10}$ or $2\sqrt{3}$ b. $3\sqrt{5}$ or $5\sqrt{2}$ c. $10\sqrt{2}$ or $4\sqrt{5}$ d. $2\sqrt{10}$ or $\sqrt{20}$
24. **WE36** Simplify each of the following.
- a. $6\sqrt{2} + 3\sqrt{2} - 7\sqrt{2}$ b. $4\sqrt{5} - 6\sqrt{5} - 2\sqrt{5}$
c. $-3\sqrt{3} - 7\sqrt{3} + 4\sqrt{3}$ d. $-9\sqrt{6} + 6\sqrt{6} + 3\sqrt{6}$
25. Simplify each of the following.
- a. $10\sqrt{11} - 6\sqrt{11} + \sqrt{11}$ b. $\sqrt{7} + \sqrt{7}$
c. $4\sqrt{2} + 6\sqrt{2} + 5\sqrt{3} + 2\sqrt{3}$ d. $10\sqrt{5} - 2\sqrt{5} + 8\sqrt{6} - 7\sqrt{6}$
26. Simplify each of the following.
- a. $5\sqrt{10} + 2\sqrt{3} + 3\sqrt{10} + 5\sqrt{3}$ b. $12\sqrt{2} - 3\sqrt{5} + 4\sqrt{2} - 8\sqrt{5}$
c. $6\sqrt{6} + \sqrt{2} - 4\sqrt{6} - \sqrt{2}$ d. $16\sqrt{5} + 8 + 7 - 11\sqrt{5}$
27. **WE37** Simplify each of the following.
- a. $\sqrt{8} + \sqrt{18} - \sqrt{32}$ b. $\sqrt{45} - \sqrt{80} + \sqrt{5}$
c. $-\sqrt{12} + \sqrt{75} - \sqrt{192}$ d. $\sqrt{7} + \sqrt{28} - \sqrt{343}$
e. $\sqrt{24} + \sqrt{180} + \sqrt{54}$
28. Simplify each of the following.
- a. $\sqrt{12} + \sqrt{20} - \sqrt{125}$ b. $2\sqrt{24} + 3\sqrt{20} - 7\sqrt{8}$
c. $3\sqrt{45} + 2\sqrt{12} + 5\sqrt{80} + 3\sqrt{108}$ d. $6\sqrt{44} + 4\sqrt{120} - \sqrt{99} - 3\sqrt{270}$
e. $2\sqrt{32} - 5\sqrt{45} - 4\sqrt{180} + 10\sqrt{8}$
29. **MC** Choose the correct answer from the given options.
- a. $\sqrt{2} + 6\sqrt{3} - 5\sqrt{2} - 4\sqrt{3}$ is equal to:
A. $-5\sqrt{2} + 2\sqrt{3}$ **B.** $-3\sqrt{2} + 23$ **C.** $6\sqrt{2} + 2\sqrt{3}$ **D.** $-4\sqrt{2} + 2\sqrt{3}$ **E.** $4\sqrt{2} + 2\sqrt{3}$
- b. $6 - 5\sqrt{6} + 4\sqrt{6} - 8$ is equal to:
A. $-2 - \sqrt{6}$ **B.** $14 - \sqrt{6}$ **C.** $-2 + \sqrt{6}$ **D.** $-2 - 9\sqrt{6}$ **E.** $2 + 9\sqrt{6}$
- c. $4\sqrt{8} - 6\sqrt{12} - 7\sqrt{18} + 2\sqrt{27}$ is equal to:
A. $-7\sqrt{5}$ **B.** $29\sqrt{2} - 18\sqrt{3}$ **C.** $-13\sqrt{2} - 6\sqrt{3}$ **D.** $-13\sqrt{2} + 6\sqrt{3}$ **E.** $7\sqrt{5}$
- d. $2\sqrt{20} + 5\sqrt{24} - \sqrt{54} + 5\sqrt{45}$ is equal to:
A. $19\sqrt{5} + 7\sqrt{6}$ **B.** $9\sqrt{5} - 7\sqrt{6}$ **C.** $-11\sqrt{5} + 7\sqrt{6}$ **D.** $-11\sqrt{5} - 7\sqrt{6}$ **E.** $-9\sqrt{5} - 7\sqrt{6}$

Understanding

30. **MC** Choose the correct answer from the given options.
- a. $\sqrt{1000}$ is equal to:
A. 31.6228 **B.** $50\sqrt{2}$ **C.** $50\sqrt{10}$ **D.** $10\sqrt{10}$ **E.** $100\sqrt{10}$
- b. $\sqrt{80}$ in simplest form is equal to:
A. $4\sqrt{5}$ **B.** $2\sqrt{20}$ **C.** $8\sqrt{10}$ **D.** $5\sqrt{16}$ **E.** $2 + 16\sqrt{5}$
- c. Choose which of the following surds is in simplest form.
A. $\sqrt{60}$ **B.** $\sqrt{147}$ **C.** $\sqrt{105}$ **D.** $\sqrt{117}$ **E.** $\sqrt{20}$
- d. Choose which of the following surds is **not** in simplest form.
A. $\sqrt{102}$ **B.** $\sqrt{110}$ **C.** $\sqrt{116}$ **D.** $\sqrt{118}$ **E.** $\sqrt{105}$

31. **MC** Choose the correct answer from the given options.

a. $6\sqrt{5}$ is equal to:

- A. $\sqrt{900}$ B. $\sqrt{30}$ C. $\sqrt{150}$ D. $\sqrt{180}$ E. $\sqrt{1800}$

b. Choose the expression out of the following that is **not** equal to the rest.

- A. $\sqrt{128}$ B. $2\sqrt{32}$ C. $8\sqrt{2}$ D. $64\sqrt{2}$ E. $6\sqrt{2}$

c. Choose the expression out of the following that is **not** equal to the rest.

- A. $4\sqrt{4}$ B. $2\sqrt{16}$ C. 8 D. 16 E. 32

d. $5\sqrt{48}$ is equal to:

- A. $80\sqrt{3}$ B. $20\sqrt{3}$ C. $9\sqrt{3}$ D. $21\sqrt{3}$ E. $10\sqrt{3}$

32. Reduce each of the following to simplest form.

- a. $\sqrt{675}$ b. $\sqrt{1805}$ c. $\sqrt{1792}$ d. $\sqrt{578}$

33. Reduce each of the following to its simplest form.

- a. $\sqrt{a^2c}$ b. $\sqrt{bd^4}$ c. $\sqrt{h^2jk^2}$ d. $\sqrt{f^3}$

34. Write the following sequences of numbers in order from smallest to largest.


- a. $6\sqrt{2}$, 8, $2\sqrt{7}$, $3\sqrt{6}$, $4\sqrt{2}$, $\sqrt{60}$
 b. $\sqrt{6}$, $2\sqrt{2}$, $\sqrt{2}$, 3, $\sqrt{3}$, 2, $2\sqrt{3}$

Reasoning

35. The formula for calculating the speed of a car before it brakes in an emergency is $v = \sqrt{20d}$, where v is the speed in m/s and d is the braking distance in metres.

Calculate the speed of a car before braking if the braking distance is 32.50 m.

Write your answer as a surd in its simplest form.



36. A gardener wants to divide their vegetable garden into 10 equal squares, each of area exactly 2 m^2 .

- Calculate the exact side length of each square. Explain your reasoning.
- Determine the side length of each square, correct to 2 decimal places.
- The gardener wishes to group the squares in their vegetable garden next to each other so as to have the smallest perimeter possible for the whole garden.
 - Determine how they should arrange the squares.
 - Explain why the exact perimeter of the vegetable patch is $14\sqrt{2}$ m.
 - Calculate the perimeter, correct to 2 decimal places.



37. Explain why $\sqrt{a^3b^2}$ can be simplified to $ab\sqrt{a}$.


38. Determine the smallest values that a and b can have, given that they are both natural numbers, that $1 < a < b$, and also that \sqrt{ab} is not a surd.

Problem solving

39. Kyle wanted a basketball court in his backyard, but he could not fit a full-sized court in his yard. He was able to get a rectangular court laid with a width of $6\sqrt{2}$ m and a length of $3\sqrt{10}$ m. Calculate the area of the basketball court and represent it in its simplest surd form.
40. A netball team went on a pre-season training run. They completed 10 laps of a triangular course that had side lengths of $(200\sqrt{3} + 50)$ m, $(50\sqrt{2} + 75\sqrt{3})$ m and $(125\sqrt{2} - 18)$ m. Determine the distance they ran, in its simplest surd form.



41. The area of a square is x cm². Explain whether the side length of the square would be a rational number.
42. To calculate the length of the hypotenuse of a right-angled triangle, use the formula $c^2 = a^2 + b^2$.



- a. Calculate the length of the hypotenuse for triangles with other side lengths as given in the following pairs of values.
- i. $\sqrt{5}, \sqrt{8}$ ii. $\sqrt{2}, \sqrt{7}$ iii. $\sqrt{15}, \sqrt{23}$
- b. Describe any pattern in your answers to part a.
- c. Calculate the length of the hypotenuse (without calculations) for triangles with the following side lengths.
- i. $\sqrt{1000}, \sqrt{500}$ ii. $\sqrt{423}, \sqrt{33}$ iii. $\sqrt{124}, \sqrt{63}$
- d. Your friend wrote down the following explanation.

$$\sqrt{b} + \sqrt{c} = \sqrt{a}$$


Is this answer correct?

If not, explain what the correct answer should be.

LESSON

1.10 Review




1.10.1 Topic summary



1.10.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
1.2	I can identify real numbers that are natural, integer, rational or irrational.			
	I can recall and use correct symbols for the sets of real, natural, integer, rational or irrational numbers.			
	I can write integers and decimals as rational number fractions.			
	I can write recurring decimals using dot notation.			
	I can explain the difference between recurring and terminating decimals.			
1.3	I can round decimals to a given number of decimal places.			
	I can identify how many significant figures a decimal number has.			
	I can round decimals to a given number of significant figures.			
1.4	I can identify the base and the index of a number written in index form.			
	I can recall the first three index laws.			
	I can use the first three index laws to simplify expressions involving indices.			
1.5	I can apply the Fourth, Fifth and Sixth index laws to raise a power to another power.			
	I can apply the first six index laws to simplify expressions involving indices.			
1.6	I can apply the Seventh Index Law to evaluate expressions using negative indices.			
	I can apply the first seven index laws to simplify expressions involving indices.			
1.7	I can identify numbers that are written in scientific notation.			
	I can write numbers in scientific notation as decimal numbers.			
	I can write decimal numbers using scientific notation.			
	I can represent decimals in exponential form.			

(continued)

(continued)


1.8	I can write square and cube roots using the radical symbol.			
	I can use fractional indices to represent square roots and cube roots.			
	I can simplify expressions that use square roots and cube roots.			
1.9	I can identify when a square root forms a surd.			
	I can place surds in their approximate position on a number line using geometric constructions.			
	I can simplify expressions containing surds.			

1.10.3 Project

Concentric squares

Consider a set of squares drawn around a central point on a grid, as shown.

These squares can be regarded as concentric squares, because the central point of each square is the point labelled **X**. The diagram shows four labelled squares drawn on one-centimetre grid paper.



- Use the diagram above to complete the following table, leaving your answers in simplest surd form (if necessary). The first square has been completed for you.


Square	1	2	3	4
Side length (cm)	2			
Diagonal length (cm)	$2\sqrt{2}$			
Perimeter (cm)	8			
Area (cm ²)	4			

Observe the patterns in the table and answer the following questions.

2. What would be the side length of the tenth square of this pattern?
3. What would be the length of the diagonal of this tenth square?

Consider a different arrangement of these squares on one-centimetre grid paper, as shown.

These squares all still have a central point, labelled Y.



4. Use the diagram to complete the following table for these squares, leaving your answers in simplest surd form if necessary. The first square has been done for you.

Square	1	2	3	4
Side length (cm)	$\sqrt{2}$			
Diagonal length (cm)	2			
Perimeter (cm)	$4\sqrt{2}$			
Area (cm ²)	2			


5. What would be the side length of the tenth square in this pattern of squares?
6. What would be the length of the diagonal of the tenth square?

Use the two diagrams above to answer the following questions.



7. In which of the two arrangements does the square have the greater side length?
8. Which of the two arrangements shows the squares with the greater diagonal length?
9. Compare the area of a square (your choice) in the first diagram with the area of the corresponding square in the second diagram.
10. Compare the perimeter of the same two corresponding squares in each of the two diagrams.
11. Examine the increase in area from one square to the next in these two diagrams. What increase in area would you expect from square 7 to square 8 in each of the two diagrams?

Consider a set of concentric circles around a centre labelled **Z**, drawn on one-centimetre grid paper, as shown.

12. Investigate the change in circumference of the circles, moving from one circle to the next (from smallest to largest). Write a general formula for this increase in circumference in terms of π .
13. Write a general formula in terms of π and r that can be used to calculate the increase in area from one circle to the next (from smallest to largest).



on Resources

-  **eWorkbook** Topic 1 Workbook (worksheets, code puzzle and project) (ewbk-2001)
-  **Interactivities** Crossword (int-0698)
Sudoku puzzle (int-3202)

Exercise 1.10 Review questions

learn **on**

Fluency

1. **MC** Choose which of the given numbers are rational: $\sqrt{\frac{6}{12}}$, $\sqrt{0.81}$, 5, -3.26 , $0.\dot{5}$, $\frac{\pi}{5}$, $\sqrt{\frac{3}{12}}$.

A. $\sqrt{0.81}$, 5, -3.26 , $0.\dot{5}$ and $\sqrt{\frac{3}{12}}$	B. $\sqrt{\frac{6}{12}}$ and $\frac{\pi}{5}$
C. $\sqrt{\frac{6}{12}}$, $\sqrt{0.81}$ and $\sqrt{\frac{3}{12}}$	D. 5, -3.26 and $\sqrt{\frac{6}{12}}$
E. $\sqrt{\frac{6}{12}}$, 5 and $\sqrt{0.81}$	
2. For each of the following, state whether the number is rational or irrational and give the reason for your answer.

a. $\sqrt{12}$	b. $\sqrt{121}$	c. $\frac{2}{9}$	d. $0.\dot{6}$	e. $\sqrt[3]{0.08}$
----------------	-----------------	------------------	----------------	---------------------
3. **MC** Choose which of the numbers in the set $\{3\sqrt{2}, 5\sqrt{7}, 9\sqrt{4}, 6\sqrt{10}, 7\sqrt{12}, 12\sqrt{64}\}$ are surds.

A. $9\sqrt{4}$, $12\sqrt{64}$	B. $3\sqrt{2}$ and $7\sqrt{12}$ only
C. $3\sqrt{2}$, $5\sqrt{7}$ and $6\sqrt{10}$ only	D. $3\sqrt{2}$, $5\sqrt{7}$, $6\sqrt{10}$ and $7\sqrt{12}$
E. $9\sqrt{4}$ and $7\sqrt{12}$	
4. State which of the following are not irrational numbers.

a. $\sqrt{7}$	b. $\sqrt{8}$	c. $\sqrt{81}$	d. $\sqrt{361}$
---------------	---------------	----------------	-----------------

5. **MC** Choose which of the following represents the number 0.000 67 written using scientific notation.
A. 67×10^{-5} **B.** 0.67×10^{-3} **C.** 6.7×10^{-4} **D.** 6.7×10^{-5} **E.** 6.7×10^{-3}
6. **MC** Choose which of the following is a simplification of the expression $\sqrt{250}$.
A. $25\sqrt{10}$ **B.** $5\sqrt{10}$ **C.** $10\sqrt{5}$ **D.** $5\sqrt{50}$ **E.** $10\sqrt{25}$
7. Simplify each of the following.
a. $b^7 \times b^3$ **b.** $m^9 \times m^2$ **c.** $k^3 \times k^5$
d. $f^2 \times f^8 \times f^4$ **e.** $h^4 \times h^5 \times h$
8. Simplify each of the following.
a. $2q^5 \times 3q^2 \times q^{10}$ **b.** $5w^3 \times 7w^{12} \times w^{14}$ **c.** $2e^2 p^3 \times 6e^3 p^5$
d. $5a^2 b^4 \times 3a^8 b^5 \times 7a^6 b^8$ **e.** $a^5 \div a^2$
9. Simplify each of the following.
a. $t^5 \div t$ **b.** $r^{19} \div r^{12}$ **c.** $p^8 \div p^5$ **d.** $\frac{f^{17}}{f^{12}}$ **e.** $\frac{y^{100}}{y^{10}}$
10. Simplify each of the following.
a. $\frac{m^{24}}{m^{14}}$ **b.** $\frac{g^4 \times g^5}{g^2}$ **c.** $\frac{x^6 \times x^2 \times x}{x^8}$ **d.** $\frac{d^6 \times d^7 \times d^2}{d^8}$ **e.** $\frac{t^7 \times t \times t^3}{t^2 \times t^4}$
11. Simplify each of the following.
a. $\frac{p^5 \times p^3 \times p \times p^4}{p^2 \times p^4 \times p^2}$ **b.** $\frac{16k^{13}}{21} \div \frac{8k^9}{42}$ **c.** $\frac{22b^{15}}{c} \div \frac{2b^8}{c^6}$
d. $\frac{9d^8}{16e^{10}} \div \frac{2d^{10}}{e^{16}}$ **e.** $\frac{2a^3 \times ab}{4 \times a \times b}$
12. Simplify each of the following.
a. 5^0 **b.** 12^0 **c.** 345^0 **d.** q^0 **e.** r^0
13. Simplify each of the following.
a. ab^0 **b.** $3w^0$ **c.** $5q^0 - 2q^0$ **d.** $100s^0 + 99t^0$ **e.** $a^7 b^0$
14. Simplify each of the following.
a. $v^{10} w^0$ **b.** prt^0 **c.** $a^9 b^4 c^0$
d. $j^8 k^0 m^3$ **e.** $4e^2 f^0 - 36(a^2 b^3)^0$
15. Simplify each of the following.
a. $2x^0 \times 3xy^2$ **b.** $-8(18x^2 y^4 z^6)^0$
c. $15 - 12x \left(\frac{3x}{8}\right)^0$ **d.** $-4p^0 \times 6(q^2 r^3)^0 \div 8(-12q^2)^0$
e. $3(6w^0)^2 \div 2(5w^5)^0$
16. Raise each of the following to the given power.
a. $(b^4)^2$ **b.** $(a^8)^3$ **c.** $(k^7)^{10}$ **d.** $(j^{100})^2$

17. **MC** $\left(\frac{4b^4}{d^2}\right)^3$ is equal to:

A. $\frac{4b^3}{d^3}$

B. $\frac{12b^{12}}{d^6}$

C. $\frac{64b^{12}}{d^6}$

D. $\frac{64b^7}{d^5}$

E. $\frac{64b^{16}}{d^5}$

18. Raise each of the following to the given power.

a. $(a^5b^2)^3$

b. $(m^7n^{12})^2$

c. $(st^6)^3$

d. $(qp^{30})^{10}$

19. Write each of the following with positive indices.

a. a^{-1}

b. k^{-4}

c. $4m^3 \div 2m^7$

d. $7x^3y^{-4} \times 6x^{-3}y^{-1}$

20. Write each of the following using a negative index.

a. $\frac{1}{x}$

b. $\frac{2}{y^4}$

c. $z \div z^4$

d. $45p^2q^{-4} \times 3p^{-5}q$

21. Simplify each of the following.

a. $\sqrt{100}$

b. $\sqrt{36}$

c. $\sqrt{a^2}$

d. $\sqrt{b^2}$

e. $\sqrt{49f^4}$

22. Simplify each of the following.

a. $\sqrt[3]{27}$

b. $\sqrt[3]{1000}$

c. $\sqrt[3]{x^3}$

d. $\sqrt[3]{8d^3}$

e. $\sqrt[3]{64f^6g^3}$

Problem solving

23. If your body produces about 2.0×10^{11} red blood cells each day, determine how many red blood cells your body has produced after one week.

24. If your body produces about 1.0×10^{10} white blood cells each day, determine how many white blood cells your body produces each hour.

25. If Earth is approximately 1.496×10^8 km from the Sun, and Mercury is about 5.8×10^7 km from the Sun, evaluate the distance from Mercury to Earth.

26. Calculate the closest distance between Mars and Saturn if Mars is 2.279×10^8 km from the Sun and Saturn is 1.472×10^9 km from the Sun.



27. Consider the numbers 6×10^9 , 3.4×10^7 and 1.5×10^9 .

a. Place these numbers in order from smallest to largest.

b. If the population of Oceania is 3.4×10^7 and the world population is 6×10^9 , determine the percentage of the world population that is represented by the population of Oceania. Give your answer to 2 decimal places.

c. Explain what you need to multiply 1.5×10^9 by to get 6×10^9 .

d. Assume the surface area of Earth is approximately $1.5 \times 10^9 \text{ km}^2$, and that the world's population is about 6×10^9 . If we divided Earth up into equal portions, determine how much surface area each person in the world would get.

28. The total area of the state of Victoria is $2.27 \times 10^5 \text{ km}^2$. If the total area of Australia is $7.69 \times 10^6 \text{ km}^2$, calculate the percentage of the total area of Australia that is occupied by the state of Victoria. Write your answer correct to two decimal places.

29. A newly discovered colony of bees contains 2.05×10^8 bees. If 0.4% of these bees are estimated to be queen bees, calculate how many queen bees live in the colony.



30. Third-generation fibre optics that can carry up to 10 trillion (1×10^{12}) bits of data per second along a single strand of fibre have recently been tested. Based on the current rate of improvement in this technology, the amount of data that one strand of fibre can carry is tripling every 6 months. This trend is predicted to continue for the next 20 years. Calculate how many years it will take for the transmission speed along a single strand of fibre to exceed 6×10^{16} bits per second.

31. Mach speed refers to the speed of sound, where Mach 1 is the speed of sound (about 343 metres per second (m/s)). Mach 2 is double the speed of sound, Mach 3 is triple the speed of sound, and so on. Earth has a circumference of about $4.0 \times 10^4 \text{ km}$.

a. The fastest recorded aircraft speed is 11 270 kilometres per hour (km/h).

Express this as a Mach speed, correct to 1 decimal place.

b. Create and complete a table for three different speeds (Mach 1, 2 and 3) with columns labeled 'Mach number', 'Speed in m/s' and 'Speed in km/h'.

c. Calculate how long it would take something travelling at Mach 1, 2 and 3 to circle Earth. Give your answer in hours, correct to 2 decimal places.

d. Evaluate how long it would take for the world's fastest aircraft to circle Earth. Give your answer in hours, correct to 2 decimal places.

e. Evaluate how many times the world's fastest aircraft could circle Earth in the time it takes an aircraft travelling at Mach 1, 2 and 3 to circle Earth once. Give your answers correct to 2 decimal places.

f. Explain the relationship between your answers in part e and their corresponding Mach value.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 1 Number skills and index laws

1.1 Pre-test

1. True
2. Recurring decimal
3. a. True b. True
4. 3
5. C
6. B
7. $\sqrt{5}$
8. C
9. 7.439
10. 7.44
11. C
12. B
13. 1.672×10^{-27}

14. $\frac{4\sqrt{3}}{3}$

15. $\frac{5\sqrt{2}}{2}$

1.2 Real numbers

1. a. $\frac{15}{1}$ b. $\frac{-8}{1}$ c. $\frac{8}{3}$ d. $\frac{-41}{8}$
2. a. $\frac{4}{1}$ b. $\frac{73}{10}$ c. $\frac{2}{1000}$ d. $\frac{872}{10}$
3. a. $\frac{0}{1}$ b. $\frac{156}{100}$ c. $\frac{3612}{1000}$ d. $\frac{-8}{100}$
4. a. 0.555 555 55 b. 0.515 151 51
c. 0.511 111 11 d. 6.031 313 1
e. 5.183 183 1
5. a. -7.024 444 4 b. 8.912 491 2
c. 5.123 434 3 d. 5.123 412 3
e. 3.002 020 2
6. a. 0.5̇ c. 0.27̇ e. 2.18̇
b. 0.428571̇ d. -1.74̇
7. a. 0.073̇ b. -3.83̇ c. 0.46̇
d. 0.46̇ e. 0.046̇
8. a. 15
b. -3, 0, 15
c. -3, $\frac{-3}{7}$, 0, 2.3̇, 2 $\frac{3}{5}$, 15
d. 2.3̇ and $-\frac{3}{7}$
9. 2.1̇, 2.12̇1̇, 2.12̇1̇, 2.12̇, 2.12̇
10. Statements a and b are true.
11. Every integer a can be written as $\frac{a}{1}$ and therefore every integer is rational.

12. a. \mathbb{Q}^-
b. \mathbb{N}^- implies a set of negative natural numbers. As the natural numbers include only positive numbers, this set would not exist.
13. Yes, because the division of an integer by another integer will always result in a rational number.
14. a. $\frac{1}{3}$, $\frac{2}{3}$, $\frac{1}{6}$ b. 6
15. Answers will vary. Sample responses are shown.
a. $\frac{1}{9}$ b. $\frac{10}{99}$ c. $\frac{100}{999}$ d. $\frac{1000}{9999}$
16. \$3330.20
17. 657 bottles of water per year

1.3 Rounding numbers and significant figures

1. a. 1.571 b. 2.236 c. 3.873
d. 16.435 e. 5.111 f. 5.156
2. a. 5.152 b. 137.358 c. 0.429
d. 0.077 e. 2.429 f. 1.000
3. a. 4.000 b. 0.254 c. 0.025
d. 0.001 e. 5.000 f. 2342.156
4. a. 2 b. 3 c. 4 d. 3
5. a. 4 b. 4 c. 2 d. 1
6. a. 3 b. 4 c. 2 d. 3
7. a. 1.5708 b. 2.2361 c. 3.8730
d. 16.435 e. 5.1111 f. 5.1556
8. a. 5.1515 b. 137.36 c. 0.428 57
d. 0.076 923 e. 2.4286 f. 1.0000
9. a. 6.5813 b. 4.0000 c. 0.253 64
d. 0.025 400 e. 0.000 913 60 f. 5.0000
10. 3.1416, 3.14159, 3.141593, 3.1415927
11. a. Yes b. No c. Yes d. No
12. a. False b. True c. False d. True
13. a. False b. True c. False d. True
14. a. True b. False c. False d. True
15. A number of answers are possible. Example answer: 2.004.
16. The 5th digit (5) causes the 4th digit to round up from 9 to 10, which causes the 3rd digit to round up from 9 to 10, which causes the 2nd digit to round up from 2 to 3.
17. a. 2 decimal places b. 3.1416
18. a. 153.86 cm²
b. 153.9380 cm²
c. 153.94 cm²
d. Yes, because 3.14 is used as an estimate and is not the value of π as used on the calculator.
19. a. 65 449.8470 cm³
b. 9
c. Yes, any number with a finite number of decimal places is a rational number.
20. 13 000

1.4 Review of index laws

1. a. $2^2 \times 3$ b. $2^3 \times 3^2$ c. 3×5^2
 d. $2^4 \times 3 \times 5$ e. $2^7 \times 5$ f. $2^3 \times 5^2 \times 7^2$
2. a. $20p^{11}$ b. $6x^8$ c. $56y^{10}$
 d. $21p^8$ e. $84t^6$ f. $30q^{15}$
3. a. $6a^6e^7$ b. $8p^6h^{12}$ c. $6k^{14}$ d. 3^{12}
4. a. $80m^9$ b. $6g^3h^6$ c. $30p^6q^9$ d. $48u^9w^7$
5. a. $27d^{11}y^{17}$ b. $42b^{14}c^9$ c. $24r^{16}s^{18}$ d. $60h^{38}v^{20}$
6. a. $3p^4$ b. $6r^4$ c. $9a^3$ d. $3b^6$
7. a. $20r^4$ b. $9q$ c. $\frac{130}{d^2}$ d. $7t^4$
8. a. $\frac{3p^5}{2}$ b. $\frac{8b^5}{3}$ c. $\frac{5m^{10}n^6}{6}$ d. $\frac{9x^8y}{4}$
9. a. $3j^5f^3$ b. $\frac{4p^2rs}{3}$ c. $\frac{9a^5b^3c}{2}$ d. $\frac{20f^6g^2h^4}{3}$
10. a. 1 b. 6 c. 1 d. 1 e. 5
11. a. 1 b. 1 c. 2 d. 2 e. 3
12. a. 1 b. 0 c. 14 d. 6 e. 6
13. a. 1 b. 2 c. 2 d. 2 e. 2
14. a. $\frac{h^2}{2}$ b. $\frac{q^4}{5}$ c. $\frac{n^3}{5}$ d. v^2 e. $2x^6$
15. a. $\frac{1}{x^3}$ b. $\frac{1}{m^8}$ c. $\frac{1}{4m^6}$ d. $\frac{2}{x^2}$
16. a. $2x^2$ b. $24t^6$ c. $\frac{1}{2y^5}$ d. $\frac{7y^3}{4x^5}$
17. a. $\frac{2}{5m^3n^4}$ b. $\frac{2}{n^2}$ c. $\frac{2y}{x}$ d. $\frac{c^4}{a^4}$
18. a. 6 b. 9 c. 18 d. 11 e. 15 f. 6
19. Sample responses can be found in the worked solutions in the online resources.
20. a. B b. D c. D
21. a. A b. D
22. Sample responses can be found in the worked solutions in the online resources.
23. a. 5^9
 b. Sample responses can be found in the worked solutions in the online resources.
24. a. 5^{16}
 b. Sample responses can be found in the worked solutions in the online resources.
25. a. 1
 b. 1
 c. 0
 d. Sample responses can be found in the worked solutions in the online resources.
26. a. $\Delta = 14$
 b. Answers will vary, but $\Delta + O + \diamond$ must sum to 12.
 Possible answers include: $\Delta = 3, O = 2, \diamond = 7; \Delta = 1, O = 3, \diamond = 8; \Delta = 4, O = 4, \diamond = 4; \Delta = 5, O = 1, \diamond = 6.$

27. a. The repeating pattern is 1, 3, 9, 7.
 b. 3
28. a. The repeating pattern is 4, 6.
 b. 4
29. a. Sample responses can be found in the worked solutions in the online resources.
 b. i. 8 ii. 1 iii. 2
30. 2^{2n+3}

1.5 Raising powers

1. a. e^6 b. f^{80} c. p^{100} d. r^{144} e. $8a^6$
2. a. a^8b^{12} b. p^5q^{15} c. $g^{30}h^{20}$
 d. $81w^{36}q^8$ e. $49e^{10}r^4q^8$
3. a. p^8q^6 b. $r^{15}w^9$ c. $b^{10}n^{18}$
 d. $j^{18}g^{12}$ e. $8a^3b^4$
4. a. q^4r^{20} b. $h^{24}j^{16}$ c. $a^{21}f^{16}$
 d. $t^{10}u^8$ e. $i^{15}j^{12}$
5. a. $\frac{9b^8}{d^6}$ b. $\frac{25h^{20}}{4j^4}$ c. $\frac{8k^{15}}{27t^{24}}$ d. $\frac{49p^{18}}{64q^{44}}$
6. a. $\frac{125y^{21}}{27z^{39}}$ b. $\frac{256a^{12}}{2401c^{20}}$ c. $\frac{-64k^6}{343m^{18}}$ d. $\frac{16g^{28}}{81h^{44}}$
7. a. 2^{20} b. t^{33} c. a^{21} d. e^{66}
8. a. g^{39} b. $12a^{14}$ c. $216d^{27}$ d. $40\,000r^{14}$
9. B
10. B
11. D
12. a. a^6 b. m^4 c. n^3
 d. b^8 e. f^{17} f. g^6
13. a. p^9 b. y^2 c. c^{20}
 d. f^7 e. k^{14} f. p^{16}
14. a. 5
 b. 6
 c. 30
 d. Sample responses can be found in the worked solutions in the online resources.
15. a. i. 1 ii. -1 iii. -1 iv. 1
 b. $(-1)^{\text{even}} = 1(-1)^{\text{odd}} = -1$. Sample responses can be found in the worked solutions in the online resources.
16. Danni is correct. Explanations will vary but should involve $(-x)(-x) = (-x)^2 = x^2$ and $-x^2 = -1 \times x^2 = -x^2$.
17. a. 5^{12}
 b. Sample responses can be found in the worked solutions in the online resources.
18. a. i. $x = \frac{5}{3}$ ii. $x = \frac{5}{3}$ iii. $x = \frac{5}{3}$
 b. When equating the powers, $3x = 5$.
19. Answers will vary. Possible answers are 4096 and 262144.
20. $10^8 \times 10^8 \times 10^8 = (10^8)^3$ atoms
21. a. 4^1 b. 7^2

22. a. 2^1 b. 3^4 c. $\frac{1}{5^2}$ d. 2^2
 e. $\frac{1}{2^2}$ f. $\frac{1}{2}$ g. $\frac{1}{2}$ h. $\frac{1}{7}$

23. a. i. 3 ii. 4 iii. 8 iv. 11
 b. i. 3 ii. 4 iii. 8 iv. 11

c. Raising a number to a power of one-half is the same as calculating the square root of that number.

1.6 Negative indices

1. a. $3^5 = 243$, $3^4 = 81$, $3^3 = 27$, $3^2 = 9$, $3^1 = 3$, $3^0 = 1$,
 $3^{-1} = \frac{1}{3}$, $3^{-2} = \frac{1}{9}$, $3^{-3} = \frac{1}{27}$, $3^{-4} = \frac{1}{81}$, $3^{-5} = \frac{1}{243}$
 b. $5^4 = 625$, $5^3 = 125$, $5^2 = 25$, $5^1 = 5$, $5^0 = 1$, $5^{-1} = \frac{1}{5}$,
 $5^{-2} = \frac{1}{25}$, $5^{-3} = \frac{1}{125}$, $5^{-4} = \frac{1}{625}$
 c. $10^4 = 10000$, $10^3 = 1000$, $10^2 = 100$, $10^1 = 10$, $10^0 = 1$,
 $10^{-1} = \frac{1}{10}$, $10^{-2} = \frac{1}{100}$, $10^{-3} = \frac{1}{1000}$, $10^{-4} = \frac{1}{10000}$

2. a. $\frac{1}{32}$ b. $\frac{1}{27}$ c. $\frac{1}{4}$ d. $\frac{1}{100}$

3. a. $\frac{1}{125}$ b. 7 c. $\frac{4}{3}$ d. $\frac{16}{9}$

4. a. 27 b. $\frac{2}{3}$ c. $\frac{16}{81}$ d. $\frac{49}{4}$

5. a. $\frac{1}{x^4}$ b. $\frac{1}{y^5}$ c. $\frac{1}{z}$

d. $\frac{a^2}{b^3}$ e. $\frac{1}{m^2n^3}$

6. a. $\frac{1}{m^2n^3}$ b. x^2y^2 c. $5x^3$

d. $\frac{w^5}{x^2}$ e. x^2y^2

7. a. $\frac{a^2c}{b^3d^4}$ b. $\frac{a^2d^3}{b^2c^2}$ c. $\frac{10y}{x^2}$

d. $\frac{x}{3}$ e. $\frac{1}{m^3x^2}$

8. a. $\frac{1}{a^5}$ b. m^5 c. $\frac{1}{m^7}$

d. $\frac{14}{x}$ e. 6 f. $\frac{5a}{b^3}$

9. a. $\frac{1}{x^3}$ b. $\frac{6}{x^5y^3}$ c. $50x^3$

d. 1 e. $\frac{50}{a^5}$ f. $10a^4$

10. a. $\frac{-32}{w^3}$ b. $\frac{16}{m^4}$ c. $\frac{27m^6}{n^{12}}$

d. $\frac{1}{a^6b^{15}}$ e. a^2b^6 f. $\frac{25}{a^2}$

11. a. $\frac{1}{x^5}$ b. $\frac{1}{x^{11}}$ c. x^{11}

d. x^5 e. $\frac{6c^4}{d^2}$

12. a. $\frac{2}{a^6}$ b. $\frac{5}{m}$ c. $\frac{1}{b}$

d. $\frac{1}{a^3b^2}$ e. $\frac{c^2}{a^4}$

13. a. $\frac{1}{16a}$ b. $\frac{1}{m^3}$ c. $\frac{3}{2t^9}$

d. t^3 e. $\frac{m^2}{n^3}$

14. a. 2^0 b. 2^3 c. 2^5 d. 2^6 e. 2^{-3} f. 2^{-5}

15. a. 4^0 b. 4^1 c. 4^3 d. 4^{-1} e. 4^{-2} f. 4^{-3}

16. a. 10^0 b. 10^1 c. 10^4 d. 10^{-1} e. 10^{-2} f. 10^{-5}

17. a. $3^{-4} = \frac{1}{3^4}$

b. Answers will vary, but should mention that if you are dividing, the power in the numerator is lower than that in the denominator. Sample responses can be found in the worked solutions in the online resources.

c. Sample responses can be found in the worked solutions in the online resources.

18. Answers will vary, but should mention that the negative 13 means the decimal point is moved 13 places to the left of 3. Using scientific notation allows the number to be expressed more concisely. Sample responses can be found in the worked solutions in the online resources.

19. a. No. The equivalent expression with positive indices is $\frac{x^6 + 1}{x^3}$.

b. No. The equivalent expression with positive indices is $\frac{(xy)^2}{(x+y)^2}$.

20. a. No. The equivalent expression with positive indices is $\frac{x^{13} - 1}{x^5}$.

b. No. The correct equivalent expression is $\frac{1}{x^6 - x^3}$.

21. a. 0.048 b. 7600 c. 0.000 29 d. 8.1

22. a. 3 b. -1 c. 2 d. -4

23. a. 2^{19} b. 3^{20}

24. a. $\frac{6}{5}$ b. $\left(\frac{3}{2}\right)^{200}$

1.7 Scientific notation

1. a. 5.00×10^3 b. 4.31×10^2
 c. 3.8×10^1 d. 3.5×10^5

2. a. 1 000 000 000 b. 392.673
 c. 591.1 d. 5100

3. a. 734 000 b. 7 141 400
 c. 35.1 d. 80 500

4. a. 614 b. 6140
c. 61 400 d. 351.8
5. a. 7.25×10^1 b. 7.25×10^2
c. 7.25×10^3 d. 7.25×10^8
6. a. 4.379×10^1 b. 5.317×10^3 c. 2.590×10^2
d. 1.101×10^2 e. 1.632×10^6 f. 1.000×10^6
7. a. 0.2 b. 0.004 c. 0.0007 d. 0.03
8. a. 0.082 73 b. 0.072 95
c. 0.002 914 2 d. 0.000 037 53
9. a. 0.000 529 b. 0.000 033 333
c. 0.000 000 002 625 d. 0.000 000 000 000 001 273
10. a. 7×10^{-1} b. 5×10^{-3}
c. 3×10^{-7} d. 1×10^{-11}
11. a. $2 \times 10^{-1} + 3 \times 10^{-2} + 1 \times 10^{-3}$
b. $3 \times 10^{-3} + 6 \times 10^{-4} + 2 \times 10^{-5}$
c. 7×10^{-4}
d. $6 \times 10^{-2} + 3 \times 10^{-3}$
12. a. 6.73×10^{-3} b. 1.43×10^{-1} c. 6.83×10^{-5}
d. 5.12×10^{-9} e. 5.09×10^{-2} f. 1.25×10^{-2}
13. a. 5.27×10^2 , 8.31×10^2 , 9.718×10^2 , 3.27×10^3
b. 4.09×10^2 , 4.37×10^2 , 7.943×10^2 , 7.95×10^2
c. 2.7×10^{-3} , 9.29×10^{-3} , 5.251×10^{-2} , 5.31×10^{-2}
d. 2.7×10^{-3} , 7.13×10^{-2} , 8.31×10^2 , 3.27×10^3
14. a. 0.000 000 000 000 000 000 000 019 94 g
b. 1.994×10^{-17} g
c. 5.015×10^{23} atoms
15. 3844 hours \approx 160.17 days
16. 333 333.33
17. The neutron is heavier by 2.3×10^{-30} kg.
18. a. 2.1×10^7 b. 1.29×10^8
19. a. 8 389 000 bytes b. 8.389×10^6 bytes
20. 0.000 000 2 N/m
21. a. i. Sample responses can be found in the worked solutions in the online resources.
ii. Sample responses can be found in the worked solutions in the online resources.
b. i. 5.36×10^7 ii. 5.36×10^7
c. The answers are the same because 2.95×10^3 is very small compared with 5.36×10^7 .
22. 8.75×10^{-7} is such a small amount (0.000 000 875) that, when it is added to 2.39×10^{-3} , it doesn't affect the value when given to 3 significant figures.
23. a. i. 385 440 km
ii. 78 489 600 km
iii. 1 278 960 000 km
iv. 5 772 840 000 km
b. i. 385 000 km
ii. 78 500 000 km
iii. 1 280 000 000 km
iv. 5 770 000 000 km

- c. i. 3.85×10^5 km
ii. 7.85×10^7 km
iii. 1.28×10^9 km
iv. 5.77×10^9 km


24. a. i. 31 536 000 s ii. 3.1536×10^7 s
b. i. 9 460 800 000 000 km ii. 9.4608×10^{12} km
c. i. 4.068×10^{13} km ii. 46 440 000 years
25. a. 5.972×10^{21} b. 4
26. a. 9.1094×10^{-31}
b. 1.6725×10^{-27}
c. 1.6726×10^{-27}
d. It is important to work with the original amounts and leave rounding until the end of a calculation so that the answer is accurate.

1.8 Square roots and cube roots

1. a. $15^{\frac{1}{2}}$ b. $m^{\frac{1}{2}}$ c. $t^{\frac{1}{3}}$ d. $(w^2)^{\frac{1}{3}}$
2. a. 7 b. 2 c. 3 d. 5
3. a. 10 b. 8 c. 4
d. 1000 e. 100 f. 9
4. a. m b. b c. $6t^2$ d. mn^2
5. a. $5t^2$ b. xy^2 c. a^2m^{10} d. $6y^2$
6. a. $4x^2y^2$ b. $5ab^2c^3$ c. b^7 d. b^3
7. B
8. D
9. A
10. a. $3^{\frac{1}{2} + \frac{1}{2}} = 3^1$ b. $\sqrt{3}$ c. $a^{\frac{1}{n}}$ d. 2
11. a. $z^{2.5} = z^{\frac{5}{2}} = (z^5)^{\frac{1}{2}} = \sqrt{z^5}$
b. No, it is the tenth root: $z^{0.3} = z^{\frac{3}{10}} = (z^3)^{\frac{1}{10}} = \sqrt[10]{z^3}$.
12. Prithvi is correct: $\sqrt{x^2} = x^{\frac{1}{2} \times 2} = x^1 = x$; x can be a positive or negative number.
13. $(-2^3)^{\frac{1}{3}} = -2$; answers will vary, but should include that we cannot take the fourth root of a negative number. Sample responses can be found in the worked solutions in the online resources.
14. $\sqrt{1871} \approx 43.25$
 $42^2 = 1764$
 $43^2 = 1849$
He was 43 years old in 1849. Therefore, he was born in $1849 - 43 = 1806$.
15. a. $d = t^{\frac{2}{3}}$ b. $t = d^{\frac{3}{2}}$
16. $\frac{16}{81}$
17. a. There are 2 possible solutions, because the square root of a number always has a positive and a negative answer.
b. $\frac{5}{3}, 1$


1.9 Surds (extending)

1. a. 1 b. 2 c. 0
 d. $\frac{1}{3}$ e. $\frac{5}{4}$
2. a. 0.4 b. 20 c. 100
 d. $\frac{2}{5}$ e. 1.2
3. a. 4.5 b. 1000 c. 0.03
 d. 16 e. $\frac{11}{7}$
4. a. 9 b. -9 c. 11 d. -21
 e. 2 b. 4 c. 7 d. 3
 e. 4 b. 5 c. -7 d. -3



8. c and d
9. c
10. a. $\sqrt{21}$ b. $-\sqrt{21}$ c. $2\sqrt{6}$
 d. $6\sqrt{7}$ e. $10\sqrt{14}$
11. a. $12\sqrt{7}$ b. $9\sqrt{7}$ c. $22\sqrt{10}$
 d. $22\sqrt{3}$ e. 3
12. a. $15\sqrt{10}$ b. 24 c. 6
 d. 11 e. 30
13. a. $\sqrt{3}$ b. $-\sqrt{2}$ c. $\sqrt{6}$ d. $\sqrt{5}$
14. a. $3\sqrt{3}$ b. $\frac{3\sqrt{6}}{2}$ c. $15\sqrt{2}$ d. $\frac{5\sqrt{5}}{3}$
15. a. $-2\sqrt{5}$ b. $\sqrt{3}$ c. $\sqrt{3}$ d. $\sqrt{7}$
16. a. $2\sqrt{5}$ b. $2\sqrt{2}$ c. $3\sqrt{2}$ d. 7
17. a. $\sqrt{30}$ b. $5\sqrt{2}$ c. $2\sqrt{7}$ d. $6\sqrt{3}$
18. a. $12\sqrt{2}$ b. $4\sqrt{3}$ c. $10\sqrt{5}$ d. $9\sqrt{2}$
19. a. $4\sqrt{2}$ b. $15\sqrt{3}$ c. 48 d. $35\sqrt{2}$ e. $20\sqrt{6}$
20. a. $10\sqrt{3}$ b. $4\sqrt{42}$ c. $72\sqrt{2}$ d. $27\sqrt{5}$ e. $132\sqrt{2}$
21. a. $\sqrt{12}$ b. $\sqrt{175}$ c. $\sqrt{108}$ d. $\sqrt{80}$ e. $\sqrt{384}$
22. a. $\sqrt{90}$ b. $\sqrt{32}$ c. $\sqrt{720}$ d. $\sqrt{600}$ e. $\sqrt{338}$

23. a. $2\sqrt{3}$ b. $5\sqrt{2}$ c. $10\sqrt{2}$ d. $2\sqrt{10}$
24. a. $2\sqrt{2}$ b. $-4\sqrt{5}$ c. $-6\sqrt{3}$ d. 0
25. a. $5\sqrt{11}$ b. $2\sqrt{7}$
 c. $10\sqrt{2} + 7\sqrt{3}$ d. $8\sqrt{5} + \sqrt{6}$
26. a. $8\sqrt{10} + 7\sqrt{3}$ b. $16\sqrt{2} - 11\sqrt{5}$
 c. $2\sqrt{6}$ d. $5\sqrt{5} + 15$
27. a. $\sqrt{2}$ b. 0 c. $-5\sqrt{3}$
 d. $-4\sqrt{7}$ e. $5\sqrt{6} + 6\sqrt{5}$
28. a. $2\sqrt{3} - 3\sqrt{5}$ b. $4\sqrt{6} + 6\sqrt{5} - 14\sqrt{2}$
 c. $29\sqrt{5} + 22\sqrt{3}$ d. $9\sqrt{11} - \sqrt{30}$
 e. $28\sqrt{2} - 39\sqrt{5}$
29. a. D b. A c. D d. A
30. a. D b. A c. C d. C
31. a. D b. D c. D d. B
32. a. $15\sqrt{3}$ b. $19\sqrt{5}$ c. $16\sqrt{7}$ d. $17\sqrt{2}$
33. a. $a\sqrt{c}$ b. $d^2\sqrt{b}$ c. $hk\sqrt{j}$ d. $f\sqrt{f}$
34. a. $2\sqrt{7}$, $4\sqrt{2}$, $3\sqrt{6}$, $\sqrt{60}$, 8, $6\sqrt{2}$
 b. $\sqrt{2}$, $\sqrt{3}$, 2, $\sqrt{6}$, $2\sqrt{2}$, 3, $2\sqrt{3}$
35. $5\sqrt{26}$
36. a. $\sqrt{2}$ m
 b. 1.41 m
 c. i. Any of the arrangements shown would work.



- ii. The length of each square is $\sqrt{2}$ m and there are 14 lengths around each shape.
- iii. 19.80 m
37. $\sqrt{a^3b^2} = \sqrt{a^3} \times \sqrt{b^2}$
 $= \sqrt{a^2} \times \sqrt{a} \times \sqrt{b^2}$
 $= a \times \sqrt{a} \times b$
 $= ab\sqrt{a}$
38. $a = 2$ and $b = 8$.
39. $36\sqrt{5} \text{ m}^2$
40. $(1750\sqrt{2} + 2750\sqrt{3} + 320) \text{ m}$
41. The side length will be rational if x is a perfect square. If x is not a perfect square, the side length will be a surd.
42. a. i. $\sqrt{13}$ ii. 3 iii. $\sqrt{38}$
 b. For side lengths \sqrt{a} and \sqrt{b} , the length of the hypotenuse is $\sqrt{a+b}$.
 c. i. $\sqrt{1500} = 10\sqrt{15}$
 ii. $2\sqrt{114}$
 iii. $\sqrt{187}$
 d. No. The correct answer is $\sqrt{b+a} = \sqrt{c}$.

Project

1. Square	1	2	3	4
Side length (cm)	2	4	6	8
Diagonal length (cm)	$2\sqrt{2}$	$4\sqrt{2}$	$6\sqrt{2}$	$8\sqrt{2}$
Perimeter (cm)	8	16	24	32
Area (cm²)	4	16	36	64

2. 20 cm
3. $20\sqrt{2}$ cm

4. Square	1	2	3	4
Side length (cm)	$\sqrt{2}$	$2\sqrt{2}$	$3\sqrt{2}$	$4\sqrt{2}$
Diagonal length (cm)	2	4	6	8
Perimeter (cm)	$4\sqrt{2}$	$8\sqrt{2}$	$12\sqrt{2}$	$16\sqrt{2}$
Area (cm²)	2	8	18	32

5. $10\sqrt{2}$ cm
6. 20 cm
7. First arrangement
8. First arrangement
9. The squares in the first diagram are twice the area of each corresponding square in the second diagram.
10. The squares in the first diagram have a perimeter that is $\sqrt{2}$ times the perimeter of each corresponding square in the second diagram.
11. An increase of 60 cm^2 in the first diagram and an increase of 30 cm^2 for the second diagram.
12. 2π
13. $\pi(2r + 1)$

1.10 Review questions

1. A
2. a. Irrational, because it is equal to a non-recurring and non-terminating decimal.
b. Rational, because it can be expressed as a whole number.
c. Rational, because it is given in a rational form.
d. Rational, because it is a recurring decimal.
e. Irrational, because it is equal to a non-recurring and non-terminating decimal.
3. D
4. c and d
5. C
6. B
7. a. b^{10} b. m^{11} c. k^8
d. f^{14} e. h^{10}
8. a. $6q^{17}$ b. $35w^{29}$ c. $12e^5p^8$
d. $105a^{16}b^{17}$ e. a^3
9. a. t^4 b. r^7 c. p^3
d. f^5 e. y^{90}

10. a. m^{10} b. g^7 c. x d. d^7 e. t^5
11. a. p^5 b. $4k^4$ c. $11b^7c^5$ d. $\frac{9e^6}{32d^2}$ e. $\frac{a^3}{2}$
12. a. 1 b. 1 c. 1 d. 1 e. 1
13. a. a b. 3 c. 3 d. 199 e. a^7
14. a. v^{10} b. pr c. a^9b^4 d. j^8m^3 e. $4e^2 - 36$
15. a. $6xy^2$ b. -8 c. $15 - 12x$
d. -3 e. 54
16. a. b^8 b. a^{24} c. k^{70} d. j^{200}
17. C
18. a. $a^{15}b^6$ b. $m^{14}n^{24}$
c. s^3t^{18} d. $q^{10}p^{300}$
19. a. $\frac{1}{a}$ b. $\frac{1}{k^4}$ c. $\frac{2}{m^4}$ d. $\frac{42}{y^5}$
20. a. x^{-1} b. $2y^{-4}$
c. z^{-3} d. $135p^{-3}q^{-3}$
21. a. 10 b. 6 c. a d. b e. $7f^2$
22. a. 3 b. 10 c. x d. $2d$ e. $4f^2g$
23. 1.4×10^{12}
24. 4.167×10^8
25. 9.16×10^7 km
26. 1.2441×10^9 km
27. a. $3.4 \times 10^7, 1.5 \times 10^9, 6 \times 10^9$
b. 0.57%
c. 4
d. 0.25 km^2
28. 2.95%
29. 820 000 queen bees
30. At the beginning of the 5th year, or in 5.5 years.
31. a. Mach 9.1

Mach number	Speed in m/s	Speed in km/h
1	343	1234.8
2	686	2469.6
3	1029	3704.4

- c. At Mach 1 it would take 32.39 hours.
At Mach 2 it would take 16.20 hours.
At Mach 3 it would take 10.79 hours.
d. 3.55 hours
e. The fastest aircraft could circle Earth 9.12 times in the time an aircraft travelling at Mach 1 could circle Earth once.
The fastest aircraft could circle Earth 4.56 times in the time an aircraft travelling at Mach 2 could circle Earth once.
The fastest aircraft could circle Earth 3.04 times in the time an aircraft travelling at Mach 3 could circle Earth once.
f. (Number of times to circle Earth) \times (Mach value) ≈ 9.1

2 Algebra

LESSON SEQUENCE

2.1 Overview	72
2.2 Using pronumerals	74
2.3 Algebra in worded problems	83
2.4 Simplification of algebraic expressions	88
2.5 Expanding brackets	95
2.6 Difference of two squares and perfect squares	104
2.7 Further expansions	109
2.8 The highest common factor	112
2.9 The highest common binomial factor	118
2.10 Solving worded problems	122
2.11 Review	131



LESSON

2.1 Overview

2.1.1 Why learn this?

Most people will tell you that if you want a career in fields like engineering, science, finance and software development, you need to study and do well in algebra at high school and university. Without algebra we wouldn't have landed on the Moon or be able to enjoy the technological marvels we do today, like our smartphones.

Learning algebra in school helps you to develop critical thinking skills. These skills can help with things like problem solving, logic, pattern recognition and reasoning.

Consider these situations:

- You want to work out the cost of a holiday to Japan at the current exchange rate.
- You're driving to a petrol station to refuel your car, but you only have \$30 in your pocket. You want to know how much petrol you can afford.
- You're trying to work out whether a new phone plan that charges \$5 per day over 24 months for the latest smartphone is worth it.



Working out the answers to these kinds of real-life questions is a direct application of algebraic thinking.

Understanding algebra can help you to make reasoned and well-considered financial and life decisions. It can help you make independent choices and assist you with many everyday tasks. Algebra is an important tool that helps us connect with — and make sense of — the world around us.

Hey students! Bring these pages to life online



Watch videos



Engage with interactivities



Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 2.1 Pre-test

1. **MC** Select the coefficient of x in the expression $\frac{-x^3}{3} + x^2 + 2ax + b$, where a and b are constants.

A. 1 B. $-\frac{1}{3}$ C. 2 D. $2a$ E. b

2. Evaluate the following expression if $x = -3$, $y = 1$ and $z = -2$.

$$2x^2 - y^2 - xyz$$

3. Determine the value of m in the following equation if $E = 25$ and $v = 5$.

$$E = \frac{1}{2}mv^2$$

4. **MC** Choose the expression out of the following that is not equivalent.

A. $x - y + z$ B. $z - y + x$ C. $x - (y - z)$ D. $z + x - y$ E. $z - (x + y)$

5. **MC** A bag of chips costs x cents. Select the correct expression for the cost of 6 bags of chips if each bag is discounted by 50 cents.

A. $6(x - 50)$ B. $6(x - 0.5)$ C. $6x - 50$ D. $6x - 0.5$ E. $50 - 6x$

6. Expand and simplify the following expressions.

a. $4(m + 3n) - (2m - n)$

b. $-x(y + 3) + y(5 - x)$

7. Expand and simplify the expression $(3q - r)(2q + r)$.

8. **MC** Students were asked to expand $-2(3x - 4)$. They gave the following answers.

Student 1: $-6x - 8$

Student 2: $-6x + 6$

Student 3: $6x - 8$

Student 4: $-6x + 8$

Student 5: $-5x + 6$

Choose the student who has the correct answer.

A. Student 1 B. Student 2 C. Student 3 D. Student 4 E. Student 5

9. State whether 23^2 can be expanded using the following method.

$$(20 + 3)^2 = 20^2 + 2 \times 20 \times 3 + 3^2$$

10. State whether the following statement is True or False.

$(a + 3)^2$ expanded is equal to $a^2 + 9$

11. Expand and simplify the following expression.

$$(2b - 5)^2 - (b + 4)(b - 3)$$

12. Factorise the following expressions.

a. $8p^2 + 20p$

b. $25x^2 - 10y$

c. $6xy + 8x^2y - 3xy^2$

13. Factorise the following expressions.

a. $7a(b + 3) - (b + 3)$

b. $10xy + 5x - 4y - 2$

14. A box has the following dimensions: x , $2x - 1$ and $3x + 2$.

Determine the correct expression for calculating the total surface area of the box.

15. Factorise the expression $x^2 + 3x - 4$.

LESSON

2.2 Using pronumerals

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify the number of terms in an algebraic expression
- identify the variables and coefficients of the terms in an expression
- identify the constant terms in an expression
- evaluate an expression or formula by substituting values for each variable.

2.2.1 Expressions

eles-4588

- A **pronumeral** is a letter or symbol that stands for a number. Algebra is the branch of mathematics in which we use and manipulate pronumerals.
- In algebra an **expression** is a group of **terms** in which each term is separated by a + sign or a – sign.
- An expression with two terms is called a **binomial**. An expression with one term is called a **monomial**.
- Here are some important things to note about the expression $5x - 3y + 2$:
 - The expression has three terms and is called a **trinomial**.
 - The pronumerals x and y may take different values and are called **variables**.
 - The term 2 has only one value and is called a **constant**.
 - The **coefficient** (the number in front) of x is 5, and the coefficient of y is -3 .

Algebraic expressions

An algebraic expression is a group of terms that are separated by + or – signs.

Consider the following expression: $5x^2 - 6y + 13x + 8 + 2y^3$.

- This expression is made up of five terms.
- The pronumerals/variables used in this expression are x and y .
- The coefficients of the terms are (in order): 5, -6 , 13 and 2.
- The coefficient of x is 13.
- The smallest coefficient is -6 .
- The constant is 8, it does not have a pronumeral.

WORKED EXAMPLE 1 Identifying coefficients of terms in an expression

For the expression $6x - 3xy + z + 2 + x^2z + \frac{y^2}{7}$, determine:

- a. the number of terms
- b. the coefficient of the second term
- c. the coefficient of the last term
- d. the constant term
- e. the term with the smallest coefficient
- f. the coefficient of x^2z .

THINK

- a. Count the number of terms.
- b. The second term is $-3xy$. The number part is the coefficient.
- c. The last term is $\frac{y^2}{7}$. This can be rewritten as $\frac{1}{7} \times y^2$.
- d. The constant is the term with no pronumeral.
- e. Identify the smallest coefficient and write the whole term to which it belongs.
- f. x^2z can be written as $1x^2z$.

WRITE

- a. There are six terms.
- b. The coefficient of the second term is -3 .
- c. The coefficient of the last term is $\frac{1}{7}$.
- d. The constant term is 2.
- e. The term with the smallest coefficient is $-3xy$.
- f. The coefficient of x^2z is 1.



2.2.2 Substitution

eles-4589

- We can evaluate (find the value of) an algebraic expression if we replace the pronumerals with their known values.
- This process is called substitution.
- Consider the expression $4x + 3y$. If we substitute the values $x = 2$ and $y = 5$, then the expression can be evaluated like this:

$$\begin{aligned}4 \times 2 + 3 \times 5 &= 8 + 15 \\ &= 23\end{aligned}$$

It is more common to use brackets for substitution instead of multiplication signs. For example:

$$\begin{aligned}4x + 3y &= 4(2) + 3(5) \\ &= 8 + 15 \\ &= 23\end{aligned}$$

WORKED EXAMPLE 2 Substituting values into terms

Evaluate the following terms for the values $x = 4$ and $x = -4$.

a. $3x^2$

b. $-x^2$

c. $\sqrt{5-x}$

THINK

- a. 1. Substitute 4 for x .
- 2. Only the x term is squared.
- 3. Evaluate.

WRITE

- a. If $x = 4$ $3x^2 = 3(4)^2$
 $= 3 \times 16$
 $= 48$

4. Repeat for $x = -4$.

$$\begin{aligned}\text{If } x = -4: \quad 3x^2 &= 3(-4)^2 \\ &= 3 \times 16 \\ &= 48\end{aligned}$$

b. 1. Substitute 4 for x .

$$\begin{aligned}\text{b. If } x = 4: \quad -x^2 &= -(4)^2 \\ &= -16\end{aligned}$$

2. Evaluate.

3. Repeat for $x = -4$.

$$\begin{aligned}\text{If } x = -4: \quad -x^2 &= -(-4)^2 \\ &= -16\end{aligned}$$

c. 1. Substitute 4 for x .

$$\begin{aligned}\text{c. If } x = 4: \quad \sqrt{5-x} &= \sqrt{5-4} \\ &= \sqrt{1} \\ &= 1\end{aligned}$$

2. Evaluate.

3. Repeat for $x = -4$.

$$\begin{aligned}\text{If } x = -4 \quad \sqrt{5-x} &= \sqrt{5-(-4)} \\ &= \sqrt{5+4} \\ &= \sqrt{9} \\ &= 3\end{aligned}$$

WORKED EXAMPLE 3 Evaluating expressions using substitution

If $x = 3$ and $y = -2$, evaluate the following expressions.

a. $3x + 2y$

b. $5xy - 3x + 1$

c. $2x^2 + y^2$

THINK

a. 1. Write the expression.

2. Substitute $x = 3$ and $y = -2$.

3. Evaluate.

b. 1. Write the expression.

2. Substitute $x = 3$ and $y = -2$.

3. Evaluate.

c. 1. Write the expression.

2. Substitute $x = 3$ and $y = -2$.

3. Evaluate.

WRITE

$$\begin{aligned}\text{a. } 3x + 2y &= 3(3) + 2(-2) \\ &= 9 - 4 \\ &= 5\end{aligned}$$


$$\begin{aligned}\text{b. } 5xy - 3x + 1 &= 5(3)(-2) - 3(3) + 1 \\ &= -30 - 9 + 1 \\ &= -38\end{aligned}$$

$$\begin{aligned}\text{c. } 2x^2 + y^2 &= 2(3)^2 + (-2)^2 \\ &= 18 + 4 \\ &= 22\end{aligned}$$

TI | THINK

a-c. On a Calculator page, complete the entry lines as:
 $3x + 2y | x = 3$
 and $y = -2$
 $5x \times y - 3x + 1 |$
 $x = 3$ and $y = -2$
 $2x^2 + y^2 | x = 3$
 and $y = -2$

Press ENTER after each entry.


WRITE

If $x = 3$ and $y = -2$, then
 a. $3x + 2y = 5$
 b. $5xy - 3x + 1 = -38$
 c. $2x^2 + y^2 = 22$.

CASIO | THINK

a-c. On the Main screen, complete the entry lines as:
 $3x + 2y | x = 3 |$
 $y = -2$
 $5x \times y - 3x + 1 |$
 $x = 3 | y = -2$
 $2x^2 + y^2 | x = 3 |$
 $y = -2$

Press EXE after each entry line.

WRITE

If $x = 3$ and $y = -2$, then
 a. $3x + 2y = 5$
 b. $5xy - 3x + 1 = -38$
 c. $2x^2 + y^2 = 22$.



2.2.3 Substitution into formulas

eles-4590

- A **formula** is a mathematical rule. Formulas are usually written using pronumerals.
- For example, the formula for the area of a rectangle is given by:

$$A = lw$$

where A represents the area of the rectangle, l represents the length and w represents the width.

- If a rectangular kitchen tile has length $l = 20$ cm and width $w = 15$ cm, we can substitute these values into the formula to find its area.

$$\begin{aligned} A &= lw \\ &= 20 \times 15 \\ &= 300 \text{ cm}^2 \end{aligned}$$



- If the area of a kitchen tile is 400 cm^2 and its width is 55 cm, then we can substitute these values into the formula to find its length.

$$\begin{aligned} A &= lw \\ 400 &= l \times 55 \\ l &= \frac{400}{55} \\ &= \frac{80}{11} \\ &\approx 7.3 \text{ cm (to 1 d.p.)} \end{aligned}$$

WORKED EXAMPLE 4 Substituting values into a formula

The formula for the voltage in an electrical circuit can be found using the formula known as Ohm's Law:

$$V = IR$$

where I = current (in amperes)

R = resistance (in ohms)

V = voltage (in volts).

a. Calculate V when:

i. $I = 2$ amperes, $R = 10$ ohms

ii. $I = 20$ amperes, $R = 10$ ohms.

b. Calculate I when $V = 300$ volts and $R = 600$ ohms.

THINK

a. i. 1. Write the formula.

2. Substitute $I = 2$ and $R = 10$.

3. Evaluate and write the answer using the correct units.

ii. 1. Write the formula.

2. Substitute $I = 20$ and $R = 10$.

3. Evaluate and write the answer using the correct units.

b. 1. Write the formula.

2. Substitute $V = 300$ and $R = 600$.

3. Evaluate and write the answer using the correct units.

WRITE

a. i. $V = IR$

$$= (2)(10)$$

$$= 20$$

The voltage is 20 volts.

ii. $V = IR$

$$= (20)(10)$$

$$= 200$$

The voltage is 200 volts.

b. $V = IR$

$$300 = I(600)$$

$$I = \frac{300}{600}$$

$$= 0.5$$

The current is 0.5 amperes.

Note: Even if we know nothing about volts, amperes and ohms, we can still do this calculation.

DISCUSSION

Which letters (pronumerals) should you avoid using when writing algebraic expressions?

Resources



eWorkbook

Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)



Video eLesson

Substitution 2 (eles-1892)



Interactivities

Individual pathway interactivity: Using pronumerals (int-4480)

Substituting positive and negative numbers (int-3765)

Individual pathways

PRACTISE

1, 4, 7, 10, 13, 14, 17, 18, 25

CONSOLIDATE

2, 5, 8, 11, 15, 19, 20, 21, 26

MASTER

3, 6, 9, 12, 16, 22, 23, 24, 27

Fluency

1. Determine the coefficient of each of the following terms.

a. $3x$ b. $-2m$ c. $\frac{x}{3}$ d. $-\frac{at}{4}$

2. Evaluate the coefficient of each of the following terms.

a. $5y$ b. $-2pq$ c. $\frac{x^2}{6}$ d. $-\frac{2mn^2}{7}$

3. Determine the coefficient of each of the following terms.

a. $15xy$ b. $-\frac{2pq}{5}$ c. $\frac{7w^2z^4}{2}$ d. $-\frac{pq^3}{7}$

4. **WE1** For each expression below, determine:

- i. the number of terms
- ii. the coefficient of the first term
- iii. the constant term
- iv. the term with the smallest coefficient.

a. $5x^2 + 7x + 8$
 b. $-9m^2 + 8m - 6$
 c. $5x^2y - 7x^2 + 8xy + 5$
 d. $9ab^2 - 8a - 9b^2 + 4$

5. For each expression below, determine:

- i. the number of terms
- ii. the coefficient of the first term
- iii. the constant term
- iv. the term with the smallest coefficient.

a. $4a - 2 + 9a^2b^2 - 3ac$
 b. $5s + s^2t + 9 + 12t - 3u$
 c. $-m + 8 + 5n^2m + m^2 + 2n$
 d. $7c^2d + 5d^2 + 14 - 3cd^2 - 2e$

6. For each expression below, determine:

- i. the number of terms
- ii. the coefficient of the x term
- iii. the constant term
- iv. the term with the smallest coefficient.

a. $-10x^2 + 3x - 7$
 b. $4y^2 + 6xy - 6x^2 + 7 - 3x$
 c. $10x^5 - 4x^{-3} + 2x^2 - 5x + 8$
 d. $6 - y^3 + 6x - 7y^5 + 10y - 15xy$

7. **WE2** Evaluate the following expressions for $x = 0$, $x = 2$ and $x = -2$.
- a. $5x + 2$ b. $x^2 - x$ c. $x^2 + 3x - 1$ d. $\frac{x}{2}$
8. Calculate the value of the following expressions for $x = 0$, $x = 3$ and $x = -3$.
- a. $-5x + 4$ b. $x^2 - 2x + 1$ c. $x(x + 4)$ d. $\frac{8}{x + 1}$
9. Determine the value of the following expressions for $x = 0$, $x = 1$ and $x = -2$.
- a. $\sqrt{x + 3}$ b. $\frac{2x^2 + 7x}{3}$ c. $(x + 3)(x - 4)$ d. $\frac{2x + 3}{5}$
10. **MC** Answer the following questions.
- a. If $x = -3$, then state the value of $-5x - 3$.
- A. -18 B. 12 C. 30 D. 18 E. -56
- b. Choose the expression that is a trinomial.
- A. $3x^2$ B. $x + 5 + 2$ C. $x^2 - 7$ D. $x^2 + x + 2$ E. $4x^3$
- c. Choose the value of x for which the expression $\sqrt{(x - 6)}$ cannot be evaluated.
- A. 2 B. 6 C. 7 D. 10 E. 16
- d. In the expression $5x^2 - 3xy + 0.5x - 0.3y - 5$, identify the smallest coefficient.
- A. 5 B. -3 C. 0.5 D. -0.3 E. -2
11. **WE3** Determine the value of the following expressions if $x = 2$, $y = -1$ and $z = 3$.
- a. $2x$ b. $3xy$ c. $2y^2z$
- d. $\frac{14}{x}$ e. $6(2x + 3y - z)$ f. $x^2 - y^2 + xyz$
12. If $x = 4$ and $y = -3$, evaluate the following expressions.
- a. $4x + 3y$ b. $3xy - 2x + 4$ c. $x^2 - y^2$

Understanding

13. **WE4** The change in the voltage in an electrical circuit can be found using the formula known as Ohm's Law $V = IR$, where $I =$ current (in amperes), $R =$ resistance (in ohms) and $V =$ voltage (in volts).
- a. Calculate V when:
- i. $I = 4$, $R = 8$ ii. $I = 25$, $R = 10$.
- b. Calculate R when:
- i. $V = 100$, $I = 25$ ii. $V = 90$, $I = 30$.
14. Evaluate each of the following by substituting the given values into each formula.
- a. If $A = bh$, calculate the value of A when $b = 5$ and $h = 3$.
- b. If $d = \frac{m}{v}$, determine the value of d when $m = 30$ and $v = 3$.
- c. If $A = \frac{1}{2}xy$, calculate the value of A when $x = 18$ and $y = 2$.
- d. If $A = \frac{1}{2}(a + b)h$, determine the value of A when $h = 10$, $a = 7$ and $b = 2$.
- e. If $V = \frac{AH}{3}$, determine the value of V when $A = 9$ and $H = 10$.

15. Calculate each of the following by substituting the given values into each formula.
- If $v = u + at$, calculate the value of v when $u = 4$, $a = 3.2$ and $t = 2.1$.
 - If $t = a + (n - 1)d$, determine the value of t when $a = 3$, $n = 10$ and $d = 2$.
 - If $A = \frac{1}{2}(x + y)h$, determine the value of A when $x = 5$, $y = 9$ and $h = 3.2$.
 - If $A = 2b^2$, calculate the value of A when $b = 5$.
 - If $y = 5x^2 - 9$, determine the value of y when $x = 6$.
16. Calculate each of the following by substituting the given values into each formula.
- If $y = x^2 - 2x + 4$, determine the value of y when $x = 2$.
 - If $a = -3b^2 + 5b - 2$, calculate the value of a when $b = 4$.
 - If $s = ut + \frac{1}{2}at^2$, determine the value of s when $u = 0.8$, $t = 5$ and $a = 2.3$.
 - If $F = \frac{mp}{r^2}$, calculate the value of F correct to 2 decimal places, when $m = 6.9$, $p = 8$ and $r = 1.2$.
 - If $C = \pi d$, determine the value of C correct to 2 decimal places if $d = 11$.

Reasoning


17. The area of a triangle is given by the formula $A = \frac{1}{2}bh$, where b is the length of the base and h is the perpendicular height of the triangle.
- Show that the area is 12 cm^2 when $b = 6 \text{ cm}$ and $h = 4 \text{ cm}$.
 - Evaluate h if $A = 24 \text{ cm}^2$ and $b = 4 \text{ cm}$.
18. Using $E = F + V - 2$, where F is the number of faces on a prism, E is the number of edges and V is the number of vertices, calculate:
- E if $F = 5$ and $V = 7$
 - F if $E = 10$ and $V = 2$.
19. The formula to convert degrees Fahrenheit (F) to degrees Celsius (C) is $C = \frac{5}{9}(F - 32)$.
- Determine the value of C when $F = 59$.
 - Show that when Celsius (C) is 15, Fahrenheit (F) is 59.



20. The length of the hypotenuse of a right-angled triangle (c) can be found using the formula $c = \sqrt{a^2 + b^2}$, where a and b are the lengths of the other 2 sides.
- Calculate c when $a = 3$ and $b = 4$.
 - Show that when $a = 5$ and $c = 13$, $b = 12$.
21. The kinetic energy (E) of an object is found by using the formula $E = \frac{1}{2}mv^2$, where m is the mass and v is the velocity of the object.
- Calculate E when $m = 3$ and $v = 3.6$.
 - Show that when $E = 25$ and $v = 5$, $m = 2$.

22. If the volume of a prism (V) is given by the formula $V = AH$, where A is the area of the cross-section and H is the height of the prism, calculate:

- a. V when $A = 7 \text{ cm}^2$ and $H = 9 \text{ cm}$
- b. H when $V = 120 \text{ cm}^3$ and $A = 30 \text{ cm}^2$.



23. The volume of a cylinder (v) is given by $v = \pi r^2 h$, where r is the radius in centimetres and h is the height of the cylinder in centimetres.

- a. Determine v correct to 2 decimal places if $r = 7$ and $h = 3$.
- b. Determine h correct to 2 decimal places if $v = 120$ and $r = 2$.

24. The surface area of a cylinder (S) is given by $S = 2\pi r(r + h)$, where r is the radius of the circular end and h is the height of the cylinder.

- a. Calculate S (to 2 decimal places) if $r = 14$ and $h = 5$.
- b. Show that for a cylinder of surface area 240 units^2 and radius 5 units, the height is 2.64 units, correct to 2 decimal places.

Problem solving

25. a. In a magic square, every row, column and diagonal adds to the same number. This number is called the magic number for that square. Complete the unfinished magic square by first finding an algebraic expression for the magic number.
- b. Make three different magic squares by substituting different values for g and d in the completed magic square from part a.
- c. Calculate the magic number for each magic square.

	$2d + 2g$	
	$d + g$	
	0	$d + 2g$

26. A flat, rectangular board is built by gluing together a number of square pieces of the same size. The board is m squares wide and n squares long. In terms of m and n , write expressions for:

- a. the total number of squares
- b. the number of completely surrounded squares
- c. the perimeter of the figure, given that each square is 1 unit by 1 unit.

27. Determine which of the following eight expressions is **not** equivalent to the others.

$x - y + z$	$z - y + x$
$z - (y - x)$	$-y + x + z$
$z + x - y$	$x - (y - z)$
$y - (z - x)$	$x + z - y$

LESSON

2.3 Algebra in worded problems

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- assign pronumerals to the unknown quantities in a worded problem
- convert the worded statements presented in a question into algebraic expressions.

2.3.1 Algebra in worded problems

eles-4591

- In order to make use of the skills you learn in algebra, it is useful to be able to take a real-life problem and convert it into an algebraic expression.
- The starting point for any worded problem is to define the unknown quantities in the question by assigning appropriate pronumerals to those unknown quantities.

Key language used in worded problems

In order to be able to turn a worded problem into an algebraic expression, it helps to look out for the following kinds of words.

- **Words for addition:** sum, altogether, add, more than, and, in total
- **Words for subtraction:** difference, less than, take away, take off, fewer than
- **Words for multiplication:** product, groups of, times, of, for each, double, triple
- **Words for division:** quotient, split into, halve, thirds

WORKED EXAMPLE 5 Converting worded problems into algebraic expressions

Write an algebraic expression for each of the following, choosing an appropriate pronumeral if necessary.

- The number that is 6 more than Ben's age.
- The total value of a bundle of \$10 notes.
- The total cost of 8 adults' and 3 children's train tickets.
- The product of a and w .
- w less than a .

THINK

- Ben's age is the unknown quantity.
Note: You must not say, 'Let Ben = b ', because b is a number.
 - Write an expression for Ben's age plus 6.
- The number of notes is the unknown quantity.
 - The value is 10 times the number of notes.
Write the expression for the value of the notes.
- The cost of tickets is the unknown quantity.
 - The adults' tickets cost $8 \times a$ and the children's Tickets cost $3 \times c$. Write the expression for the total cost of the tickets.

WRITE

- Let Ben's age = b years.
 $b + 6$
- Let n = the number of \$10 notes
Total value = $10n$ dollars
- Let an adult's ticket cost a dollars and a child's ticket cost c dollars.
Total cost = $8a + 3c$ dollars.

d. 'Product' means to multiply.

Write the expression.

d. $a \times w$

e. In this expression, w is subtracted from a .

Write the expression.

e. $a - w$

WORKED EXAMPLE 6 Solving a worded problem

Kot has a block of chocolate that is made up of 40 smaller pieces.

a. If he breaks off x pieces to give to Akira, evaluate how much of the block remains.

b. Kot then eats $\frac{1}{5}$ of the remaining chocolate for dessert. Determine how much he ate.

c. Finally Kot gives a third of what remains to his sister Temar. Calculate how much Temar receives if the original amount Kot gave to Akira was 10 pieces.

THINK

a. 1. The unknown quantity is the amount passed to Akira, which is x .

2. Write an expression for x less than 40.

b. 1. The unknown quantity is still x .

2. Using the answer to **a**, write an expression for $\frac{1}{5}$ of what is left. *Note:* 'of' is a word that means multiplication.

c. 1. The unknown quantity is still x from part **a**.

2. Use your answers to **a** and **b** to write an expression for the amount left over after Kot eats $\frac{1}{5}$ of the remaining chocolate.

3. Multiply that expression by $\frac{1}{3}$.

4. Substitute the value of $x = 10$ into the expression to find the final answer. Write the answer in words.

WRITE

a. $x =$ number of pieces given to Akira

Amount remaining = $40 - x$

b. Amount eaten = $\frac{1}{5}$ of amount remaining

Amount eaten = $\frac{1}{5} \times (40 - x)$

Amount eaten = $\frac{1}{5}(40 - x)$

c. Amount left = $40 - x - \frac{1}{5}(40 - x)$

Amount passed to Temar is $\frac{1}{3}$ of what is left over after Kot's dessert.

Temar's amount = $\frac{1}{3} \left(40 - x - \frac{1}{5}(40 - x) \right)$

When $x = 10$

= $\frac{1}{3} \left(40 - 10 - \frac{1}{5}(40 - 10) \right)$

= $\frac{1}{3} \left(30 - \frac{1}{5}(30) \right)$

= $\frac{1}{3}(30 - 6)$

= $\frac{1}{3}(24)$

= 8

Temar receives 8 pieces of chocolate from Kot.



eWorkbook Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)



Interactivity Individual pathway interactivity: Algebra in worded problems (int-4481)

Exercise 2.3 Algebra in worded problems

2.3 Quick quiz **on**

2.3 Exercise

Individual pathways

PRACTISE

1, 4, 5, 6, 10, 14

CONSOLIDATE

2, 7, 8, 11, 15, 16

MASTER

3, 9, 12, 13, 17, 18

Fluency

- Iam studies five more subjects than Xan. Determine how many subjects Iam studies if:
 - Xan studies six subjects
 - Xan studies x subjects
 - Xan studies y subjects.
- Eva and Juliette walk home from school together. Eva's home is 2 km further from school than Juliette's home. Calculate how far Eva walks if Juliette's home is:
 - 1.5 km from school
 - x km from school.
- Lisa watched television for 2.5 hours today. Calculate how many hours she will watch tomorrow if she watches:
 - 1.5 hours more than she watched today
 - t hours more than she watched today
 - y hours fewer than she watched today.
- Samir ran d km today. Calculate the distance he will cover tomorrow if he runs:
 - x km less than he ran today
 - y km more than he ran today
 - 4 km more than double the distance that he ran today.



Understanding

- WE5** Write an algebraic expression for each of the following questions.
 - If it takes 10 minutes to iron a single shirt, calculate how long it would take to iron all of Anthony's shirts.
 - Ross has 30 dollars more than Nick. If Nick has N dollars, calculate how much money Ross has.
 - In a game of Aussie Rules football Luciano kicked 4 more goals than he kicked behinds.
 - Determine how many behinds Luciano scored, if g is the number of goals kicked.
 - Calculate the number of points Luciano scored.
(Note: 1 goal scores 6 points, 1 behind scores 1 point.)

6. Jeff and Chris play Aussie Rules football for opposing teams. Jeff's team won when the two teams played each other.

- a. Calculate how many points Jeff's team scored if they kicked:
- 14 goals and 10 behinds
 - x goals and y behinds.
- b. Calculate how many points Chris's team scored if his team kicked:
- 10 goals and 6 behinds
 - p goals and q behinds.

- c. Calculate how many points Jeff's team won by if:

- Chris's team scored 10 goals and 6 behinds, and Jeff's team scored 14 goals and 10 behinds
- Chris's team scored p goals and q behinds, and Jeff's team scored x goals and y behinds.

(Note: 1 goal scores 6 points, 1 behind scores 1 point.)

7. Yvonne's mother gives her x dollars for each school subject she passes. If she passes y subjects, determine how much money Yvonne receives.

8. Rosya buys a bag containing x Smarties.

- If they divide them equally among n people, calculate how many each person gets.
- If they keep half of the Smarties for themselves and divide the remaining Smarties equally among n people, calculate how many Smarties each person gets.

9. A piece of licorice is 30 cm long.

- If Ameer cuts d cm off, calculate how much licorice remains.
- If Ameer cuts off $\frac{1}{4}$ of the remaining licorice, calculate how much licorice has been cut off.
- Determine how much licorice remains after the two cuts that Ameer has made.



Reasoning

10. One-quarter of a class of x students plays tennis at the weekend. One-sixth of the class plays tennis and also swims at the weekend.

- Write an expression to represent the number of students playing tennis at the weekend.
- Write an expression to represent the number of students playing tennis and also swimming at the weekend.
- Show that the number of students playing only tennis at the weekend is $\frac{x}{12}$.

11. During a 24-hour period, Vanessa uses her computer for c hours. Her brother Darren uses it for $\frac{1}{7}$ of the remaining time.

- Determine the length of time for which Darren used the computer.
- Show that the total number of hours that Vanessa and Darren use the computer during a 24-hour period can be expressed as $\frac{6c + 24}{7}$.

12. Dizem had a birthday party last weekend and invited n friends. The table shown indicates the number of friends at Dizem's party at certain times during the evening. Everybody arrived by 8.30 pm and everyone left the party by 11 pm.
- Show that 1 person arrived between 7.00 pm and 7.30 pm.
 - Determine the start and end times during which the most friends arrived.
 - Calculate how many friends were invited but did not arrive.
 - Show that 24 friends were invited in total.
 - Determine the start and end times during which the most friends were present at the party.

Time	Number of friends
7.00 pm	$n - 24$
7.30 pm	$n - 23$
8.00 pm	$n - 8$
8.30 pm	$n - 5$
9.00 pm	$n - 5$
9.30 pm	$n - 7$
10.00 pm	$n - 12$
10.30 pm	$n - 18$
11.00 pm	$n - 24$



13. **WE6** Ty earns \$ y per week working at his part-time job.
- If he spends \$30 a week on his phone, calculate how much money he has left.
 - If he saves two-thirds of the remaining money for rent each week, calculate how much money is left.
 - If, after paying for his phone, saving for rent and spending \$100 on food, Ty still has \$200 left over, determine how much he earns in a week.

Problem solving

14. A child builds a pyramid out of building blocks using the pattern shown.
- Determine a rule that gives the number of blocks on the bottom layer, b , for a tower that is h blocks high.
 - If the child wants the tower to be 10 blocks high, determine how many blocks they should begin with on the bottom layer.



15. A shop owner is ordering vases from a stock catalogue. Red vases cost \$4.20 and clear vases cost \$5.70. If she buys 25 vases for \$120, determine how many of each type of vase she bought.
16. A father is 4 times the age of his son. In 4 years time the father will be 3 times the age of his son. Determine how old the father is now.
17. If a circle has 3 equally spaced dots (A, B and C) on its circumference, only 1 triangle can be formed by joining the dots. Investigate to determine the number of triangles that could be formed by 4, 5 or more equally spaced dots on the circumference.



From your investigation, can you see a pattern in the number of triangles (t) formed for a circle with d equally spaced dots on its circumference?

Determine whether it is possible to express this as a formula.

18. Answer the following questions, showing full working.

a. Transpose the equation $T = \frac{x-3}{x-1}$ to make x the subject.

b. Evaluate T when $x = 1$.

c. Evaluate x when $T = 1$.

LESSON

2.4 Simplification of algebraic expressions

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify whether two terms are like terms
- simplify algebraic expressions by adding and subtracting like terms
- simplify algebraic expressions that involve multiplication and division of multiple algebraic terms.

2.4.1 Addition and subtraction of like terms

eles-4592

- Simplifying an expression involves writing it in a form with the least number of terms, without \times or \div signs, and with any fractions expressed in their simplest form.
- **Like terms** have identical pronumeral parts (including the power), but may have different coefficients.
For example:
 - $5y$ and $10y$ are like terms, but $5y^2$ and $10y$ are not like terms.
 - $3mn^2$ and $-4mn^2$ are like terms, but $3m^2n$ and $-2mn^2$ are not like terms.
- If a term contains more than one pronumeral, the convention is to write the pronumerals in alphabetical order. This makes it easier to identify like terms.
For example:
 - $3cb^2a$ and $-7b^2ca$ are like terms and should be written as $3ab^2c$ and $-7ab^2c$ to make this easier to identify.
- We can simplify like terms by adding (or subtracting) the coefficients to form a single term. We cannot simplify any terms that are not like terms.
For example:
 - $10xy + 3xy - 7xy = 6xy$.

WORKED EXAMPLE 7 Simplifying algebraic expressions

Simplify the following expressions.

a. $x + 2x + 3x - 5$

b. $9a^2b + 5ba^2 + ab^2$

c. $-12 - 4c^2 + 10 + 5c^2$

THINK

- a. 1. Write the expression.
2. Collect the like terms x , $2x$ and $3x$.
- b. 1. Write the expression.
2. Collect the like terms $9a^2b$ and $5ba^2$.
- c. 1. Write the expression.
2. Collect the like terms $4c^2$ and $5c^2$.
3. Simplify the expression.

WRITE

- a. $x + 2x + 3x - 5$
 $= 6x - 5$
- b. $9a^2b + 5ba^2 + ab^2$
 $= 14a^2b + ab^2$
- c. $-12 - 4c^2 + 10 + 5c^2$
 $= -12 + 10 - 4c^2 + 5c^2$
 $= -2 + c^2$ or $c^2 - 2$

2.4.2 Multiplication of algebraic terms

eles-4593

- Any number of algebraic terms can be multiplied together to produce a single term.
- Numbers can be multiplied in any order, so it is easiest to multiply the coefficients first, then multiply the pronumerals in alphabetical order.
- Remember that, when multiplying pronumerals, we also add their indices together.

Note: $x = x^1$

$$\begin{aligned} 4ab \times 3a^2b \times 2a^3 &= 4 \times a \times b \times 3 \times a^2 \times b \times 2 \times a^3 \\ &= 4 \times 3 \times 2 \times a \times a^2 \times a^3 \times b \times b \\ &= 24 \times a^6 \times b^2 \\ &= 24a^6b^2 \end{aligned}$$

WORKED EXAMPLE 8 Multiplication of algebraic terms

Simplify the following expressions.

a. $4a \times 2b \times a$

b. $7ax \times (-6bx) \times (-2abx)$

THINK

- a. 1. Write the expression.
 2. Rearrange the expression, writing the coefficients first.
 3. Multiply the coefficients and pronumerals separately.
- b. 1. Write the expression.
 2. Rearrange the expression, writing the coefficients first.
 3. Multiply the coefficients and pronumerals separately.

WRITE


a. $4a \times 2b \times a$
 $= 4 \times 2 \times a \times a \times b$
 $= 8a^2b$

b. $7ax \times (-6bx) \times (-2abx)$
 $= 7 \times (-6) \times (-2) \times a \times a \times b \times b \times x \times x \times x$
 $= 84a^2b^2x^3$

TI | THINK

- a–b. In a new problem, on a Calculator page, complete the entry lines as:
 $4a \times 2b \times a$ $7a \times x \times$
 $-6b \times x \times -2a \times b \times x$
- Press ENTER after each entry line.

WRITE


a–b. 

a. $4a \times 2b \times a = 8a^2b$
 b. $7a \times x \times -6b \times x \times -2a \times b \times x = 84a^2b^2x^3$

CASIO | THINK

- a–b. On the Main screen, complete the entry lines as:
 $4a \times 2b \times a$
 $7a \times x \times -6b \times x \times -2a \times b \times x$
- Press EXE after each entry line.

WRITE

a–b. 

a. $4a \times 2b \times a = 8a^2b$
 b. $7a \times x \times -6b \times x \times -2a \times b \times x = 84a^2b^2x^3$

2.4.3 Division of algebraic terms

eles-4594

- When dividing two algebraic terms, first rewrite the terms in fraction form.
- Simplify the fraction by dividing through by the highest common factor of the coefficients and cancelling down the pronumerals.
- Remember that when dividing pronumerals we subtract their indices.

$$\begin{aligned}
 36x^3yz^2 \div (40x^2y^2) &= \frac{36x^3yz^2}{40x^2y^2} \\
 &= \frac{\overset{3}{\cancel{36}} \times x \times \cancel{x} \times \cancel{x} \times y \times z \times z}{\underset{10}{\cancel{40}} \times \cancel{x} \times \cancel{x} \times y \times y} \\
 &= \frac{9xz^2}{10y}
 \end{aligned}$$

WORKED EXAMPLE 9 Dividing algebraic terms

Simplify the following terms.

a. $\frac{12xy}{4xz}$

b. $(-8ab^2) \div (-16a^2b)$

THINK

a. 1. Write the term.

2. Cancel the common factors.

3. Write the answer.

b. 1. Write the term as a fraction.

2. Cancel the common factors.

3. Simplify and write the answer.

WRITE

a. $\frac{12xy}{4xz}$

$$\begin{aligned}
 &= \frac{\overset{3}{\cancel{12}} \times \cancel{x} \times y}{\underset{1}{\cancel{4}} \times x \times z} \\
 &= \frac{3y}{z}
 \end{aligned}$$




b. $\frac{-8ab^2}{-16a^2b}$

$$\begin{aligned}
 &= \frac{\overset{1}{\cancel{-8}} \times \overset{1}{\cancel{a}} \times \cancel{b^2} \times b}{\underset{2}{\cancel{-16}} \times \underset{a}{\cancel{a^2}} \times \underset{1}{\cancel{b}}} \\
 &= \frac{1}{2} \times \frac{1}{a} \times \frac{b}{1} \\
 &= \frac{b}{2a}
 \end{aligned}$$

DISCUSSION

Is the expression ab the same as ba ? Explain your answer.



-  **eWorkbook** Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)
-  **Video eLesson** Simplification of expressions (eles-1884)
-  **Interactivities** Individual pathway interactivity: Simplification of algebraic expressions (int-4482)
 - Simplifying expressions (int-3771)
 - Multiplying variables (int-3772)
 - Dividing expressions with variables (int-3773)

Exercise 2.4 Simplification of algebraic expressions

2.4 Quick quiz **on**

2.4 Exercise

Individual pathways

PRACTISE

1, 3, 9, 10, 11, 14, 19, 22, 25, 26

CONSOLIDATE

2, 4, 5, 7, 12, 15, 17, 20, 23, 27, 28

MASTER

6, 8, 13, 16, 18, 21, 24, 29, 30

Fluency

1. For each of the following terms, select the terms listed in brackets that are like terms.

<p>a. $6ab$ ($7a, 8b, 9ab, -ab, 4a^2b^2$)</p> <p>c. $3az$ ($3ay, -3za, -az, 3z^2a, 3a^2z$)</p>	<p>b. $-x$ ($3xy, -xy, 4x, 4y, -yx$)</p> <p>d. x^2 ($2x, 2x^2, 2x^3, -2x, -x^2$)</p>
--	--

2. For each of the following terms, select the terms listed in brackets that are like terms.

<p>a. $2x^2y$ ($xy, -2xy, -2xy^2, -2x^2y, -2x^2y^2$)</p> <p>b. $3x^2y^5$ ($3xy, 3x^5y^2, 3x^4y^3, -x^2y^5, -3x^2y^5$)</p> <p>c. $5x^2p^3w^5$ ($-5x^3w^5p^3, p^3x^2w^5, 5xp^3w^5, -5x^2p^3w^5, w^5p^2x^3$)</p> <p>d. $-x^2y^5z^4$ ($-xy^5, -y^2z^5x^4, -x+y+z, 4y^5z^4x^2, -2x^2z^4y^5$)</p>	
---	--

3. **WE7** Simplify the following expressions.

<p>a. $5x + 2x$</p> <p>d. $13q - 2q$</p>	<p>b. $3y + 8y$</p> <p>e. $17r - 9r$</p>	<p>c. $7m + 12m$</p> <p>f. $-x + 4x$</p>
--	--	--

4. Simplify the following expressions.

<p>a. $5a + 2a + a$</p> <p>d. $14p - 3p + 5p$</p>	<p>b. $9y + 2y - 3y$</p> <p>e. $2q^2 + 7q^2$</p>	<p>c. $7x - 2x + 8x$</p> <p>f. $5x^2 - 2x^2$</p>
---	--	--

5. Simplify the following expressions.

<p>a. $6x^2 + 2x^2 - 3y$</p> <p>d. $-5m^2 + 5m - 4m + 15$</p>	<p>b. $3m^2 + 2n - m^2$</p> <p>e. $12a^2 + 3b + 4b^2 - 2b$</p>	<p>c. $-2g^2 - 4g + 5g - 12$</p> <p>f. $6m + 2n^2 - 3m + 5n^2$</p>
---	--	--

6. Simplify the following expressions.

<p>a. $3xy + 2y^2 + 9yx$</p> <p>d. $4m^2n + 3n - 3m^2n + 8n$</p>	<p>b. $3ab + 3a^2b + 2a^2b - ab$</p> <p>e. $-3x^2 - 4yx^2 - 4x^2 + 6x^2y$</p>	<p>c. $9x^2y - 3xy + 7yx^2$</p> <p>f. $4 - 2a^2b - ba^2 + 5b - 9a^2$</p>
--	---	--

7. **MC** Choose which of the following is a simplification of the expression $18p - 19p$.

A. p	B. $-p$	C. p^2	D. -1	E. $p - 1$
---------------	----------------	-----------------	----------------	-------------------

8. **MC** Choose which of the following is a simplification of the expression $5x^2 - 8x + 6x - 9$.
- A. $3x - 9$ B. $3x^2 - 9$ C. $5x^2 + 2x - 9$
 D. $5x^2 - 2x - 9$ E. $-3x^2 + 6x - 9$
9. **MC** Choose which of the following is a simplification of the expression $12a - a + 15b - 14b$.
- A. $11a + b$ B. 12 C. $11a - b$
 D. $13a + b$ E. $-12a - b$
10. **MC** Choose which of the following is a simplification of the expression $-7m^2n + 5m^2 + 3 - m^2 + 2m^2n$.
- A. $-9m^2n + 4m^2 + 3$ B. $-9m^2n + 8$ C. $-5m^2n - 4m^2 + 3$
 D. $-5m^2n + 4m^2 + 3$ E. $5m^2n + 4m^2 + 3$
11. **WE8** Simplify the following.
- a. $3m \times 2n$ b. $4x \times 5y$ c. $2p \times 4q$ d. $5x \times (-2y)$ e. $3y \times (-4x)$
12. Simplify the following.
- a. $(-3m) \times (-5n)$ b. $5a \times 2a$ c. $3mn \times 2p$ d. $(-6ab) \times b$ e. $(-5m) \times (-2mn)$
13. Simplify the following.
- a. $(-6a) \times 3ab$ b. $(-3xy) \times (-5xy) \times 2x$ c. $4pq \times (-p) \times 3q^2$
 d. $4c \times (-7cd) \times 2c$ e. $(-3a^2) \times (-5ab^3) \times 2ab^4$
14. **WE9** Simplify the following.
- a. $\frac{6x}{2}$ b. $\frac{9m}{3}$ c. $\frac{12y}{6}$ d. $\frac{8m}{2}$ e. $12m \div 3$
15. Simplify the following.
- a. $14x \div 7$ b. $-21x \div 3$ c. $-32m \div 8$ d. $\frac{4m}{8}$ e. $\frac{6x}{18}$
16. Simplify the following.
- a. $\frac{8mn}{18n}$ b. $\frac{6ab}{12a^2b}$ c. $\frac{28xyz}{14x}$
 d. $\frac{2x^2yz}{8xz}$ e. $-7xy^2z^2 \div (11xyz)$

Understanding

17. Simplify the following.
- a. $5x \times 4y \times 2xy$ b. $7xy \times 4ax \times 2y$ c. $\frac{6x^2y}{12y^2}$ d. $\frac{-15x^2ab}{12b^2x^2}$
18. Simplify the following.
- a. $\frac{2p^3q^2}{p^3q^2}$ b. $(-4a) \times (-5ab^2) \times 2a$
 c. $(-a) \times 4ab \times 2ba \times b$ d. $2a \times 2a \times 2a \times 2a$
19. Jim buys m pens at p cents each and n books at q dollars each.
- a. Calculate how much Jim spends in:
 i. dollars ii. cents.
 b. Calculate how much change Jim will have if he starts with \$20.
20. At a local discount clothing store 4 shirts and 3 pairs of shorts cost \$138 in total. If a pair of shorts costs 2.5 times as much as a shirt, calculate the cost of each kind of clothing.

21. Anthony and Jamila are taking their 3 children to see a movie at the cinema. The total cost for the 2 adults and 3 children is \$108. If an adult's ticket is 1.5 times the cost of a child's, calculate the cost of 1 child's ticket plus 1 adult's ticket.

Reasoning

22. Class 9A were given an algebra test. One of the questions is shown below.

Simplify the following expression: $\frac{3ab}{2} \times \frac{4ac}{6b} \times 7c$.

Sean, who is a student in class 9A, wrote his answer as $\frac{12aabc}{12b} \times 7c$. Explain why Sean's answer is incorrect, and write the correct answer.

23. Using the appropriate method to divide fractions, simplify the following expression.

$$\frac{15a^2b^2}{16c^4} \div \left(\frac{5ac^2}{4b^3} \times \frac{a}{2b^4} \right)$$

24. Answer the following questions.



a. Simplify the expression $\frac{n}{(n+1)} \times \frac{(n+1)}{(n+2)}$.

b. Simplify the expression $\frac{n}{(n+1)} \times \frac{(n+1)}{(n+2)} \times \frac{(n+2)}{(n+3)}$.



c. Use the results from parts a and b to evaluate $\frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} \times \dots \times \frac{99}{100}$.

Problem solving

25. a. Fill in the empty bricks in the following pyramids. The expression in each brick is obtained by adding the expressions in the two bricks below it.




- b. Fill in the empty bricks in the following pyramids. The expression in each brick is obtained by multiplying the expressions in the two bricks below it.




26. For the triangle shown:

- a. write an expression for the perimeter of the triangle
- b. show that the triangle's perimeter can be simplified to $\frac{43}{21}x$
- c. calculate the cost (to the nearest dollar) to frame the triangle using timber that costs \$4.50 per metre, if x equals 2 metres.




27. For the rectangle shown:



- write an expression for the perimeter of the rectangle
 - show that the expression for the perimeter can be simplified to $6\frac{4}{7}w$
 - calculate the cost (to the nearest cent) to create a wire frame for the rectangle using wire that costs \$1.57 per metre, if w equals seven metres.
28. A rectangular chocolate block has dimensions x and $(4x - 7)$.



- Write an expression for the perimeter (P) in the form $P = \square x + \square(4x - 7)$.
- Expand and simplify this expression.
- Calculate the perimeter of the chocolate block when $x = 3$.
- Explain why x cannot equal 1.

29. A doghouse in the shape of a rectangular prism is to be reinforced with steel edging along its outer edges including the base.



The cost of the steel edging is \$1.50 per metre. All measurements are in centimetres.

- Write an expression for the total length of the straight edges of the frame.
 - Expand the expression.
 - Evaluate the cost of steel needed when $x = 80$ cm.
30. A tile manufacturer produces tiles that have the side lengths shown. All measurements are in centimetres.



- Write an expression for the perimeter of each shape.
- Evaluate the value of x for which the perimeter of the triangular tile is the same as the perimeter of the square tile.

LESSON

2.5 Expanding brackets

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- apply the distributive law to expand and simplify an expression with a single set of brackets
- apply the distributive law to expand and simplify expressions containing two or more sets of brackets.

2.5.1 The Distributive Law

eles-4595

- There are two ways of calculating the area of the rectangle shown.
 1. The rectangle can be treated as a single shape with a length of 4 cm and a width of 5 + 3 cm.


$$\begin{aligned}A &= l \times w \\ &= 4(5 + 3) \\ &= 4 \times 8 \\ &= 32 \text{ cm}^2\end{aligned}$$



2. The areas of the two smaller rectangles can be added together.

$$\begin{aligned}A &= 4 \times 5 + 4 \times 3 \\ &= 20 + 12 \\ &= 32 \text{ cm}^2\end{aligned}$$

- The length of the rectangle shown is a and its width is $b + c$.



- This rectangle's area can be found in two different ways.

1. $A = l \times w$
 $= a(b + c)$

This expression is described as **factorised** because it shows one number (or factor) multiplied by another. The two factors are a and $(b + c)$.

2. The areas of the two small rectangles can be added together.

$$A = ab + ac$$

This expression is described as **expanded**, which means that it is written without brackets.

- $a(b + c) = ab + ac$ is called the **Distributive Law**.

The Distributive Law

In order to expand a single set of brackets we apply the distributive law, which states that:

$$a(b + c) = a \times b + a \times c = ab + ac$$

- A helpful way to expand brackets is to draw arrows between factors as you work out each multiplication.

$$x(x - y) = x^2 \dots$$

$$x(x - y) = x^2 - xy$$

WORKED EXAMPLE 10 Expanding a single set of brackets

Use two different methods to calculate the value of the following.

a. $7(5 + 15)$

b. $10(9 - 1)$

THINK

a. 1. Method 1: Work out the brackets first, then evaluate.

2. Method 2: Expand the brackets first, then evaluate.

b. 1. Method 1: Work out the brackets first, then evaluate.

2. Method 2: Expand the brackets first, then evaluate

WRITE

a. $7(5 + 15)$
 $= 7 \times 20$
 $= 140$

$7(5 + 15)$
 $= 7 \times 5 + 7 \times 15$
 $= 35 + 105$
 $= 140$

b. $10(9 - 1)$
 $= 10 \times 8$
 $= 80$

$10(9 - 1)$
 $= 10 \times 9 + 10 \times -1$
 $= 90 - 10$
 $= 80$

WORKED EXAMPLE 11 Expanding using arrows

Expand the following expressions.

a. $5(x + 3)$

b. $-4y(2x - w)$

c. $3x(5 - 6x + 2y)$

THINK

a. 1. Draw arrows to help with the expansion.

2. Simplify and write the answer.

b. 1. Draw arrows to help with the expansion.

2. Simplify and write the answer.

c. 1. Draw arrows to help with the expansion.

2. Simplify and write the answer.

WRITE

a. $5(x + 3)$
 $= 5 \times x + 5 \times 3$
 $= 5x + 15$

b. $-4y(2x - w)$
 $= -4y \times 2x - 4y \times (-w)$
 $= -8xy + 4wy$

c. $3x(5 - 6x + 2y)$
 $= 3x \times 5 + 3x \times (-6x) + 3x \times 2y$
 $= 15x - 18x^2 + 6xy$

eles-4596 **2.5.2 Expanding and simplifying**

- When a problem involves **expanding** that is more complicated, it is likely that the like terms will need to be collected and simplified after you have expanded the brackets.
- When solving these more complicated problems, expand all sets of brackets first, then simplify any like terms that result from the expansion.

WORKED EXAMPLE 12 Expanding and simplifying

Expand and simplify the following expressions by collecting like terms.

a. $4(x - 4) + 5$

b. $x(y - 2) + 5x$

c. $-x(y - 2) + 5x$

d. $7x - 6(y - 2x)$

THINK

a. 1. Expand the brackets.

2. Simplify and write the answer.

b. 1. Expand the brackets.

2. Simplify and write the answer.

c. 1. Expand the brackets.

2. Simplify and write the answer. (*Note: There are no like terms.*)

d. 1. Expand the brackets.

2. Simplify and write the answer.

WRITE

a. $4(x - 4) + 5$
 $= 4x - 16 + 5$
 $= 4x - 11$

b. $x(y - 2) + 5x$
 $= xy - 2x + 5x$
 $= xy + 3x$

c. $-x(y - 2) + 5x$
 $= -xy + xz + 5x$

d. $7x - 6(y - 2x)$
 $= 7x - 6y + 12x$
 $= 19x - 6y$

WORKED EXAMPLE 13 Expanding two sets of brackets and simplifying

Expand and simplify the following expressions.

a. $5(x + 2y) + 6(x - 3y)$

b. $5x(y - 2) - y(x + 3)$

THINK

a. 1. Expand each set of brackets.

2. Simplify and write the answer.

b. 1. Expand each set of brackets.

2. Simplify and write the answer.

WRITE


a. $5(x + 2y) + 6(x - 3y)$
 $= 5x + 10y + 6x - 18y$
 $= 11x - 8y$

b. $5x(y - 2) - y(x + 3)$
 $= 5xy - 10x - xy - 3y$
 $= 4xy - 10x - 3y$

2.5.3 Expanding binomial factors

eles-4597

- Remember that a **binomial** is an expression containing two terms, for example $x + 3$ or $2y - z^2$. In this section we will look at how two binomials can be multiplied together.
- The rectangle shown has length $a + b$ and width $c + d$.



- There are two ways of finding the area of the large rectangle.
 - $A = l \times w$
 $= (a + b) \times (c + d)$
 $= (a + b)(c + d)$
 This is a factorised expression in which the two factors are $(a + b)$ and $(c + d)$.
 - The areas of the four small rectangles can be added together.
 $A = ac + ad + bc + bd$
 So $(a + b)(c + d) = ac + ad + bc + bd$
- There are several methods that can be helpful when remembering how to expand binomial factors. One commonly used method is **FOIL**. Each of the letters in FOIL stand for:
 - F**irst — multiply the first term in each bracket.
 - O**uter — multiply the two outer terms of each bracket.
 - I**nner — multiply the two inner terms of each bracket.
 - L**ast — multiply the last term of each bracket.

FOIL – First Outer Inner Last

$$\begin{array}{c}
 \text{F} \quad \text{O} \\
 \text{I} \quad \text{L} \\
 (a + b)(c + d) \\
 (a + b)(c + d) = ac + ad + bc + bd
 \end{array}$$

Expanding binomial factors

Using FOIL, the expanded product of two binomial factors is given by:

$$(a + b)(c + d) = a \times c + a \times d + b \times c + b \times d = ac + ad + bc + bd$$

It is expected that the result will have 4 terms.

Note: It may be possible to simplify like terms after expanding.

WORKED EXAMPLE 14 Expanding and simplifying two binomial factors

Expand and simplify each of the following expressions.

a. $(x - 5)(x + 3)$

b. $(x + 2)(x + 3)$

c. $(2x + 2)(2x + 3)$

THINK

a. 1. Expand the brackets using FOIL.

2. Simplify the expression by collecting like terms.

b. 1. Expand the brackets using FOIL.

2. Simplify the expression by collecting like terms.

c. 1. Expand the brackets using FOIL.

2. Simplify the expression by collecting like terms.

WRITE

a.

$$(x - 5)(x + 3)$$

$$= x^2 + 3x - 5x - 15$$

$$= x^2 - 2x - 15$$

b.

$$(x + 2)(x + 3)$$

$$= x^2 + 3x + 2x + 6$$

$$= x^2 + 5x + 6$$

c.

$$(2x + 2)(2x + 3)$$

$$= 4x^2 + 6x + 4x + 6$$

$$= 4x^2 + 10x + 6$$

TI | THINK

a–c. In a new problem, on a Calculator page, press:

- MENU
- 3: Algebra 3
- 3: Expand 3.

Complete the entry lines as: expand $((x - 5) \times (x + 3))$
expand $((x + 2) \times (x + 3))$
expand $((2x + 2) \times (2x + 3))$

Press ENTER after each entry.
Note: Remember to include the multiplication sign between the brackets.

WRITE

a–c.



a. $(x - 5)(x + 3)$
 $= x^2 - 2x - 15$
b. $(x + 2)(x + 3)$
 $= x^2 + 5x + 6$
c. $(2x + 2)(2x + 3)$
 $= 4x^2 + 10x + 6$

CASIO | THINK

a–c. On the Main screen, press:


- Action
- Transformation
- Expand

Complete the entry lines as:
expand $((x - 5)(x + 3))$
expand $((x + 2)(x + 3))$
expand $((2x + 2)(2x + 3))$

Press EXE after each entry line.

WRITE




a–c.



a. $(x - 5)(x + 3) = x^2 - 2x - 15$
b. $(x + 2)(x + 3) = x^2 + 5x + 6$
c. $(2x + 2)(2x + 3) = 4x^2 + 10x + 6$

DISCUSSION

Explain why, when expanded, $(x + y)(2x + y)$ gives the same result as $(2x + y)(x + y)$.

-  **eWorkbook** Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)
-  **Video eLessons** Expanding brackets (eles-1888)
Expansion of binomial expressions (eles-1908)
-  **Interactivities** Individual pathway interactivity: Expanding brackets (int-4483)
Expanding binomial factors (int-6033)
Expanding brackets (int-6034)
Like terms (int-6035)

Exercise 2.5 Expanding brackets

2.5 Quick quiz **on**

2.5 Exercise

Individual pathways

PRACTISE

1, 2, 7, 9, 12, 15, 18, 22, 24, 26, 30, 33

CONSOLIDATE

3, 4, 6, 10, 13, 17, 20, 21, 25, 27, 31, 34

MASTER

5, 8, 11, 14, 16, 19, 23, 28, 29, 32, 35, 36

Fluency

1. **WE10** Use two different methods to calculate the value of the following expressions.

a. $8(10 - 2)$	b. $11(99 + 1)$	c. $-5(3 + 1)$	d. $7(100 - 1)$
----------------	-----------------	----------------	-----------------
2. **WE11** Expand the following expressions. For the first two examples, draw a diagram to represent the expression.

a. $3(x + 2)$	b. $4(x + 3)$	c. $4(x + 1)$	d. $7(x - 1)$
---------------	---------------	---------------	---------------
3. Expand the following expressions.

a. $-3(p - 2)$	b. $-(x - 1)$	c. $3(2b - 4)$	d. $8(3m - 2)$
----------------	---------------	----------------	----------------
4. Expand each of the following. For the first two examples, draw a diagram to represent the expression.

a. $x(x + 2)$	b. $a(a + 5)$	c. $x(4 + x)$	d. $m(7 - m)$
---------------	---------------	---------------	---------------
5. Expand each of the following.

a. $2x(y + 2)$	b. $-3y(x + 4)$	c. $-b(3 - a)$	d. $-6a(5 - 3a)$
----------------	-----------------	----------------	------------------
6. **WE12** Expand and simplify the following expressions by collecting like terms.

a. $2(p - 3) + 4$	b. $5(x - 5) + 8$	c. $-7(p + 2) - 3$
d. $-4(3p - 1) - 1$	e. $6x(x - 3) - 2x$	
7. Expand and simplify the following expressions by collecting like terms.

a. $2m(m + 5) - 3m$	b. $3x(p + 2) - 5$	c. $4y(y - 1) + 7$
d. $-4p(p - 2) + 5p$	e. $5(x - 2y) - 3y - x$	
8. Expand and simplify the following expressions by collecting like terms.

a. $2m(m - 5) + 2m - 4$	b. $-3p(p - 2q) + 4pq - 1$	c. $-7a(5 - 2b) + 5a - 4ab$
d. $4c(2d - 3c) - cd - 5c$	e. $6p + 3 - 4(2p + 5)$	

9. **WE13** Expand and simplify the following expressions.

- a. $2(x + 2y) + 3(2x - y)$ b. $4(2p + 3q) + 2(p - 2q)$ c. $7(2a + 3b) + 4(a + 2b)$
d. $5(3c + 4d) + 2(2c + d)$ e. $-4(m + 2n) + 3(2m - n)$

10. Expand and simplify the following expressions.

- a. $-3(2x + y) + 4(3x - 2y)$ b. $-2(3x + 2y) + 3(5x + 3y)$ c. $-5(4p + 2q) + 2(3p + q)$
d. $6(a - 2b) - 5(2a - 3b)$ e. $5(2x - y) - 2(3x - 2y)$

11. Expand and simplify the following expressions.

- a. $4(2p - 4q) - 3(p - 2q)$ b. $2(c - 3d) - 5(2c - 3d)$ c. $7(2x - 3y) - (x - 2y)$
d. $-5(p - 2q) - (2p - q)$ e. $-3(a - 2b) - (2a + 3b)$

12. Expand and simplify the following expressions.

- a. $a(b + 2) + b(a - 3)$ b. $x(y + 4) + y(x - 2)$ c. $c(d - 2) + c(d + 5)$
d. $p(q - 5) + p(q + 3)$ e. $3c(d - 2) + c(2d - 5)$

13. Expand and simplify the following expressions.

- a. $7a(b - 3) - b(2a + 3)$ b. $2m(n + 3) - m(2n + 1)$ c. $4c(d - 5) + 2c(d - 8)$
d. $3m(2m + 4) - 2(3m + 5)$ e. $5c(2d - 1) - (3c + cd)$

14. Expand and simplify the following expressions.

- a. $-3a(5a + b) + 2b(b - 3a)$
b. $-4c(2c - 6d) + d(3d - 2c)$
c. $6m(2m - 3) - (2m + 4)$
d. $7x(5 - x) + 6(x - 1)$
e. $-2y(5y - 1) - 4(2y + 3)$

15. **MC** Choose the expression that is the equivalent of $3(a + 2b) + 2(2a - b)$.

- A. $5a + 6b$ B. $7a + 4b$ C. $5(3a + b)$ D. $7a + 8b$ E. $5a + 2b$

16. **MC** Choose the expression that is the equivalent of $-3(x - 2y) - (x - 5y)$.

- A. $-4x + 11y$ B. $-4x - 11y$ C. $4x + 11y$ D. $4x + 7y$ E. $-4x - 7y$

17. **MC** Choose the expression that is the equivalent of $2m(n + 4) + m(3n - 2)$.

- A. $3m + 4n - 8$ B. $5mn + 4m$ C. $5mn + 10m$ D. $5mn + 6m$ E. $3mn + 4m$

18. **WE14** Expand and simplify each of the following expressions.

- a. $(a + 2)(a + 3)$ b. $(x + 4)(x + 3)$ c. $(y + 3)(y + 2)$ d. $(m + 4)(m + 5)$ e. $(b + 2)(b + 1)$

19. Expand and simplify each of the following expressions.

- a. $(p + 1)(p + 4)$ b. $(a - 2)(a + 3)$ c. $(x - 4)(x + 5)$ d. $(m + 3)(m - 4)$ e. $(y + 5)(y - 3)$

20. Expand and simplify each of the following expressions.

- a. $(y - 6)(y + 2)$ b. $(x - 3)(x + 1)$ c. $(x - 3)(x - 4)$ d. $(p - 2)(p - 3)$ e. $(x - 3)(x - 1)$

21. Expand and simplify each of the following expressions.

- a. $(2a + 3)(a + 2)$ b. $(c - 6)(4c - 7)$ c. $(7 - 2t)(5 - t)$ d. $(2 + 3t)(5 - 2t)$
e. $(7 - 5x)(2 - 3x)$ f. $(5x - 2)(5x - 2)$

22. Expand and simplify each of the following expressions.

- a. $(x + y)(z + 1)$ b. $(2x + y)(z + 4)$ c. $(3p + q)(r + 1)$
d. $(a + 2b)(a + b)$ e. $(2c + d)(c - 3d)$ f. $(x + y)(2x - 3y)$


23. Expand and simplify each of the following expressions.

- a. $(4p - 3q)(p + q)$ b. $(a + 2b)(b + c)$ c. $(3p - 2q)(1 - 3r)$
d. $(4x - y)(3x - y)$ e. $(p - q)(2p - r)$ f. $(5 - 2j)(3k - 1)$

24. **MC** Choose the expression that is the equivalent of $(4 - y)(7 + y)$.
A. $28 - y^2$ **B.** $28 - 3y + y^2$ **C.** $28 - 3y - y^2$ **D.** $11 - 2y$ **E.** $28 - 11y - y^2$
25. **MC** Choose the expression that is the equivalent of $(2p + 1)(p - 5)$.
A. $2p^2 - 5$ **B.** $2p^2 - 11p - 5$ **C.** $2p^2 - 9p - 5$ **D.** $2p^2 - 6p - 5$ **E.** $2p^2 - 4p - 5$

Understanding



26. Expand the following expressions using FOIL, then simplify.
a. $(x + 3)(x - 3)$ **b.** $(x + 5)(x - 5)$ **c.** $(x + 7)(x - 7)$
d. $(x - 1)(x + 1)$ **e.** $(x - 2)(x + 2)$ **f.** $(2x - 1)(2x + 1)$
27. Expand the following expressions using FOIL, then simplify.
a. $(x + 1)(x + 1)$ **b.** $(x + 2)(x + 2)$ **c.** $(x + 8)(x + 8)$
d. $(x - 3)(x - 3)$ **e.** $(x - 5)(x - 5)$ **f.** $(x - 9)(x - 9)$
28. Simplify the following expressions.
a. $2.1x(3x + 4.7y) - 3.1y(1.4x + y)$ **b.** $(2.1x - 3.2y)(2.1x + 3.2y)$
c. $(3.4x + 5.1y)^2$
29. For the box shown, calculate the following in expanded form:



- a.** the total surface area **b.** the volume.

Reasoning

30. For each of the following shapes:
i. write down the area in factor form
ii. expand and simplify the expression
iii. discuss any limitations on the value of x .



31. Show that the following is true.

$$(a - x)(a + x) - 2(a - x)(a - x) - 2x(a - x) \\ = -(a - x)^2$$

32. A series of incorrect expansions of $(x + 8)(x - 3)$ are shown below. For each of these incorrect expansions, explain the mistake that has been made.
a. $x^2 + 11x + 24$ **b.** $x^2 + 5x + 24$ **c.** $x^2 - 3x$

Problem solving

33. Three students gave the following incorrect answers when expanding $-5(3x - 20)$.

i. $-5x - 20$

ii. $-8x + 25$

iii. $15x - 100$

- a. Explain the errors made by each student.
b. Determine the correct answer.

34. In a test, a student expanded brackets and obtained the following answers. Identify and correct the student's errors and write the correct expansions.

a. $-2(a - 5) = -2a - 10$

b. $2b(3b - 1) = 6b^2 - 1$

c. $-2(c - 4) = 2c + 8$

35. Three students' attempts at expanding $(3x + 4)(2x + 5)$ are shown.

Student A

$$\begin{aligned}(3x+4)(2x+5) \\ = 3x \times 2x + 3x \times 5 + 4 \times 2x + 4 \times 5 \\ = 6x + 15x + 8x + 20 \\ = 29x + 20 \\ = 49x\end{aligned}$$

Student B

$$\begin{aligned}(3x+4)(2x+5) \\ = 3x \times 2x + 4 \times 2x + 4 \times 5 \\ = 6x^2 + 8x + 20\end{aligned}$$

Student C

$$\begin{aligned}(3x+4)(2x+5) \\ = 3x \times 2x + 3x \times 5 + 4 \times 2x + 4 \times 5 \\ = 6x^2 + 15x + 8x + 20 \\ = 6x^2 + 23x + 20\end{aligned}$$

- a. State which student's work is correct.
b. Copy each of the incorrect answers into your workbook and correct the mistakes in each one as though you were their teacher.
36. Rectangular floor mats have an area of $(x^2 + 2x - 15)$ cm².
- a. The length of a mat is $(x + 5)$ cm. Determine an expression for the width of this mat.
b. If the length of a mat is 70 cm, evaluate its width.
c. If the width of a mat is 1 m, evaluate its length.

LESSON

2.6 Difference of two squares and perfect squares

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- recognise and use the difference of two squares rule to quickly expand and simplify binomial products with the form $(a + b)(a - b)$
- recognise and use the perfect square rule to quickly expand and simplify binomial products with the form $(a \pm b)^2$.

2.6.1 Difference of two squares

eles-4598

- Consider the expansion of $(x + 4)(x - 4)$:

$$(x + 4)(x - 4) = x^2 - 4x + 4x - 16 = x^2 - 16$$

- Now consider the expansion of $(x - 6)(x + 6)$:

$$(x - 6)(x + 6) = x^2 + 6x - 6x - 36 = x^2 - 36$$

- In both cases the middle two terms cancel each other out, leaving two terms, both of which are perfect squares.
- The two terms that are left are the first value squared minus the second value squared. This is where the phrase **difference of two squares** originates.
- In both cases the binomial terms can be written in the form $(x + a)$ and $(x - a)$. If we can recognise expressions that have this form, we can use the pattern above to quickly expand those expressions. For example: $(x + 12)(x - 12) = x^2 - 12^2 = x^2 - 144$

Difference of two squares

The difference of two squares rule is used to quickly expand certain binomial products, as long as they are in the forms shown below:

- $(a + b)(a - b) = a^2 - b^2$
- $(a - b)(a + b) = a^2 - b^2$

Because the two binomial brackets are being multiplied, the order of the brackets does not affect the final result.

WORKED EXAMPLE 15 Expanding using difference of two squares

Use the difference of two squares rule to expand and simplify each of the following expressions.

a. $(x + 8)(x - 8)$

b. $(6 - 3)(6 + x)$

c. $(2x - 3)(2x + 3)$

d. $(3x + 5)(5 - 3x)$

THINK

- a. 1. Write the expression.
2. This expression is in the form $(a + b)(a - b)$, so the difference of two squares rule can be used. Expand using the formula.

WRITE

a. $(x + 8)(x - 8)$
 $= x^2 - 8^2$
 $= x^2 - 64$

b. 1. Write the expression.

2. This expression is in the form $(a + b)(a - b)$, so the difference of two squares rule can be used. Expand using the formula.

Note: $36 - x^2$ is not the same as $x^2 - 36$.

c. 1. Write the expression.

2. This expression is in the form $(a - b)(a + b)$, so the difference of two squares rule can be used. Expand using the formula.

Note: $(2x)^2$ and $2x^2$ are not the same. In this case $a = 2x$, so $a^2 = (2x)^2$.

d. 1. Write the expression.

2. The difference of two squares rule can be used if we rearrange the terms, since $3x + 5 = 5 + 3x$. Expand using the formula.

$$\text{b. } (6 - x)(6 + x)$$

$$= 6^2 - x^2$$

$$= 36 - x^2$$

$$\text{c. } (2x - 3)(2x + 3)$$

$$= (2x)^2 - 3^2$$

$$= 4x^2 - 9$$

$$\text{d. } (3x + 5)(5 - 3x)$$

$$(5 + 3x)(5 - 3x)$$

$$= 5^2 - (3x)^2$$


$$= 25 - 9x^2$$

2.6.2 Perfect squares

eles-4599

- A **perfect square** is the result of the square of a whole number. $1 \times 1 = 1$, $2 \times 2 = 4$ and $3 \times 3 = 9$, showing that 1, 4 and 9 are all perfect squares.
- Similarly, $(x + 3)(x + 3) = (x + 3)^2$ is a perfect square because it is the result of a binomial factor multiplied by itself.
- Consider the diagram illustrating $(x + 3)^2$. What shape is it?
- The area is given by $x^2 + 3x + 3x + 9 = x^2 + 6x + 9$.
- We can see from the diagram that there are two squares produced (x^2 and $3^2 = 9$) and two rectangles that are identical to each other ($2 \times 3x$).
- Compare this with the expansion of $(x + 6)^2$:

$$(x + 6)(x + 6) = x^2 + 6x + 6x + 36 = x^2 + 12x + 36$$



- A pattern begins to emerge after comparing these two expansions. The square of a binomial equals the square of the first term, plus double the product of the two terms plus the square of the second term. For example: $(x + 10)(x + 10) = x^2 + 2 \times 10 \times x + 10^2 = x^2 + 20x + 100$

Perfect squares

The rule for the expansion of the square of a binomial is given by:

- $(a + b)^2 = (a + b)(a + b) = a^2 + 2ab + b^2$
- $(a - b)^2 = (a - b)(a - b) = a^2 - 2ab + b^2$

We can use the above rules to quickly expand binomial products that are presented as squares.

WORKED EXAMPLE 16 Expanding perfect squares

Use the rules for expanding perfect binomial squares to expand and simplify the following.

a. $(x + 1)(x + 1)$

b. $(x - 2)^2$

c. $(2x + 5)^2$

d. $(4x - 5y)^2$

THINK

a. 1. This expression is the square of a binomial.

2. Apply the formula for perfect squares:

$$(a + b)^2 = a^2 + 2ab + b^2$$

b. 1. This expression is the square of a binomial.

2. Apply the formula for perfect squares:

$$(a - b)^2 = a^2 - 2ab + b^2.$$

c. 1. This expression is the square of a binomial.

2. Apply the formula for perfect squares:

$$(a + b)^2 = a^2 + 2ab + b^2.$$

d. 1. This expression is the square of a binomial.

2. Apply the formula for perfect squares:

$$(a - b)^2 = a^2 - 2ab + b^2.$$

WRITE

a. $(x + 1)(x + 1)$

$$= x^2 + 2 \times x \times 1 + 1^2$$

$$= x^2 + 2x + 1$$

b. $(x - 2)^2$

$$= (x - 2)(x - 2)$$

$$= x^2 - 2 \times x \times 2 + 2^2$$

$$= x^2 - 4x + 4$$

c. $(2x + 5)^2$

$$= (2x)^2 + 2 \times 2x \times 5 + 5^2$$

$$= 4x^2 + 20x + 25$$

d. $(4x - 5y)^2$

$$= (4x)^2 - 2 \times 4x \times 5y + (5y)^2$$

$$= 16x^2 - 40xy + 25y^2$$

DISCUSSION

How could you represent $(x - 3)^2$ on a diagram?

on Resources



eWorkbook Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)



Interactivities Individual pathway interactivity: Difference of two squares and perfect squares (int-4484)

Difference of two squares (int-6036)

2.6 Quick quiz

2.6 Exercise

Individual pathways

■ PRACTISE

1, 3, 5, 8, 13, 16, 20, 21

■ CONSOLIDATE

2, 6, 9, 11, 14, 17, 22, 23

■ MASTER

4, 7, 10, 12, 15, 18, 19, 24, 25

Fluency

- WE15** Use the difference of two squares rule to expand and simplify each of the following.

a. $(x+2)(x-2)$ b. $(y+3)(y-3)$ c. $(m+5)(m-5)$ d. $(a+7)(a-7)$
- Use the difference of two squares rule to expand and simplify each of the following.

a. $(x+6)(x-6)$ b. $(p-12)(p+12)$ c. $(a+10)(a-10)$ d. $(m-11)(m+11)$
- Use the difference of two squares rule to expand and simplify each of the following.

a. $(2x+3)(2x-3)$ b. $(3y-1)(3y+1)$ c. $(5d-2)(5d+2)$ d. $(7c+3)(7c-3)$ e. $(2+3p)(2-3p)$
- Use the difference of two squares rule to expand and simplify each of the following.

a. $(d-9x)(d+9x)$ b. $(5-12a)(5+12a)$ c. $(3x+10y)(3x-10y)$
 d. $(2b-5c)(2b+5c)$ e. $(10-2x)(2x+10)$
- WE16** Use the rule for the expansion of the square of a binomial to expand and simplify each of the following.

a. $(x+2)(x+2)$ b. $(a+3)(a+3)$ c. $(b+7)(b+7)$ d. $(c+9)(c+9)$
- Use the rule for the expansion of the square of a binomial to expand and simplify each of the following.

a. $(m+12)^2$ b. $(n+10)^2$ c. $(x-6)^2$ d. $(y-5)^2$
- Use the rule for the expansion of the square of a binomial to expand and simplify each of the following.

a. $(9-c)^2$ b. $(8+e)^2$ c. $2(x+y)^2$ d. $(u-v)^2$
- Use the rule for expanding perfect binomial squares rule to expand and simplify each of the following.

a. $(2a+3)^2$ b. $(3x+1)^2$ c. $(2m-5)^2$ d. $(4x-3)^2$
- Use the rule for expanding perfect binomial squares to expand and simplify each of the following.

a. $(5a-1)^2$ b. $(7p+4)^2$ c. $(9x+2)^2$ d. $(4c-6)^2$
- Use the rule for expanding perfect binomial squares to expand and simplify each of the following.

a. $(5+3p)^2$ b. $(2-5x)^2$ c. $(9x-4y)^2$ d. $(8x-3y)^2$

Understanding

- Use the difference of two squares rule to expand and simplify each of the following.

a. $(x+3)(x-3)$ b. $(2x+3)(2x-3)$ c. $(7x-4)(7x+4)$
 d. $(2x+7y)(2x-7y)$ e. $(x^2+y^2)(x^2-y^2)$
- Expand and simplify the following perfect squares.

a. $(4x+5)^2$ b. $(7x-3y)^2$ c. $(5x^2-2y)^2$
 d. $2(x-y)^2$ e. $\left(\frac{2}{x}+4x\right)^2$
- A square has a perimeter of $4x+12$. Calculate its area.


14. Francis has fenced off a square in her paddock for spring lambs. The area of the paddock is $(9x^2 + 6x + 1) \text{ m}^2$.
Using pattern recognition, determine the side length of the paddock in terms of x .
15. A square has an area of $x^2 + 18x + 81$. Determine an expression for the perimeter of this square.

Reasoning

16. Show that $a^2 - b^2 = (a + b)(a - b)$ is true for each of the following.
- a. $a = 5, b = 4$ b. $a = 9, b = 1$ c. $a = 2, b = 7$ d. $a = -10, b = -3$
17. Lin has a square bedroom. Her sister Tasneem has a room that is 1 m shorter in length than Lin's room, but 1 m wider.
- a. Show that Lin has the larger bedroom.
b. Determine how much bigger Lin's bedroom is than Tasneem's bedroom.
18. Expand each of the following pairs of expressions.
- a. i. $(x - 4)(x + 4)$ and $(4 - x)(4 + x)$
ii. $(x - 11)(x + 11)$ and $(11 - x)(11 + x)$
iii. $(2x - 9)(2x + 9)$ and $(9 - 2x)(9 + 2x)$
b. State what you notice about the answers to the pairs of expansions above.
c. Explain how this is possible.
19. Answer the following questions.
- a. Expand $(10k + 5)^2$.
b. Show that $(10k + 5)^2 = 100k(k + 1) + 25$.
c. Using part b, evaluate 25^2 and 85^2 .

Problem solving

20. A large square has been subdivided into 2 squares and 2 rectangles.
- a. Write formulas for the areas of these 4 pieces, using the dimensions a and b marked on the diagram.
b. Write an equation that states that the area of the large square is equal to the combined area of its 4 pieces. Do you recognise this equation?



21. Expand each of the following pairs of expressions.
- a. i. $(x - 3)^2$ and $(3 - x)^2$
ii. $(x - 15)^2$ and $(15 - x)^2$
iii. $(3x - 7)^2$ and $(7 - 3x)^2$
b. State what you notice about the answers to the pairs of expansions above.
c. Explain how this is possible.
22. Use the perfect squares rule to quickly evaluate the following.
- a. 27^2 b. 33^2 c. 39^2 d. 47^2
23. Allen is creating a square deck with a square pool installed in the middle of it. The side length of the deck is $(2x + 3) \text{ m}$ and the side length of the pool is $(x - 2) \text{ m}$. Evaluate the area of the decking around the pool.
24. Ram wants to create a rectangular garden that is 10 m longer than it is wide. Write an expression for the area of the garden in terms of x , where x is the average length of the two sides.

25. The expansion of perfect squares $(a + b)^2 = a^2 + 2ab + b^2$ and $(a - b)^2 = a^2 - 2ab + b^2$ can be used to simplify some arithmetic calculations. For example:

$$\begin{aligned} 97^2 &= (100 - 3)^2 \\ &= 100^2 - 2 \times 100 \times 3 + 3^2 \\ &= 9409 \end{aligned}$$

Use this method to evaluate the following.

- a. 103^2 b. 62^2 c. 997^2 d. 1012^2 e. 53^2 f. 98^2

LESSON

2.7 Further expansions

LEARNING INTENTION

At the end of this lesson you should be able to:

- expand multiple sets of brackets and simplify the result.

2.7.1 Expanding multiple sets of brackets


eles-4600

- When expanding expressions with more than two sets of brackets, expand all brackets first before simplifying the like terms.
- It is important to be careful with signs, particularly when subtracting one expression from another.

For example:

$$\begin{aligned} 3x(x + 6) - (x + 5)(x - 3) &= 3x^2 + 18x - (x^2 + 5x - 3x - 15) \\ &= 3x^2 + 18x - (x^2 + 2x - 15) \\ &= 3x^2 + 18x - x^2 - 2x + 15 \\ &= 2x^2 + 16x + 15 \end{aligned}$$

- When expanding this, since the entire expression $(x + 5)(x - 3)$ is being subtracted from $3x(x + 6)$, its expanded form needs to be kept inside its own set of brackets. This helps us to remember that, as each term is being subtracted, the sign of each term will switch when opening up the brackets.
- When expanding, it can help to treat the expression as though it has a -1 at the front, as shown.

$$\begin{aligned} -(x^2 + 2x - 15) &= -1(x^2 + 2x - 15) \\ &= -x^2 - 2x + 15 \end{aligned}$$


WORKED EXAMPLE 17 Expanding and simplifying multiple sets of brackets

Expand and simplify each of the following expressions.

a. $(x + 3)(x + 4) + 4(x - 2)$

THINK

a. 1. Expand each set of brackets.

2. Simplify by collecting like terms.

b. 1. Expand and simplify each pair of brackets.

Because the second expression is being subtracted, keep it in a separate set of brackets.

2. Subtract all of the second result from the

first result. Remember that $-(x^2 + x - 2) = -1(x^2 + x - 2)$. Simplify by collecting like terms.

b. $(x - 2)(x + 3) - (x - 1)(x + 2)$

WRITE

a. $(x + 3)(x + 4) + 4(x - 2)$

$$= x^2 + 4x + 3x + 12 + 4x - 8$$

$$= x^2 + 11x + 4$$

b. $(x - 2)(x + 3) - (x - 1)(x + 2)$

$$= x^2 + 3x - 2x - 6 - (x^2 + 2x - x - 2)$$

$$= x^2 + x - 6 - (x^2 + x - 2)$$

$$= x^2 + x - 6 - x^2 - x + 2$$

$$= -4$$

WORKED EXAMPLE 18 Further expanding with multiple sets of brackets

Consider the expression $(4x + 3)(4x + 3) - (2x - 5)(2x - 5)$.

a. Expand and simplify this expression.

b. Apply the difference of perfect squares rule to verify your answer.

THINK

a. 1. Expand each set of brackets.

2. Simplify by collecting like terms.

b. 1. Rewrite as the difference of two squares.

2. Apply the difference of two squares rule, where $a = (4x + 3)$ and $b = (2x - 5)$.

3 Simplify, then expand the new brackets.

4. Simplify by collecting like terms.

WRITE

a. $(4x + 3)(4x + 3) - (2x - 5)(2x - 5)$

$$= 16x^2 + 12x + 12x + 9 - (4x^2 - 10x - 10x + 25)$$

$$= 16x^2 + 24x + 9 - (4x^2 - 20x + 25)$$

$$= 16x^2 + 24x + 9 - 4x^2 + 20x - 25$$

$$= 12x^2 + 44x - 16$$

b. $(4x + 3)(4x + 3) - (2x - 5)(2x - 5)$

$$= (4x + 3)^2 - (2x - 5)^2$$

$$= ((4x + 3) + (2x - 5))((4x + 3) - (2x - 5))$$

$$= (6x - 2)(2x + 8)$$

$$= 12x^2 + 48x - 4x - 16$$

$$= 12x^2 + 44x - 16$$

DISCUSSION

On a diagram, how would you show $(m + 2)(m + 3) - (m + 2)(m + 4)$?



Exercise 2.7 Further expansions

2.7 Quick quiz **on**

2.7 Exercise

Individual pathways

PRACTISE

1, 4, 8, 11, 14

CONSOLIDATE

2, 5, 7, 9, 12, 15

MASTER

3, 6, 10, 13, 16

Fluency

1. **WE17** Expand and simplify each of the following expressions.

a. $(x + 3)(x + 5) + (x + 2)(x + 3)$

b. $(x + 4)(x + 2) + (x + 3)(x + 4)$

c. $(x + 5)(x + 4) + (x + 3)(x + 2)$

d. $(x + 1)(x + 3) + (x + 2)(x + 4)$

2. Expand and simplify each of the following expressions.

a. $(p - 3)(p + 5) + (p + 1)(p - 6)$

b. $(a + 4)(a - 2) + (a - 3)(a - 4)$

c. $(p - 2)(p + 2) + (p + 4)(p - 5)$

d. $(x - 4)(x + 4) + (x - 1)(x + 20)$

3. Expand and simplify each of the following expressions.

a. $(y - 1)(y + 3) + (y - 2)(y + 2)$

b. $(d + 7)(d + 1) + (d + 3)(d - 3)$

c. $(x + 2)(x + 3) + (x - 4)(x - 1)$

d. $(y + 6)(y - 1) + (y - 2)(y - 3)$

4. Expand and simplify each of the following expressions.

a. $(x + 2)^2 + (x - 5)(x - 3)$

b. $(y - 1)^2 + (y + 2)(y - 4)$

c. $(p + 2)(p + 7) + (p - 3)^2$

d. $(m - 6)(m - 1) + (m + 5)^2$

5. Expand and simplify each of the following expressions.

a. $(x + 3)(x + 5) - (x + 2)(x + 5)$

b. $(x + 5)(x + 2) - (x + 1)(x + 2)$

c. $(x + 3)(x + 2) - (x + 4)(x + 3)$

d. $(m - 2)(m + 3) - (m + 2)(m - 4)$

6. Expand and simplify each of the following expressions.

a. $(b + 4)(b - 6) - (b - 1)(b + 2)$

b. $(y - 2)(y - 5) - (y + 2)(y + 6)$

c. $(p - 1)(p + 4) - (p - 2)(p - 3)$

d. $(x + 7)(x + 2) - (x - 3)(x - 4)$

7. Expand and simplify each of the following expressions.

a. $(m + 3)^2 - (m + 4)(m - 2)$

b. $(a - 6)^2 - (a - 2)(a - 3)$

c. $(p - 3)(p + 1) - (p + 2)^2$

d. $(x + 5)(x - 4) - (x - 1)^2$

Understanding

8. **WE18** Consider the expression $(x + 3)(x + 3) - (x - 5)(x - 5)$.

a. Expand and simplify this expression.

b. Apply the difference of perfect squares rule to verify your answer.

9. Consider the expression $(4x - 5)(4x - 5) - (x + 2)(x + 2)$.
- Expand and simplify this expression.
 - Apply the difference of perfect squares rule to verify your answer.
10. Consider the expression $(3x - 2y)(3x - 2y) - (y - x)(y - x)$.
- Expand and simplify this expression.
 - Apply the difference of perfect squares rule to verify your answer.

Reasoning

11. Determine the value of x for which $(x + 3) + (x + 4)^2 = (x + 5)^2$ is true.
12. Show that $(p - 1)(p + 2) + (p - 3)(p + 1) = 2p^2 - p - 5$.
13. Show that $(x + 2)(x - 3) - (x + 1)^2 = -3x - 7$.

Problem solving

14. Answer the following questions.
- Show that $(a^2 + b^2)(c^2 + d^2) = (ac - bd)^2 + (ad + bc)^2$.
 - Using part **a**, write $(2^2 + 1^2)(3^2 + 4^2)$ as the sum of two squares and evaluate.
15. Answer the following questions.
- Expand $(x^2 + x - 1)^2$.
 - Show that $(x^2 + x - 1)^2 = (x - 1)x(x + 1)(x + 2) + 1$.
- c. i.** Evaluate $4 \times 3 \times 2 \times 1 + 1$.
- ii.** Determine the value of x if $4 \times 3 \times 2 \times 1 + 1 = (x - 1)x(x + 1)(x + 2) + 1$.
16. Answer the following questions.
- Expand $(a + b)(d + e)$.
 - Expand $(a + b + c)(d + e + f)$. Draw a diagram to illustrate your answer.

LESSON

2.8 The highest common factor

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- determine the highest common factor for a group of terms
- factorise expressions by dividing out the highest common factor from each term.



2.8.1 Factorising expressions

eles-4601

- Factorising** a number or term means writing it as the product of a pair of its factors. For example, $18ab$ could be factorised as $3a \times 6b$.
- Factorising an algebraic expression is the opposite process to expanding.
- $3(x + 6)$ is the product of 3 and $(x + 6)$ and is the factorised form of $3x + 18$.
- In this section we will start with the expanded form of an expression, for example $6x + 12$, and work to re-write it in factorised form, in this case $6(x + 2)$.

WORKED EXAMPLE 19 Factorising terms

Complete the following factorisations.

a. $15x = 5x \times$ _____

b. $15x^2 = 3x \times$ _____

c. $-18ab^2 = 6a \times$ _____

THINK

a. Divide $15x$ by $5x$.

b. Divide $15x^2$ by $3x$.

c. Divide $-18ab^2$ by $6a$.

WRITE

a. $\frac{15x}{5x} = 3$

So, $15x = 5x \times 3$.

b. $\frac{15x^2}{3x} = 5x$

So, $15x^2 = 3x \times 5x$.

c. $\frac{-18ab^2}{6a} = -3b^2$

So, $-18ab^2 = 6a \times (-3b^2)$.

- The **highest common factor (HCF)** of two or more numbers is the largest factor that divides into all of them. The highest common factor of 18 and 27 is 9, since it is the biggest number that divides evenly into both 18 and 27.
- We can determine the HCF of two or more algebraic terms by determining the highest common factor of the coefficients, as well as any pronumerals that are common to all terms. For example, the HCF of $30xyz$, $25x^2z$ and $15xz^2$ is $5xz$. It can help to write each term in expanded form so that you can determine all of the common pronumerals.

WORKED EXAMPLE 20 Determining the highest common factor of two terms

Determine the highest common factor for each of the following pairs of terms.

a. $25a^2b$ and $10ab$

b. $3xy$ and $-3xz$

THINK

a. 1. Write $25a^2b$ in expanded form.

2. Write $10ab$ in expanded form.

3. Write the HCF.

b. 1. Write $3xy$ in expanded form.

2. Write $-3xz$ in expanded form.

3. Write the HCF.

WRITE

a. $25a^2b = 5 \times 5 \times a \times a \times b$

$10ab = 2 \times 5 \times a \times b$

HCF = $5ab$

b. $3xy = 3 \times x \times y$

$-3xz = -1 \times 3 \times x \times z$

HCF = $3x$



eles-4602

2.8.2 Factorising expressions by determining the highest common factor

- To factorise an expression such as $15x^2yz + 12xy^2$, determine the highest common factor of both terms. Writing each term in expanded form can help identify what is common to both terms.

$$\begin{aligned}15x^2yz + 12xy^2 &= 3 \times 5 \times x \times x \times y \times z + 3 \times 4 \times x \times y \times y \\ &= 3 \times x \times y \times 5 \times x \times z + 3 \times x \times y \times 4 \times y\end{aligned}$$

- We can see that $3xy$ is common to both terms. To complete the factorisation, place $3xy$ on the outside of a set of brackets and simplify what is left inside the brackets.

$$= 3 \times x \times y(5 \times x \times z + 4 \times y)$$

$$= 3xy(5xz + 4y)$$

- By removing the HCF from both terms, we have factorised the expression.
- The answer can always be checked by expanding out the brackets and making sure it produces the original expression.

WORKED EXAMPLE 21 Factorising by first determining the HCF

Factorise each expression by first determining the HCF.

- $5x + 15y$
- $-14xy - 7y$
- $6x^2y + 9xy^2$

THINK

1. The HCF is 5.
 2. Divide each term by 5 to determine the binomial.
 3. Check the answer by expanding.
1. The HCF is $7y$ or $-7y$, but $-7y$ makes things a little simpler.
 2. Divide each term by $-7y$ to determine the binomial.
 3. Check the answer by expanding.
1. The HCF is $3xy$.
 2. Divide each term by $3xy$ to determine the binomial.
 3. Check the answer by expanding.

WRITE

$$\text{a. } 5x + 15y = 5(\quad)$$

$$= 5(x + 3y)$$

$$5(x + 3y) = 5x + 15y \text{ (correct)}$$

$$\text{b. } -14xy - 7y = -7y(\quad)$$

$$= -7y(2x + 1)$$

$$-7y(2x + 1) = -14xy - 7y \text{ (correct)}$$

$$\text{c. } 6x^2y + 9xy^2 = 3xy(\quad)$$

$$= 3xy(2x + 3y)$$

$$3xy(2x + 3y) = 6x^2y + 9xy^2 \text{ (correct)}$$

TI | THINK


a-c. In a new problem, on a Calculator page, press:

- MENU
- 3: Algebra 3
- 2: Factor 2.

Complete the entry lines as:
 factor(5x + 15y)
 factor(-14x × y - 7y)
 factor(6x² × y + 9x × y²)

Press ENTER after each entry.

WRITE



- a. $5x + 15y = 5(x + 3y)$
 b. $-14xy - 7y = -7y(2x + 1)$
 c. $6x^2y + 9xy^2 = 3xy(2x + 3y)$

CASIO | THINK


a-c. On the Main screen, press:

- Action
- Transformation
- Factor.

Complete the entry lines as:
 factor(5x + 15y)
 factor(-14x × y - 7y)
 factor(6x² × y + 9x × y²)

Press EXE after each entry.

WRITE



- a. $(5x + 15y) = 5(x + 3y)$
 b. $(-14xy - 7y) = -7y(2x + 1)$
 c. $(6x^2y + 9xy^2) = 3xy(2x + 3y)$

DISCUSSION

How do you find the factors of terms within algebraic expressions?

on Resources

- eWorkbook** Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)
- Video elesson** Factorisation (eles-1887)
- Interactivities** Individual pathway interactivity: The highest common factor (int-4486)
Highest common factor (int-6037)

Exercise 2.8 The highest common factor

learn on

2.8 Quick quiz

2.8 Exercise

Individual pathways

■ PRACTISE

1, 3, 7, 11, 13, 16, 21, 23, 26

■ CONSOLIDATE

2, 4, 6, 8, 10, 14, 17, 18, 22, 24, 27

■ MASTER

5, 9, 12, 15, 19, 20, 25, 28

Fluency

1. **WE19** Complete each of the following factorisations by writing in the missing factor.

a. $8a = 4 \times \underline{\hspace{2cm}}$

b. $8a = 2a \times \underline{\hspace{2cm}}$

c. $12x^2 = 4x \times \underline{\hspace{2cm}}$

d. $-12x^2 = 3x^2 \times \underline{\hspace{2cm}}$

e. $3x^2 = x \times \underline{\hspace{2cm}}$

f. $15a^2b = ab \times \underline{\hspace{2cm}}$

2. Complete each of the following factorisations by writing in the missing factor.
- a. $12x = -4 \times \underline{\hspace{2cm}}$ b. $10mn = 10n \times \underline{\hspace{2cm}}$ c. $10mn = -10 \times \underline{\hspace{2cm}}$
d. $a^2b^2 = ab \times \underline{\hspace{2cm}}$ e. $30x^2 = 10x \times \underline{\hspace{2cm}}$ f. $-15mn^2 = -3m \times \underline{\hspace{2cm}}$
3. **WE20** Determine the highest common factor (HCF) of each of the following.
- a. 4 and 12 b. 6 and 15 c. 10 and 25 d. 24 and 32 e. 12, 15 and 21
4. Determine the highest common factor (HCF) of each of the following.
- a. 25, 50 and 200 b. 17 and 23 c. $6a$ and $12ab$
d. $14xy$ and $21xz$ e. $60pq$ and $30q$
5. Determine the highest common factor (HCF) of each of the following.
- a. $50cde$ and $70fgh$ b. $6x^2$ and $15x$ c. $6a$ and $9c$
d. $5ab$ and 25 e. $3x^2y$ and $4x^2z$
6. **MC** Choose which of the following pairs has a highest common factor of $5m$.
- A. $2m$ and $5m$ B. $5m$ and m C. $25mn$ and $15lm$
D. $20m$ and $40m$ E. $5m$ and lm
7. **WE21** Factorise each of the following expressions.
- a. $4x + 12y$ b. $5m + 15n$ c. $7a + 14b$ d. $7m - 21n$ e. $-8a - 24b$ f. $8x - 4y$
8. Factorise each of the following expressions.
- a. $-12p - 2q$ b. $6p + 12pq + 18q$ c. $32x + 8y + 16z$
d. $16m - 4n + 24p$ e. $72x - 8y + 64pq$ f. $15x^2 - 3y$
9. Factorise each of the following expressions.
- a. $5p^2 - 20q$ b. $5x + 5$ c. $56q + 8p^2$
d. $7p - 42x^2y$ e. $16p^2 + 20q + 4$ f. $12 + 36a^2b - 24b^2$
10. Factorise each of the following expressions.
- a. $9a + 21b$ b. $4c + 18d^2$ c. $12p^2 + 20q^2$ d. $35 - 14m^2n$ e. $25y^2 - 15x$
11. Factorise each of the following expressions.
- a. $16a^2 + 20b$ b. $42m^2 + 12n$ c. $63p^2 + 81 - 27y$
d. $121a^2 - 55b + 110c$ e. $10 - 22x^2y^3 + 14xy$
12. Factorise each of the following expressions.
- a. $18a^2bc - 27ab - 90c$ b. $144p + 36q^2 - 84pq$ c. $63a^2b^2 - 49 + 56ab^2$
d. $22 + 99p^3q^2 - 44p^2r$ e. $36 - 24ab^2 + 18b^2c$
13. Factorise each of the the following expressions.
- a. $-x + 5$ b. $-a + 7$ c. $-b + 9$ d. $-2m - 6$ e. $-6p - 12$ f. $-4a - 8$
14. Factorise each of the following expressions.
- a. $-3n^2 + 15m$ b. $-7x^2y^2 + 21$ c. $-7y^2 - 49z$
d. $-12p^2 - 18q$ e. $-63m + 56$ f. $-12m^3 - 50x^3$
15. Factorise each of the following expressions.
- a. $-9a^2b + 30$ b. $-15p - 12q$ c. $-18x^2 + 4y^2$
d. $-3ab + 18m - 21$ e. $-10 - 25p^2 - 45q$ f. $-90m^2 + 27n + 54p^3$
16. Factorise each of the following expressions.
- a. $a^2 + 5a$ b. $14q - q^2$ c. $18m + 5m^2$
d. $6p + 7p^2$ e. $7n^2 - 2n$ f. $7p - p^2q + pq$

17. Factorise each of the following expressions.

a. $xy + 9y - 3y^2$

b. $5c + 3c^2d - cd$

c. $3ab + a^2b + 4ab^2$

d. $2x^2y + xy + 5xy^2$

e. $5p^2q^2 - 4pq + 3p^2q$

f. $6x^2y^2 - 5xy + x^2y$

18. Factorise each of the following expressions.

a. $5x^2 + 15x$

b. $24m^2 - 6m$

c. $32a^2 - 4a$

d. $-2m^2 + 8m$

e. $-5x^2 + 25x$

f. $-7y^2 + 14y$

19. Factorise each of the following expressions.

a. $-3a^2 + 9a$

b. $-12p^2 - 2p$

c. $-26y^2 - 13y$


d. $4m - 18m^2$

e. $-6t + 36t^2$

f. $-8p - 24p^2$

Understanding

20. A large billboard display is in the shape of a rectangle as shown. The billboard has 3 regions (A, B, C) with dimensions in terms of x , as shown.



a. Calculate the total area of the billboard. Give your answer in factorised form.

b. Determine an expression for the area of each region of the billboard. Write the expression in its simplest form.

21. Consider the expression $3ab^2 + 18ab + 27a$.

a. Factorise this expression by taking out the highest common factor from each term.

b. Use the rule for perfect squares to fully factorise this expression.

22. Consider the expression $x^2(x+4) + 8x(x+4) + 16(x+4)$.

a. Factorise this expression by taking out the highest common factor from each term.

b. Use the rule for perfect squares to fully factorise this expression.

Reasoning

23. A question on Marcia's recent Algebra test was, 'Using factorisation, simplify the following expression: $a^2(a-b) - b^2(a-b)$ '. Marcia's answer was $(a+b)(a-b)^2$. If Marcia used the difference of two squares rule to get her solution, explain the steps she took to get that answer.

24. Prove that, as long as $a \neq 0$, then $(x+a)^2 \neq x^2 + a^2$.

25. Using the fact that any positive number, n , can be written as $(\sqrt{n})^2$, factorise the following expressions using the difference of two squares rule.

a. $x^2 - 13$

b. $4x^2 - 17$

c. $(x+3)^2 - 10$

Problem solving

26. Evaluate $\frac{x^2 + 2xy + y^2}{4x^2 - 4y^2} \div \frac{x^2 + xy}{8xy - 8x^2}$.

27. It has been said that, for any 2 numbers, the product of their LCM and HCF is the same as the product of the 2 numbers themselves. Show whether this is true.

28. a. Factorise $36x^2 - 100y^2$ by first taking out the common factor and then using the difference of two squares rule.

b. Factorise $36x^2 - 100y^2$ by first using the difference of two squares rule and then taking out the common factor.

c. Comment on whether you got the same answer for parts a and b.

LESSON

2.9 The highest common binomial factor

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- factorise expressions by taking out a common binomial factor
- factorise expressions by grouping terms.

2.9.1 The common binomial factor

eles-4603

- When factorising an expression, we look for the highest common factor(s) first.
- It is possible for the HCF to be a binomial expression.
- Consider the expression $7(a - b) + 8x(a - b)$. The binomial expression $(a - b)$ is a common factor to both terms. Factorising this expression looks like this:

$$\begin{aligned}7(a - b) + 8x(a - b) &= 7 \times (a - b) + 8x \times (a - b) \\ &= (a - b)(7 + 8x)\end{aligned}$$

WORKED EXAMPLE 22 Factorising by finding a binomial common factor

Factorise each of the following expressions.

a. $5(x + y) + 6b(x + y)$

b. $2b(a - 3b) - (a - 3b)$

Note: In both of these expressions the HCF is a binomial factor.

THINK

a. 1. The HCF is $(x + y)$.

2. Divide each term by $(x + y)$ to determine the binomial.

b. 1. The HCF is $(a - 3b)$.

2. Divide each term by $(a - 3b)$ to determine the binomial.

WRITE

a. $\frac{5(x + y)}{x + y} = 5, \frac{6b(x + y)}{x + y} = 6b$

Therefore,

$$\begin{aligned}5(x + y) + 6b(x + y) \\ = (x + y)(5 + 6b)\end{aligned}$$

b. $\frac{2b(a - 3b)}{a - 3b} = 2b, \frac{-1(a - 3b)}{a - 3b} = -1$

Therefore,

$$\begin{aligned}2b(a - 3b) - (a - 3b) \\ = 2b(a - 3b) - 1(a - 3b) \\ = (a - 3b)(2b - 1)\end{aligned}$$

2.9.2 Factorising by grouping in pairs

eles-4604

- If an algebraic expression has four terms and no common factors in any of its terms, it may be possible to group the terms in pairs and find a common factor in each pair.
- Consider the expression $10x + 15 - 6ax - 9a$.
- We can attempt to factorise by grouping the first two terms and the last two terms:

$$\begin{aligned}10x + 15 - 6ax - 9a &= 5 \times 2x + 5 \times 3 - 3a \times 2x - 3a \times 3 \\ &= 5(2x + 3) - 3a(2x + 3)\end{aligned}$$

- Once a common factor has been taken out from each pair of terms, a common binomial factor will appear. This common binomial factor can also be factorised out.

$$5(2x + 3) - 3a(2x + 3) = (2x + 3)(5 - 3a)$$

- Thus the expression $10x + 15 - 6ax - 9a$ can be factorised to become $(2x + 3)(5 - 3a)$.
- It is worth noting that it doesn't matter which terms are paired up first — the final result will still be the same.

$$\begin{aligned} 10x + 15 - 6ax - 9a &= 10x - 6ax + 15 - 9a \\ &= 2x \times 5 + 2x \times (-3a) + 3 \times 5 + 3 \times (-3a) \\ &= 2x(5 - 3a) + 3(5 - 3a) \\ &= (5 - 3a)(2x + 3) \\ &= (2x + 3)(5 - 3a) \end{aligned}$$

WORKED EXAMPLE 23 Factorising by grouping in pairs

Factorise each of the following expressions by grouping the terms in pairs.

a. $5a + 10b + ac + 2bc$

b. $x - 3y + ax - 3ay$

c. $5p + 6q + 15pq + 2$

THINK

a. 1. Write the expression.

2. Take out the common factor $a + 2b$.

b. 1. Write the expression.

2. Take out the common factor $x - 3y$.

c. 1. Write the expression.

2. There are no simple common factors. Write the terms in a different order.

3. Take out the common factor $1 + 3q$.
Note: $1 + 3q = 3q + 1$.

WRITE

a. $5a + 10b + ac + 2bc$

$$5a + 10b = 5(a + 2b)$$

$$ac + 2bc = c(a + 2b)$$

$$= 5(a + 2b) + c(a + 2b)$$

$$= (a + 2b)(5 + c)$$

b. $x - 3y + ax - 3ay$

$$x - 3y = 1(x - 3y)$$

$$ax - 3ay = a(x - 3y)$$

$$= 1(x - 3y) + a(x - 3y)$$

$$= (x - 3y)(1 + a)$$

c. $5p + 6q + 15pq + 2$

$$= 5p + 15pq + 6q + 2$$

$$5p + 15pq = 5p(1 + 3q)$$

$$6q + 2 = 2(3q + 1)$$

$$= 5p(1 + 3q) + 2(3q + 1)$$

$$= 5p(1 + 3q) + 2(1 + 3q)$$

$$= (1 + 3q)(5p + 2)$$

DISCUSSION

How do you factorise expressions with four terms?



eWorkbook Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)



Interactivities Individual pathway interactivity: The highest common binomial factor (int-4487)
Common binomial factor (int-6038)

Exercise 2.9 The highest common binomial factor

2.9 Quick quiz



2.9 Exercise

Individual pathways

PRACTISE

1, 3, 6, 9, 12

CONSOLIDATE

2, 4, 7, 10, 13

MASTER

5, 8, 11, 14

Fluency

- WE22** Factorise each of the following expressions.

a. $2(a + b) + 3c(a + b)$ b. $4(m + n) + p(m + n)$ c. $7x(2m + 1) - y(2m + 1)$
 d. $4a(3b + 2) - b(3b + 2)$ e. $z(x + 2y) - 3(x + 2y)$
- Factorise each of the following expressions.

a. $12p(6 - q) - 5(6 - q)$ b. $3p^2(x - y) + 2q(x - y)$ c. $4a^2(b - 3) + 3b(b - 3)$
 d. $p^2(q + 2p) - 5(q + 2p)$ e. $6(5m + 1) + n^2(5m + 1)$
- WE23** Factorise each of the following expressions by grouping the terms in pairs.

a. $xy + 2x + 2y + 4$ b. $ab + 3a + 3b + 9$ c. $xy - 4y + 3x - 12$
 d. $2xy + x + 6y + 3$ e. $3ab + a + 12b + 4$ f. $ab - 2a + 5b - 10$
- Factorise each of the following expressions by grouping the terms in pairs.

a. $m - 2n + am - 2an$ b. $5 + 3p + 15a + 9ap$ c. $15mn - 5n - 6m + 2$
 d. $10pq - q - 20p + 2$ e. $6x - 2 - 3xy + y$ f. $16p - 4 - 12pq + 3q$
- Factorise each of the following expressions by grouping the terms in pairs.

a. $10xy + 5x - 4y - 2$ b. $6ab + 9b - 4a - 6$ c. $5ab - 10ac - 3b + 6c$
 d. $4x + 12y - xz - 3yz$ e. $5pr + 10qr - 3p - 6q$ f. $ac - 5bc - 2a + 10b$

Understanding

- Simplify the following expressions using factorising.


a. $\frac{ax + 2ay + 3az}{bx + 2by + 3bz}$ b. $\frac{10(3x - 4) + 2y(3x - 4)}{7a(10 + 2y) - 5(10 + 2y)}$
- Use factorising by grouping in pairs to simplify the following expressions.

a. $\frac{3x + 6 + xy + 2y}{6 + 2y + 18x + 6xy}$ b. $\frac{5xy + 10x + 3ay + 6a}{15bx - 10x + 9ab - 6a}$
- Use factorising by grouping in pairs to simplify the following expressions.

a. $\frac{6x^2 + 15xy - 4x - 10y}{6xy + 4x + 15y^2 + 10y}$ b. $\frac{mp + 4mq - 4np - 16nq}{mp + 4mq + 4np + 16nq}$

Reasoning


9. Using the method of rectangles to expand, show how $a(m + n) + 3(m + n)$ equals $(a + 3)(m + n)$.
10. Fully factorise $6x + 4x^2 + 6x + 9$ by grouping in pairs. Discuss what you noticed about this factorisation.
11. a. Write out the product $5(x + 2)(x + 3)$ and show that it also corresponds to the diagram shown.



- b. Explain why $5(x + 2)(x + 3)$ is equivalent to $(5x + 10)(x + 3)$. Use bracket expansion and a labelled diagram to support your answer.
- c. Explain why $5(x + 2)(x + 3)$ is equivalent to $(x + 2)(5x + 15)$. Use bracket expansion and a labelled diagram to support your answer.

Problem solving

12. A series of five squares of increasing size and a list of five area formulas are shown.
- Use factorisation to calculate the side length that correlates to each area formula.
 - Using the area given and the side lengths found, match the squares below with the appropriate algebraic expression of their area.



$$A_a = x^2 + 6x + 9$$

$$A_b = x^2 + 10x + 25$$

$$A_c = x^2 + 16x + 64$$

$$A_d = x^2 - 6x + 9$$

$$A_e = x^2 + 12x + 36$$

- c. If $x = 5$ cm, use the formula given to calculate the area of each square.
13. Fully factorise both sets and brackets in the expression $(-12xy + 27x + 8y - 18) - (-8xy + 18x + 12y - 27)$ using grouping in pairs, then factorise the result.
14. The area formulas shown relate to either squares or rectangles.
- $9s^2 + 48s + 64$
 - $25s^2 - 4$
 - $s^2 + 4s + 3$
 - $4s^2 - 28s - 32$
- Without completing any algebraic operations, examine these formulas and work out which ones belong to squares and which ones belong to rectangles. Explain your answer.
 - Factorise each formula and classify it as a square or rectangle. Check your classifications against your answer to part a.

LESSON

2.10 Solving worded problems

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- write an algebraic expression for worded problems
- apply the skills covered in this topic to solve worded problems.

2.10.1 Converting worded problems into algebraic expressions

eles-4605

- It is important to be able to convert worded problems (or 'real-world problems') into algebraic expressions. The key words that were identified in lesson 2.3 can help with this process.
- Drawing diagrams is a useful strategy for understanding and solving worded problems.
- If a worded problem uses units in its questions, make sure that any answers also use those units.

WORKED EXAMPLE 24 Converting worded problems

A rectangular swimming pool measures 30 m by 20 m.
A path around the edge of the pool is x m wide on each side.




- Determine the area of the pool.
- Write an expression for the area of the pool plus the area of the path.
- Write an expression for the area of the path.
- If the path is 1.5 m wide, calculate the area of the path.

THINK

1. Construct a drawing of the pool.

WRITE/DRAW



2. Calculate the area of the pool.

$$\begin{aligned}\text{Area} &= \text{length} \times \text{width} \\ &= 20 \times 30 \\ &= 600 \text{ m}^2\end{aligned}$$

1. Write an expression for the total length from one edge of the path to the other. Write another expression for the total width from one edge of the path to the other.

$$\begin{aligned}\text{Length} &= 30 + x + x \\ &= 30 + 2x \\ \text{Width} &= 20 + x + x \\ &= 20 + 2x\end{aligned}$$

2. Area = length \times width

$$\begin{aligned}\text{Area} &= \text{length} \times \text{width} \\ &= (30 + 2x)(20 + 2x) \\ &= 600 + 60x + 40x + 4x^2 \\ &= (600 + 100x + 4x^2) \text{ m}^2\end{aligned}$$

c. Determine an expression for the area of the path by subtracting the area of the pool from the total area of the pool and the path combined.

$$\begin{aligned} \text{c. Area of path} &= \text{total area} - \text{area of pool} \\ &= 600 + 100x + 4x^2 - 600 \\ &= (100x + 4x^2) \text{ m}^2 \end{aligned}$$

d. Substitute 1.5 for x in the expression you have found for the area of the path.


$$\begin{aligned} \text{d. When } x = 1.5, \\ \text{Area of path} &= 100(1.5) + 4(1.5)^2 \\ &= 159 \text{ m}^2 \end{aligned}$$

- The algebraic expression found in part c of Worked example 24 allows us to calculate the area of the path for any given width.

WORKED EXAMPLE 25 Writing expressions for worded problems

Suppose that the page of a typical textbook is 24 cm high by 16 cm wide. Each page has margins of x cm at the top and bottom, and margins of y cm on the left and right.


- Write an expression for the height of the section of the page that lies inside the margins.
- Write an expression for the width of the section of the page that lies inside the margins.
- Write an expression for the area of the section of the page that lies inside the margins.
- Show that, if the margins on the left and right are doubled, the area within the margins is reduced by $(48y - 4xy)$ cm.
- If the margins at the top and the bottom of the page are 1.5 cm and the margins on the left and right of the page are 1 cm, calculate the size of the area that lies within the margins.



THINK

- Construct a drawing that shows the key dimensions of the page and its margins.

WRITE/DRAW



- The total height of the page (24 cm) is effectively reduced by x cm at the top and x cm at the bottom.

$$\begin{aligned} \text{Height} &= 24 - x - x \\ &= (24 - 2x) \text{ cm} \end{aligned}$$

b. The total width of the page (16 cm) is reduced by y cm on the left and y cm on the right.

$$\begin{aligned}\text{b. Width} &= 16 - y - y \\ &= (16 - 2y) \text{ cm}\end{aligned}$$

c. The area of the page that lies within the margins is the product of the width and height.

$$\begin{aligned}\text{c. Area}_1 &= (24 - 2x)(16 - 2y) \\ &= (384 - 48y - 32x + 4xy) \text{ cm}^2\end{aligned}$$

d. 1. If the left and right margins are doubled they both become $2y$ cm. Determine the new expression for the new width of the page.

$$\begin{aligned}\text{d. Width} &= 16 - 2y - 2y \\ &= (16 - 4y) \text{ cm}\end{aligned}$$

2. Determine the new expression for the reduced area.

$$\begin{aligned}\text{Area}_2 &= (24 - 2x)(16 - 4y) \\ &= 24 \times 16 + 24 \times -4y - 2x \times 16 - 2x \times -2y \\ &= (384 - 96y - 32x + 8xy) \text{ cm}^2\end{aligned}$$

3. Determine the difference in area by subtracting the reduced area obtained in part d from the original area obtained in part c.

$$\begin{aligned}\text{Difference in area} &= \text{Area}_1 - \text{Area}_2 \\ &= 384 - 48y - 32x + 4xy \\ &\quad - (384 - 96y - 32x + 8xy) \\ &= 384 - 48y - 32x + 4xy \\ &\quad - 384 + 96y + 32x - 8xy \\ &= 48y - 4xy\end{aligned}$$

So the amount by which the area is reduced is $(48y - 4xy) \text{ cm}^2$.

e. Using the area found in c, substitute 1.5 for x and 1 for y . Solve the expression.

$$\begin{aligned}\text{e. Area} &= (384 - 48y - 32x + 4xy) \text{ cm}^2 \\ &= 384 - 48(1) - 32(1.5) + 4(1.5)(1) \\ &= 294 \text{ cm}^2\end{aligned}$$

DISCUSSION

How can you use algebraic skills in real-life situations?

Resources



eWorkbook Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)



Interactivity Individual pathway interactivity: Solving worded problems (int-4488)

Individual pathways

PRACTISE

1, 2, 6, 9, 12

CONSOLIDATE







3, 5, 7, 10, 13

MASTER

4, 8, 11, 14, 15

Fluency

1. Answer the following for each shape shown.
 - i. Determine an expression for the perimeter.
 - ii. Determine the perimeter when $x = 5$.
 - iii. Determine an expression for the area. If necessary, simplify the expression by expanding.
 - iv. Determine the area when $x = 5$.




Understanding

2. **WE24** A rectangular swimming pool measures 50 m by 25 m. A path around the edge of the pool is x m wide on each side.
 - a. Determine the area of the pool.
 - b. Write an expression for the area of the pool plus the area of the path.
 - c. Write an expression for the area of the path.
 - d. If the path is 2.3 m wide, calculate the area of the path.
 - e. If the area of the path is 200 m^2 , write an equation that can be solved to calculate the width of the path.



3. **WE25** The pages of a book are 20 cm high by 15 cm wide. The pages have margins of x cm at the top and bottom and margins of y cm on the left and right.
 - a. Write an expression for the height of the section of the pages that lie inside the margins.
 - b. Write an expression for the width of the section of the pages that lie inside the margins.
 - c. Write an expression for the area of the section of the pages that lie inside the margins.
 - d. Show that, if the margins on the left and right are doubled, the area between the margins is reduced by $(40y - 4xy)$ cm.




4. A rectangular book cover is 8 cm long and 5 cm wide.
- Calculate the area of the book cover.
 - If the length of the book cover is increased by v cm, write an expression for its new length.
 - If the width of the book cover is increased by v cm, write an expression for its new width.
 - Write an expression for the new area of the book cover. Expand this expression.
 - Calculate the increased area of the book cover if $v = 2$ cm.
 - If the length of the book cover is decreased by d cm, write an expression for its new length.
 - If the width of the book cover is decreased by d cm, write an expression for its new width.
 - Write an expression for the new area of the book cover. Expand this expression.
 - Calculate the new area of the book cover if $d = 2$ cm.
 - If the length of the book cover is made x times longer, write an expression for its new length.
 - If the width of the book cover is increased by x cm, write an expression for its new width.
 - Write an expression for the new area of the book cover. Expand this expression.
 - Calculate the new area of the book cover if $x = 5$ cm.
5. A square has sides of length $5x$ m.
- Write an expression for its perimeter.
 - Write an expression for its area.
 - If the square's length is decreased by 2 m, write an expression for its new length.
 - If the square's width is decreased by 3 m, write an expression for its new width.
 - Write an expression for the square's new area. Expand this expression.
 - Calculate the square's area when $x = 6$ m.
6. A rectangular sign has a length of $2x$ cm and a width of x cm.
- Write an expression for the sign's perimeter.
 - Write an expression for the sign's area.
 - If the sign's length is increased by y cm, write an expression for its new length.
 - If the sign's width is decreased by y cm, write an expression for its new width.
 - Write an expression for the sign's new area and expand.
 - Calculate the sign's area when $x = 4$ cm and $y = 3$ cm using your expression.
7. A square has a side length of x cm.
- Write an expression for its perimeter.
 - Write an expression for its area.
 - If the square's side length is increased by y cm, write an expression for its new side length.
 - Write an expression for the square's new perimeter. Expand this expression.
 - Calculate the square's perimeter when $x = 5$ cm and $y = 9$ cm.
 - Write an expression for the square's new area and expand.
 - Calculate the square's area when $x = 3.2$ cm and $y = 4.6$ cm.

8. A swimming pool with length $(4p + 2)$ m and width $3p$ m is surrounded by a path of width p m.


Write the following in expanded form.

- An expression for the perimeter of the pool.
- An expression for the area of the pool.
- An expression for the length of the pool and path.
- An expression for the width of the pool and path.
- An expression for the perimeter of the pool and path.
- An expression for the area of the pool and path.
- An expression for the area of the path.
- The area of the path when $p = 2$ m.




Reasoning

9. The Body Mass Index (B) is used as an indicator of whether or not a person is in a healthy weight range for their height. It can be calculated using the formula $B = \frac{m}{h^2}$, where m is the person's mass in kilograms and h is the person's height in metres. Use this formula to answer the following questions, correct to 1 decimal place.
- Calculate Samir's Body Mass Index if they weigh 85 kg and are 1.75 m tall.
 - A person is considered to be in a healthy weight range if their Body Mass Index is between 21 to 25 inclusive. Comment on Samir's weight for a person of their height.
 - Calculate the Body Mass Index, correct to 1 decimal place, for each of the following people:
 - Sammara, who is 1.65 m tall and has a mass of 52 kg
 - Nimco, who is 1.78 m tall and has a mass of 79 kg
 - Manuel, who is 1.72 m tall and has a mass of 65 kg.
 - The healthy Body Mass Index range for children changes up until adulthood. These graphs show the Body Mass Index for boys and girls aged from 2 to 20 years of age. Study these graphs carefully and decide on possible age ranges (between 2 and 20) for Sammara, Nimco and Manuel if their BMI was in the satisfactory range. Give the age ranges in half-year intervals.




10. The figure shown is a rectangle. For the purpose of this activity we are going to call this rectangle an 'algebra rectangle'.




The length of each algebra rectangle is x cm and the width is 1 cm.

These algebra rectangles are put together to form larger rectangles in one of two ways.

Long algebra rectangle (3-long)



Tall algebra rectangle (3-tall)



To calculate the perimeter of each of these algebra rectangles:

- perimeter of 3-long algebra rectangle $P = (x + x + x + x + x + x) + (1 + 1) = 6x + 2$
 - perimeter of 3-tall algebra rectangle $P = (x + x) + (1 + 1 + 1 + 1 + 1 + 1) = 2x + 6$.
- a. Using the expressions for the perimeters of a 3-long algebra rectangle and a 3-tall algebra rectangle as a basis, find the perimeter of each of the following algebra rectangles and add them to the table shown.

Type of algebra rectangle	Perimeter	Type of algebra rectangle	Perimeter
1-long		1-tall	
2-long		2-tall	
3-long	$6x + 2$	3-tall	$2x + 6$
4-long		4-tall	
5-long		5-tall	

- b. Can you see the pattern? Based on this pattern, calculate the perimeter of a 20-long algebra rectangle and a 20-tall algebra rectangle.
- c. Determine the values of x that will result in a tall algebra rectangle that has a larger perimeter than a long algebra rectangle.
11. Can you evaluate 997^2 without a calculator in less than 90 seconds? It is possible to work out the answer using long multiplication, but it would take a fair amount of time and effort. Mathematicians are always looking for quick and simple ways of solving problems.

What if we consider the expanding formula that produces the difference of two squares?

$$(a + b)(a - b) = a^2 - b^2$$

Adding b^2 to both sides gives $(a + b)(a - b) + b^2 = a^2 - b^2 + b^2$.


Simplifying and swapping sides gives $a^2 = (a + b)(a - b) + b^2$.

We can use this formula, combined with the fact that multiplying by 1000 is an easy operation, to evaluate 997^2 .


- a. If $a = 997^2$, determine what the value of b should be so that $(a + b)$ equals 1000.
- b. Substitute these a and b values into the formula to evaluate 997^2 .
- c. Try the above method to evaluate the following.
- 995^2
 - 990^2

Problem solving

12. In the picture of the phone shown, the dimensions are in cm.




- a. Use the information in the picture to write an expression for the area of:
- the viewing screen
 - the entire front face of the phone.
- b. Your friend has a phone that is 4 cm longer than the one shown, but also 1 cm narrower. Write expressions in expanded form for:
- the length and width of your friend's phone
 - the area of your friend's phone.
- c. If the phone in the picture is 5 cm wide, use your answer to part b to calculate the area of the front face of your friend's phone.
13. A proposed new flag for Australian schools will have the Australian flag in the top left-hand corner. The dimensions of this new flag are given in metres.



- a. Write an expression in factorised form for the area of:
- the Australian flag section of the proposed flag
 - the whole area of the proposed flag.
- b. Use the answers from part a to write the area of the Australian flag as a fraction of the school flag. Simplify this fraction.
- c. Use the fraction from part b to express the area of the Australian flag as a percentage of the proposed school flag.
- d. Use the formula for the percentage of the area taken up by the Australian flag to find the percentages for the following suggested widths for the proposed school flag.
- 4 m
 - 4.5 m
 - 4.8 m
- e. If the percentage of the school flag taken up by the Australian flag measures the importance a school places on Australia, determine what can be said about the three suggested flag widths.

14. A new game has been created by students for the school fair. To win the game you need to hit the shaded region of the target shown with 5 darts.




- Write an expression to calculate the area of the shaded region of the game board.
 - Calculate the area of the shaded region of the game board if $R = 7.5$ cm and $x = 4$ cm.
 - Show that $R = \sqrt{\frac{A + x^2}{\pi}}$ by transposing the formula found in part a.
 - If $A = 80 \text{ cm}^2$ and $x = 3$ cm, calculate the value of R .
 - The students found that the best size for the game board is when $R = 10$ and $x = 5$. For a board of this size, determine the percentage of the total board that is shaded.
15. Cubic expressions are the expanded form of expressions with three linear factors. The expansion process for three linear factors is called trinomial expansion.
- Explain how the area model can be altered to show that binomial expansion can also be altered to become a model for trinomial expansion.
 - Investigate the powers in cubic expressions by expanding the following expressions.
 - $(3x + 2)(-x + 1)(x + 1)$
 - $(x + 5)(2x - 2)(4x - 8)$
 - $(3 - x)(x + 8)(5 - x)$
 - Describe the patterns in the powers in a cubic expression.

LESSON

2.11 Review




2.11.1 Topic summary



2.11.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.


(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
2.2	I can identify the number of terms in an algebraic expression.			
	I can identify the variables, constant terms and coefficients of the terms in an expression.			
	I can evaluate an expression or formula by substituting values for each variable.			
2.3	I can assign pronumerals to the unknown quantities in a worded problem.			
	I can convert the worded statements presented in a question into algebraic expressions.			
2.4	I can identify like terms.			
	I can simplify algebraic expressions by adding and subtracting like terms.			
	I can simplify algebraic expressions that involve multiplication and division of multiple algebraic terms.			
2.5	I can use the distributive law to expand and simplify an expression with a single set of brackets.			
	I can use the distributive law to expand and simplify expressions containing two or more sets of brackets.			
2.6	I can quickly expand and simplify binomial products using the rule for difference of two squares.			
	I can quickly expand and simplify binomial products using the rule for perfect squares.			
2.7	I can expand multiple sets of brackets and simplify the result.			
2.8	I can determine the highest common factor for a group of terms.			
	I can factorise expressions by dividing out the highest common factor from each term.			
2.9	I can factorise an expression by taking out a common binomial factor			
	I can factorise expressions by grouping terms in pairs.			
2.10	I can write an algebraic expression for a worded problem and use it to solve that problem.			

2.11.3 Project

Quilt squares


People all over the world are interested in quilt making, which involves stitching pieces of fabric together and inserting stuffing between layers of stitched-together fabric. When making a quilt, the fabric can be arranged and sewn in a variety of ways to create attractive geometric designs. Because of the potential for interesting designs in quilt-making, quilts are often used as decorative objects.



Medini is designing a quilt. She is sewing pieces of differently coloured fabric together to make a block, then copying the block and sewing the blocks together in a repeated pattern.

Making your own quilt

A scaled diagram of the basic block that Medini is using to make her quilt is shown. The letters indicate the colours of the fabric that make up the block: yellow, black and white. The yellow and white pieces are square, while the black pieces are rectangular. The finished blocks are sewn together in rows and columns.





Trace or copy the basic block shown onto a sheet of paper. Repeat this process until you have 9 blocks. Colour in each section and cut out all 9 blocks.

1. Place all 9 blocks together to form a 3×3 square. Draw and colour a scaled diagram of your result.
2. Describe the feature created by arranging the blocks in the manner described in question 1. Observe the shapes created by the different colours.

Medini sold her design to a company that now manufactures quilts made from 100 of these blocks. Each quilt covers an area of 1.44 m^2 . Each row and column has the same number of blocks. Answer the following, ignoring any seam allowances.

3. Calculate the side length of each square block.
4. If the entire quilt has an area of 1.44 m^2 , what is the area of each block?
5. Determine the dimensions of the yellow, black and white pieces of fabric of each block.
6. Calculate the area of the yellow, black and white pieces of fabric of each block.
7. Determine the total area of each of the three different colours required to construct this quilt.
8. Due to popular demand, the company that manufactures these quilts now makes them in different sizes. A customer can specify the approximate quilt area, the three colours they want and the number of blocks in the quilt, but the quilt must be either square or rectangular. Come up with a general formula that would let the company quickly work out the areas of the three coloured fabrics in each block. Give an example of this formula. Draw a diagram on a separate sheet of paper to illustrate your formula.

Resources

-  **eWorkbook** Topic 2 Workbook (worksheets, code puzzle and project) (ewbk-2002)
-  **Interactivities** Crossword (int-0699)
Sudoku puzzle (int-3203)

Fluency

1. **MC** Select the coefficient of the second term in the expression $-5xy^2 + 2y + 8y^2 + 6$.
A. -5 **B.** 2 **C.** 8 **D.** 6 **E.** 5
2. **MC** Jodie had \$55 in her purse. She spent \$ x . Select the expression for the amount of money she has left.
A. $\$(x - 55)$ **B.** $\$(x + 55)$ **C.** $\$55x$ **D.** $\$(55 - x)$ **E.** $\$\left(\frac{55}{x}\right)$
3. **MC** $\frac{3x}{5} - \frac{x}{4}$ can be simplified to:
A. $\frac{7x}{20}$ **B.** $\frac{2x}{20}$ **C.** $\frac{2x}{1}$ **D.** $\frac{4x}{20}$ **E.** $\frac{x}{20}$
4. **MC** $\frac{3}{p} \div \frac{6}{p}$ is equal to:
A. 2 **B.** $\frac{1}{2}$ **C.** $12p$ **D.** $12p^2$ **E.** $\frac{1}{2p^2}$
5. **MC** The volume of a sphere, V , is given by the formula $V = \frac{4\pi r^3}{3}$, where r is the radius of the sphere. If $r = 3$, select the volume of the sphere from the following options.
A. 25.12 **B.** 38.37 **C.** 64.05 **D.** 113.10 **E.** 4.19
6. **MC** Choose the equivalent of $6 - 4x(x + 2) + 3x$.
A. $6 - 4x^2 - 5x$ **B.** $6 + 4x^2 + 5x$ **C.** $6 - 4x^2 + 5x$ **D.** $6 + 4x^2 + 5x$ **E.** $6 + 4x^2 - 5x$
7. **MC** Choose the equivalent of $(3 - a)(3 + a)$.
A. $9 + a^2$ **B.** $9 - a^2$ **C.** $3 + a^2$ **D.** $3 - a^2$ **E.** $9 - a$
8. **MC** Choose the equivalent of $(2y + 5)^2$.
A. $4y^2 + 20y + 25$ **B.** $4y^2 + 10y + 25$ **C.** $2y^2 + 20y + 25$ **D.** $2y^2 + 20y + 5$ **E.** $2y^2 + 10y + 25$
9. **MC** Select what $6(a + 2b) - x(a + 2b)$ equals when it is factorised.
A. $6 - x(a + 2b)$ **B.** $(6 - x)(a + 2b)$ **C.** $6(a + 2b - x)$ **D.** $(6 + x)(a - 2b)$ **E.** $(6 + x)(a + 2b)$
10. **a.** For the expression $-8xy^2 + 2x + 8y^2 - 5$:
 i. state the number of terms
 ii. state the coefficient of the first term
 iii. state the constant term
 iv. state the term with the lowest coefficient.
b. Write expressions for the following, where x and y represent the following numbers.
 i. A number 8 more than y .
 ii. The difference between x and y .
 iii. The sum of x and y .
 iv. A number that is 7 times the product of x and y .
 v. The number given when 2 times x is subtracted from 5 times y .

11. a. Leo receives x dollars for each car he washes. If he washes y cars, calculate how much he earn.
 b. A piece of rope is 24 m long.
 i. If Rawiri cuts k m off, calculate how much is left.
 ii. After Rawiri has cut k m off, he divides the rest of the rope into three pieces of equal length. Calculate the length of each piece.
12. Simplify the following expressions by collecting like terms.
- a. $8p + 9p$ b. $5y^2 + 2y - 4y^2$ c. $9s^2t - 12s^2t$
 d. $11c^2d - 2cd + 5dc^2$ e. $n^2 - p^2q - 3p^2q + 6$ f. $8ab + 2a^2b^2 - 5a^2b^2 + 7ab$
13. Simplify the following expressions.
- a. $6a \times 2b$ b. $2ab \times b$ c. $2xy \times 4yx$ d. $\frac{4x}{12}$ e. $18 \div (4b)$
14. Expand the following expressions.
- a. $5(x + 3)$ b. $-(y + 5)$ c. $-x(3 - 2x)$ d. $-4x(2m + 1)$
15. Expand and simplify the following expressions by collecting like terms.
- a. $3(x - 2) + 9$ b. $-2(5m - 1) - 3$
 c. $4m(m - 3) + 3m - 5$ d. $7p - 2 - (3p + 4)$
16. Expand and simplify the following expressions.
- a. $3(a + 2b) + 2(3a + b)$ b. $-4(2x + 3y) + 3(x - 2y)$
 c. $2m(n + 6) - m(3n + 1)$ d. $-2x(3 - 2x) - (4x - 3)$
17. Expand and simplify the following expressions.
- a. $(x + 4)(x + 5)$ b. $(m - 2)(m + 1)$ c. $(3m - 2)(m - 5)$ d. $(2a + b)(a - 3b)$
18. Expand and simplify the following expressions.
- a. $(x + 4)(x - 4)$ b. $(9 - m)(9 + m)$ c. $(x + y)(x - y)$ d. $(1 - 2a)(1 + 2a)$
19. Expand and simplify the following expressions.
- a. $(x + 5)^2$ b. $(m - 3)^2$ c. $(4x + 1)^2$ d. $(2 - 3y)^2$
20. Expand and simplify the following expressions.
- a. $(x + 2)(x + 1) + (x + 3)(x + 2)$ b. $(m + 7)(m - 2) + (m + 3)^2$
 c. $(x + 6)(x + 2) - (x + 3)(x - 1)$ d. $(b - 7)^2 - (b - 3)(b - 4)$
21. Expand and simplify the following expressions.
- a. $(x + 2)^2 + (x + 3)^2$ b. $(x - 2)^2 - (x - 3)^2$ c. $(x + 4)^2 - (x - 4)^2$
22. If $y = 5x^2 + 2x - 1$, determine the value of y when:
 a. $x = 2$
 b. $x = 5$.
23. The volume (V) of each of the following paint tins is given by the formula $V = \pi r^2 h$, where r is the radius and h is the height of the cylinder. Determine the value of V (to 2 decimal places) when:
 a. $r = 7$ cm and $h = 2$ cm
 b. $r = 9.3$ cm and $h = 19.8$ cm.



24. Factorise each of the following expressions by determining common factors.

a. $6x + 12$

b. $6x^2 + 12x^2y$

c. $8a^2 - 4b$

d. $16x^2 - 24xy$

e. $-2x - 4$

f. $b^2 - 3b + 4bc$

25. Simplify each of the following expressions using common factor techniques.

a. $5(x + y) - 4a(x + y)$

b. $7a(b + 5c) - 6c(b + 5c)$

c. $15x(d + 2e) + 25xy(d + 2e)$

d. $2x + 2y + ax + ay$

e. $6xy + 4x - 6y - 4$

f. $pq - r + p - rq$

Problem solving

26. A rectangular rug has a length of $3x$ cm and a width of x cm.

a. Write an expression for the rug's perimeter.

b. Write an expression for the rug's area.

c. The rug's side length is increased by y cm.

i. Write an expression for its new side length.

ii. Write an expression for the rug's new perimeter. Expand this expression.

iii. Calculate the rug's perimeter when $x = 90$ cm and $y = 30$ cm.

iv. Write an expression for the rug's new area. Expand this expression.

v. Calculate the rug's area when $x = 90$ cm and $y = 30$ cm.

27. A rectangular garden bed has a length of 15 m and a width of 8 m. It is surrounded by a path with a width of p m.

a. Write down the total area of the garden bed and path in factorised form.


b. Expand the expression you wrote down in part a.

c. Calculate the area of the path in terms of p .

d. Write an equation that can be solved to find the width of the path if the area of the path is 200 m^2 .

28. Calculate the circumference of a circle whose area is $\pi \left(\frac{x}{y}\right) \text{ cm}^2$. Give your answer in terms of π .

29. The large sign shown appears in a parking lot at the entrance to car park 5. It has a uniform width, with dimensions shown.



Write an algebraic expression for the area of the front of the sign.

30. At the end of a lesson involving boxes of Smarties, the Smarties left over are shared equally among the class. There are 6 boxes full of Smarties left over and 42 Smarties that are no longer in boxes.
- Write an expression for the total number of Smarties left over.
(Hint: Let n = the number of Smarties in a box.)
 - Write the expression from part **a** in factorised form.
 - If there are only three people in the class, write an expression for the number of Smarties each person receives.
 - If there are x people in the class, write an expression for the number of Smarties each person receives.



31. When entering numbers into an electronic device, or even writing numbers down, we can often make mistakes. A common type of mistake is called a transposition error. This is when two digits are written in the reverse order. For example, the number 2869 could be written as 8269, 2689 or 2896.



A common rule for checking if these mistakes have been made is to see if the difference between the correct number and the recorded number is a multiple of 9. If it is, a transposition error has occurred.

We can use algebraic expressions to check this rule. Let the digit in the thousands position be represented by a , the digit in the hundreds position by b , the digit in the tens position by c , and the digit in the ones position by d .

In this way the real number can be represented as $1000a + 100b + 10c + d$.

- If the digits in the ones position and the tens position were written in the reverse order, the number would be $1000a + 100b + 10d + c$. The difference between the correct number and the incorrect one would then be $1000a + 100b + 10c + d - (1000a + 100b + 10d + c)$.

- Simplify this expression.
- Is the expression a multiple of nine? Explain your answer.

- If a transposition error had occurred with the tens and hundreds positions, the incorrect number would be $1000a + 100c + 10b + d$.

Perform the procedure shown in part **a** to work out whether the difference between the correct number and the incorrect one is a multiple of 9.

- Consider a transposition error in the thousands and hundreds positions. Explain whether the difference between the two numbers is a multiple of 9.
- Comment on this rule for transposition errors.
- The price marked on the tag of a CD in a music store is x dollars and y cents. Its value is less than \$100. The person on the counter makes a mistake and enters a value of y dollars and x cents into the cash register.

Show whether the multiple of 9 rule for checking transposition errors applies in this case.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 2 Algebra

2.1 Pre-test

- D
- $2x^2 - y^2 - xyz = 11$
- $m = 2$
- E
- A
- a. $2m + 13n$ b. $-2xy - 3x + 5y$
- $6q^2 + qr - r^2$
- D
- Yes
- False
- $3b^2 - 21b + 37$
- a. $4p(2p + 5)$ b. $5(5x^2 - 2y)$ c. $xy(6 + 8x - 3y)$
- a. $(7a - 1)(b + 3)$ b. $(5x - 2)(2y + 1)$
- $2x(2x - 1) + 2x(3x + 2) + 2(3x + 2)(2x - 1)$
- $(x + 4)(x - 1)$

2.2 Using pronumerals

- a. 3 b. -2 c. $\frac{1}{3}$ d. $-\frac{1}{4}$
- a. 5 b. -2 c. $\frac{1}{6}$ d. $-\frac{2}{7}$
- a. 15 b. $-\frac{2}{5}$ c. $-\frac{7}{2}$ d. $-\frac{1}{7}$
- a. i. 3 ii. 5 iii. 8 iv. $5x^2$
 b. i. 3 ii. -9 iii. -6 iv. $-9m^2$
 c. i. 4 ii. 5 iii. 5 iv. $-7x^2$
 d. i. 4 ii. 9 iii. 4 iv. $-9b^2$
- a. i. 4 ii. 4 iii. -2 iv. $-3ac$
 b. i. 5 ii. 5 iii. 9 iv. $-3u$
 c. i. 5 ii. -1 iii. 8 iv. $-m$
 d. i. 5 ii. 7 iii. 14 iv. $-3cd^2$
- a. i. 3 ii. 3 iii. -7 iv. $-10x^2$
 b. i. 5 ii. -3 iii. 7 iv. $-6x^2$
 c. i. 5 ii. -5 iii. 8 iv. $-5x$
 d. i. 6 ii. 6 iii. 6 iv. $-15xy$
- a. 2, 12, -8 b. 0, 2, 6
 c. -1, 9, -3 d. 0, 1, -1
- a. 4, -11, 19 b. 1, 4, 16
 c. 0, 21, -3 d. 8, 2, -4
- a. $\sqrt{3}$, 2, 1 b. 0, 3, -2
 c. -12, -12, -6 d. $\frac{3}{5}$, 1, $-\frac{1}{5}$
- a. B b. D c. A d. B
- a. 4 b. -6 c. 6
 d. 7 e. -12 f. -3
- a. 7 b. -40 c. 7
- a. i. 32 ii. 250
 b. i. 4 ii. 3

- a. 15 b. 10 c. 18
 d. 45 e. 30
- a. 10.72 b. 21 c. 22.4
 d. 50 e. 171
- a. 4 b. -30 c. 32.75
 d. 38.33 e. 34.56
- a. Sample responses can be found in the worked solutions in the online resources.
 b. 12 cm
- a. 10 b. 10
- a. 15
 b. Sample responses can be found in the worked solutions in the online resources.
- a. 5
 b. Sample responses can be found in the worked solutions in the online resources.
- a. 19.44
 b. Sample responses can be found in the worked solutions in the online resources.
- a. 63 cm^3 b. 4 cm
- a. 461.81 b. 9.55
- a. 1671.33 b. 2.64
- a.

d	$2d + 2g$	g
$2g$	$d + g$	$2d$
$2d + g$	0	$d + 2g$

Magic number = $3d + 3g$

- a. Sample responses can be found in the worked solutions in the online resources.
- a. mn
 b. $(m - 2)(n - 2)$
 c. $(2m + 2n)$ units
- $y - (z - x)$

2.3 Algebra in worded problems

- a. 11 b. $x + 5$ c. $y + 5$
- a. 3.5 km b. $(x + 2)$ km
- a. 4 b. $2.5 + t$ c. $2.5 - y$ hours
- a. $d - x$ km b. $d + y$ km c. $2d + 4$ km
- a. $10n$, where n = number of shirts
 b. $N + 30$, where N = the number of Nick's dollars
 c. i. $g - 4$ ii. $7g - 4$
- a. i. 94 ii. $6x + y$
 b. i. 66 ii. $6p + q$
 c. i. 28 ii. $6x + y - 6p - q$
- xy dollars
- a. $\frac{x}{n}$ b. $\frac{x}{2n}$
- a. $(30 - d)$ cm b. $\frac{30 - d}{4}$ cm c. $\frac{3(30 - d)}{4}$ cm
- a. $\frac{x}{4}$ b. $\frac{x}{6}$ c. $\frac{x}{4} - \frac{x}{6} = \frac{x}{12}$

11. a. $\frac{24-c}{7}$ hours b. $\left(c + \frac{24-c}{7}\right) = \frac{6c+24}{7}$

12. a. 1 b. 7:30 pm and 8:00 pm
 c. 5 d. 24
 e. 8:30 pm and 9:00 pm

13. a. $\$(y-30)$
 b. $\left(\frac{1}{3}(y-30)\right)$
 c. \$930

14. a. $b = 2h - 1$ b. 19 blocks

15. 15 red vases and 10 clear vases

16. The father is 32 years old.

17. 4 dots = 4 triangles
 5 dots = 10 triangles
 6 dots = 20 triangles
 7 dots = 35 triangles
 $t = \frac{d!}{6(d-3)!}$ where $d!$ is the product of all positive integers less than or equal to d .

18. a. $x = \frac{T-3}{T-1}$
 b, c. Answer is undefined.

2.4 Simplification of algebraic expressions

1. a. $9ab, -ab$ b. $4x$
 c. $-3za, -az$ d. $2x^2, -x^2$
2. a. $-2x^2y$ b. $-x^2y^5, -3x^2y^5$
 c. $p^3x^2w^5, -5x^2p^3w^5$ d. $4y^5z^4x^2, -2x^2z^4y^5$
3. a. $7x$ b. $11y$ c. $19m$ d. $11q$ e. $8r$ f. $3x$
4. a. $8a$ b. $8y$ c. $13x$ d. $16p$ e. $9q^2$ f. $3x^2$
5. a. $8x^2 - 3y$ b. $2m^2 + 2n$
 c. $-2g^2 + g - 12$ d. $-5m^2 + m + 15$
 e. $12a^2 + b + 4b^2$ f. $3m + 7n^2$
6. a. $12xy + 2y^2$ b. $2ab + 5a^2b$
 c. $16x^2y - 3xy$ d. $m^2n + 11n$
 e. $-7x^2 + 2x^2y$ f. $-3a^2b - 9a^2 + 5b + 4$

7. B

8. D

9. A

10. D

11. a. $6mn$ b. $20xy$ c. $8pq$
 d. $-10xy$ e. $-12xy$

12. a. $15mn$ b. $10a^2$ c. $6mnp$
 d. $-6ab^2$ e. $10m^2n$

13. a. $-18a^2b$ b. $30x^3y^2$ c. $-12p^2q^3$
 d. $-56c^3d$ e. $30a^4b^7$

14. a. $3x$ b. $3m$ c. $2y$ d. $4m$ e. $4m$

15. a. $2x$ b. $-7x$ c. $-4m$ d. $\frac{m}{2}$ e. $\frac{x}{3}$

16. a. $\frac{4m}{9}$ b. $\frac{1}{2a}$ c. $2yz$

d. $\frac{xy}{4}$ e. $\frac{-7yz}{11}$

17. a. $40x^2y^2$ b. $56ax^2y^2$

c. $\frac{x^2}{2y}$ d. $\frac{-5a}{4b}$

18. a. 2 b. $40a^3b^2$

c. $-8a^3b^3$ d. $16a^4$

19. a. i. $(0.01mp + nq)$ dollars ii. $(mp + 100nq)$ cents

b. $20 - (0.01mp + nq)$ dollars

20. Shirt = \$12 each
 Shorts = \$30 each


21. \$45

22. The correct answer is $7a^2c^2$. Sample responses can be found in the worked solutions in the online resources.


23. $\frac{3b^9}{2c^6}$

24. a. $\frac{n}{(n+2)}$ b. $\frac{n}{(n+3)}$ c. $\frac{1}{100}$


25. a. i.




ii.



b. i.



ii.



26. a. $P = \frac{x}{3} + x + \frac{5x}{7}$

b. Sample responses can be found in the worked solutions in the online resources.

c. \$18

27. a. $P = 2w + \frac{32w}{7}$

b. Sample responses can be found in the worked solutions in the online resources.

c. \$72.22

28. a. $P = 2x + 2(4x - 7)$

b. $P = 10x - 14$

c. 16 units

d. If $x = 1$ the rectangle will have a negative value for the perimeter, which is not possible.

29. a. $l = 4(x + 40) + 8(x + 20)$

b. $l = 12x + 320$

c. \$19.20

30. a. $P_1 = (10x + 12)$ cm b. $x = 2$
 $P_2 = 16x$ cm

2.5 Expanding brackets

1. a. 64 b. 1100
 c. -20 d. 693
2. a. $3x + 6$
- | | | |
|---|------|-----|
| 3 | $3x$ | 6 |
| | x | 2 |
- b. $4x + 12$
- | | | |
|---|------|------|
| 4 | $4x$ | 12 |
| | x | 3 |
- c. $4x + 4$ d. $7x - 7$
3. a. $-3p + 6$ b. $-x + 1$
 c. $6b - 12$ d. $24m - 16$
4. a. $x^2 + 2x$
- | | | |
|---|-------|------|
| x | x^2 | $2x$ |
| | x | 2 |
- b. $a^2 + 5a$
- | | | |
|---|-------|------|
| a | a^2 | $5a$ |
| | a | 5 |
- c. $4x + x^2$ d. $7m - m^2$
5. a. $2xy + 4x$ b. $-3xy - 12y$
 c. $-3b + ab$ d. $-30a + 18a^2$
6. a. $2p - 2$ b. $5x - 17$ c. $-7p - 17$
 d. $-12p + 3$ e. $6x^2 - 20x$
7. a. $2m^2 + 7m$ b. $3px + 6x - 5$
 c. $4y^2 - 4y + 7$ d. $-4p^2 + 13p$
 e. $4x - 13y$
8. a. $2m^2 - 8m - 4$ b. $-3p^2 + 10pq - 1$
 c. $-30a + 10ab$ d. $7cd - 12c^2 - 5c$
 e. $-2p - 17$
9. a. $8x + y$ b. $10p + 8q$ c. $18a + 29b$
 d. $19c + 22d$ e. $2m - 11n$
10. a. $6x - 11y$ b. $9x + 5y$ c. $-14p - 8q$
 d. $-4a + 3b$ e. $4x - y$
11. a. $5p - 10q$ b. $-8c + 9d$ c. $13x - 19y$
 d. $-7p + 11q$ e. $-5a + 3b$
12. a. $2ab + 2a - 3b$ b. $2xy + 4x - 2y$
 c. $2cd + 3c$ d. $2pq - 2p$
 e. $5cd - 11c$
13. a. $5ab - 21a - 3b$ b. $5m$
 c. $6cd - 36c$ d. $6m^2 + 6m - 10$
 e. $9cd - 8c$
14. a. $-15a^2 + 2b^2 - 9ab$
 b. $-8c^2 + 3d^2 + 22cd$
 c. $12m^2 - 20m - 4$
 d. $-7x^2 + 41x - 6$
 e. $-10y^2 - 6y - 12$
15. B
16. A

17. D
18. a. $a^2 + 5a + 6$ b. $x^2 + 7x + 12$
 c. $y^2 + 5y + 6$ d. $m^2 + 9m + 20$
 e. $b^2 + 3b + 2$
19. a. $p^2 + 5p + 4$ b. $a^2 + a - 6$
 c. $x^2 + x - 20$ d. $m^2 - m - 12$
 e. $y^2 + 2y - 15$
20. a. $y^2 - 4y - 12$ b. $x^2 - 2x - 3$
 c. $x^2 - 7x + 12$ d. $p^2 - 5p + 6$
 e. $x^2 - 4x + 3$
21. a. $2a^2 + 7a + 6$ b. $4c^2 - 31c + 42$
 c. $2t^2 - 17t + 35$ d. $-6t^2 + 11t + 10$
 e. $15x^2 - 31x + 14$ f. $25x^2 - 20x + 4$
22. a. $xz + x + yz + y$ b. $2xz + 8x + yz + 4y$
 c. $3pr + 3p + qr + q$ d. $a^2 + 3ab + 2b^2$
 e. $2c^2 - 5cd - 3d^2$ f. $2x^2 - xy - 3y^2$
23. a. $4p^2 + pq - 3q^2$ b. $ab + ac + 2b^2 + 2bc$
 c. $3p - 9pr - 2q + 6qr$ d. $12x^2 - 7xy + y^2$
 e. $2p^2 - pr - 2pq + qr$ f. $15k - 5 - 6jk + 2j$
24. C
25. C
26. a. $x^2 - 9$ b. $x^2 - 25$ c. $x^2 - 49$
 d. $x^2 - 1$ e. $x^2 - 4$ f. $4x^2 - 1$
27. a. $x^2 + 2x + 1$ b. $x^2 + 4x + 4$
 c. $x^2 + 16x + 64$ d. $x^2 - 6x + 9$
 e. $x^2 - 10x + 25$ f. $x^2 - 18x + 81$
28. a. $6.3x^2 + 5.53xy - 3.1y^2$
 b. $4.41x^2 - 10.24y^2$
 c. $11.56x^2 + 34.68xy + 26.01y^2$
29. a. Surface area = $38x^2 + 14x - 6$
 b. Volume = $12x^3 + 5x^2 - 3x$
30. a. i. $((x + 3y)(4x - y))m^2$
 ii. $(4x^2 + 11xy - 3y^2)cm^2$
 iii. Both brackets must be positive; therefore, $x > -3y$, $x > \frac{y}{4}$
- b. i. $\frac{(2x - 1)(x + 5)}{2}$
 ii. $\frac{2x^2 + 9x - 5}{2}$
 iii. Sample responses can be found in the worked solutions in the online resources.
31. Sample responses can be found in the worked solutions in the online resources.
32. a. Negative sign ignored.
 b. Negative sign ignored.
 c. Distributive law not used.
33. a. i. The student did not multiply both terms.
 ii. The student used addition instead of multiplication.
 iii. The student did not change negative and positive signs.
 b. $100 - 15x$
34. a. $-2(a - 5) = 2a - 10$ is incorrect because the student did not change the multiplied negative signs for $-2 \times (-5)$ to a positive sign. The correct answer is $-2a + 10$.

- b. $2b(3b - 1) = 6b^2 - 1$ is incorrect because the student did not multiply -1 and $2b$ together. The correct answer is $6b^2 - 2b$.
- c. $-2(c - 4) = 2c + 8$ is incorrect because the student left out the negative sign when multiplying -2 and c . The correct answer is $-2c + 8$.
35. a. Student C
- b. Corrections to the students' answers are shown in bold.
Student A:
 $(3x + 4)(2x + 5)$
 $= 3x \times 2x + 3x \times 5 + 4 \times 2x + 4 \times 5$
 $= 6x^2 + 23x + 20$
 (Also, $29x + 20$ does not equal $49x$.)
 Student B:
 $(3x + 4)(2x + 5)$
 $= 3x \times 2x + 3x \times 5 + 4 \times 2x + 4 \times 5$
 $= 6x^2 + 15x + 8x + 20$
 $= 6x^2 + 23x + 20$
36. a. $w = (x - 3)$ cm
 b. $w = 62$ cm
 c. $l = 108$ cm

2.6 Difference of two squares and perfect squares

1. a. $x^2 - 4$ b. $y^2 - 9$ c. $m^2 - 25$ d. $a^2 - 49$
2. a. $x^2 - 36$ b. $p^2 - 144$ c. $a^2 - 100$ d. $m^2 - 121$
3. a. $4x^2 - 9$ b. $9y^2 - 1$ c. $25d^2 - 4$
 d. $49c^2 - 9$ e. $4 - 9p^2$
4. a. $d^2 - 81x^2$ b. $25 - 144a^2$ c. $9x^2 - 100y^2$
 d. $4b^2 - 25c^2$ e. $100 - 4x^2$
5. a. $x^2 + 4x + 4$ b. $a^2 + 6a + 9$
 c. $b^2 + 14b + 49$ d. $c^2 + 18c + 81$
6. a. $m^2 + 24m + 144$ b. $n^2 + 20n + 100$
 c. $x^2 - 12x + 36$ d. $y^2 - 10y + 25$
7. a. $81 - 18c + c^2$ b. $64 + 16e + e^2$
 c. $2x^2 + 4xy + 2y^2$ d. $u^2 - 2uv + v^2$
8. a. $4a^2 + 12a + 9$ b. $9x^2 + 6x + 1$
 c. $4m^2 - 20m + 25$ d. $16x^2 - 24x + 9$
9. a. $25a^2 - 10a + 1$ b. $49p^2 + 56p + 16$
 c. $81x^2 + 36x + 4$ d. $16c^2 - 48c + 36$
10. a. $25 + 30p + 9p^2$ b. $4 - 20x + 25x^2$
 c. $81x^2 - 72xy + 16y^2$ d. $64x^2 - 48xy + 9y^2$
11. a. $x^2 - 9$ b. $4x^2 - 9$ c. $49x^2 - 16$
 d. $4x^2 - 49y^2$ e. $x^4 - y^4$
12. a. $16x^2 + 40x + 25$ b. $49x^2 - 42xy + 9y^2$
 c. $25x^4 - 20x^2y + 4y^2$ d. $2x^2 - 4xy + 2y^2$
 e. $\frac{4}{x^2} + 16 + 16x^2$
13. $(x^2 + 6x + 9)$ units²
14. $(3x + 1)$ m
15. Perimeter = $4x + 36$
16. Sample responses can be found in the worked solutions in the online resources.

17. a. Sample responses can be found in the worked solutions in the online resources.
 b. Lin's bedroom is larger by 1 m^2 .
18. a. i. $x^2 - 16$ and $16 - x^2$ iii. $4x^2 - 81$ and $81 - 4x^2$
 ii. $x^2 - 121$ and $121 - x^2$
- b. The answers to the pairs of expansions are the same, except that the negative and positive signs are reversed.
- c. This is possible because when a negative number is multiplied by a positive number, it becomes negative. When expanding a DOTS in which the expressions have different signs, the signs will be reversed.
19. a. $100k^2 + 100k + 25$
- b. $(10k + 5)^2 = 100 \times k \times k + 100 \times k + 25$
 $= 100k(k + 1) + 25$
- c. $25^2 = (10 \times 2 + 5)^2$
 Let $k = 2$.
 $25^2 = 100k(k + 1) + 25$
 $= 100 \times 2 \times (2 + 1) + 25$
 $= 625$
 $85^2 = (10 \times 8 + 5)^2$
 Let $k = 8$.
 $85^2 = 100k(k + 1) + 25$
 $= 100 \times 8 \times (8 + 1) + 25$
 $= 7225$
20. a. $A_1 = a^2$ units²
 $A_2 = ab$ units²
 $A_3 = ab$ units²
 $A_4 = b^2$ units²
- b. $A = a^2 + ab + ab + b^2$
 $= a^2 + 2ab + b^2$
 This is the equation for perfect squares.
21. a. i. $x^2 - 6x + 9$ and $9 - 6x + x^2$
 ii. $x^2 - 30x + 225$ and $225 - 30x + x^2$
 iii. $9x^2 - 42x + 49$ and $49 - 42x + 9x^2$
- b. The answers to the pairs of expansions are the same.
- c. This is possible because when a negative number is multiplied by itself, it becomes positive. When expanding a perfect square in which the two expressions are the same, the negative signs cancel out and result in the same answer.
22. a. 729 b. 1089 c. 1521 d. 2209
23. $(3x^2 + 16x + 5)$ m²
24. $(x^2 - 25)$ m²
25. a. 10609 b. 3844 c. 994 009
 d. 1 024 144 e. 2809 f. 9604

2.7 Further expansions

1. a. $2x^2 + 13x + 21$ b. $2x^2 + 13x + 20$
 c. $2x^2 + 14x + 26$ d. $2x^2 + 10x + 11$
2. a. $2p^2 - 3p - 21$ b. $2a^2 - 5a + 4$
 c. $2p^2 - p - 24$ d. $2x^2 + 19x - 36$
3. a. $2y^2 + 2y - 7$ b. $2d^2 + 8d - 2$
 c. $2x^2 + 10$ d. $2y^2$

4. a. $2x^2 - 4x + 19$
c. $2p^2 + 3p + 23$
5. a. $x + 5$
c. $-2x - 6$
6. a. $-3b - 22$
c. $8p - 10$
7. a. $4m + 17$
c. $-6p - 7$
8. a. $16x - 16$
b. Sample responses can be found in the worked solutions in the online resources.
9. a. $15x^2 - 44x + 21$
b. Sample responses can be found in the worked solutions in the online resources.
10. a. $8x^2 - 10xy + 3y^2$
b. Sample responses can be found in the worked solutions in the online resources.
11. $x = -6$
12. Sample responses can be found in the worked solutions in the online resources.
13. Sample responses can be found in the worked solutions in the online resources.
14. a. Sample responses can be found in the worked solutions in the online resources.
b. $a = 2, b = 1, c = 3, d = 4$
 $(2^2 + 1^2)(3^2 + 4^2) = (2 \times 3 - 1 \times 4)^2 + (2 \times 4 - 1 \times 3)^2$
 $= 4 + 121$
 $= 125$
15. a. $x^4 + 2x^3 - x^2 - 2x + 1$
b. Sample responses can be found in the worked solutions in the online resources.
c. i. 25
ii. $x = 2$
16. a. $ad + ae + bd + be$
b. $(a + b + c)(d + e + f) = ad + ae + bd + be + cd + ce + af + bf + cf$

	a	b	c
d	ad	bd	cd
e	ae	be	ce
f	af	bf	cf

2.8 The highest common factor

1. a. $2a$
d. -4
2. a. $-3x$
d. ab
3. a. 4
4. a. 25
5. a. 10
6. C
7. a. $4(x + 3y)$
c. $7(a + 2b)$
e. $-8(a + 3b)$
- b. 4
e. $3x$
b. m
e. $3x$
- c. $3x$
f. $15a$
c. $-mn$
f. $5n^2$
d. 8
e. 3
c. 6a
d. $7x$
e. $30q$
d. 5
e. x^2
b. $5(m + 3n)$
d. $7(m - 3n)$
f. $4(2x - y)$
8. a. $-2(6p + q)$
c. $8(4x + y + 2z)$
e. $8(9x - y + 8pq)$
9. a. $5(p^2 - 4q)$
c. $8(7q + p^2)$
e. $4(4p^2 + 5q + 1)$
10. a. $3(3a + 7b)$
c. $4(3p^2 + 5q^2)$
e. $5(5y^2 - 3x)$
11. a. $4(4a^2 + 5b)$
c. $9(7p^2 + 9 - 3y)$
e. $2(5 - 11x^2y^3 + 7xy)$
12. a. $9(2a^2bc - 3ab - 10c)$
c. $7(9a^2b^2 - 7 + 8ab^2)$
e. $6(6 - 4ab^2 + 3b^2c)$
13. a. $-(x - 5)$
d. $-2(m + 3)$
14. a. $-3(n^2 - 5m)$
d. $-6(2p^2 + 3q)$
15. a. $-3(3a^2b - 10)$
c. $-2(9x^2 - 2y^2)$
e. $-5(2 + 5p^2 + 9q)$
16. a. $a(a + 5)$
d. $p(6 + 7p)$
17. a. $y(x + 9 - 3y)$
c. $ab(3 + a + 4b)$
e. $pq(5pq - 4 + 3p)$
18. a. $5x(x + 3)$
d. $-2m(m - 4)$
19. a. $-3a(a - 3)$
d. $2m(2 - 9m)$
20. a. $2(x + 3)(4x + 1)$
b. $A = x(x + 3)$
B = $(5x + 2)(x + 3)$
C = $2x(x + 3)$
21. a. $3a(b^2 + 6b + 9)$
b. $3a(b + 3)^2$
22. a. $(x + 4)(x^2 + 8x + 16)$
b. $(x + 4)^3$
23. Sample responses can be found in the worked solutions in the online resources.
24. Sample responses can be found in the worked solutions in the online resources.
25. a. $(x + \sqrt{13})(x - \sqrt{13})$
b. $(2x - \sqrt{17})(2x + \sqrt{17})$
c. $(x + 3 + \sqrt{10})(x + 3 - \sqrt{10})$
26. -2
27. True. Sample responses can be found in the worked solutions in the online resources.
28. a. $4(3x - 5y)(3x + 5y)$
b. $4(3x - 5y)(3x + 5y)$
c. Yes, the answers are the same.
- b. $6(p + 2pq + 3q)$
d. $4(4m - n + 6p)$
f. $3(5x^2 - y)$
b. $5(x + 1)$
d. $7(p - 6x^2y)$
f. $12(1 + 3a^2b - 2b^2)$
b. $2(2c + 9d^2)$
d. $7(5 - 2m^2n)$
b. $6(7m^2 + 2n)$
d. $11(11a^2 - 5b + 10c)$
b. $12(12p + 3q^2 - 7pq)$
d. $11(2 + 9p^3q^2 - 4p^2r)$
c. $-(b - 9)$
f. $-4(a + 2)$
c. $-7(y^2 + 7z)$
f. $-2(6m^3 + 25x^3)$
b. $-3(5p + 4q)$
d. $-3(ab - 6m + 7)$
f. $-9(10m^2 - 3n - 6p^3)$
c. $m(18 + 5m)$
f. $p(7 - pq + q)$
b. $c(5 + 3cd - d)$
d. $xy(2x + 1 + 5y)$
f. $xy(6xy - 5 + x)$
b. $6m(4m - 1)$
c. $4a(8a - 1)$
f. $-7y(y - 2)$
c. $-13y(2y + 1)$
f. $-8p(1 + 3p)$

2.9 The highest common binomial factor

3. a. $(a + b)(2 + 3c)$ b. $(m + n)(4 + p)$
 c. $(2m + 1)(7x - y)$ d. $(3b + 2)(4a - b)$
 e. $(x + 2y)(z - 3)$
4. a. $(6 - q)(12p - 5)$ b. $(x - y)(3p^2 + 2q)$
 c. $(b - 3)(4a^2 + 3b)$ d. $(q + 2p)(p^2 - 5)$
 e. $(5m + 1)(6 + n^2)$
5. a. $(y + 2)(x + 2)$ b. $(b + 3)(a + 3)$
 c. $(x - 4)(y + 3)$ d. $(2y + 1)(x + 3)$
 e. $(3b + 1)(a + 4)$ f. $(b - 2)(a + 5)$
6. a. $(m - 2n)(1 + a)$ b. $(5 + 3p)(1 + 3a)$
 c. $(3m - 1)(5n - 2)$ d. $(10p - 1)(q - 2)$
 e. $(3x - 1)(2 - y)$ f. $(4p - 1)(4 - 3q)$
7. a. $(2y + 1)(5x - 2)$ b. $(2a + 3)(3b - 2)$
 c. $(b - 2c)(5a - 3)$ d. $(x + 3y)(4 - z)$
 e. $(p + 2q)(5r - 3)$ f. $(a + 5b)(c - 2)$
8. a. $\frac{a}{b}$ b. $\frac{3x - 4}{7a - 5}$
 7. a. $\frac{x + 2}{2 + 6x}$ b. $\frac{y + 2}{3b - 2}$
 8. a. $\frac{3x - 2}{3y + 2}$ b. $\frac{m - 4n}{m + 4n}$

9. Sample responses can be found in the worked solutions in the online resources.

10. $(2x + 3)^2$, this is a perfect square.

11. a. $5(x + 2)(x + 3) = 5(x^2 + 5x + 6) = 5x^2 + 25x + 30$

	x	3	x	3	x	3	x	3	x	3
x	x^2	3x	x^2	3x	x^2	3x	x^2	3x	x^2	3x
2	2x	6	2x	6	2x	6	2x	6	2x	6

b. $5(x + 2)(x + 3) = (5 \times x + 2 \times 5)(x + 3)$
 $= (5x + 10)(x + 3)$

	x	2	x	2	x	2	x	2	x	2
x	x^2	2x	x^2	2x	x^2	2x	x^2	2x	x^2	2x
3	3x	6	3x	6	3x	6	3x	6	3x	6

c. $5(x + 2)(x + 3) = (5 \times x + 3 \times 5)(x + 2)$
 $= (5x + 15)(x + 2)$

	5x	15
x	$5x^2$	15x
2	10x	30

12. a. Side length_a = $x + 3$
 Side length_b = $x + 5$
 Side length_c = $x + 8$
 Side length_d = $x - 3$
 Side length_e = $x + 6$
- b. $A_a = \text{II} = (x + 3)^2$
 $A_b = \text{III} = (x + 5)^2$
 $A_c = \text{V} = (x + 8)^2$
 $A_d = \text{I} = (x - 3)^2$
 $A_e = \text{IV} = (x + 6)^2$

c. $A_a = \text{II} = 64 \text{ cm}^2$
 $A_b = \text{III} = 100 \text{ cm}^2$
 $A_c = \text{V} = 169 \text{ cm}^2$
 $A_d = \text{I} = 4 \text{ cm}^2$
 $A_e = \text{IV} = 121 \text{ cm}^2$

13. $-(x + 1)(4y - 9)$

14. a. i. Square, because it is a perfect square.
 ii. Rectangle, because it is a DOTS.
 iii. Rectangle, because it is a trinomial.
 iv. Rectangle, because it is a trinomial.
- b. i. $(3s + 8)^2$ ii. $(5s + 2)(5s - 2)$
 iii. $(s + 1)(s + 3)$ iv. $4(s - 8)(s + 1)$

2.10 Solving worded problems

1. a. i. $12x$ ii. 60 iii. $9x^2$ iv. 225
 b. i. $4x + 8$ ii. 28
 iii. $x^2 + 4x + 4$ iv. 49
 c. i. $16x - 4$ ii. 76
 iii. $16x^2 - 8x + 1$ iv. 361
 d. i. $10x + 2$ ii. 52
 iii. $6x^2 + 2x$ iv. 160
 e. i. $12x - 2$ ii. 58
 iii. $5x^2 - 13x - 6$ iv. 54
 f. i. $20x + 18$ ii. 118
 iii. $21x^2 + 42x$ iv. 735
2. a. 1250 m^2
 b. $(1250 + 150x + 4x^2) \text{ m}^2$
 c. $(150x + 4x^2) \text{ m}^2$
 d. 366.16 m^2
 e. $150x + 4x^2 = 200$
3. a. $(20 - 2x) \text{ cm}$
 b. $(15 - 2y) \text{ cm}$
 c. $(300 - 40y - 30x + 4xy) \text{ cm}^2$
 d. Sample responses can be found in the worked solutions in the online resources.
4. a. 40 cm^2
 b. i. $(8 + v) \text{ cm}$
 ii. $(5 + v) \text{ cm}$
 iii. $(v^2 + 13v + 40) \text{ cm}^2$
 iv. 70 cm^2
 c. i. $(8 - d) \text{ cm}$
 ii. $(5 - d) \text{ cm}$
 iii. $(40 - 13d + d^2) \text{ cm}^2$
 iv. 18 cm^2
 d. i. $8x \text{ cm}^2$ ii. $(5 + x) \text{ cm}$
 iii. $(8x^2 + 40x) \text{ cm}^2$ iv. 400 cm^2
5. a. $20x \text{ m}$ b. $25x^2 \text{ m}^2$
 c. i. $(5x - 2) \text{ m}$
 ii. $(5x - 3) \text{ m}$
 iii. $(25x^2 - 25x + 6) \text{ m}^2$
 iv. 756 m^2
6. a. $6x \text{ cm}$
 b. $2x^2 \text{ cm}^2$

- c. i. $(2x + y)$ cm
 ii. $(x - y)$ cm
 iii. $(2x^2 - xy - y^2)$ cm²
 iv. 11 cm²

7. a. $4x$ cm
 b. x^2 cm²

- c. i. $(x + y)$ cm
 ii. $(4x + 4y)$ cm
 iii. 56 cm
 iv. $(x^2 + 2xy + y^2)$ cm²
 v. 60.84 cm²

8. a. $(14p + 4)$ m b. $(12p^2 + 6p)$ m²
 c. $(6p + 2)$ m d. $5p$ m
 e. $(22p + 4)$ m f. $(30p^2 + 10p)$ m²
 g. $(18p^2 + 4p)$ m² h. 80 m²

9. a. 27.8
 b. Samir is not within the healthy weight range.
 c. i. 19.1 ii. 24.9 iii. 22.0
 d. Sammara: 9 to 20 years of age; Nimco: 17.5 to 20 years of age; Manuel: 13.5 to 20 years of age

10. a.

Type of algebra	Perimeter	Type of algebra rectangle	Perimeter
1-long	$2x + 2$	1-tall	$2x + 2$
2-long	$4x + 2$	2-tall	$2x + 4$
3-long	$6x + 2$	3-tall	$2x + 6$
4-long	$8x + 2$	4-tall	$2x + 8$
5-long	$10x + 2$	5-tall	$2x + 10$

- b. Perimeter: long $40x + 2$, tall $2x + 40$
 c. $0 < x < 1$

11. a. 3
 b. 994009
 c. i. 990025 ii. 980100

12. a. i. $A_{\text{screen}} = x(x + 2) = x^2 + 2x$
 ii. $A_{\text{phone 1}} = (x + 1)(x + 7) = x^2 + 8x + 7$
 b. i. $l = x + 11$, $w = x$
 ii. $A_{\text{phone 2}} = x(x + 11) = x^2 + 11x$
 c. $x = 4$ cm, $A_{\text{phone 2}} = 60$ cm²

13. a. i. $x(x - 2)$ m² ii. $x(2x)$ m²
 b. $\frac{x - 2}{2x}$
 c. $\frac{50(x - 2)}{x}\%$
 d. i. 25% ii. 27.8% iii. 29.17%

e. The third flag places the most importance on Australia.

14. a. $A = \pi R^2 - x^2$
 b. $A = 160.71$ cm²
 c. Sample responses can be found in the worked solutions in the online resources.
 d. $R = 5.32$ cm
 e. 92.0%

15. a. Sample responses can be found in the worked solutions in the online resources.

- b. i. $-3x^3 - 2x^2 + 3x + 2$
 ii. $8x^3 + 16x^2 - 104x + 80$
 iii. $x^3 - 49x + 120$

c. In a cubic expression there are four terms with descending powers of x and ascending values of the pronumeral.
 For example, $(x + a)^3 = x^3 + 3ax^2 + 3a^2x + a^3$, where the powers of x are descending and the values of a are ascending.

Project

- Solution not required.
- Students need to describe the feature created by arranging the blocks.
- 0.12 m
- 0.0144 m²
- Yellow: $0.03 \text{ m} \times 0.03 \text{ m}$
 Black: $0.03 \text{ m} \times 0.06 \text{ m}$
 White: $0.06 \text{ m} \times 0.06 \text{ m}$
- Yellow: 0.0009 m^2
 Black: 0.0018 m^2
 White: 0.0036 m^2
- Yellow: 0.0036 m^2
 Black: 0.0072 m^2
 White: 0.0036 m^2
- Students are required to derive a general formula that would allow the company to determine the area of the three coloured fabrics in each block. They should give an example and illustrate it with the help of a diagram.

2.11 Review questions

- B
- D
- A
- B
- D
- A
- B
- A
- B
- a. i. 4
 ii. -8
 iii. -5
 iv. $-8xy^2$
 b. i. $y + 8$
 ii. $x - y$
 iii. $x + y$
 iv. $7xy$
 v. $5y - 2x$
- a. $\$xy$
 b. i. $(24 - k)$ m
 ii. $\frac{24 - k}{3}$ m

12. a. $17p$
 c. $-3s^2t$
 e. $n^2 - 4p^2q + 6$
13. a. $12ab$
 d. $\frac{x}{3}$
14. a. $5x + 15$
 c. $-3x + 2x^2$
15. a. $3x + 3$
 c. $4m^2 - 9m - 5$
16. a. $9a + 8b$
 c. $-mn + 11m$
17. a. $x^2 + 9x + 20$
 c. $3m^2 - 17m + 10$
18. a. $x^2 - 16$
 c. $x^2 - y^2$
19. a. $x^2 + 10x + 25$
 c. $16x^2 + 8x + 1$
20. a. $2x^2 + 8x + 8$
 c. $6x + 15$
21. a. $2x^2 + 10x + 13$
 c. $16x$
22. a. 23
23. a. 307.88 cm^3
24. a. $6(x + 2)$
 c. $4(2a^2 - b)$
 e. $-2(x + 2)$
25. a. $(5 - 4a)(x + y)$
 c. $5x(3 + 5y)(d + 2e)$
 e. $2(x - 1)(3y + 2)$
26. a. $8x \text{ cm}$
 b. $3x \text{ cm}^2$
 c. i. $(3x + y) \text{ cm}$
 ii. $(8x + 2y) \text{ cm}$
 iii. 780 cm
 iv. $(3x^2 + xy) \text{ cm}^2$
 v. $27\,000 \text{ cm}^2$
27. a. $(8 + 2p)(15 + 2p) \text{ m}^2$
 c. $(4p^2 + 46p) \text{ m}^2$
28. $\frac{2\pi x}{y} \text{ cm}$
29. $5xy - 4y^2$
30. a. $6n + 42$
 c. $2(n + 7)$
31. a. i. $9(c - d)$
 ii. Yes, this is a multiple of 9 because the number that multiplies the brackets is 9.
- b. $y^2 + 2y$
 d. $16c^2d - 2cd$
 f. $15ab - 3a^2b^2$
- b. $2ab^2$
 e. $\frac{9}{2b}$
- b. $-y - 5$
 d. $-8mx - 4x$
- b. $-10m - 1$
 d. $4p - 6$
- b. $-5x - 18y$
 d. $4x^2 - 10x + 3$
- b. $m^2 - m - 2$
 d. $2a^2 - 5ab - 3b^2$
- b. $81 - m^2$
 d. $1 - 4a^2$
- b. $m^2 - 6m + 9$
 d. $4 - 12y + 9y^2$
- b. $2m^2 + 11m - 5$
 d. $-7b + 37$
- b. $2x - 5$
- b. 134
- b. 5379.98 cm^3
- b. $6x^2(1 + 2xy)$
 d. $8x(2x - 3y)$
 f. $b(b - 3 + 4c)$
- b. $(7a - 6c)(b + 5c)$
 d. $(2 + a)(x + y)$
 f. $(p - r)(q + 1)$
- b. $90(b - c) - 90$ is a multiple of 9, so the difference between the correct and incorrect numbers is a multiple of 9.
- c. $900(a - b)$ is a multiple of 9, so the difference between the correct and incorrect numbers is a multiple of 9.
- d. If two adjacent digits are transposed, the difference between the correct number and the transposed number is a multiple of 9.
- e. Let the correct price of the CD be $\$ab.cd$. This can be written as $10a + b + 0.1c + 0.01d$. The cashier enters $\$cd.ab$ as the price. This can be written as $10c + d + 0.1$. The difference is $10a + b + 0.1c + 0.01d - (10c + d + 0.1a + 0.01b)$. This is equal to $9.9a + 0.99b - 9.9c - 0.99d$, which is a multiple of 9. This means that the rule for checking transposition errors applies in this case.

3 Linear equations

LESSON SEQUENCE

3.1 Overview	148
3.2 Solving linear equations and inequations	150
3.3 Solving linear equations with brackets	161
3.4 Solving linear equations with pronumerals on both sides	165
3.5 Solving problems with linear equations	169
3.6 Rearranging formulas	175
3.7 Review	181



LESSON

3.1 Overview

3.1.1 Why learn this?

Mathematical equations are all around us. Learning how to solve equations lets us work out unknown values. Being able to manipulate and solve equations has many uses in many fields, including chemistry, medicine, economics and commerce, to name just a few.

Being able to solve linear equations is a useful skill for many everyday tasks. Linear equations can be used when converting temperature from Celsius to Fahrenheit, when working out how to balance chemical equations, or when converting between different currencies. Linear equations can also be used when working out fuel consumption for a road trip, deciding on the right dosage of medicine, or budgeting for a holiday.

Any calculation of rates will use some kind of linear equation. Linear equations can also be used to help predict future trends involving growth or decay. An understanding of the underlying principles for solving linear equations can also be applied to the solution of other mathematical models.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 3.1 Pre-test

- Solve the linear equation $\frac{x}{3} - 2 = -1$.
- Solve the linear equation $0.4x - 2.6 = 6.2$.
- MC** Select which of the following equations has the solution $x = 2$.
 A. $x + 3 = -5$ B. $3 - x = 5$ C. $3(x + 1) = 2x + 5$
 D. $\frac{3}{x} = 6$ E. $x^2 - x = 1$
- Solve the following linear equations.
 a. $\frac{3x + 1}{4} = -5$ b. $\frac{3 - z}{2} = -4$
- Solve the equation $-3(2a - 4) = -12$.
- Dylan is solving the equation $4(y + 3) = -10$. State whether it is true or false that Dylan will calculate the correct value of y if he subtracts 3 from both sides of the equation then divides both sides by 4.
- MC** Select the correct value for m in the equation $0.4(m - 8) = 0.6$.
 A. $m = -6.5$ B. $m = 6.5$ C. $m = 7.8$
 D. $m = 8.2$ E. $m = 9.5$
- MC** Select the first step to solve the equation $2x + 1 = 11 - 3x$.
 A. Add $2x$ to both sides. B. Add 1 to both sides. C. Add $3x$ to both sides.
 D. Add 11 to both sides. E. Subtract $3x$ from both sides.
- Solve the following equations.
 a. $5(a + 2) = 4a + 12$ b. $0.2(b - 6) = 5.2 + b$
- MC** Choose the equation that matches the statement 'subtracting 6 from 5 multiplied by a certain number gives a result of -8 '.
 A. $6 - 5x = -8$ B. $5x - 6 = -8$ C. $5 - 6x = -8$
 D. $6x - 5 = -8$ E. $-6 - 5x = -8$
- The cost of renting a car is given by $c = 75d + 0.4k$, where d = number of days rented and k = number of kilometres driven. Toni has \$340 to spend on a car for three days. Calculate the total distance she can travel.
- Solve the value of x in the statement 'x is multiplied by 5 and then 4 is subtracted. The result is the same as three times x minus 10'.
- MC** Select the correct equation in which $\frac{y + b}{x + b} = m$ is rearranged to make x the subject.
 A. $y = \frac{4x + 3}{x}$ B. $x = \frac{m - b}{y + b}$ C. $x = \frac{y + b}{m} + b$
 D. $x = \frac{y + b - bm}{m}$ E. $x = \frac{y}{m}$



14. **MC** Select the correct equation in which $x(y - 4) = y + 3$ is rearranged to make y the subject.

A. $y = \frac{4x + 3}{x}$

B. $y = \frac{4x + 3}{x - 1}$

C. $y = 3x + 3$

D. $y = \frac{y + 3}{x} + 4$

E. $y = y + 3 - 4x$

15. Rearrange $\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$ to make:

a. a the subject

b. b the subject.

LESSON

3.2 Solving linear equations and inequations

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- recognise inverse operations
- solve multistep equations and inequations with pronumerals on one side of the equation
- solve equations and inequations with algebraic fractions.

3.2.1 What is a linear equation?

eles-4689

- An **equation** is a mathematical statement that contains an equals sign (=).
- The **expression** on the left-hand side of the equals sign has the same value as the expression on the right-hand side of the equals sign.
- Solving a **linear equation** means determining a value for the pronumeral that makes the statement true.
- ‘Doing the same thing’ to both sides of an equation (also known as applying the **inverse operation**) ensures that the expressions on either side of the equals sign remain equal, or **balanced**.

WORKED EXAMPLE 1 Using substitution to check equations

For each of the following equations, determine whether $x = 10$ is a solution.

a. $\frac{x + 2}{3} = 6$

b. $2x + 3 = 3x - 7$

c. $x^2 - 2x = 9x - 10$

THINK

a. 1. Substitute 10 for x in the left-hand side of the equation.

2. Write the right-hand side.

3. Is the equation true? Does the left-hand side equal the right-hand side?

4. State whether $x = 10$ is a solution.

WRITE

$$\begin{aligned} \text{a. LHS} &= \frac{x + 2}{3} \\ &= \frac{10 + 2}{3} \\ &= \frac{12}{3} \\ &= 4 \end{aligned}$$

$$\text{RHS} = 6$$

$$\text{LHS} \neq \text{RHS}$$

$x = 10$ is not a solution.

b. 1. Substitute 10 for x in the left-hand side of the equation.

$$\begin{aligned} \text{b. LHS} &= 2x + 3 \\ &= 2(10) + 3 \\ &= 23 \end{aligned}$$

2. Substitute 10 for x in the right-hand side of the equation.

$$\begin{aligned} \text{RHS} &= 3x - 7 \\ &= 3(10) - 7 \\ &= 23 \end{aligned}$$

3. Is the equation true? Does the left-hand side equal the right-hand side?

$$\text{LHS} = \text{RHS}$$

4. State whether $x = 10$ is a solution.

$x = 10$ is a solution.

c. 1. Substitute 10 for x in the left-hand side of the equation.

$$\begin{aligned} \text{c. LHS} &= x^2 - 2x \\ &= 10^2 - 2(10) \\ &= 100 - 20 \\ &= 80 \end{aligned}$$

2. Substitute 10 for x in the right-hand side of the equation.

$$\begin{aligned} \text{RHS} &= 9x - 10 \\ &= 9(10) - 10 \\ &= 90 - 10 \\ &= 80 \end{aligned}$$

3. Is the equation true? Does the left-hand side equal the right-hand side?

$$\text{LHS} = \text{RHS}$$

4. State whether $x = 10$ is a solution.

$x = 10$ is a solution.

3.2.2 Solving multistep equations

eles-4690

- If an equation performs two operations on a pronumeral, it is known as a **two-step equation**.
- If an equation performs more than two operations on a pronumeral, it is known as a **multistep equation**.
- To solve multistep equations, first determine the order in which the operations were performed.
- Once the order of operations is determined, perform the inverse of those operations, in the reverse order, to both sides of the equation. This will keep the equation balanced.

Inverse operations

The inverse operation of an operation has the effect of undoing the original operation.

Operation	Inverse operation
+	-
-	+
×	÷
÷	×

- Each inverse operation must be performed one step at a time.
- These steps can be applied to any equation with two or more steps, as shown in the worked examples that follow.

WORKED EXAMPLE 2 Solving equations using inverse operations

Solve the following linear equations.

a. $3x + 1 = 11$

b. $1 - \frac{x}{3} = 11$

c. $\frac{2x}{3} + \frac{x}{4} = -11$

d. $\frac{-x - 0.3}{2} = -1.1$

THINK

- a. 1. Subtract 1 from both sides of the equation.
2. Divide both sides by 3.
3. Write the value for x .
Note: It is preferable to write fractions in improper form rather than as mixed numbers.

- b. 1. Subtract 1 from both sides of the equation.

2. Multiply both sides by -3 .

3. Write the value for x .

- b. 1. Subtract 1 from both sides.

2. Multiply both sides by 3.

3. Divide both sides by -1 .

4. Write the value for x .

- c. 1. Add like terms to simplify the left-hand side of the equation. To do this, you will need to determine the lowest common denominator.

2. Multiply both sides by 12.

3. Divide both sides by 11.

4. Write the value for x .

WRITE

a. $3x + 1 = 11$
 $3x + 1 - 1 = 11 - 1$

$$\frac{3x}{3} = \frac{10}{3}$$

$$x = \frac{10}{3}$$

b. $1 - \frac{x}{3} = 11$

$$1 - \frac{x}{3} - 1 = 11 - 1$$

$$\frac{-x}{3} \times (-3) = 10 \times (-3)$$

$$x = -30$$

b. $1 - \frac{x}{3} = 11$

$$1 - \frac{x}{3} - 1 = 11 - 1$$

$$\frac{-x}{3} \times 3 = 10 \times 3$$

$$\frac{-x}{-1} = \frac{30}{-1}$$

$$x = -30$$

c. $\frac{2x}{3} + \frac{x}{4} = -11$

$$\frac{2x}{3} \times \frac{4}{4} + \frac{x}{4} \times \frac{3}{3} = -11$$

$$\frac{8x + 3x}{12} = -11$$

$$\frac{11x}{12} \times 12 = -11 \times 12$$

$$\frac{11x}{11} = \frac{-132}{11}$$

$$x = -12$$

d. 1. Multiply both sides of the equation by 2.

2. Add 0.3 to both sides.

3. Divide both sides by -1 .

4. Write the value for x .

$$\begin{aligned} \text{d.} \quad & \frac{-x-0.3}{2} = -1.1 \\ & \frac{-x-0.3}{2} \times 2 = -1.1 \times 2 \end{aligned}$$

$$-x - 0.3 + 0.3 = -2.2 + 0.3$$

$$\frac{-x}{-1} = \frac{-1.9}{-1}$$

$$x = 1.9$$

TI | THINK

a-d. On a Calculator page, press:

- MENU
- 3: Algebra
- 1: Solve

Complete the entry lines as:

$$\text{solve}(3x + 1 = 11, x)$$


$$\text{solve}\left(1 - \frac{x}{3} = 11, x\right)$$

$$\text{solve}\left(\frac{2x}{3} + \frac{x}{4} = -11, x\right)$$

$$\text{solve}\left(\frac{-x-0.3}{2} = -1.1, x\right)$$

Press ENTER after each entry.

DISPLAY/WRITE



a. $3x + 1 = 11$
 $x = \frac{10}{3}$

b. $1 - \frac{x}{3} = 11$
 $x = -30$

c. $\frac{2x}{3} + \frac{x}{4} = -11$
 $x = -12$

d. $\frac{-x-0.3}{2} = -1.1$
 $x = 1.9$

CASIO | THINK

a-d. On a standard Main screen, press:

- Action
- Advanced
- Solve

Complete the entry lines as:

$$\text{solve}(3x + 1 = 11)$$


$$\text{solve}\left(1 - \frac{x}{3} = 11\right)$$

$$\text{solve}\left(\frac{2x}{3} + \frac{x}{4} = -11\right)$$

$$\text{solve}\left(\frac{-x-0.3}{2} = -1.1\right)$$

Press EXE after each entry.
Note: As x is the only variable in each equation there is no need to specify 'solve for x ', so the ' x ' at the end of each entry line may be left out. This only applies if x is the variable.

DISPLAY/WRITE



a. $3x + 1 = 11$
 $x = \frac{10}{3}$

b. $1 - \frac{x}{3} = 11$
 $x = -30$

c. $\frac{2x}{3} + \frac{x}{4} = -11$
 $x = -12$

d. $\frac{-x-0.3}{2} = -1.1$
 $x = 1.9$



3.2.3 Algebraic fractions with the pronumeral in the denominator

eles-4691

- If an equation uses fractions and the pronumeral is in the denominator, there is an extra step involved in determining the solution.
- Consider the following equation.

$$\frac{4}{x} = \frac{3}{2}$$

- In order to solve this equation, first multiply both sides by x .

$$\frac{4}{x} \times x = \frac{3}{2} \times x$$

$$4 = \frac{3x}{2}$$

or $\frac{3x}{2} = 4$

- The pronumeral is now in the numerator, and the equation is easy to solve.

$$\frac{3x}{2} = 4$$

$$3x = 8$$

$$x = \frac{8}{3}$$

- Alternatively, the equation can be solved using the equivalent ratios method.

$$\frac{4}{x} = \frac{3}{2}$$

- This equation can be written as the equivalent ratio.

$$\frac{x}{4} = \frac{2}{3}$$

- Now the equation can be solved by multiplying both sides by 4.

$$\frac{x}{4} \times 4 = \frac{2}{3} \times 4$$

$$x = \frac{8}{3}$$

WORKED EXAMPLE 3 Solving equations with pronumerals in the denominator

Solve each of the following linear equations.

a. $\frac{3}{a} = \frac{4}{5}$

THINK

a. 1. Write the equation

2. Multiply both sides by a .

3. Multiply both sides by 5.

4. Divide both sides by 4.

b. 1. Write the equation.

2. Multiply both sides by b .

3. Divide both sides by -2 .

b. $\frac{5}{b} = -2$

WRITE

a. $\frac{3}{a} = \frac{4}{5}$

$$3 = \frac{4a}{5}$$

$$15 = 4a$$

$$a = \frac{15}{4}$$

b. $\frac{5}{b} = -2$

$$5 = -2b$$

$$\frac{5}{-2} = b \text{ or } \frac{-5}{2} = b$$

3.2.4 Solving linear inequations

eles-3869

- Compared to an equation, an inequation is a statement of inequality such as $x > 9$ (x is greater than 9).
- The solution to an inequation has many values.

Inequations on a number line

The following table displays four inequations and their representation on a number line.

Mathematical statement	English statement	Number line diagram
$x > -4$	x is greater than -4	
$x \geq -4$	x is greater than or equal to -4	
$x < -4$	x is less than -4	
$x \leq -4$	x is less than or equal to -4	

- Note that an open circle does not include the number and a solid circle does include the number.

To solve inequations:

- Imagine that the inequality sign is replaced with an equal sign.
- Solve the inequation as if it was an equation and keep the original inequality sign unless there is a special case as outlined below.

WORKED EXAMPLE 4 Solving linear inequations

Solve each of the following linear inequations and display the solution on a number line.

a. $x + 7 \leq 19$

b. $3x - 12 > 9$

THINK

a. 1. To get x by itself, subtract 7 from both sides and keep the inequality the same.

2. Display the answer on a number line.

3. Write the answer.

b. 1. Firstly, get $3x$ by itself, by adding 12 to both sides and keep the inequality the same.

WRITE/DRAW

a. $x + 7 \leq 19$

$$x + 7 - 7 \leq 19 - 7$$

$$x \leq 12$$



$$x \leq 12$$

b. $3x - 12 > 9$

$$3x - 12 + 12 > 9 + 12$$

$$3x > 21$$


2. To get x by itself divide both sides by 3.

$$3x > 21$$

$$\frac{3x}{3} > \frac{21}{3}$$

$$x > 7$$

3. Display the answer on a number line.



4. Write the answer.

$$x > 7$$

The special case – multiply or divide both sides of the inequation by a negative number

- When multiplying or dividing by a negative number, change the direction of the inequality sign.

WORKED EXAMPLE 5 Solving linear inequation with negatives

Solve the following linear inequation.

a. $-4x - 16 \leq 20$

b. $-1.6x + 4.8 \leq 12.2$

THINK

a. 1. To get $-4x$ by itself, add 16 to both sides and keep the inequality the same.

2. To get x by itself, divide both sides by -4 . Since dividing by a negative number, change the direction of the inequality sign.

3. Write the answer.

b. 1. To get $-1.6x$ by itself, subtract 4.8 from both sides and keep the inequality the same.

2. To get x by itself, divide both sides by -1.6 . Since dividing by a negative number, change the direction of the inequality sign.

3. Write the answer.

WRITE

a.
$$\begin{aligned} -4x - 16 &\leq 20 \\ -4x - 16 + 16 &\leq 20 + 16 \\ -4x &\leq 36 \end{aligned}$$

$$-4x \leq 36$$

$$\frac{-4x}{-4} \geq \frac{36}{-4}$$

$$x \geq -9$$

$$x \geq -9$$

b.
$$\begin{aligned} -1.6x + 4.8 &\leq 12.2 \\ -1.6x + 4.8 - 4.8 &\leq 12.2 - 4.8 \\ -1.6x &\leq 7.4 \end{aligned}$$

$$-1.6x \leq 7.4$$


$$\frac{-1.6x}{-1.6} \geq \frac{7.4}{-1.6}$$


$$x \geq -4.625$$

$$x \geq -4.625$$

on Resources

 **eWorkbook** Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)

 **Video eLesson** Solving linear equations (eles-1895)

 **Interactivities** Individual pathway interactivity: Solving linear equations and inequations (int-4489)
Using algebra to solve problems (int-3805)

3.2 Quick quiz **on**

3.2 Exercise

Individual pathways

PRACTISE

1, 3, 6, 8, 12, 16, 20, 21, 25, 28, 31

CONSOLIDATE

2, 4, 9, 10, 13, 14, 17, 19, 22, 23, 26, 29, 32

MASTER

5, 7, 11, 15, 18, 24, 27, 30, 33, 34

Fluency

1. **WE1** For each of the following equations, determine whether $x = 6$ is a solution.

a. $x + 3 = 7$

b. $2x - 5 = 7$

c. $x^2 - 2 = 38$

d. $\frac{6}{x} + x = 7$

e. $\frac{2(x+1)}{7} = 2$

f. $3 - x = 9$

2. For each of the following equations, determine whether $x = 6$ is a solution.

a. $x^2 + 3x = 39$

b. $3(x + 2) = 5(x - 4)$

c. $x^2 + 2x = 9x - 6$

d. $x^2 = (x + 1)^2 - 14$

e. $(x - 1)^2 = 4x + 1$

f. $5x + 2 = x^2 + 4$

3. **WE2a** Solve each of the following linear equations. Check your answers by substitution.

a. $x - 43 = 167$

b. $x + 286 = 516$

c. $58 + x = 81$

d. $209 - x = 305$

e. $5x = 185$

f. $60x = 1200$

4. **WE2b** Solve each of the following linear equations. Check your answers by substitution.

a. $5x = 250$

b. $\frac{x}{23} = 6$

c. $\frac{x}{17} = 26$

d. $\frac{x}{9} = 27$

e. $y - 16 = -31$

f. $5.5 + y = 7.3$

5. **WE2c** Solve each of the following linear equations. Check your answers by substitution.

a. $y - 7.3 = 5.5$

b. $6y = 14$

c. $0.9y = -0.05$

d. $\frac{y}{5} = 4.3$

e. $\frac{y}{7.5} = 23$

f. $\frac{y}{8} = -1.04$

6. Solve each of the following linear equations.

a. $2y - 3 = 7$

b. $2y + 7 = 3$

c. $5y - 1 = 0$

d. $6y + 2 = 8$

e. $7 + 3y = 10$

f. $8 + 2y = 12$

7. Solve each of the following linear equations.

a. $15 = 3y - 1$

b. $-6 = 3y - 1$

c. $6y - 7 = 140$

d. $4.5y + 2.3 = 7.7$

e. $0.4y - 2.7 = 6.2$

f. $600y - 240 = 143$

8. Solve each of the following linear equations.

a. $3 - 2x = 1$

b. $-3x - 1 = 5$

c. $-4x - 7 = -19$

d. $1 - 3x = 19$

e. $-5 - 7x = 2$

f. $-8 - 2x = -9$

9. Solve each of the following linear equations.

a. $9 - 6x = -1$

b. $-5x - 4.2 = 7.4$

c. $2 = 11 - 3x$

d. $-3 = -6x - 8$

e. $-1 = 4 - 4x$

f. $35 - 13x = -5$

10. Solve each of the following linear equations.

a. $7 - x = 8$

b. $8 - x = 7$

c. $5 - x = 5$

d. $5 - x = 0$

e. $15.3 = 6.7 - x$

f. $5.1 = 4.2 - x$

11. Solve each of the following linear equations.

a. $9 - x = 0.1$

b. $140 - x = 121$

c. $-30 - x = -4$

d. $-5 = -6 - x$

e. $-x + 1 = 2$

f. $-2x - 1 = 0$

12. Solve each of the following linear equations.

a. $\frac{x}{4} + 1 = 3$

b. $\frac{x}{3} - 2 = -1$

c. $\frac{x}{8} = \frac{1}{2}$

d. $-\frac{x}{3} = 5$

e. $5 - \frac{x}{2} = -8$

f. $4 - \frac{x}{6} = 11$

13. Solve each of the following linear equations.

a. $\frac{2x}{3} = 6$

b. $\frac{5x}{2} = -3$

c. $-\frac{3x}{4} = -7$

d. $-\frac{8x}{3} = 6$

e. $\frac{2x}{7} = -2$

f. $-\frac{3x}{10} = -\frac{1}{5}$

14. Solve each of the following linear equations.

a. $\frac{z-1}{3} = 5$

b. $\frac{z+1}{4} = 8$

c. $\frac{z-4}{2} = -4$

d. $\frac{6-z}{7} = 0$

e. $\frac{3-z}{2} = 6$

f. $\frac{-z-50}{22} = -2$

15. Solve each of the following linear equations.

a. $\frac{z-4.4}{2.1} = -3$

b. $\frac{z+2}{7.4} = 1.2$

c. $\frac{140-z}{150} = 0$

d. $\frac{-z-0.4}{2} = -0.5$

e. $\frac{z-6}{9} = -4.6$

f. $\frac{z+65}{73} = 1$

16. Solve each of the following linear equations.

a. $\frac{5x+1}{3} = 2$

b. $\frac{2x-5}{7} = 3$

c. $\frac{3x+4}{2} = -1$

d. $\frac{4x-13}{9} = -5$

e. $\frac{4-3x}{2} = 8$

f. $\frac{1-2x}{6} = -10$

17. Solve each of the following linear equations.

a. $\frac{-5x-3}{9} = 3$

b. $\frac{-10x-4}{3} = 1$

c. $\frac{4x+2.6}{5} = 8.8$

d. $\frac{5x-0.7}{-0.3} = -3.1$

e. $\frac{1-0.5x}{4} = -2.5$

f. $\frac{-3x-8}{14} = \frac{1}{2}$

18. **WE3** Solve each of the following linear equations.

a. $\frac{2}{x} = \frac{1}{2}$

b. $\frac{3}{x} = 7$

c. $\frac{-4}{x} = \frac{7}{2}$

d. $\frac{5}{x} = \frac{-3}{4}$

e. $\frac{0.4}{x} = \frac{9}{2}$

f. $\frac{8}{x} = 1$

19. Solve each of the following linear equations.

a. $\frac{-4}{x} = \frac{2}{3}$

b. $\frac{-6}{x} = \frac{-4}{5}$

c. $\frac{1.7}{x} = \frac{1}{3}$

d. $\frac{6}{x} = -1$

e. $\frac{4}{x} = \frac{-15}{22}$

f. $\frac{50}{x} = \frac{-35}{43}$

20. Solve each of the following linear equations.

a. $3a + 7 = 4$

b. $5 - b = -5$

c. $4c - 4.4 = 44$

d. $\frac{d-4}{67} = 0$

e. $5 - 3e = -10$

f. $\frac{2f}{3} = 8$

21. **WE4** Solve each of the following linear inequations and display the solution on a number line.

a. $x + 12 \leq 28$

b. $x - 3 > 14$

22. Solve each of the following linear inequations.

a. $2x + 6 \leq 24$

b. $5x - 28 > 22$

23. **WE5** Solve the following linear inequations.

a. $-5x - 35 \leq 30$

b. $-8x + 21 < -35$

24. Solve the following linear inequations.

a. $-2.5x - 15 \leq 20$

b. $-1.4x - 13.1 < -38.3$

Understanding

25. Write each of the following worded statements as a mathematical sentence and then solve for the unknown value.

a. Seven is added to the product of x and 3, which gives the result of 4.

b. Four is divided by x and this result is equivalent to $\frac{2}{3}$.

c. Three is subtracted from x and this result is divided by 12 to give 25.

26. Driving lessons are usually quite expensive, but a discount of \$15 per lesson is given if a family member belongs to the automobile club.

If 10 lessons cost \$760 after the discount, calculate the cost of each lesson before the discount.

27. Anton lives in Australia. His friend Utan lives in the USA. Anton's home town of Horsham experienced one of the hottest days on record with a temperature of 46.7°C .

Utan said that his home town had experienced a day hotter than that, with the temperature reaching 113°F .

The formula for converting Celsius to Fahrenheit is $F = \frac{9}{5}C + 32$.

State whether Utan was correct. Show full working.

Reasoning

28. If the expression $\frac{12}{x-4}$ always results in a positive integer value, explain how you would determine the possible values for x .
29. Santo solved the linear equation $9 = 5 - x$. His second step was to divide both sides by -1 . Trudy, his mathematics buddy, said she multiplied both sides by -1 . Explain why they are both correct.
30. Determine the mistake in the following working and explain what is wrong.


$$\begin{aligned}\frac{x}{5} - 1 &= 2 \\ x - 1 &= 10 \\ x &= 11\end{aligned}$$

Problem solving

31. Sweet-tooth Sami goes to the corner store and buys an equal number of 25-cent and 30-cent lollies for a total of \$16.50. Determine the amount of lollies he bought.



32. In a cannery, cans are filled by two machines that together produce 16 000 cans during an 8-hour shift. If the newer machine of the two produces 340 more cans per hour than the older machine, evaluate the number of cans produced by each machine in an eight-hour shift.
33. General admission to a music festival is \$55 for an adult ticket, \$27 for a child and \$130 for a family of two adults and two children.
- Evaluate how much you would save by buying a family ticket instead of two adult and two child tickets.
 - Determine if it is worthwhile buying a family ticket if a family has only one child.
34. A teacher asks her students to determine the value of n in the diagram shown. Use your knowledge of linear equations to solve this problem.



LESSON

3.3 Solving linear equations with brackets

LEARNING INTENTION

At the end of this lesson you should be able to:

- solve equations involving brackets.

3.3.1 Linear equations with brackets

eles-4692

- Consider the equation $3(x + 5) = 18$. There are two methods for solving this equation.

Method 1:

Start by dividing both sides by 3.

$$\begin{aligned}\frac{3(x+5)}{3} &= \frac{18}{3} \\ x+5 &= 6 \\ x &= 1\end{aligned}$$

Method 2:

Start by expanding the brackets.

$$\begin{aligned}3(x+5) &= 18 \\ 3x+15 &= 18 \\ 3x &= 3 \\ x &= 1\end{aligned}$$

- In this case, method 1 works well because 3 divides exactly into 18.
- Now try the equation $7(x + 2) = 10$.

Method 1:

Start by dividing both sides by 7.

$$\begin{aligned}\frac{7(x+2)}{7} &= \frac{10}{7} \\ x+2 &= \frac{10}{7} \\ x &= \frac{10}{7} - 2 \\ x &= -\frac{4}{7}\end{aligned}$$

Method 2:

Start by expanding the brackets.

$$\begin{aligned}7(x+2) &= 10 \\ 7x+14 &= 10 \\ 7x &= -4 \\ x &= \frac{-4}{7}\end{aligned}$$

- In this case, method 2 works well because it avoids the addition or subtraction of fractions.
- For each equation you need to solve, try both methods and choose the method that works best.

WORKED EXAMPLE 6 Solving equations with brackets

Solve each of the following linear equations.

a. $7(x - 5) = 28$

b. $6(x + 3) = 7$

THINK

1. 7 is a factor of 28, so it will be easier to divide both sides by 7.
2. Add 5 to both sides.
3. Write the value of x .

WRITE

a. $7(x - 5) = 28$

$$\frac{7(x-5)}{7} = \frac{28}{7}$$
$$x - 5 + 5 = 4 + 5$$
$$x = 9$$

b. 1. 6 is not a factor of 7, so it will be easier to expand the brackets first.

2. Subtract 18 from both sides.

3. Divide both sides by 6.

b. $6(x + 3) = 7$
 $6x + 18 = 7$

$$6x + 18 - 18 = 7 - 18$$

$$6x = -11$$

$$x = -\frac{11}{6}$$

TI | THINK


a-b. In a new problem, on a Calculator page, press:

- MENU
- 3: Algebra
- 1: Solve

Complete the entry lines as:

solve
 $(7(x - 5) = 28, x)$
 solve
 $(6(x + 3) = 7, x)$
 Press ENTER after each entry.

DISPLAY/WRITE



a. $x = 9$
 b. $x = -\frac{11}{6}$

CASIO | THINK


a-b. On the Main screen, press:

- Action
- Advanced
- Solve

Complete the entry lines as:

solve
 $(7(x - 5) = 28)$
 solve
 $(6(x + 3) = 7)$
 Press EXE after each entry.

DISPLAY/WRITE



a. $x = 9$
 b. $x = -\frac{11}{6}$

on Resources



eWorkbook Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)



Interactivities Individual pathway interactivity: Solving linear equations with brackets (int-4490)
 Linear equations with brackets (int-6039)

3.3 Quick quiz **on**

3.3 Exercise

Individual pathways

PRACTISE

1, 3, 5, 11, 13, 16

CONSOLIDATE

2, 7, 9, 12, 14, 17

MASTER

4, 6, 8, 10, 15, 18

Fluency

1. **WE6** Solve each of the following linear equations.

a. $5(x - 2) = 20$

b. $4(x + 5) = 8$

c. $6(x + 3) = 18$

d. $5(x - 41) = 75$

e. $8(x + 2) = 24$

f. $3(x + 5) = 15$

2. Solve each of the following linear equations.

a. $5(x + 4) = 15$

b. $3(x - 2) = -12$

c. $7(x - 6) = 0$

d. $-6(x - 2) = 12$

e. $4(x + 2) = 4.8$

f. $16(x - 3) = 48$

3. Solve each of the following linear equations.

a. $6(b - 1) = 1$

b. $2(m - 3) = 3$

c. $2(a + 5) = 7$

d. $3(m + 2) = 2$

e. $5(p - 2) = -7$

f. $6(m - 4) = -8$

4. Solve each of the following linear equations.

a. $-10(a + 1) = 5$

b. $-12(p - 2) = 6$

c. $-9(a - 3) = -3$

d. $-2(m + 3) = -1$

e. $3(2a + 1) = 2$

f. $4(3m + 2) = 5$

5. Solve each of the following linear equations.

a. $9(x - 7) = 82$

b. $2(x + 5) = 14$

c. $7(a - 1) = 28$

d. $4(b - 6) = 4$

e. $3(y - 7) = 0$

f. $-3(x + 1) = 7$

6. Solve each of the following linear equations.

a. $-6(m + 1) = -30$

b. $-4(y + 2) = -12$

c. $-3(a - 6) = 3$

d. $-2(p + 9) = -14$

e. $3(2m - 7) = -3$

f. $2(4p + 5) = 18$

7. Solve each of the following linear equations. Round the answers to 3 decimal places where appropriate.

a. $2(y + 4) = -7$

b. $0.3(y + 8) = 1$

c. $4(y + 19) = -29$

d. $7(y - 5) = 25$

e. $6(y + 3.4) = 3$

f. $7(y - 2) = 8.7$

8. Solve each of the following linear equations. Round the answers to 3 decimal places where appropriate.

a. $1.5(y + 3) = 10$

b. $2.4(y - 2) = 1.8$

c. $1.7(y + 2.2) = 7.1$

d. $-7(y + 2) = 0$

e. $-6(y + 5) = -11$

f. $-5(y - 2.3) = 1.6$

9. **MC** Select the best first step for solving the equation $7(x - 6) = 23$.

A. Add 6 to both sides.

B. Subtract 7 from both sides.

C. Divide both sides by 23.

D. Expand the brackets.

E. Multiply both sides by 7.

10. **MC** Select which one of the following is closest to the solution for the equation $84(x - 21) = 782$.

A. $x = 9.31$

B. $x = 9.56$

C. $x = 30.31$

D. $x = -11.69$

E. $x = 30.32$

Understanding

11. In 1974 a mother was 6 times as old as her daughter. If the mother turned 50 in the year 2000, calculate the year in which the mother's age was double her daughter's age.
12. New edging is to be placed around a rectangular children's playground. The width of the playground is x metres and the length is 7 metres longer than the width.
- Write down an expression for the perimeter of the playground. Write your answer in factorised form.
 - If the amount of edging required is 54 m, determine the dimensions of the playground.



Reasoning

13. Explain the two possible methods for solving equations in factorised form.
14. Juanita is solving the equation $2(x - 8) = 10$. She performs the following operations to both sides of the equation in order: $+ 8, \div 2$. Explain why Juanita will not be able to determine the correct value of x using her order of inverse operations, then solve the equation correctly.
15. As your first step to solve the equation $3(2x - 7) = 18$, you are given three options:
- Expand the brackets on the left-hand side.
 - Add 7 to both sides.
 - Divide both sides by 3.
- Explain which of these options is your least preferred.

Problem solving

16. Five times the sum of four and an unknown number is equal to 35. Evaluate the value of the unknown number.
17. Oscar earns \$55 more than Josue each week, but Hector earns three times as much as Oscar. If Hector earns \$270 a week, determine how much Oscar and Josue earn each week.
18. A school wants to hire a bus to travel to a football game. The bus will take 28 passengers, and the school will contribute \$48 towards the cost of the trip. The price of each ticket is \$10. If the hiring of the bus is $\$300 + 10\%$ of the total cost of all the tickets, evaluate the cost per person.



LESSON

3.4 Solving linear equations with pronumerals on both sides

LEARNING INTENTION

At the end of this lesson you should be able to:

- solve equations by collecting pronumerals on one side of the equation before solving.



3.4.1 Linear equations with pronumerals on both sides

eles-4693

- If pronumerals occur on both sides of an equation, the first step in solving the equation is to move all of the pronumerals so that they appear on only one side of the equation.
- When solving equations, it is important to remember that whatever you do to one side of an equation you must also do to the other side.

WORKED EXAMPLE 7 Solving equations with pronumerals on both sides

Solve each of the following linear equations.

a. $5y = 3y + 3$

c. $3(x + 1) = 14 - 2x$

b. $7x + 5 = 2 - 4x$

d. $2(x + 3) = 3(x + 7)$

THINK

a. 1. $3y$ is smaller than $5y$. This means it is easiest to subtract $3y$ from both sides of the equation.

2. Divide both sides by 2.

b. 1. $-4x$ is smaller than $7x$. This means it is easiest to add $4x$ to both sides of the equation.

2. Subtract 5 from both sides.

3. Divide both sides by 11.

c. 1. Expand the brackets.

2. $-2x$ is smaller than $3x$. This means it is easiest to add $2x$ to both sides of the equation.

WRITE

a.
$$\begin{aligned} 5y &= 3y + 3 \\ 5y - 3y &= 3y + 3 - 3y \\ 2y &= 3 \end{aligned}$$

$$\begin{aligned} \frac{2y}{2} &= \frac{3}{2} \\ y &= \frac{3}{2} \end{aligned}$$

b.
$$\begin{aligned} 7x + 5 &= 2 - 4x \\ 7x + 5 + 4x &= 2 - 4x + 4x \\ 11x + 5 &= 2 \end{aligned}$$

$$\begin{aligned} 11x + 5 - 5 &= 2 - 5 \\ 11x &= -3 \end{aligned}$$

$$\begin{aligned} \frac{11x}{11} &= \frac{-3}{11} \\ x &= \frac{-3}{11} \end{aligned}$$

c.
$$\begin{aligned} 3(x + 1) &= 14 - 2x \\ 3x + 3 &= 14 - 2x \\ 3x + 3 + 2x &= 14 - 2x + 2x \\ 5x + 3 &= 14 \end{aligned}$$

3. Subtract 3 from both sides.

$$5x + 3 - 3 = 14 - 3$$

$$5x = 11$$

4. Divide both sides by 5.

$$\frac{5x}{5} = \frac{11}{5}$$

$$x = \frac{11}{5}$$

d. 1. Expand the brackets.

d.

$$2(x + 3) = 3(x + 7)$$

$$2x + 6 = 3x + 21$$

2. $2x$ is smaller than $3x$. This means it is easiest to subtract $2x$ from both sides of the equation.

$$2x + 6 - 2x = 3x + 21 - 2x$$

$$6 = x + 21$$

3. Subtract 21 from both sides.

$$6 - 21 = x + 21 - 21$$

$$-15 = x$$

4. Write the answer with the pronumeral written on the left-hand side.

$$x = -15$$

TI | THINK

a-d. In a new problem, on a Calculator page, press:

- MENU
- 3: Algebra
- 1: Solve

Complete the entry lines as:

solve ($5y = 3y + 3, y$)


solve ($7x + 5 = 2 - 4x, x$)

solve ($3(x + 1) = 14 - 2x, x$)

solve ($2(x + 3) = 3(x + 7), x$)

Press ENTER after each entry.

DISPLAY/WRITE



- a. $y = \frac{3}{2}$
- b. $x = -\frac{3}{11}$
- c. $x = \frac{11}{5}$
- d. $x = -15$

CASIO | THINK

a-d. On the Main screen, press:

- Action
- Advanced
- Solve

Complete the entry lines as:

solve ($5y = 3y + 3, y$)


solve ($7x + 5 = 2 - 4x, x$)

solve ($3(x + 1) = 14 - 2x, x$)

solve ($2(x + 3) = 3(x + 7), x$)

Press EXE after each entry. *Note:* If the variable is *not* x , it must be specified.




DISPLAY/WRITE



- a. $y = \frac{3}{2}$
- b. $x = -\frac{3}{11}$
- c. $x = \frac{11}{5}$
- d. $x = -15$

DISCUSSION

Draw a diagram that could represent $2x + 4 = 3x + 1$.

-  **eWorkbook** Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)
-  **Video eLesson** Solving linear equations with pronumerals on both sides (eles-1901)
-  **Interactivity** Individual pathway interactivity: Solving linear equations with pronumerals on both sides (int-4491)

Exercise 3.4 Solving linear equations with pronumerals on both sides

3.4 Quick quiz **on**

3.4 Exercise

Individual pathways

PRACTISE

1, 3, 5, 8, 12, 14, 17

CONSOLIDATE

2, 6, 10, 13, 15, 18

MASTER

4, 7, 9, 11, 16, 19, 20

Fluency

1. **WE7a** Solve each of the following linear equations.

a. $5y = 3y - 2$	b. $6y = -y + 7$	c. $10y = 5y - 15$
d. $25 + 2y = -3y$	e. $8y = 7y - 45$	f. $15y - 8 = -12y$
2. Solve each of the following linear equations.

a. $7y = -3y - 20$	b. $23y = 13y + 200$	c. $5y - 3 = 2y$
d. $6 - 2y = -7y$	e. $24 - y = 5y$	f. $6y = 5y - 2$
3. **MC** Select the first step for solving the equation $3x + 5 = -4 - 2x$.

A. Add $3x$ to both sides.	B. Add 5 to both sides.	C. Add $2x$ to both sides.
D. Subtract $2x$ from both sides.	E. Divide both sides by 3.	
4. **MC** Select the first step for solving the equation $6x - 4 = 4x + 5$.

A. Subtract $4x$ from both sides.	B. Add $4x$ to both sides.	C. Subtract 4 from both sides.
D. Add 5 to both sides.	E. Divide both sides by 6.	
5. **WE7b** Solve each of the following linear equations.

a. $2x + 3 = 8 - 3x$	b. $4x + 11 = 1 - x$	c. $x - 3 = 6 - 2x$
d. $4x - 5 = 2x + 3$	e. $3x - 2 = 2x + 7$	f. $7x + 1 = 4x + 10$
6. Solve each of the following linear equations.

a. $5x + 3 = x - 5$	b. $6x + 2 = 3x + 14$	c. $2x - 5 = x - 9$
d. $10x - 1 = -2x + 5$	e. $7x + 2 = -5x + 2$	f. $15x + 3 = 7x - 3$
7. Solve each of the following linear equations.

a. $x - 4 = 3x + 8$	b. $2x + 9 = 7x - 1$	c. $-2x + 7 = 4x + 19$
d. $-3x + 2 = -2x - 11$	e. $11 - 6x = 18 - 5x$	f. $6 - 9x = 4 + 3x$
8. **MC** Determine the solution to $5x + 2 = 2x + 23$.

A. $x = 3$	B. $x = -3$	C. $x = 5$	D. $x = 7$	E. $x = -5$
------------	-------------	------------	------------	-------------
9. **MC** Determine the solution to $3x - 4 = 11 - 2x$.

A. $x = 15$	B. $x = 7$	C. $x = 3$	D. $x = 5$	E. $x = -3$
-------------	------------	------------	------------	-------------

10. **WE7c,d** Solve each of the following.

a. $5(x - 2) = 2x + 5$

b. $7(x + 1) = x - 11$

c. $2(x - 8) = 4x$

d. $3(x + 5) = x$

e. $6(x - 3) = 14 - 2x$

f. $9x - 4 = 2(3 - x)$

11. Solve each of the following.

a. $4(x + 3) = 3(x - 2)$

b. $5(x - 1) = 2(x + 3)$

c. $8(x - 4) = 5(x - 6)$

d. $3(x + 6) = 4(2 - x)$

e. $2(x - 12) = 3(x - 8)$

f. $4(x + 11) = 2(x + 7)$

Understanding

12. Aamir's teacher gave him the following algebra problem and told him to solve it.

$$3x + 7 = x^2 + k = 7x + 15$$


Suggest how you can help Aamir calculate the value of k .

13. A classroom contained an equal number of boys and girls. Six girls left to play hockey, leaving twice as many boys as girls in the classroom.

Determine the original number of students present.

Reasoning

14. Express the information in the diagram shown as an equation, then show that $n = 29$ is the solution.



15. The block shown has a width of 1 unit and length of $(x + 1)$ units.



a. Draw two rectangles with different sizes, each using 3 of the blocks shown.

b. Show that the areas of both of the rectangles you have drawn using the blocks shown are the same. Explain how the areas of these rectangles relate to expanding brackets.

16. Explain what the difficulty is when trying to solve the equation $4(3x - 5) = 6(4x + 2)$ without expanding the brackets first.

Problem solving

17. This year Tom is 4 times as old as his daughter. In 5 years' time he will be only 3 times as old as his daughter. Determine the age of Tom and his daughter now.

18. If you multiply an unknown number by 6 and then add 5, the result is 7 less than the unknown number plus 1, multiplied by 3. Evaluate the unknown number.



19. You are looking into getting a business card printed for your new game store. A local printing company charges \$250 for the materials and an hourly rate for labour of \$40.



- a. If h is the number of hours of labour required to print the cards, construct an equation for the cost of the cards, C .
- b. You have budgeted \$1000 for this printing job. Determine the number of hours of labour that you can afford. Give your answer to the nearest minute.
- c. The company estimates that it can print 1000 cards an hour. Determine the number of cards that you can afford to get printed with your current budget.
- d. An alternative to printing is photocopying. The company charges 15 cents per side for the first 10 000 cards, then 10 cents per side for additional cards. Determine the cheaper option to get 18 750 single-sided cards made. Determine how much cheaper this is than the other option.
20. A local games arcade offers its regular customers the following deal. For a monthly fee of \$40, players can play 25 \$2 games of their choice. Extra games cost \$2 each. After a player has played 50 games in a month, all further games are \$1.
- a. If Iman has \$105 to spend in a month, determine the number of games she can play if she takes up the special deal.
- b. Determine the amount of money Iman will save by taking up the special deal compared to playing the same number of games at \$2 a game.



LESSON

3.5 Solving problems with linear equations

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- define a pronumeral and write an equation to represent a problem
- solve a problem and answer the question (or questions) posed in the problem.



3.5.1 Solving problems with linear equations

eles-4694

- When pronumerals are not provided in a problem, you need to introduce a pronumeral to the problem.
- When translating a worded problem into an equation, make sure that the order of the operations is presented correctly.

WORKED EXAMPLE 8 Translating worded expressions into linear equations

Write linear equations for each of the following statements, using x to represent the unknown value. Solve the equations.

- Three more than seven times a certain number is zero. Determine the number.
- The sum of three consecutive integers is 102. Determine the three numbers.
- The difference between two numbers is 11. Determine the numbers if, when eight is added to each of the two numbers, the larger result will be double the value of the smaller result.

THINK

1. Let x be the number.
2. 7 times the number is $7x$. Increasing the value of $7x$ by 3 gives $7x + 3$. This expression equals 0. Write the equation.

3. Solve the equation using inverse operations.

4. Write the value for x .

1. Let x be the first number.
2. The next two consecutive integers are $(x + 1)$ and $(x + 2)$. The sum of all three integers equals 102. Write the equation.

3. Solve the equation using inverse operations.

4. Write the value for x and the next 2 integers.

1. Let x be the larger number.
2. The difference between the 2 numbers is 11. This means the smaller number is 11 less than x . Write this as $(x - 11)$.
3. Increasing both numbers by 8 makes the larger number $(x + 8)$ and the smaller number $(x - 11 + 8)$.
The larger number becomes double the smaller number.

4. Solve the equation using inverse operations.

5. Write the value of x and the other number.

WRITE

- a. $x =$ unknown number

$$7x + 3 = 0$$

$$7x + 3 - 3 = 0 - 3$$

$$\frac{7x}{7} = \frac{-3}{7}$$

$$x = \frac{-3}{7}$$

- b. $x =$ smallest number

$$x + (x + 1) + (x + 2) = 102$$

$$3x + 3 - 3 = 102 - 3$$

$$\frac{3x}{3} = \frac{99}{3}$$

$$x = 33$$

The three numbers are 33, 34 and 35.

- c. $x =$ larger number

$$(x - 11) = \text{smaller number}$$

$$(x + 8) = 2(x - 11 + 8)$$

$$x + 8 = 2(x - 3)$$

$$x + 8 - 8 = 2x - 6 - 8$$

$$x - x = 2x - 14 - x$$

$$0 + 14 = x - 14 + 14$$

$$x = 14$$

The two numbers are 14 and 3.

WORKED EXAMPLE 9 Solving problems using linear equations

A taxi charges \$3.60 plus \$1.38 per kilometre for any trip in Melbourne. If Elena's taxi ride cost \$38.10, calculate the distance she travelled.



THINK

1. The distance travelled by Elena has to be found. Define the pronumeral.
2. It costs 1.38 to travel 1 kilometre, so the cost to travel x kilometres = $1.38x$. The fixed cost is \$3.60. Write an expression for the total cost.
3. Let the total cost = 38.10.
4. Solve the equation.
5. Write the solution in words.

WRITE

Let x = the distance travelled (in kilometres).

$$\text{Total cost} = 3.60 + 1.38x$$

$$3.60 + 1.38x = 38.10$$

$$3.60 + 1.38x = 38.10$$

$$1.38x = 34.50$$

$$x = \frac{34.50}{1.38}$$

$$= 25$$

Elena travelled 25 kilometres.

WORKED EXAMPLE 10 Solving problems using linear equations

In a basketball game, Hao scored 5 more points than Seve. If they scored a total of 27 points between them, calculate how many points each of them scored.



THINK

1. Define the pronumeral.
2. Hao scored 5 more than Seve. Write an expression for Hao's score.
3. Hao and Seve scored a total of 27 points. Write this as an equation.
4. Solve the equation.

WRITE

Let Seve's score be x .

Hao's score is $x + 5$.

$$x + (x + 5) = 27$$

$$2x + 5 = 27$$

$$2x = 22$$

$$x = 11$$

5. Since $x = 11$, Seve's score is 11. Substitute this into the expression to work out Hao's score.
6. Write the answer in words.

$$\begin{aligned}\text{Hao's score} &= x + 5 \\ &= 11 + 5 \\ &= 16\end{aligned}$$

Seve scored 11 points and Hao scored 16 points.

WORKED EXAMPLE 11 Solving problems using linear equations

A collection of 182 marbles is owned by four friends. Pat has twice the number of marbles that Quentin has, and Rachel has 20 fewer marbles than Pat. If Sam has two-thirds the number of marbles that Rachel has, determine who has the second-largest number of marbles.

THINK

- Let p represent the number of marbles Pat has.

Pat has twice the number of marbles that Quentin has.
Rachel has 20 fewer marbles than Pat.
Sam has $\frac{2}{3}$ of the number of marbles that Rachel has.
- Let the sum of all marbles equal 182.
- Simplify the left-hand side of the equation by adding like terms.
- Solve using inverse operations.
- Calculate the number of marbles that each friend has.
- Write down the second-highest number of marbles and the name of the person with that number of marbles.

WRITE

$$\begin{aligned}p &= \text{number of Pat's marbles} \\ \left(\frac{p}{2}\right) &= \text{number of Quentin's marbles} \\ p - 20 &= \text{number of Rachel's marbles} \\ \frac{2}{3}(p - 20) &= \text{number of Sam's marbles} \\ p + \left(\frac{p}{2}\right) + (p - 20) + \frac{2}{3}(p - 20) &= 182 \\ p + \frac{p}{2} + p + \frac{2p}{3} - 20 - \frac{40}{3} &= 182 \\ \frac{6p}{6} + \frac{3p}{6} + \frac{6p}{6} + \frac{4p}{6} - \frac{120}{6} - \frac{80}{6} &= 182 \\ \frac{19p - 200}{6} \times 6 &= 182 \times 6 \\ 19p - 200 + 200 &= 1092 + 200 \\ \frac{19p}{19} &= \frac{1292}{19} \\ p &= 68 \\ \text{Pat has } 68 \text{ marbles.} \\ \text{Quentin has } \frac{68}{2} &= 34 \text{ marbles.} \\ \text{Rachel has } 68 - 20 &= 48 \text{ marbles.} \\ \text{Sam has } \frac{2}{3} \times 48 &= 32 \text{ marbles.} \\ \text{Rachel has the second-highest number of} \\ \text{marbles. She has } 48 \text{ marbles.}\end{aligned}$$

DISCUSSION

Why is it important to define the pronumeral when using a linear equation to solve a problem?

on Resources



eWorkbook Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)



Interactivity Individual pathway interactivity: Solving problems with linear equations (int-4492)

Exercise 3.5 Solving problems with linear equations

learnon

3.5 Quick quiz **on**

3.5 Exercise

Individual pathways

■ PRACTISE

1, 3, 7, 11, 12, 16

■ CONSOLIDATE

2, 4, 8, 9, 13, 17

■ MASTER

5, 6, 10, 14, 15, 18

Fluency

- WE8** Write linear equations for each of the following statements, using x to represent the unknown, without solving the equations.
 - When 3 is added to a certain number, the answer is 5.
 - Subtracting 9 from a certain number gives a result of 7.
 - Multiplying a certain number by 7 gives 24.
 - A certain number divided by 5 gives a result of 11.
 - Dividing a certain number by 2 equals -9 .
- Write linear equations for each of the following statements, using x to represent the unknown, without solving the equations.
 - Subtracting 3 from 5 times a certain number gives a result of -7 .
 - When a certain number is subtracted from 14 and the result is then multiplied by 2, the final result is -3 .
 - When 5 is added to 3 times a certain number, the answer is 8.
 - When 12 is subtracted from 2 times a certain number, the result is 15.
 - The sum of 3 times a certain number and 4 is divided by 2. This gives a result of 5.
- MC** Select the equation that matches the following statement.
A certain number, when divided by 2, gives a result of -12 .
 - $x = \frac{-12}{2}$
 - $2x = -12$
 - $\frac{x}{2} = -12$
 - $\frac{x}{12} = -2$
 - $\frac{2}{x} = 12$
- MC** Select the equation that matches the following statement.
Dividing 7 times a certain number by -4 equals 9.
 - $\frac{x}{-4} = 9$
 - $\frac{-4x}{7} = 9$
 - $\frac{7+x}{-4} = 9$
 - $\frac{7x}{-4} = 9$
 - $\frac{7x}{-4} = -9$
- MC** Select the equation that matches the following statement.
Subtracting twice a certain number from 8 gives 12.
 - $2x - 8 = 12$
 - $8 - 2x = 12$
 - $2 - 8x = 12$
 - $8 - (x + 2) = 12$
 - $2(8 - x) = 12$


6. **MC** Select the equation that matches the following statement.

When 15 is added to a quarter of a number, the answer is 10.

A. $15 + 4x = 10$ B. $10 = \frac{x}{4} + 15$ C. $\frac{x + 15}{4} = 10$ D. $15 + \frac{4}{x} = 10$ E. $15 = \frac{x}{4} + 10$


Understanding

7. When a certain number is added to 3 and the result is multiplied by 4, the answer is the same as when that same number is added to 4 and the result is multiplied by 3. Determine the number.
8. **WE9** John is three times as old as his son Jack. The sum of their ages is 48. Calculate how old John is.
9. In one afternoon's shopping Seedevi spent half as much money as Georgia, but \$6 more than Amy. If the three of them spent a total of \$258, calculate how much Seedevi spent.
10. The rectangular blocks of land shown have the same area. Determine the dimensions of each block and use these dimensions to calculate the area.



Reasoning

11. **WE10** A square pool is surrounded by a paved area that is 2 metres wide. If the area of the paving is 72 m^2 , determine the length of the pool. Show all working.



12. Maria is paid \$11.50 per hour plus \$7 for each jacket that she sews. If she earned \$176 for a particular eight-hour shift, determine the number of jackets she sewed during that shift.
13. Mai hired a car for a fee of \$120 plus \$30 per day. Casey's rate for his car hire was \$180 plus \$26 per day. If their final cost and rental period is the same for both Mai and Casey, determine the rental period.
14. **WE11** The cost of producing an album on CD is quoted as \$1200 plus \$0.95 per CD. If Maya has a budget of \$2100 for her debut album, determine the number of CDs she can make.



15. Joseph wants to have some flyers for his grocery business delivered. Post Quick quotes a price of \$200 plus 50 cents per flyer, and Fast Box quotes \$100 plus 80 cents per flyer.
- If Joseph needs to order 1000 flyers, determine the distributor that would be cheaper to use.
 - Determine the number of flyers that Joseph need to get delivered for the cost to be the same for either distributor.

Problem solving

16. A certain number is multiplied by 8 and then 16 is subtracted. The result is the same as 4 times the original number minus 8. Evaluate the number.

17. Carmel sells three different types of healthy drinks: herbal, vegetable and citrus fizz. In an hour she sells 4 herbal, 3 vegetable and 6 citrus fizz drinks for \$60.50.

In the following hour she sells 2 herbal, 4 vegetable and 3 citrus fizz drinks. In the third hour she sells 1 herbal, 2 vegetable and 4 citrus.

The total amount in cash sales for the three hours is \$136.50.


Carmel made \$7 less in the third hour than she did in the second hour of sales.

Evaluate Carmel's sales in the fourth hour if she sells 2 herbal, 3 vegetable and 4 citrus fizz drinks.



18. A rectangular swimming pool is surrounded by a path that is enclosed by a pool fence. All measurements given are in metres and are not to scale in the diagram shown.

- Write an expression for the area of the entire fenced-off section.
- Write an expression for the area of the path surrounding the pool.
- If the area of the path surrounding the pool is 34 m^2 , evaluate the dimensions of the swimming pool.
- Determine the fraction of the fenced-off area taken up by the pool.



LESSON

3.6 Rearranging formulas

LEARNING INTENTION

At the end of this lesson you should be able to:

- rearrange formulas with two or more pronumerals to make only one of the pronumerals the subject.



3.6.1 Rearranging formulas

eles-4695

- Formulas are equations that are used for specific purposes. They are generally written in terms of two or more pronumerals (also known as 'variables').
- One pronumeral is usually written on the left of the equals sign. This pronumeral is called the **subject**.
- Solving a linear equation requires determining a value for the pronumeral (or pronumerals), but rearranging formulas does not. For this reason formulas are defined as **literal equations**.

Inverse operations

When rearranging formulas, use inverse operations in reverse order, just as you do when you are solving linear equations.

For example, $C = 2\pi r$ is a formula describing the relationship of the circumference of a circle (C) to its radius (r). To make r the subject of this formula, the following steps are required:

Divide both sides by 2π .	$C = 2\pi r$ $\frac{C}{2\pi} = \frac{2\pi r}{2\pi}$ $\frac{C}{2\pi} = r$
Now r is the subject.	$r = \frac{C}{2\pi}$

DISCUSSION

In pairs, share some formulas that you know. Discuss each of the formulas by thinking about why we need them, how and when they are used in the real world, and why rearranging the pronumerals in these formulas can be helpful.

WORKED EXAMPLE 12 Rearranging formulas

Rearrange each formula to make x the subject.

a. $y = kx + m$

b. $6(y + 1) = 7(x - 2)$

THINK

a. 1. Subtract m from both sides.

2. Divide both sides by k .

3. Rewrite the equation so that x is on the left-hand side.

b. 1. Expand the brackets.

WRITE

a.
$$y = kx + m$$
$$y - m = kx$$

$$\frac{y - m}{k} = \frac{kx}{k}$$

$$\frac{y - m}{k} = x$$

$$x = \frac{y - m}{k}$$

b.
$$6(y + 1) = 7(x - 2)$$
$$6y + 6 = 7x - 14$$

2. Add 14 to both sides.

$$6y + 6 + 14 = 7x - 14 + 14$$

$$6y + 20 = 7x$$

3. Divide both sides by 7.

$$\frac{6y + 20}{7} = \frac{7x}{7}$$

$$\frac{6y + 20}{7} = x$$

4. Rewrite the equation so that x is on the left-hand side.

$$x = \frac{6y + 20}{7}$$

TI | THINK

a-b. In a new problem, on a Calculator page, press:


- MENU
- 3: Algebra
- 1: Solve

Complete the entry lines as:

solve
 $(y = k \times x + m, x)$
 solve $(6(y + 1) = 7(x - 2), x)$

Press ENTER after each entry.

DISPLAY/WRITE



$$x = \frac{y - m}{k}$$

$$x = \frac{6y + 20}{7}$$

CASIO | THINK

a-b. On the Main screen, press:


- Action
- Advanced
- Solve

Complete the entry lines as:

solve
 $(y = k \times x + m, x)$
 solve $(6(y + 1) = 7(x - 2), x)$

Press EXE after each entry.

DISPLAY/WRITE



$$x = \frac{y - m}{k}$$

$$x = \frac{6y + 20}{7}$$

WORKED EXAMPLE 13 Solving literal equations

a. If $g = 6d - 3$, solve for d .

b. Solve for v , given that $a = \frac{v - u}{t}$.

THINK

a. 1. Add 3 to both sides.

2. Divide both sides by 6.

3. Rewrite the equation so that d is on the left-hand side.

WRITE

a. $g = 6d - 3$
 $g + 3 = 6d$

$$\frac{g + 3}{6} = \frac{6d}{6}$$

$$\frac{g + 3}{6} = d$$

$$d = \frac{g + 3}{6}$$

b. 1. Multiply both sides by t .

$$\begin{aligned} \text{b. } a &= \frac{v-u}{t} \\ a \times t &= \frac{v-u}{t} \times t \\ at &= v-u \end{aligned}$$

2. Add u to both sides.

$$at + u = v$$

3. Rewrite the equation so that v is on the left-hand side.

$$v = at + u$$

on Resources



eWorkbook Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)



Interactivities Individual pathway interactivity: Rearranging formulas (int-4493)
Rearranging formulas (int-6040)

Exercise 3.6 Rearranging formulas

learn on

3.6 Quick quiz

on

3.6 Exercise

Individual pathways

PRACTISE

1, 3, 6, 8, 11

CONSOLIDATE

2, 4, 7, 9, 12

MASTER

5, 10, 13

Fluency

1. **WE12** Rearrange each formula to make x the subject.

a. $y = ax$

b. $y = ax + b$

c. $y = 2ax - b$

d. $y + 4 = 2x - 3$

e. $6(y + 2) = 5(4 - x)$

f. $x(y - 2) = 1$

2. Rearrange each formula to make x the subject.

a. $x(y - 2) = y + 1$

b. $5x - 4y = 1$

c. $6(x + 2) = 5(x - y)$

d. $7(x - a) = 6x + 5a$

e. $5(a - 2x) = 9(x + 1)$

f. $8(9x - 2) + 3 = 7(2a - 3x)$

3. **WE13** For each of the following, make the variable shown in brackets the subject of the formula.

a. $g = 4P - 3$ (P)

b. $f = \frac{9c}{5}$ (c)

c. $f = \frac{9c}{5} + 32$ (c)

d. $v = u + at$ (t)

4. For each of the following, make the variable shown in brackets the subject of the formula.

a. $d = b^2 - 4ac$ (c)

b. $m = \frac{y-k}{h}$ (y)

c. $m = \frac{y-a}{x-b}$ (a)

d. $m = \frac{y-a}{x-b}$ (x)

5. For each of the following, make the variable shown in brackets the subject of the formula.

a. $C = \frac{2\pi}{r}$ (r)

b. $f = ax + by$ (x)

c. $s = ut + \frac{1}{2}at^2$ (a)

d. $F = \frac{GMm}{r^2}$ (G)

Understanding

6. The cost to rent a car is given by the formula $C = 50d + 0.2k$, where d = the number of days rented and k = the number of kilometres driven. Lin has \$300 to spend on car rental for her four-day holiday.



Calculate the distance that she can drive on her holiday.

7. A cyclist pumps up a bike tyre that has a slow leak.



The volume of air (in cm^3) after t minutes is given by the formula $V = 24\,000 - 300t$.

a. Determine the volume of air in the tyre when it is first filled.

b. Write out the equation for the tyre's volume and solve it to work out how long it would take the tyre to go completely flat.

Reasoning

8. The total surface area of a cylinder is given by the formula $T = 2\pi r^2 + 2\pi rh$, where $r =$ radius and $h =$ height. A car manufacturer wants its engines' cylinders to have a radius of 4 cm and a total surface area of 400 cm^2 . Show that the height of the cylinder is approximately 11.92 cm, correct to 2 decimal places. (*Hint: Express h in terms of T and r .*)



9. If $B = 3x - 6xy$, write x as the subject. Explain the process you followed by showing your working.
10. The volume, V , of a sphere can be calculated using the formula $V = \frac{4}{3}\pi r^3$, where r is the radius of the sphere.



Explain how you would work out the radius of a spherical ball that has the capacity to hold 5 litres of air.

Problem solving

11. Use algebra to show that $\frac{1}{v} = \frac{1}{u} - \frac{1}{f}$ can also be written as $u = \frac{fv}{v+f}$.

12. Consider the formula $d = \sqrt{b^2 - 4ac}$. Rearrange the formula to make a the subject.

13. Evaluate the values for a and b such that:

$$\frac{4}{x+1} - \frac{3}{x+2} = \frac{ax+b}{(x+1)(x+2)}$$

LESSON

3.7 Review

3.7.1 Topic summary

LINEAR EQUATIONS

Inverse operations

- Solving a linear equation means determining a value for the pronumeral that makes the statement true.
- Solving equations requires inverse operations.
- The inverse operation has the effect of undoing the original operation.

Operation	Inverse operation
+	-
-	+
×	÷
÷	×

- To keep an equation balanced, inverse operations are done on both sides of the equation.

e.g.

$$3x + 1 = 11$$

$$3x + 1 - 1 = 11 - 1$$

$$\frac{3x}{3} = \frac{10}{3}$$

$$x = \frac{10}{3}$$

Equations with brackets

- The number in front of the brackets indicates the multiple of the expression inside the brackets.
- Brackets can be removed by dividing both sides by the multiple.

e.g.

$$3(x + 5) = 18$$

$$\frac{3(x + 5)}{3} = \frac{18}{3}$$

$$(x + 5) = 6$$

- Alternatively, brackets can be removed by expanding (multiplying) out the expression.

e.g.

$$3(x + 5) = 18$$

$$3x + 15 = 18$$

Equations and inequations with pronumerals on both sides

- Pronumerals must be moved so that they appear only on one side using inverse operations.
- Once this is done, you can solve the equation.

e.g.

$$5y = 3y + 3$$

$$5y - 3y = 3y + 3 - 3y$$

$$2y = 3$$

$$y = \frac{3}{2}$$

e.g.

$$3x + 2 < 14$$

$$3x + 2 - 2 < 14 - 2$$

$$3x < 12$$

$$x < 4$$

Pronumeral in the denominator

- The pronumeral needs to be expressed as a numerator.
- This can be done by multiplying both sides of the equation by the pronumeral.

e.g.

$$\frac{4}{x} \times x = \frac{3}{2} \times x$$

- Alternatively, you can flip both sides of the equation.

e.g.

$$\frac{4}{x} = \frac{3}{2}$$

becomes $\frac{x}{4} = \frac{2}{3}$.

Worded problems

- Worded problems should be translated into equations.
 - Pronumerals must be defined in the answer if this is not already done in the question.
- e.g. The equation for the worded problem 'When 6 is subtracted from a certain number, the result is 15' is 'x - 6 = 15'.

Rearranging formulas

- The subject of the formula is the pronumeral that is written by itself.
 - The difference between rearranging formulas and solving linear equations is that rearranging formulas does not require a value for the pronumeral to be found.
- e.g. Rearrange the following to make x the subject:

$$y = kx + m$$

$$y - m = kx$$




$$y - \frac{m}{k} = \frac{kx}{k}$$

$$y - \frac{m}{k} = x$$

3.7.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
3.2	I can recognise inverse operations.			
	I can solve multistep equations and inequations with pronumerals on one side of the equation.			
	I can solve equations and inequations with algebraic fractions.			
3.3	I can solve equations involving brackets.			
3.4	I can solve equations by collecting pronumerals on one side of an equation before solving.			
3.5	I can define a pronumeral and write an equation to represent the problem.			
	I can solve a problem and answer the question (or questions) posed in the problem.			
3.6	I can rearrange formulas with two or more pronumerals to make only one of the pronumerals the subject.			

3.7.3 Project

Forensic science

Scientific studies have been conducted on the relationship between a person's height and the measurements of a variety of body parts. One study has suggested that there is a general relationship between a person's height and the humerus bone in their upper arm, and that this relationship is slightly different for men and women.

According to this study, there is a general trend indicating that $h = 3.08l + 70.45$ for men, and $h = 3.36l + 57.97$ for women, where h represents body height in centimetres and l the length of the humerus in centimetres.

Imagine the following situation.

A decomposed body has been in the bushland outside your town. A team of forensic scientists suspects that the body could be the remains of either Alice Brown or James King, both of whom have been missing for several years.

From the descriptions provided by their Missing Persons files, Alice is 162 cm tall and James is 172 cm tall. The forensic scientists hope to identify the body based on the length of the body's humerus.



1. Based on the relationship suggested by the study previously mentioned, complete both of the following tables using the equations provided. Calculate body height to the nearest centimetre.

Body height (men)

Length of humerus, l (cm)	20	25	30	35	40
Body height, h (cm)					

Body height (women)

Length of humerus, l (cm)	20	25	30	35	40
Body height, h (cm)					



2. On a piece of graph paper, draw the first quadrant of a Cartesian plane. Since the length of the humerus is the independent variable, place it on the x -axis. Place the dependent variable, body height, on the y -axis.
3. Plot the points from the two tables onto the set of axes drawn in question 2. Join the points with straight lines, using different colours to represent men and women.
4. Describe the shape of the two graphs.
5. Measure the length of your humerus. Use your graph to predict your height. How accurate is the measurement?
6. The two lines of your graph will intersect if extended. At what point does this occur? Comment on this value.



The forensic scientists measured the length of the humerus of the decomposed body and found it to be 33 cm.

7. Using methods covered in this activity, is it more probable that the body is that of Alice or of James? Justify your decision with mathematical evidence.

on Resources

- eWorkbook** Topic 3 Workbook (worksheets, code puzzle and project) (ewbk-2003)
- Interactivities** Crossword (int-0700)
- Sudoku puzzle (int-3204)

Exercise 3.7 Review questions

learn on

Fluency

1. **MC** Select the linear equation represented by the sentence ‘When a certain number is multiplied by 3, the result is 5 times that certain number plus 7’.
- A. $3x + 7 = 5x$ B. $5(x + 7) = 3x$ C. $5x + 7 = 3x$
D. $5x = 3x + 7$ E. $3x = 7x + 5$
2. **MC** Calculate the solution to the equation $\frac{x}{3} = 5$.
- A. $x = -15$ B. $x = 15$ C. $x = 1\frac{2}{3}$ D. $x = 3$ E. $x = -3$

3. **MC** Determine the solution to the equation $7 = 21 + x$.
- A. $x = 28$ B. $x = -28$ C. $x = -14$ D. $x = 14$ E. $x = 7$
4. **MC** Calculate the solution to the equation $5x + 3 = 37$.
- A. $x = 8$ B. $x = -8$ C. $x = 6.8$ D. $x = 106$ E. $x = -6.8$
5. **MC** Determine the solution to the equation $8 - 2x = 22$.
- A. $x = 11$ B. $x = 15$ C. $x = -15$ D. $x = -7$ E. $x = 7$
6. **MC** Determine the solution to the equation $4x + 3 = 7x - 33$.
- A. $x = -12$ B. $x = 12$ C. $x = \frac{36}{11}$ D. $x = \frac{30}{11}$ E. $x = \frac{11}{30}$
7. **MC** Calculate the solution to the equation $7(x - 15) = 28$.
- A. $x = 11$ B. $x = 19$ C. $x = 20$ D. $x = 6.14$ E. $x = 6$
8. **MC** Select the correct rearrangement of $y = ax + b$ in terms of x .
- A. $x = \frac{y-a}{b}$ B. $x = \frac{y-b}{a}$ C. $x = \frac{b-y}{a}$ D. $x = \frac{y+b}{a}$ E. $x = \frac{x-b}{a}$
9. Identify which of the following are linear equations.
- a. $5x + y^2 = 0$ b. $2x + 3 = x - 2$
c. $\frac{x}{2} = 3$ d. $x^2 = 1$
e. $\frac{1}{x} + 1 = 3x$ f. $8 = 5x - 2$
g. $5(x + 2) = 0$ h. $x^2 + y = -9$
i. $r = 7 - 5(4 - r)$
10. Solve each of the following linear equations.
- a. $3a = 8.4$ b. $a + 2.3 = 1.7$
c. $\frac{b}{21} = -0.12$ d. $b - 1.45 = 1.65$
e. $b + 3.45 = 0$ f. $7.53b = 5.64$
11. Solve each of the following linear equations.
- a. $\frac{2x-3}{7} = 5$ b. $\frac{5-x}{2} = -4$ c. $\frac{-3x-4}{5} = 3$
d. $\frac{6}{x} = 5$ e. $\frac{4}{x} = \frac{3}{5}$ f. $\frac{x+1.7}{2.3} = -4.1$
12. Solve each of the following linear equations.
- a. $5(x-2) = 6$ b. $7(x+3) = 40$ c. $4(5-x) = 15$
d. $6(2x+3) = 1$ e. $4(x+5) = 2x-5$ f. $3(x-2) = 7(x+4)$
13. Liz has a packet of 45 lolly snakes. She saves 21 to eat tomorrow, but she rations the remainder so that she can eat 8 snakes every hour until today's share of snakes is gone.
- a. Write a linear equation in terms of the number of hours, h , to represent this situation.
b. Determine the number of hours it will take Liz to eat today's share.

14. Solve each of the following linear equations.
- | | |
|----------------------|----------------------|
| a. $11x = 15x - 2$ | b. $3x + 4 = 16 - x$ |
| c. $5x + 2 = 3x + 8$ | d. $8x - 9 = 7x - 4$ |
| e. $2x + 5 = 8x - 7$ | f. $3 - 4x = 6 - x$ |
15. Translate the following sentences into algebraic equations. Use x to represent the number in question.
- Twice a certain number is equal to 3 minus that certain number.
 - When 8 is added to 3 times a certain number, the result is 19.
 - Multiplying a certain number by 6 equals 4.
 - Dividing 10 by a certain number equals one more than dividing that number by 6.
 - Multiply a certain number by 2, then add 5. Multiply this result by 7. This expression equals 0
 - Twice a certain distance travelled is 100 metres more than that certain distance travelled plus 50 metres.
16. Takanori decides to go on a holiday. He travels a certain distance on the first day, twice that distance on the second day, three times that distance on the third day and four times that distance on the fourth day. If Takanori's total journey is 2000 km, calculate the distance he travelled on the third day.
17. For each of the following, make the variable shown in brackets the subject of the formula.
- | | |
|--------------------------------|--------------------------|
| a. $y = 6x - 4$ (x) | b. $y = mx + c$ (x) |
| c. $q = 2(P - 1) + 2r$ (P) | d. $P = 2l + 2w$ (w) |
18. For each of the following, make the variable shown in brackets the subject of the formula.
- | | |
|------------------------------|--|
| a. $v = u + at$ (a) | b. $s = \left(\frac{u+v}{2}\right)t$ (t) |
| c. $v^2 = u^2 + 2as$ (a) | d. $2A = h(a + b)$ (b) |

Problem solving

19. Saeed is comparing two car rental companies, Golden Ace Rental Company and Silver Diamond Rental Company. Golden Ace Rental Company charges a flat rate of \$38 plus \$0.20 per kilometre. The Silver Diamond Rental Company charges a flat rate of \$30 plus \$0.32 per kilometre. Saeed plans to rent a car for three days.
- Write an equation for the cost of renting a car for three days from the Golden Ace Rental Company, in terms of the number of kilometres travelled, k .
 - Write an algebraic equation for the cost of renting a car for three days from the Silver Diamond Rental Company, in terms of the number of kilometres travelled, k .
 - Evaluate the number of kilometres Saeed would have to travel so that the cost of hiring from each company is the same.
20. Frederika has \$24 000 saved to pay for a holiday. Her travel expenses are \$5400 and her daily expenses are \$260.
- Write down an equation for the cost of her holiday if she stays for d days.
 - Determine the number of days Frederika can spend on holiday if she wants to be able to buy a new laptop for \$ 2500 when she gets back from her holidays.



21. A company that makes bottled orange juice buys their raw materials from two sources. The first source provides liquid with 6% orange juice, whereas the second source provides liquid with 3% orange juice. The company wants to make 1-litre bottles that have 5% orange juice. Let x = the amount of liquid (in litres) that the company buys from the first source.
- Write an expression for the amount of orange juice from the first supplier, given that x is the amount of liquid.
 - Write an expression for the amount of liquid from the second supplier, given that x is the amount of liquid used from the first supplier.
 - Write an expression for the amount of orange juice from the second supplier.
 - Write an equation for the total amount of orange juice in a mixture of liquids from the two suppliers, given that 1 litre of bottled orange juice will be mixed to contain 5% orange juice.
 - Determine the quantity of the first supplier's liquid that the company uses.
22. Jayani goes on a four-day bushwalk. She travels a certain distance on the first day, half of that distance on the second day, a third of that distance on the third day and a quarter of that distance on the fourth day. If Jayani's total journey is 50 km, evaluate the distance that she walked on the first day.



23. Svetlana goes on a five-day bushwalk, travelling the same relative distances as Jayani travelled in question 22 (a certain amount, then half that amount, then one third of that, then one quarter of that, and finally one fifth of that). If Svetlana's journey is also 50 km, determine the distance that she travelled on the first day.
24. An online bookstore advertises its shipping cost to Australia as a flat rate of \$20 for up to 10 books. Their major competitor offers a flat rate of \$12 plus \$1.60 per book. Determine the number of books you would have to buy (6, 7, 8, 9 or 10) for the first bookstore's shipping cost to be a better deal.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 3 Linear equations

3.1 Pre-test


- 3
- 22
- C
- a. $x = -7$ b. $z = 11$
- 4
- False
- E
- C
- a. $a = 2$ b. $b = -8$
- B
- 287.5km
- 3
- D
- B
- a. $a = \frac{bc}{(b+c)}$ b. $b = \frac{ac}{(c-a)}$

3.2 Solving linear equations and inequations


- a. No b. Yes c. No
d. Yes e. Yes f. No
- a. No b. No c. Yes
d. No e. Yes f. No
- a. $x = 210$ b. $x = 230$ c. $x = 23$
d. $x = -96$ e. $x = 37$ f. $x = 20$
- a. $x = 50$ b. $x = 138$ c. $x = 442$
d. $x = 243$ e. $y = -15$ f. $y = 1.8$
- a. $y = 12.8$ b. $y = 2\frac{1}{3}$ c. $y = -\frac{1}{18}$
d. $y = 21.5$ e. $y = 172.5$ f. $y = -8.32$
- a. $y = 5$ b. $y = -2$ c. $y = 0.2$
d. $y = 1$ e. $y = 1$ f. $y = 2$
- a. $y = 5\frac{1}{3}$ b. $y = -1\frac{2}{3}$ c. $y = 24.5$
d. $y = 1.2$ e. $y = 22.25$ f. $y = \frac{383}{600}$
- a. $x = 1$ b. $x = -2$ c. $x = 3$
d. $x = -6$ e. $x = -1$ f. $x = \frac{1}{2}$
- a. $x = 1\frac{2}{3}$ b. $x = -2.32$ c. $x = 3$
d. $x = -\frac{5}{6}$ e. $x = 1\frac{1}{4}$ f. $x = 3\frac{1}{13}$
- a. $x = -1$ b. $x = 1$ c. $x = 0$
d. $x = 5$ e. $x = -8.6$ f. $x = -0.9$
- a. $x = 8.9$ b. $x = 19$ c. $x = -26$
d. $x = -1$ e. $x = -1$ f. $x = -\frac{1}{2}$
- a. $x = 8$ b. $x = 3$ c. $x = 4$
d. $x = -15$ e. $x = 26$ f. $x = -42$

- a. $x = 9$ b. $x = -1\frac{1}{5}$ c. $x = 9\frac{1}{3}$
d. $x = -2\frac{1}{4}$ e. $x = -7$ f. $x = \frac{2}{3}$
- a. $z = 16$ b. $z = 31$ c. $z = -4$
d. $z = 6$ e. $z = -9$ f. $z = -6$
- a. $z = -1.9$ b. $z = 6.88$ c. $z = 140$
d. $z = 0.6$ e. $z = -35.4$ f. $z = 8$
- a. $x = 1$ b. $x = 13$ c. $x = -2$
d. $x = -8$ e. $x = -4$ f. $x = 30\frac{1}{2}$
- a. $x = -6$ b. $x = -\frac{7}{10}$ c. $x = 10.35$
d. $x = 0.326$ e. $x = 22$ f. $x = -5$
- a. $x = 4$ b. $x = \frac{3}{7}$ c. $x = -1\frac{1}{7}$
d. $x = -6\frac{2}{3}$ e. $x = \frac{4}{45}$ f. $x = 8$
- a. $x = -6$ b. $x = 7.5$ c. $x = 5.1$
d. $x = -6$ e. $x = -5\frac{13}{15}$ f. $x = -61\frac{3}{7}$
- a. $a = -1$ b. $b = 10$ c. $c = 12.1$
d. $d = 4$ e. $e = 5$ f. $f = 12$

21. a. $x \leq 16$



- b. $x > 17$




- a. $x \leq 9$ b. $x > 10$
- a. $x \geq -13$ b. $x > -7$
- a. $x \geq -14$ b. $x > 18$
- a. -1 b. 6 c. 303
- \$91
- No. $46.7^\circ\text{C} \approx 116.1^\circ\text{F}$
- For a positive integer, $(x-4)$ must be a factor of 12.
 $x = 5, 6, 7, 8, 10, 16$
- Sample responses can be found in the worked solutions in the online resources.
- The mistake is in the second line: the -1 should have been multiplied by 5.
- 60 lollies
- Old machine: 6640 cans; new machine: 9360 cans
- a. \$34
b. Yes, a saving of \$7
- 17


3.3 Solving linear equations with brackets

- $x = 6$
 - $x = -3$
 - $x = 0$
 - $x = 56$
 - $x = 1$
 - $x = 0$
- $x = -1$
 - $x = -2$
 - $x = 6$
 - $x = 0$
 - $x = -0.8$
 - $x = 6$
- $b = 1\frac{1}{6}$
 - $m = 4\frac{1}{2}$
 - $a = -1\frac{1}{2}$
 - $m = -1\frac{1}{3}$
 - $p = \frac{3}{5}$
 - $m = 2\frac{2}{3}$
- $a = -1\frac{1}{2}$
 - $p = 1\frac{1}{2}$
 - $a = 3\frac{1}{3}$
 - $m = -2\frac{1}{2}$
 - $a = -\frac{1}{6}$
 - $m = -\frac{1}{4}$
- $x = 16\frac{1}{9}$
 - $x = 2$
 - $a = 5$
 - $b = 7$
 - $y = 7$
 - $x = -3\frac{1}{3}$
- $m = 4$
 - $y = 1$
 - $a = 5$
 - $p = -2$
 - $m = 3$
 - $p = 1$
- $y = -7.5$
 - $y = -4.667$
 - $y = -26.25$
 - $y = 8.571$
 - $y = -2.9$
 - $y = 3.243$
- $y = 3.667$
 - $y = 2.75$
 - $y = 1.976$
 - $y = -2$
 - $y = -3.167$
 - $y = 1.98$
- D
- C
- 1990
- $[2(2x + 7)]m$
 - Width 10 m, length 17 m
- Expand out the brackets or divide by the factor.
- $x = 13$. Sample responses can be found in the worked solutions in the online resources.
- Adding 7 to both sides is the least preferred option, as it does not resolve the subtraction of 7 within the brackets.
- 3
- Oscar: \$90, Josue: \$35
- \$20

3.4 Solving linear equations with pronumerals on both sides

- $y = -1$
 - $y = 1$
 - $y = -3$
 - $y = -5$
 - $y = -45$
 - $y = \frac{8}{27}$
- $y = -2$
 - $y = 20$
 - $y = 1$
 - $y = -1\frac{1}{5}$
 - $y = 4$
 - $y = -2$
- C
- A
- $x = 1$
 - $x = -2$
 - $x = 3$
 - $x = 4$
 - $x = 9$
 - $x = 3$
- $x = -2$
 - $x = 4$
 - $x = -4$
 - $x = \frac{1}{2}$
 - $x = 0$
 - $x = -\frac{3}{4}$

- $x = -6$
 - $x = 2$
 - $x = -2$
 - $x = 13$
 - $x = -7$
 - $x = \frac{1}{6}$
- D
- C
- $x = 5$
 - $x = -3$
 - $x = -8$
 - $x = -7\frac{1}{2}$
 - $x = 4$
 - $x = \frac{10}{11}$
- $x = -18$
 - $x = 3\frac{2}{3}$
 - $x = \frac{2}{3}$
 - $x = -1\frac{3}{7}$
 - $x = 0$
 - $x = -15$
- $k = -3$
- 24
- $3(n - 36) - 98 = -11n + 200$. Sample responses can be found in the worked solutions in the online resources.
- 



- Area of first rectangle $= (x + 1) \times 3$
 $= 3x + 3$
 - Area of second rectangle $= (x + 1 + x + 1 + x + 1) \times 1$
 $= 3x + 3$
 - Both rectangles have an area of $3x + 3$.
- You cannot easily divide the left-hand side by 6 or the right-hand side by 4.
 - Daughter = 10 years, Tom = 40 years
 - The unknown number is -3 .
 - $C = 40h + 250$
 - 18 hours, 45 minutes
 - 18 750
 - The printing is cheaper by \$1375.
 - 65 games
 - \$25

3.5 Solving problems with linear equations

- $x + 3 = 5$
 - $x - 9 = 7$
 - $7x = 24$
 - $\frac{x}{5} = 11$
 - $\frac{x}{2} = -9$
- $5x - 3 = -7$
 - $2(14 - x) = -3$
 - $3x + 5 = 8$
 - $2x - 12 = 15$
 - $\frac{3x + 4}{2} = 5$
- C
- D
- B
- B
- 0

8. 36 years old
9. \$66
10. 20×15 ; 30×10 ; area = 300 square units
11. 7 m
12. 12 jackets
13. 15 days
14. 947 CDs
15. a. Post Quick (cost = \$700)
 - b. The cost is nearly the same for 333 flyers (\$366.50 and \$366.40).
16. 2
17. \$42.50
18. a. $A_{\text{fenced}} = (5x + 20) \text{ m}^2$ b. $A_{\text{path}} = (3x + 16) \text{ m}^2$

c. $l = 8 \text{ m}$, $w = 2 \text{ m}$ d. $\frac{8}{25} \text{ m}$

3.6 Rearranging formulas

1. a. $x = \frac{y}{a}$ b. $x = \frac{y-b}{a}$
 - c. $x = \frac{y+b}{2a}$ d. $x = \frac{y+7}{2}$
 - e. $x = \frac{8-6y}{5}$ f. $x = \frac{1}{y-2}$
2. a. $x = \frac{y+1}{y-2}$ b. $x = \frac{4y+1}{5}$ c. $x = -5y - 12$
 - d. $x = 12a$ e. $x = \frac{5a-9}{19}$ f. $x = \frac{14a+13}{93}$
3. a. $P = \frac{g+3}{4}$ b. $c = \frac{5f}{9}$
 - c. $c = \frac{5(f-32)}{9}$ d. $t = \frac{v-u}{a}$
4. a. $c = \frac{b^2-d}{4a}$ b. $y = hm + k$
 - c. $a = y - m(x-b)$ d. $x = \frac{y-a+mb}{m}$
5. a. $r = \frac{2\pi}{C}$ b. $x = \frac{f-by}{a}$
 - c. $a = \frac{2(s-ut)}{t^2}$ d. $G = \frac{Fr^2}{Mm}$
6. 500 km
7. a. $24\,000 \text{ cm}^3$
 - b. $t = 80 \text{ min} = 1 \text{ h } 20 \text{ min}$
8. Sample responses can be found in the worked solutions in the online resources.
9. $\frac{B}{3(1-2y)} = x$
10. radius = 10.6 cm
11. Sample responses can be found in the worked solutions in the online resources.
12. $a = \frac{b^2-d^2}{4c}$
13. $a = 1$; $b = 5$

Project


1. Body height (men)

Length of humerus, l (cm)	20	25	30	35	40
Body height, h (cm)	132	147	163	178	194

Body height (women)

Length of humerus, l (cm)	20	25	30	35	40
Body height, h (cm)	125	142	159	176	192

2. and 3.



3. Linear

4. Results will vary for each student.
5. (44.6, 207.8)
6. The height of the male body with $l = 33 \text{ cm}$:
 $h = 3.08 \times 33 + 70.45 = 172.09 \text{ cm}$
 The height of the female body with $l = 33 \text{ cm}$:
 $h = 3.36 \times 33 + 57.97 = 168.85 \text{ cm}$
 The estimated height for a male body is very close to the known height of James King; the estimated height for a female body is more different from the known height of Alice Brown. Therefore, the body is more likely to be that of James King.

3.7 Review questions

1. C
2. B
3. C
4. C
5. D
6. B
7. B
8. B
9. b, c, f, g, i
10. a. $a = 2.8$ b. $a = -0.6$
 - c. $b = -2.52$ d. $b = 3.1$
 - e. $b = -3.45$ f. $b = 0.749$

11. a. $x = 19$ b. $x = 13$ c. $x = -6\frac{1}{3}$
 d. $x = 1\frac{1}{5}$ e. $x = 6\frac{2}{3}$ f. $x = -11.13$
12. a. $x = 3\frac{1}{5}$ b. $x = 2\frac{5}{7}$
 c. $x = 1\frac{1}{4}$ d. $x = -1\frac{5}{12}$
 e. $x = -12\frac{1}{2}$ f. $x = -8\frac{1}{2}$
13. a. $8h + 21 = 45$ b. 3 hours
14. a. $x = \frac{1}{2}$ b. $x = 3$ c. $x = 3$
 d. $x = 5$ e. $x = 2$ f. $x = -1$
15. a. $2x = 3 - x$ b. $3x + 8 = 19$
 c. $6x = 4$ d. $\frac{10}{x} - 1 = \frac{x}{6}$
 e. $7(2x + 5) = 0$ f. $2x - 100 = x + 50$
16. 600 km
17. a. $x = \frac{y + 4}{6}$ b. $x = \frac{y - c}{m}$
 c. $P = \frac{q - 2r}{2} + 1$ d. $w = \frac{P - 2l}{2}$
18. a. $a = \frac{v - u}{t}$ b. $t = \frac{2s}{u + v}$
 c. $a = \frac{v^2 - u^2}{2s}$ d. $b = \frac{2A - ah}{h}$
19. a. $C_G = 114 + 0.20k$
 b. $C_S = 90 + 0.32k$
 c. 200 km
20. a. $5400 + 260d = C_H$ b. 61 days
21. a. $0.06x$
 b. $(1 - x)$
 c. $0.03(1 - x)$
 d. $0.06x + 0.03(1 - x) = 0.05$
 e. 0.667 or 66.7%
22. 24 km
23. $21\frac{123}{137} \approx 21.9$ km
24. The online bookstore's fee is a better price than their major competitor's fee for all of the options given: 6, 7, 8, 9 and 10 books.

4 Congruence and similarity

LESSON SEQUENCE

4.1 Overview	192
4.2 Ratio and scale	195
4.3 Congruent figures	204
4.4 Nets, polyhedra and Euler's formula	212
4.5 Similar figures	217
4.6 Area and volume of similar figures	229
4.7 Review	238



LESSON

4.1 Overview

4.1.1 Why learn this?

Geometry allows us to explore our world in very precise ways. It is also one of the oldest areas of mathematics. It involves the study of points, lines and angles and how they can be combined to make different shapes. Similarity and congruence are two important concepts in geometry. When trying to determine whether two shapes are exactly the same, or if they are enlargements of each other, the answers can be found by considering the sides and angles of those shapes.

When you take a photo on your phone it is very small. If you save it to your computer you can make it larger. The larger photo is an enlargement of the original photo — this is an example of how similar figures work in the everyday world.



The principle of similar triangles can be used to work out the height of tall objects by calculating the length of their shadows. This technique was extremely important in early engineering and architecture. Today, architects and designers still prepare scale diagrams before starting the building process.

In manufacturing, the products that come off an assembly line all have exactly the same shape and size. These products can be described as congruent. Designers, engineers and surveyors all use the concepts of congruence and similarity in their daily work.

Hey students! Bring these pages to life online



Watch videos



Engage with interactivities



Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets


Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 4.1 Pre-test

1. Express the ratio $1\frac{2}{3} : 2\frac{3}{5}$ in simplest form.


$$1\frac{2}{3} : 2\frac{3}{5} = \boxed{} : \boxed{}$$

2. Determine the value of x in the proportion $2 : 3 = 9 : x$
3. **MC** Select the correct symbol for congruence.
- A. =
 B. \approx
 C. \sim
 D. \cong
 E. \equiv
4. **MC** If $OB = OC$, choose which of the following congruency tests can be used to prove $\triangle AOB \cong \triangle DOC$.




- A. AAA
 B. RHS
 C. SSS
 D. ASA
 E. SAS
5. Determine the simplest ratio of $y : z$ if $3x = 2y$ and $4x = 3z$.
6. The ratio $1.4 : 0.2$ in its simplest form is $14 : 2$. State whether this statement is True or False.
7. **MC** Select the only pair of congruent triangles from the following.

- A. $\triangle ABC \cong \triangle BCD$
 B. $\triangle ABD \cong \triangle BCD$
 C. $\triangle BCD \cong \triangle DEF$
 D. $\triangle ABC \cong \triangle DEF$
 E. $\triangle ABD \cong \triangle DEF$




8. For the diagram given:
- a. name the angle that is equal to angle ABC
- b. state the congruency test that can be used to prove $\triangle ABC \cong \triangle CED$.




9. **MC** Select which similarity test can be used to prove $\triangle ABC \sim \triangle ADE$.

- A. AAA
- B. RHS
- C. SSS
- D. ASA
- E. SAS




10. **MC** $\triangle ABC$ and $\triangle CDE$ are similar. Choose which of the following statements is true.

- A. $\frac{AB}{ED} = \frac{BC}{CD}$
- B. $\frac{AB}{ED} = \frac{AC}{CD}$
- C. $\frac{AC}{EC} = \frac{BC}{CD}$
- D. $\frac{AB}{ED} = \frac{BC}{CE}$
- E. $\frac{AB}{ED} = \frac{CD}{AC}$




11. $\triangle ABC$ and $\triangle CDE$ are similar.



Determine the value of the length x correct to 1 decimal place.

12. $\triangle ABC$ and $\triangle CDE$ are similar. Determine the value of b .



13. A pair of cones are similar. The ratio of their volumes is $27 : 125$.
- The perpendicular height ratio of the two cones is $\square : \square$.
 - The areas of the base ratio of the two cones is $\square : \square$.
14. A pair of rectangles are similar. If the width of the first rectangle is 3 times the width of the other, the ratio of their areas is:
larger area : smaller area = $\square : \square$.
15. **MC** A rectangular box has a surface area of 94 cm^2 and volume of 60 cm^3 . Select the volume and surface area of a similar box that has side lengths that are twice the size of the original.
- Volume = 480 cm^3 and surface area = 752 cm^2
 - Volume = 480 cm^3 and surface area = 376 cm^2
 - Volume = 120 cm^3 and surface area = 188 cm^2
 - Volume = 240 cm^3 and surface area = 376 cm^2
 - Volume = 240 cm^3 and surface area = 752 cm^2

LESSON

4.2 Ratio and scale

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- compare two quantities of the same type using ratios
- simplify ratios
- enlarge a figure using a scale factor
- calculate and use a scale factor.

4.2.1 Ratio


eles-4727

- Ratios** are used to compare quantities of the same kind, measured in the same unit.
- The ratio '1 is to 4' can be written in two ways: as $1 : 4$ or as $\frac{1}{4}$.
- The order of the numbers in a ratio is important.
- In simplest form, a ratio is written using the smallest whole numbers possible.

WORKED EXAMPLE 1 Expressing ratios in simplest form

A lighthouse is positioned on a cliff that is 80 m high. A ship at sea is 3600 m from the base of the cliff.

- Write the following ratios in simplest form.
 - The height of the cliff to the distance of the ship from shore.
 - The distance of the ship from shore to the height of the cliff.
- Compare the distance of the ship from shore with the height of the cliff.



THINK

a. i. 1. The height and distance are in the same units (m). Write the height first and the distance second.

2. Simplify the ratio by dividing both terms by the highest common factor (80).

ii. 1. Write the distance first and the height second.

2. Simplify by dividing both terms by the highest common factor (80).

Note: Do not write $\frac{45}{1}$ as 45, because a ratio is a comparison between two numbers.

b. 1. Write the ratio 'distance of the ship from shore to height of the cliff'.

2. Write the answer.

WRITE

a. i. Height of cliff : distance of ship from shore
= 80 : 3600

$$= 1 : 45$$

ii. Distance of ship from shore : height of cliff

$$= \frac{3600}{80}$$

$$= \frac{45}{1}$$

b. 45 : 1

The distance of the ship from shore is 45 times the height of the cliff.

WORKED EXAMPLE 2 Simplifying ratios into simplest form

Express each of the following ratios in simplest form.

a. 24 : 8

b. 3.6 : 8.4

c. $1\frac{4}{9} : 1\frac{2}{3}$

THINK

a. Divide both terms by the highest common factor (8).

b. 1. Multiply both terms by 10 to obtain whole numbers.

2. Divide both terms by the highest common factor (12).

c. 1. Change both mixed numbers into improper fractions.

2. Multiply both terms by the lowest common denominator (9) to obtain whole numbers.

WRITE

a. 24 : 8

$$= 3 : 1$$

b. 3.6 : 8.4

$$= 36 : 84$$

$$= 3 : 7$$

c. $1\frac{4}{9} : 1\frac{2}{3}$

$$= \frac{13}{9} : \frac{5}{3}$$

$$= 13 : 15$$

- A **proportion** is a statement that indicates that two ratios are equal.
- A proportion can be written in two ways, for example in the format used for $4 : 7 = x : 15$ or in the format used for $\frac{4}{7} = \frac{x}{15}$.

WORKED EXAMPLE 3 Calculating a value in a proportion

Determine the value of x in the proportion $4 : 9 = 7 : x$.

THINK

1. Write the ratios as equal fractions.
2. Multiply both sides by x .
3. Solve the equation to obtain the value of x .
4. Write the answer.

WRITE

$$\frac{4}{9} = \frac{7}{x}$$

$$\frac{4x}{9} = 7$$


$$4x = 63$$

$$x = 15.75$$


4.2.2 Scale

eles-4728


- Ratios are used when creating scale drawings or maps.
- Consider the situation in which we want to enlarge a triangle ABC (the **object**) by a **scale factor** of 2 (this means we want to make it twice its size).
The following is one method that we can use.



1. Mark a point O somewhere outside the triangle and draw the lines OA, OB and OC as shown.



- Measure the length of OA and mark in the point A' (this is called the **image** of A) so that the distance OA' is twice the distance of OA .
- In the same way, mark in points B' and C' . ($OB' = 2 \times OB$, and $OC' = 2 \times OC$.)




- Joining $A'B'C'$ gives a triangle that has side lengths double those of $\triangle ABC$. $\triangle A'B'C'$ is called the image of $\triangle ABC$.

Scale factor

$$\text{Scale factor} = \frac{\text{image length}}{\text{object length}}$$

WORKED EXAMPLE 4 Enlarging a figure


Enlarge triangle ABC by a scale factor of 3, with the centre of enlargement at point O .



THINK

- Join each vertex of the triangle to the centre of enlargement (O) with straight lines, then extend them.
- Locate points A' , B' and C' along the lines so that $OA' = 3OA$, $OB' = 3OB$ and $OC' = 3OC$.
- Join points A' , B' and C' to complete the image.

DRAW



- Enlargements have the following properties.
 - The corresponding side lengths of the enlarged figure are changed in a fixed ratio (they have same ratio).
 - The corresponding angles are the same.
 - A scale factor greater than 1 produces an enlarged figure.
 - If the scale factor is a positive number less than 1, the image is smaller than the object (this means that a reduction has taken place).

WORKED EXAMPLE 5 Calculating the scale factor

A triangle PQR has been enlarged to create triangle P'Q'R'. PQ = 4 cm, PR = 6 cm, P'Q' = 10 cm and Q'R' = 20 cm. Calculate:


- the scale factor for the enlargement
- the length of P'R'
- the length of QR.

THINK

1. Draw a diagram.

WRITE/DRAW

a.



2. Identify two corresponding sides. P'Q' corresponds to PQ.

$$\begin{aligned}
 \text{Scale factor} &= \frac{\text{image length}}{\text{object length}} \\
 &= \frac{P'Q'}{PQ} \\
 &= \frac{10}{4} \\
 &= 2.5
 \end{aligned}$$

1. Apply the scale factor. $P'R' = 2.5 \times PR$

$$\begin{aligned}
 \text{b. } P'R' &= 2.5 \times PR \\
 &= 2.5 \times 6 \\
 &= 15
 \end{aligned}$$

2. Write the answer.



P'R' is 15 cm long.

1. Apply the scale factor. $Q'R' = 2.5 \times QR$

$$\begin{aligned}
 \text{c. } \frac{Q'R'}{QR} &= \frac{20 \text{ cm}}{x \text{ cm}} = 2.5 \\
 Q'R' &= 2.5 \times QR \\
 20 &= 2.5 \times QR \\
 QR &= \frac{20}{2.5} \\
 &= 8
 \end{aligned}$$

2. Write the answer.

QR is 8 cm long.

-  **eWorkbook** Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)
-  **Interactivities** Individual pathway interactivity: Ratio and scale (int-4494)
Introduction to ratios (int-3733)
Proportion (int-3735)
Scale factors (int-6041)

Exercise 4.2 Ratio and scale

4.2 Quick quiz **on**

4.2 Exercise

Individual pathways

PRACTISE

1, 3, 5, 10, 13, 16, 17, 19, 22, 26

CONSOLIDATE

2, 4, 6, 8, 11, 15, 18, 21, 23, 27, 28

MASTER

7, 9, 12, 14, 20, 24, 25, 29, 30


Fluency

1. **WE1** The horse track shown is 1200 m long and 35 m wide.

a. Write the following ratios in simplest form.

- i. Track length to track width.
- ii. Track width to track length.

b. Compare the distance of the length of the track with the width of the track.




2. A dingo perched on top of a cliff spots an emu on the ground below.

a. Write the following ratios in simplest form.

- i. Cliff height to distance between cliff base and emu.
- ii. Distance between emu and cliff base to cliff height.

b. Compare the height of the cliff with the horizontal distance between the base of the cliff and the emu.



3. **WE2a** Express each of the following ratios in simplest form.

- a. 12 : 18
- b. 8 : 56
- c. 9 : 27
- d. 14 : 35

4. Express each of the following ratios in simplest form.

- a. 16 : 60
- b. 200 : 155
- c. 32 : 100
- d. 800 : 264


5. **WE2b** Express each of the following ratios in simplest form.

- a. 1.2 : 0.2
- b. 3.9 : 4.5
- c. 9.6 : 2.4
- d. 18 : 3.6


6. Express each of the following ratios in simplest form.
- a. $1.8 : 3.6$ b. $4.4 : 0.66$ c. $0.9 : 5.4$ d. $0.35 : 0.21$
7. Express each of the following ratios in simplest form.
- a. $6 : 1.2$ b. $12.1 : 5.5$ c. $8.6 : 4$ d. $0.07 : 14$
8. **WE2c** Write each of the following ratios in the simplest form.
- a. $1\frac{1}{2} : 2$ b. $2 : 1\frac{3}{4}$ c. $1\frac{1}{3} : 2$ d. $1\frac{2}{5} : 1\frac{1}{4}$
9. Write each of the following ratios in the simplest form.
- a. $\frac{4}{7} : 2$ b. $5 : 1\frac{1}{2}$ c. $2\frac{3}{4} : 1\frac{1}{3}$ d. $3\frac{5}{6} : 2\frac{1}{2}$
10. **WE3** Determine the value of the pronumeral in each of the following proportions.
- a. $a : 15 = 3 : 5$ b. $b : 18 = 4 : 3$ c. $24 : c = 3 : 4$
11. Determine the value of the pronumeral in each of the following proportions.
- a. $e : 33 = 5 : 44$ b. $6 : f = 5 : 12$ c. $3 : 4 = g : 5$
12. Determine the value of the pronumeral in each of the following proportions.
- a. $11 : 3 = i : 8$ b. $7 : 20 = 3 : j$ c. $15 : 13 = 12 : k$

For questions 13 to 15, enlarge the figures shown by the given scale factor and the centre of enlargement marked with O. Show the image of each of these figures.


13. **WE4** Scale factor = 2



14. Scale factor = 1.5




15. Scale factor = $\frac{1}{2}$



16. **WE5** A quadrilateral ABCD is enlarged to $A'B'C'D'$. $AB = 7$ cm, $AD = 4$ cm, $A'B' = 21$ cm, $B'C' = 10.5$ cm. Determine:
- a. the scale factor for enlargement
- b. $A'D'$
- c. BC.

Understanding

17. The estimated volume of Earth's salt water is 1 285 600 000 cubic kilometres. The estimated volume of fresh water is about 35 000 000 cubic kilometres.
- Determine the ratio of fresh water to salt water (in simplest form).
 - Determine the value of x , to the nearest whole number, when the ratio found in part a. is expressed in the form $1 : x$.
18. Super strength glue comes in two tubes that contain 'Part A' and 'Part B' pastes. These pastes have to be mixed in the ratio $1 : 4$ for maximum strength. Determine how many mL of Part A should be mixed with 10 mL of Part B.
19. A recipe for a tasty cake says the butter and flour needs to be combined in the ratio $2 : 7$. Determine the amount of butter that should be mixed with 3.5 kg of flour.
20. The diagram shown lays out the floor plan of a house. The actual size of bedroom 1 is $8 \text{ m} \times 4 \text{ m}$.



- If the dimensions of bedroom 1 as it appears on the ground plan are 4 cm by 2 cm, calculate the scale factor when the actual house (object) is built from the plan (image).
- Determine the real-life dimensions of bedroom 3 if the dimensions as shown on the ground plan are $3 \text{ cm} \times 3 \text{ cm}$.
- Determine the real-life dimensions of the kitchen if the dimensions as shown on the ground plan are $2.5 \text{ cm} \times 2.5 \text{ cm}$.

Reasoning

21. Pure gold is classed as 24-carat gold. This kind of gold is too soft to use for making jewellery, so it gets combined with other metals to form an alloy. The ratio of gold to other metals in 18-carat gold is 18 : 6.

The composition of 18-carat rose gold is 75% gold, 22.25% copper and 2.75% silver.

- a. Show that the mass of silver in a 2.5-gram rose gold bracelet is 0.07 g.
b. Determine the composition of metals in a rose gold bracelet that contains 0.5 g of copper.



22. The angles of a triangle have the ratio 3 : 4 : 5. Show that the sizes of the three angles are 45° , 60° and 75° .
23. The dimensions of a rectangular box have the ratio 2 : 3 : 5. The box's volume is $21\,870\text{ cm}^3$. Show that the dimensions of the box are 18, 27 and 45 cm.
24. Tyler, Dylan and Ari invested money in the ratio 11 : 9 : 4. If their profits are shared in the ratio 17 : 13 : 6, is this fair to each person? Explain your answer.
25. It costs the same amount to buy either 5 pens or 2 pens and 6 pencils. This is also the cost of 6 sharpeners and a pencil. Show a relationship between the cost of each kind of item.

Problem solving

26. Sharnee, a tourist at Kakadu National Park, takes a picture of a 2-metre-long crocodile beside a cliff. When they develops the pictures, they can see that on the photo the crocodile is 2.5 cm long and the cliff is 8.5 cm high. Determine the actual height of the cliff in cm.



27. Evaluate the ratio of $y : z$ if $2x = 3y$ and $3x = 4z$.
28. The quantities P and Q are in the ratio 2 : 3. If P is reduced by 1, the ratio becomes $\frac{1}{2}$. Determine the values of P and Q .
29. The ratio of boys to girls among the students who signed up for a basketball competition is 4 : 3. If 3 boys drop out of the competition and 4 girls join, there will be the same number of boys and girls. Evaluate the number of students who have signed up for the basketball competition.
30. In a group of students who voted in a Year 9 class president election, the ratio of girls to boys is 2 : 3. If 10 more girls and 5 more boys had voted, the ratio would have been 3 : 4. Evaluate the number of students who voted altogether.

LESSON

4.3 Congruent figures

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- identify congruent figures
- show that two triangles are congruent using the appropriate congruency test.

4.3.1 Congruent figures

eles-4729

- **Congruent figures** are identical figures. They have exactly the same shape and size.
- Congruent figures can be superimposed exactly on top of each other using reflection, rotation or translation (or a combination of some or all of these actions).




- The symbol for congruence is \cong . When reading this symbol out loud we say, 'is congruent to'.
- For the diagrams shown, $ABC \cong A'B'C'$ and $ABCDE \cong PQRST$.
- When writing congruence statements, the vertices of each figure are named in corresponding order.


WORKED EXAMPLE 6 Identifying congruent shapes

Identify a pair of congruent shapes from the following set.


a.




b.



c.



d.



THINK

Figures **a** and **c** are identical in shape and size. They just have different orientations.

WRITE

Shape **a** \cong shape **c**

4.3.2 Testing triangles for congruence

eles-4730

- To determine whether two triangles are congruent, it is not necessary to know that all three sides and all three angles of a each triangle are equal to the corresponding sides and angles of the other triangle.
- Certain minimum conditions can guarantee that two triangles are congruent.


Side-side-side condition for congruence (SSS)

- If all of the corresponding sides of two triangles are equal, the angles opposite those corresponding sides will also be equal. This means that the two triangles are congruent.
- This is known as the *side-side-side* (SSS) condition for congruence.




Side-angle-side condition for congruence (SAS)

- If two triangles have two corresponding sides that are equal, and the angles between those corresponding sides are equal, then the two triangles are congruent.
- This is known as the *side-angle-side* (SAS) condition for congruence.



Angle-side-angle condition for congruence (ASA)

- If two triangles have two corresponding angles that are equal, and 1 side that is also equal, then the two triangles are congruent. (*Note:* The third corresponding angle will also be equal.)
- This is known as the *angle-side-angle* (ASA) condition for congruence.







Right angle-hypotenuse-side condition for congruence (RHS)

- If the hypotenuse and one other side of two right-angled triangles are equal, then the two triangles are congruent.
- This is known as the *right angle-hypotenuse-side* (RHS) condition for congruence.




Summary of congruence tests

Congruence test	Description	Abbreviation
	<ul style="list-style-type: none"> • All corresponding sides are equal in length. 	SSS (side-side-side)
	<ul style="list-style-type: none"> • Two corresponding sides are equal in length. • The corresponding angles between them are equal in size. 	SAS (side-angle-side)
	<ul style="list-style-type: none"> • Two corresponding angles are equal in size. • One corresponding side is equal in length. 	ASA (angle-side-angle)
	<ul style="list-style-type: none"> • The hypotenuse and one other corresponding side of two right-angled triangles are equal in length. 	RHS (right angle-hypotenuse-side)

WORKED EXAMPLE 7 Identifying congruent triangles

Identify which of the following triangles are congruent. Give reasons for your answer.



THINK


In all three triangles two given sides are of equal length (2 cm and 5 cm). The included angles in triangles ABC and KLM are also equal (60°). B corresponds to L, and A corresponds to M.

WRITE

$$\triangle ABC \cong \triangle KLM \text{ (SAS)}$$

WORKED EXAMPLE 8 Using congruent triangles to determine values of the pronumerals

Given that $\triangle ABD \cong \triangle CBD$, determine the values of the pronumerals in the figure shown.



THINK

1. Congruent triangles have corresponding sides that are equal in length. Side AD (marked x) corresponds to side CD.
2. Since these triangles are congruent, the corresponding angles are equal.

WRITE

$$\triangle ABD \cong \triangle CBD$$

$$AD = CD$$

$$x = 3 \text{ cm}$$

$$\angle A = \angle C$$


$$y = 40^\circ$$

$$\angle BDA = \angle BDC$$

$$z = 90^\circ$$

WORKED EXAMPLE 9 Proving that two triangles are congruent

Prove that $\triangle PQS$ is congruent to $\triangle RQS$.



THINK


1. Study the diagram and state which sides and/or angles are equal.
2. This fits the SSS condition, which proves congruence.

WRITE

$QP = QR$ (given)
 $PS = RS$ (given)
 QS is common.
 $\triangle PQS \cong \triangle RQS$ (SSS)

COLLABORATIVE TASK

On a piece of paper, draw the trapezium shown as per the dimensions given in the diagram. Working in pairs or small teams, try to divide the trapezium into four congruent trapeziums that are similar in shape to the original trapezium.

**on Resources**

eWorkbook Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)



Interactivities Individual pathway interactivity: Congruent figures (int-4495)

Congruency tests (int-3755)

Congruent figures (int-3749)

Congruent triangles (int-3754)

Exercise 4.3 Congruent figures**learn on****4.3 Quick quiz** **on****4.3 Exercise****Individual pathways****PRACTISE**

1, 4, 5, 8, 11, 12, 16, 17, 22

CONSOLIDATE

2, 6, 9, 13, 14, 18, 19, 23


MASTER

3, 7, 10, 15, 20, 21, 24, 25

Fluency

1. **WE6** Select a pair of congruent shapes from the figures shown.

a.




b.







c.



d.



2. Select a pair of congruent shapes from the figures shown.









3. Select a pair of congruent shapes from the figures shown.






Understanding




4. **MC** Select which of the following is congruent to the triangle shown.






5. **WE7** Identify which of the following triangles are congruent. Give a reason for your answer.






6. Identify which of the following triangles are congruent. Give a reason for your answer.






7. Identify which of the triangles are congruent. Give a reason for your answer.




8. Identify which of the following triangles are congruent. Give a reason for your answer.





9. Identify which of the following triangles are congruent. Give a reason for your answer.




10. Identify which of the following triangles are congruent. Give a reason for your answer.



11. **WEB** Determine the value of the pronumeral in the following pair of congruent triangles. All side lengths are in centimetres.





12. Determine the value of the pronumeral in the pair of congruent triangles shown. All side lengths are in centimetres.







13. Determine the value of the pronumerals in the pair of congruent triangles shown. All side lengths are in centimetres.



14. Determine the value of the pronumerals in the pair of congruent triangles shown. All side lengths are in centimetres.






15. Calculate the length of the side marked with the pronumeral using the following congruent triangles.




Reasoning

16. Do congruent figures have the same area? Explain your answer.
17. If two congruent triangles have a right angle, is the reason for their congruence always due to the RHS condition of congruence? Justify your answer.
18. Give an example to show that triangles with two angles of equal size and a pair of non-corresponding sides of equal length may not be congruent.
19. **WE9** For each of the following, prove that the statement accompanying the diagram is true.




20. ABCD is a trapezium with both AD and BC perpendicular to AB. If a right-angled triangle DEC is constructed with an angle $\angle ECD$ equal to 45° , prove that $\triangle EDA \cong \triangle ECB$.
21. A teacher asked their class to each draw a triangle with side lengths of 5 cm and 4 cm, and an angle of 45° that is not formed at the point joining the 5 cm and 4 cm side. Explain why the triangles drawn by every member of the class would be congruent.




Problem solving

22. Construct five congruent triangles from nine matchsticks.
23. Construct seven congruent triangles from nine matchsticks.
24. Demonstrate how the figure shown can be cut into four congruent pieces.



25. Determine the ratio of the outer (unshaded) area to the inner (shaded) area of the six-pointed star shown.



LESSON

4.4 Nets, polyhedra and Euler's formula

LEARNING INTENTIONS

At the end of this lesson students should be able to:

- identify and produce nets of various solids
- define polyhedra and identify examples of Platonic solids
- verify Euler's formula for various polyhedra.


4.4.1 Nets

eles-4731

- The **faces** forming a solid can be drawn as plane shapes, which are joined across the edges to form the solid.
- The complete set of faces forming a solid is called its net. Note that for some figures, different nets can be drawn.

WORKED EXAMPLE 10 Drawing a possible net for a cone





Draw a possible net for a cone, which has a radius of 7 cm and a slanting height of 10 cm.



THINK

1. The base of a cone is a circle of radius 7 cm.
2. The other (slanted) part of the cone when split open will form a sector of a circle of radius 10 cm.
3. If the two parts (from steps 1 and 2) are put together, the complete net of a cone is obtained.

DRAW




4.4.2 Polyhedra construction

eles-4839


- A 3-dimensional solid where each of the faces is a polygon is called a **polyhedron**. If all the faces are congruent, the solid is called a regular polyhedron or a Platonic solid.

• Platonic solids are shown below.


- A **cube**, with 6 faces, each of which is a square.




- A **tetrahedron**, with 4 faces, each of which is an equilateral triangle.




- An **octahedron**, with 8 faces, each of which is an equilateral triangle.



- A **dodecahedron**, with 12 faces, each of which is a regular pentagon.



- An **icosahedron**, with 20 faces, each of which is an equilateral triangle.



• Construction of polyhedra is conveniently done from nets.


WORKED EXAMPLE 11 Drawing a net of an octahedron

Draw a net of an octahedron.

THINK

An octahedron is a polyhedron with 8 faces, each of which is an equilateral triangle. So its net will consist of 8 equilateral triangles. Draw a possible net of an octahedron.

DRAW



4.4.3 Euler's formula

eles-4840

- **Euler's formula** shows the relationship between the number of **edges**, the number of **faces** and the number of vertices in any polyhedron. Note that a **vertex** (the singular of vertices) is a point or a corner of a shape where the straight edges meet.


Euler's formula

Euler's formula states that for any polyhedron:

$$\text{number of faces } (F) + \text{number of vertices } (V) - 2 = \text{number of edges } (E)$$
$$F + V - 2 = E$$

WORKED EXAMPLE 12

Verify Euler's formula for a tetrahedron.



THINK

1. Look at the shape of a tetrahedron and state the number of faces, vertices and edges.
2. Write Euler's formula.
3. To verify Euler's formula, show that the left-hand side (LHS) is equal to right-hand side (RHS). Substitute the values of F , V and E in the formula.
4. Write the conclusion.

WRITE

In a tetrahedron there are four faces, four vertices and six edges.

number of faces (F) + number of vertices (V) - 2 =
number of edges (E)

$$F = 4, V = 4 \text{ and } E = 6$$

$$\text{LHS} = F + V - 2$$

$$= 4 + 4 - 2$$

$$= 6$$

$$\text{RHS} = 6$$

$$\text{LHS} = \text{RHS}$$

Euler's formula holds for a tetrahedron.

on Resources



eWorkbook Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)



Interactivities Individual pathway interactivity: Nets, polyhedra and Euler's rule (int-4428)
Nets (int-3759)
Polyhedra construction (int-3760)
Euler's rule (int-3761)

Exercise 4.4 Nets, polyhedra and Euler's formula

learnon

4.4 Quick quiz **on**

4.4 Exercise

Individual pathways

PRACTISE

1, 3, 7, 11, 14

CONSOLIDATE


2, 5, 8, 10, 12, 15

MASTER

4, 6, 9, 13, 16


Fluency

1. **WE10** Draw a possible net for the cylinder shown.




2. Draw a possible net for each of the following solids.


a.



b.




c.



3. **WE11** Draw 2 different nets to form a cube.

4. Draw a net for the octahedron that is different from the one shown. Cut out your net and fold it to form the octahedron.




5. Using nets generated from geometry software or elsewhere, construct:


- a. a cube
- b. a tetrahedron
- c. a dodecahedron.

6. Construct the pyramids shown in the figures.

a.



b.



Understanding


7. Draw some 3-dimensional shapes of your choice. State how many of these shapes are polyhedra, and name them.

8. **WE12** Verify Euler's formula for the following Platonic solids.


- a. A cube.
- b. An octahedron.

9. Show that Euler's formula holds true for these solids.


a.



b.




c.




10. Make the prisms shown. Verify Euler's formula for each shape.

a.



b.




Reasoning


11. Discuss why a knowledge of 3D shapes is important.

12. Explain whether you can verify Euler's formula for the following shapes. Give reasons.


a.



b.



c.



13. A spherical scoop of ice-cream sits on a cone. Draw this shape. Explain whether you can verify Euler's formula for this shape.




Problem solving


14. Renee knows that a polyhedron has 12 faces and 8 vertices. Show how she can determine the number of edges.

15. Determine which of the following compound shapes are nets that can be folded into a 3D solid. You may wish to enlarge them onto paper, cut them out and fold them.


a.




b.



c.



16. The net forms an icosahedron as shown. Verify Euler's formula for this shape, showing your working.



LESSON

4.5 Similar figures

LEARNING INTENTIONS

At the end of this lesson you should be able to:



- identify similar figures
- determine the scale factor between similar figures
- show that two triangles are similar using the appropriate similarity test.



4.5.1 Similar figures and similarity condition

eles-4732

- **Similar figures** are identical in shape, but different in size.
- The corresponding angles in similar figures are equal in size and the corresponding sides are in the same ratio.
- The symbol used to denote similarity is \sim . When reading this symbol out loud we say, 'is similar to'.
- In the triangles shown $\triangle ABC$ is similar to $\triangle UVW$. That is, $\triangle ABC \sim \triangle UVW$.
- The ratio of side lengths is known as the scale factor.



Scale factor

$$\text{Scale factor} = \frac{\text{length of image}}{\text{length of object}}$$

- If the scale factor is > 1 , an enlargement has occurred.
- If the scale factor is < 1 , a reduction has occurred.

- The scale factor for the triangles shown is 2. Each side in UVW is twice the length of the corresponding side in ABC.
- Enlargements and reductions are transformations that create similar figures.


- The method for creating enlarged figures that is explained in lesson 4.2 can also be used to create similar figures.

Testing triangles for similarity

- As with congruent triangles, to determine whether two triangles are similar, it is not necessary to know that all pairs of corresponding sides are in the same ratio and that all corresponding angles are equal.
- Certain minimum conditions can guarantee that two triangles are similar.


Angle–angle–angle condition of similarity (AAA)

- If all of the angles of two triangles are equal, then the triangles are similar.
- This is known as the angle–angle–angle (AAA) condition for similarity.
- In the diagram shown, $\triangle ABC \sim \triangle RST$ (AAA).




Side–side–side condition for similarity (SSS)

- If two triangles have a constant ratio for all corresponding side lengths, then the two triangles are similar.
- This is known as the side–side–side (SSS) condition for similarity.
- In the diagram shown, the ratios of all corresponding side lengths are equal $\left(\frac{9}{6} = \frac{15}{10} = \frac{10.5}{7} = 1.5\right)$, therefore $\triangle ABC \sim \triangle RST$ (SSS).




Side–angle–side condition for similarity (SAS)

- If two triangles have two corresponding sides in the same ratio and the included angles of those sides are equal, then the two triangles are similar.
- This is known as the side–angle–side (SAS) condition for similarity.
- In the diagram shown, the ratio of the triangles' two corresponding side lengths are equal $\left(\frac{9}{6} = \frac{15}{10} = 1.5\right)$ and the included angles are also the same, therefore $\triangle ABC \sim \triangle RST$ (SAS).



Right angle–hypotenuse–side condition for similarity (RHS)

- If the hypotenuse and one other corresponding side of two right-angled triangles are in the same ratio, then the two triangles are similar.
- This is known as the the right angle–hypotenuse–side (RHS) condition for similarity.
- In the diagram shown, the ratio of the hypotenuses and one other pair of corresponding sides are equal $\left(\frac{12}{6} = \frac{10}{5} = 2\right)$, therefore $\triangle ABC \sim \triangle RST$ (RHS).



Summary of similarity tests

- Triangles can be checked for similarity using one of the tests described in the table shown.


Test description	Abbreviation
All corresponding angles are equal in size	AAA (angle–angle–angle)
All corresponding sides are in the same ratio	SSS (side–side–side)
Two corresponding sides are in the same ratio and the included angles are equal in size	SAS (side–angle–side)
The hypotenuse and one other corresponding side of two right-angled triangles are in the same ratio	RHS (right angle–hypotenuse–side)

Note: When using the AAA test, it is sufficient to show that two corresponding angles are equal. Since the sum of the interior angles in any triangle is 180° , the third corresponding angle will automatically be equal.


WORKED EXAMPLE 13 Identifying similar triangles

Identify a pair of similar triangles from the triangles shown. Give a reason for your answer.


a.



b.



c.



THINK

1. In each triangle we know the size of two sides and the included angle, so the SAS test can be applied. Since all included angles are equal (30°), we need to determine the ratios of the corresponding sides, looking at two triangles at a time.

2. Write the answer.

WRITE

For triangles a and b:

$$\frac{15}{10} = \frac{9}{6} = 1.5$$


For triangles a and c:

$$\frac{20}{10} = 2, \frac{15}{6} = 2.5$$

Triangle a \sim triangle b (SAS)

WORKED EXAMPLE 14 Proving that two triangles are similar

Prove that $\triangle ABC$ is similar to $\triangle EDC$.



THINK

1. AB is parallel to DE. $\angle ABC$ and $\angle EDC$ are alternate angles.
2. $\angle BAC$ and $\angle DEC$ are alternate angles.
3. The third pair of angles must be equal.
4. This proves that the triangles are similar.


WRITE

- $\angle ABC = \angle EDC$ (alternate angles)
- $\angle BAC = \angle DEC$ (alternate angles)
- $\angle BCA = \angle DCE$ (vertically opposite angles)
- $\triangle ABC \sim \triangle EDC$ (AAA)

- The ratio of the corresponding sides in similar figures can be used to calculate missing side lengths or angles in those figures.

WORKED EXAMPLE 15 Solving worded problems using similar triangles


A pole 1.5 metres high casts a shadow 3 metres long, as shown. Calculate the height of a building that casts a shadow 15 metres long at the same time of the day.



THINK

1. Represent the given information on a diagram.
 $\angle BAC = \angle EDC$; $\angle BCA = \angle ECD$
2. Triangles ABC and DEC are similar. This means the ratios of corresponding sides are the same. Write the ratios.

WRITE/DRAW



$$\triangle ABC \sim \triangle DEC \text{ (AAA)}$$

$$\frac{h}{1.5} = \frac{15}{3}$$

3. Solve the equation for h .

$$h = \frac{15 \times 1.5}{3}$$

$$= 7.5$$

4. Write the answer in words, including units.

The building is 7.5 metres high.

Digital tools to develop similar shapes

We have looked at similar shapes that get bigger with a scale factor greater than one and a similar shape that reduces in size with a scale factor greater than zero and less than one.

When a shape is enlarged or reduced:

- Angles at the vertex stay the same.
- The side lengths increase or decrease.

These similar shapes can be produced by using CAS technology, as shown.


Digital technology

As an example, we will draw a square and enlarge it with a scale factor of 3 and then reduce it with a scale factor of 0.5.


TI | THINK

1. On a Calculator page, press MENU and select:
 - 3 Add Geometry

DISPLAY/WRITE



2. Select
 - 2 View and click on 1 Graphing. Then select 6 Grid and click on 2 Dot Grid.





CASIO | THINK


1. On the Main screen tap:
 - Geometry


DISPLAY/WRITE




2. After selecting the Geometry screen:
 - Tap  to create a grid. (You might need to tap this multiple times to get the grid).




3. Click on  and select:
9 Settings and tick
Automatically label points
then click OK.




4. To draw a shape, press
MENU and select:
8 Geometry
2 Shapes
To select a rectangle, select
3 Rectangle




5. Tap on the screen where you
want one of the corners to
be, then drag to the size of
the shape you require.





6. From the Menu, click on
8 Geometry,
then select
1 Points & Lines and click
on 1 Point.
Right click on the point and
select 5 Text.




7. Add text to the screen. Text
being a numerical quantity
'3' in this case.






3. Tap:
 to select a
rectangle, or the arrow
next to it to choose other
shapes.




4. Tap on the screen where
you want one of your
corners to be, then drag
to the size of the shape you
require.



5. Tap:
 -  on the arrow as
shown.
 -  to enlarge/reduce
the size of the shape.




6. Tap the centre of the
shape you want to
enlarge/reduce.
Then type in 3 for the
scale factor and then OK.




8. To enlarge the rectangle by a factor of 3. Select from the Menu:

8 Geometry
5 Transformation
5 Dilation

Dilation of the rectangle by a factor of 3 about a point A. The diagram shows the new shape increased by a scale factor of 3.




9. Completing the same process but reducing the size of the shape by using a scale factor of 0.5.





8. The diagram shows the new shape increased by a scale factor of 3.

This shows that the angle at the vertices stay the same (90°), but the side lengths are three times longer.




9. Starting with the shape shown, complete the same process but reducing the size of the shape by using a scale factor of 0.5. Use the same process to input a scale factor of 0.5.



10. Tap:

- OK

The shape again has the same angles at the vertices (90°), but the side lengths are half their original length to produce the similar shape.



Resources

- eWorkbook** Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)
- Video eLesson** Similar triangles (eles-1925)
- Interactivity** Individual pathway interactivity: Similar figures (int-4496)

4.5 Quick quiz **on**

4.5 Exercise

Individual pathways

PRACTISE

1, 2, 6, 7, 11, 14, 15, 19, 20

CONSOLIDATE

3, 4, 8, 12, 16, 21


MASTER

5, 9, 10, 13, 17, 18, 22, 23, 24


Fluency

WE13 For questions 1 to 5, Identify the pair of similar triangles among those shown. Give reasons for your answers.


1. a.




b.




c.




2. a.




b.




c.




3. a.




b.




c.




4. a.




b.




c.




5. a.



b.








c.




Understanding

6. Name two similar triangles in each of the following figures, ensuring that vertices are listed in the correct order.




7. In the diagram shown, C is the centre of the circle. Complete this statement: $\triangle ABC$ is similar to ...



8. For the diagram shown:

a. complete the statement: $\frac{AB}{AD} = \frac{BC}{\square} = \frac{\square}{AE}$


b. determine the value of the pronumeral.




9. Using the diagram shown:

a. determine the values of h and i



b. determine the values of j and k .




10. Determine the value of the pronumeral in the diagram shown.



11. If the two triangles shown are similar, determine the values of the pronumerals x and y .





12. Determine the values of the pronumerals x and y in the diagram shown.





13. Determine the value of each pronumeral in the following triangles. Show how you arrived at your answers.



a.



b.




c.







Reasoning

14. **WE15** A ladder just touches a bench and also leans on a wall that is 4 metres high, as shown in the diagram. If the bench is 50 centimetres high and 1 metre from the base of the ladder, show that the base of the ladder is 8 metres from the wall.



15. **WE14** Prove that $\triangle ABC$ is similar to $\triangle EDC$ in each of the following.




16. Natalie is 1.5 metres tall. They cast a shadow 2 metres long at a certain time of the day.



If Alex is 1.8 metres tall, show that their shadow would be 2.4 m long at the same time of day.

17. A string 50 metres long is pegged to the ground and tied to the top of a flagpole. It just touches Maz on the top of her head. If Maz is 1.5 metres tall and 5 metres away from the point where the string is held to the ground, show that the height, h , of the flagpole is 14.37 m.



18. Using diagrams or other methods, explain whether the following statements are True or False.
- All equilateral triangles are similar.
 - All isosceles triangles are similar.
 - All right-angled triangles are similar.
 - All right-angled isosceles triangles are similar.

Problem solving

19. Paw and Thuy play tennis at night under floodlights. When Paw stands 2.5 m from the base of the floodlight, her shadow is 60 cm long.


- If Paw is 1.3 m tall, evaluate the height of the floodlight in metres, correct to 2 decimal places.
- If Thuy, who is 1.6 m tall, stands in the same place, calculate her shadow length in cm.



20. To determine the height of a flagpole, Jenna and Mia decide to measure the shadow cast by the flagpole. They place a 1 m ruler at a distance of 3 m from the base of the flagpole and measure the shadows that both the ruler and flagpole cast. Both shadows finish at the same point.


After measuring the shadow of the flagpole, Jenna and Mia calculate that the height of the flagpole is 5 m. Determine the length of the shadow cast by the flagpole, as measured by Jenna and Mia. Give your answer in metres.

21. Use the diagram shown to determine the value of a if $XZ = 8$ cm, $X'Z' = 12$ cm, $X'X = a$ cm and $XY = (a + 1)$ cm.




22. AB and CD are parallel lines in the figure shown.

- State the similar triangles.
- Determine the values of x and y .




23. PQ is the diameter of the circle shown. The circle's centre is located at S. R is any point on the circumference. T is the midpoint of PR.

- Write down everything you know about this figure.
- Explain why $\triangle PTS$ is similar to $\triangle PRQ$.
- Determine the length of TS if RQ is 8 cm.
- Determine the length of every other side, given that PT is 3 cm and the angle PRQ is a right angle.



24. For the diagram shown, show that, if the base of the triangle is raised to half of the height of the triangle, the length of the base of the newly formed triangle will be half of its original length.



LESSON

4.6 Area and volume of similar figures

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- convert between different units of length, area and volume
- determine the area of similar figures, given a scale factor of n^2
- determine the volume of similar figures, given a scale factor of n^3 .

4.6.1 Converting between units of length, area and volume

eles-4733


Units of length

- Metric units of length include millimetres (mm), centimetres (cm), metres (m) and kilometres (km).
- Length units can be converted using the chart shown.



Units of area


- **Area** is measured in square units, such as square millimetres (mm^2), square centimetres (cm^2), square metres (m^2) and square kilometres (km^2).
- Area units can be converted using the chart shown. Note, $10^2 = 100$; $100^2 = 10\,000$; $1000^2 = 1\,000\,000$



- Area units are the squares of the corresponding length units.

Units of volume

- **Volume** is measured in cubic units, such as cubic millimetres (mm^3), cubic centimetres (cm^3) and cubic metres (m^3).
- Volume units can be converted using the chart shown. Note, $10^3 = 1000$; $100^3 = 1\,000\,000$; $1000^3 = 1\,000\,000\,000$




- Volume units are the cubes of the corresponding length units.

WORKED EXAMPLE 16 Converting between units of measurements

- Convert 9 m into mm.
- Convert 150 cm^2 into mm^2 .
- A cube has a side length of 8 cm. Calculate the volume of the cube in m^3 .


THINK

1. To convert m to mm:



- To convert 9 m to mm, multiply 9 with 1000.
- Simplify and write the answer.

1. To convert cm^2 into mm^2 :



- To convert 150 cm^2 into mm^2 , multiply 150 by 10^2 (100).
- Simplify and write the answer.

1. Write the formula for the volume of a cube.
2. Substitute the value of the side length (l) in the volume formula.

WRITE

- $1\text{ m} = 1000\text{ mm}$

$$9\text{ m} = 9 \times 1000\text{ mm}$$

$$9\text{ m is } 9000\text{ mm.}$$

- $1\text{ cm}^2 = 100\text{mm}^2$


$$150\text{ cm}^2 = 150 \times 100\text{ mm}^2$$

$$150\text{ cm}^2 \text{ is } 15\,000\text{ mm}^2.$$

- Volume of a cube (V) = $l \times l \times l$, where l is the side length.

$$\begin{aligned} V &= 8 \times 8 \times 8 \\ &= 512\text{ cm}^3 \end{aligned}$$

3. The question states that the answer should be given in m^3 .
To convert cm^3 into m^3 :



$$1 \text{ cm}^3 = \frac{1}{100^3} \text{ m}^3$$

4. To convert 512 cm^3 into m^3 , multiply 512 by $\frac{1}{100^3} \text{ m}^3$.

$$512 \text{ cm}^3 = \frac{512}{100^3} \text{ m}^3$$

5. Simplify and write the answer.

The volume of the cube is $0.000 512 \text{ m}^3$.



4.6.2 Area and volume of similar figures

eles-4734

Area and surface area of similar figures

- If the side lengths of any figure are increased by a scale factor of n , then the area of similar figures increases by a scale factor of n^2 .

For example, consider the squares shown.



$$\begin{aligned} \text{Area A} &= 2 \times 2 \\ &= 4 \text{ cm}^2 \end{aligned}$$



$$\begin{aligned} \text{Area B} &= 4 \times 4 \\ &= 16 \text{ cm}^2 \end{aligned}$$




$$\begin{aligned} \text{Area C} &= 6 \times 6 \\ &= 36 \text{ cm}^2 \end{aligned}$$

- The scale factors for the side lengths and the scale factors for the areas are calculated in the table shown.

Squares	Scale factor for side length	Scale factor for area
A and B	$\frac{4}{2} = 2$	$\frac{16}{4} = 4 = 2^2$
A and C	$\frac{6}{2} = 3$	$\frac{36}{4} = 9 = 3^2$
B and C	$\frac{6}{4} = \frac{3}{2}$	$\frac{36}{16} = \frac{9}{4} = \left(\frac{3}{2}\right)^2$

- The surface area of a 3D object also increases by the square of the length scale factor. For example, consider the cubes shown.



$$\begin{aligned} \text{Surface area A} &= 6 \times 4 \\ &= 24 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{Surface area B} &= 6 \times 16 \\ &= 96 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{Surface area C} &= 6 \times 36 \\ &= 216 \text{ cm}^2 \end{aligned}$$

- The scale factors for the side lengths and the scale factors for the surface areas are calculated in the table shown.


Cubes	Scale factor for side length	Scale factor for surface area
A and B	$\frac{4}{2} = 2$	$\frac{96}{24} = 4 = 2^2$
A and C	$\frac{6}{2} = 3$	$\frac{216}{24} = 9 = 3^2$
B and C	$\frac{6}{4} = \frac{3}{2}$	$\frac{216}{96} = \frac{9}{4} = \left(\frac{3}{2}\right)^2$

Areas of similar figures

When side lengths are increased by a factor of n , the area increases by a factor of n^2 .

Volume of similar figures

- If the side lengths of any solid are increased by a scale factor of n , then the volume of similar solids increases by a scale factor of n^3 . For example, consider the cubes shown.



$$\begin{aligned} \text{Volume A} &= 2 \times 2 \times 2 \\ &= 8 \text{ cm}^3 \end{aligned}$$

$$\begin{aligned} \text{Volume B} &= 4 \times 4 \times 4 \\ &= 64 \text{ cm}^3 \end{aligned}$$

$$\begin{aligned} \text{Volume C} &= 6 \times 6 \times 6 \\ &= 216 \text{ cm}^3 \end{aligned}$$

- The scale factors for the side lengths and the scale factors for the volumes are calculated in the table shown.

Cubes	Scale factor for side length	Scale factor for volume
A and B	$\frac{4}{2} = 2$	$\frac{64}{8} = 8 = 2^3$
A and C	$\frac{6}{2} = 3$	$\frac{216}{8} = 27 = 3^3$
B and C	$\frac{6}{4} = \frac{3}{2}$	$\frac{216}{64} = \frac{27}{8} = \left(\frac{3}{2}\right)^3$

Volumes of similar figures

When side lengths are increased by a factor of n , the volume increases by a factor of n^3 .

WORKED EXAMPLE 17 Calculating area and volume of similar figures

The side lengths of a box have been increased by a factor of 3.

- Calculate the surface area of the new box if the original surface area was 94 cm^2 .
- Determine the volume of the new box if the original volume was 60 cm^3 .



THINK

- State the scale factor for side length used to increase the size of the original box.
 - The scale factor for surface area is the square of the scale factor for length.
 - Calculate the surface area of the new box.
- The scale factor for volume is the cube of the scale factor for length.
 - Calculate the volume of the new box.

WRITE

- Scale factor for side length = 3

$$\begin{aligned} \text{Scale factor for surface area} &= 3^2 \\ &= 9 \end{aligned}$$

$$\begin{aligned} \text{Surface area of new box} &= 94 \times 9 \\ &= 846 \text{ cm}^2 \end{aligned}$$

- Scale factor for volume = 3^3
 - $= 27$

$$\begin{aligned} \text{Volume of new box} &= 60 \times 27 \\ &= 1620 \text{ cm}^3 \end{aligned}$$

WORKED EXAMPLE 18 Using technology to calculate volume for similar figures

A small cone has a radius of 5 cm and a height of 15 cm.

- If the scale factor is 4, determine the dimensions of a larger similar cone.
- Giving your answers both in exact form and correct to 3 decimal places, calculate the volume of:
 - the smaller cone
 - the larger cone.
- Show that the volumes of the two cones are in the ratio 1 : 64.

THINK

- The scale factor is 4, so multiply the radius and height by 4. Write the answers.
 - Use your calculator to work out the volume of the smaller cone by substituting the following into the formula:
 $r = 5$
 $h = 15$.
 - Use your calculator to work out the volume of the larger cone (V_L) by substituting the following into the formula:
 $r = 20$
 $h = 60$.
- Use your calculator to evaluate the ratio of the volumes of the smaller cone and the larger cone.

$$\frac{V_s}{V_L} = \frac{125\pi}{8000\pi}$$

WRITE

- For the larger cone:
 radius = $5 \text{ cm} \times 4 = 20 \text{ cm}$
 height = $15 \text{ cm} \times 4 = 60 \text{ cm}$
- $$V_s = \frac{1}{3}\pi r^2 h$$

$$= \frac{1}{3}\pi \times 5^2 \times 15$$

$$= 125\pi \text{ cm}^3$$

$$\approx 392.699 \text{ cm}^3 \text{ (to 3 d.p.)}$$
 - $$V_L = \frac{1}{3}\pi r^2 h$$

$$= \frac{1}{3}\pi \times 20^2 \times 60$$

$$= 8000\pi$$


$$\approx 25\,132.741 \text{ cm}^3 \text{ (to 3 d.p.)}$$
- $$\frac{V_s}{V_L} = \frac{1}{64}$$

The volumes are in the ratio 1 : 64.

TI | THINK

- a-b.** In a new problem, on a Calculator page, complete the entry line as:
- $$\frac{1}{3} \times \pi \times 5^2 \times 15$$
- Then press ENTER.
- $$\frac{1}{3} \times \pi \times 20^2 \times 60$$
- Then press ENTER.
- To convert the answer to decimal:
- Menu
 - 2: Number
 - 1: Convert to Decimal
- Then press ENTER.

DISPLAY/WRITE



$$V_s = 125\pi \text{ cm}^3$$

$$V_s \approx 392.699 \text{ cm}^3 \text{ (to 3 d.p.)}$$

$$V_L = 8000\pi \text{ cm}^3$$

$$V_L \approx 25\,132.741 \text{ cm}^3 \text{ (to 3 d.p.)}$$

CASIO | THINK

- a-b.** On the Main screen, complete the entry line as:
- $$\frac{1}{3} \times \pi \times 5^2 \times 15$$
- Then press EXE.
- $$\frac{1}{3} \times \pi \times 20^2 \times 60$$
- Then press EXE.
- Note: Change Standard to Decimal

DISPLAY/WRITE





$$V_s = 125\pi \text{ cm}^3$$


$$V_s \approx 392.699 \text{ cm}^3 \text{ (to 3 d.p.)}$$

$$V_L = 8000\pi \text{ cm}^3$$

$$V_L \approx 25\,132.741 \text{ cm}^3 \text{ (to 3 d.p.)}$$

-  **eWorkbook** Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)
-  **Interactivities** Individual pathway interactivity: Area and volume of similar figures (int-4497)
 - Units of length (int-3779)
 - Area of similar figures (int-6043)
 - Volume and surface area of similar figures (int-6044)

Exercise 4.6 Area and volume of similar figures

4.6 Quick quiz 

4.6 Exercise

Individual pathways

PRACTISE

1, 3, 4, 8, 9, 13

CONSOLIDATE

2, 5, 7, 10, 14, 15

MASTER

6, 11, 12, 16, 17, 18

Fluency

1. **WE17a** The side lengths of the following shapes have all been increased by a factor of 3. Copy and complete the table shown.

Original surface area	Enlarged surface area
100 cm ²	a.
7.5 cm ²	b.
95 mm ²	c.
d.	918 cm ²
e.	45 m ²
f.	225 mm ²

2. A rectangular box has a surface area of 96 cm² and volume of 36 cm³. Calculate the volume and surface area of a similar box that has side lengths that are double the size of the original box.
3. **WE17b** The side lengths of the following shapes have all been increased by a factor of 3. Copy and complete the table shown.

Original volume	Enlarged volume
200 cm ³	a.
12.5 cm ³	b.
67 mm ³	c.
d.	2700 cm ³
e.	67.5 m ³
f.	27 mm ³


Understanding

4. The area of a bathroom as drawn on a house plan is 5 cm^2 . Calculate the area of the actual bathroom if the map has a scale of $1 : 100$.
5. The area of a kitchen is 25 m^2 .
 - a. **WE16** Convert 25 m^2 to cm^2 .
 - b. Calculate the area of the kitchen as drawn on a plan if the scale of the plan is $1 : 120$. (Give your answer correct to 1 decimal place.)
6. The volume of a swimming pool as it appears on its construction plan is 20 cm^3 . Determine the actual volume of the pool if the plan has a scale of $1 : 75$.
7. The total surface area of an aeroplane's wings is 120 m^3 .
 - a. Convert 120 m^2 to cm^2 .
 - b. Calculate the total surface area of the wings of a scale model of the aeroplane if the model is built using the scale $1 : 80$.



Reasoning


8. A triangle ABC maps to triangle $A'B'C'$ under an enlargement.
 $AB = 7 \text{ cm}$, $AC = 5 \text{ cm}$, $A'B' = 21 \text{ cm}$, $B'C' = 30 \text{ cm}$.
 - a. Show that the scale factor for enlargement is 3.
 - b. Determine BC .
 - c. Determine $A'C'$.
 - d. If the area of $\triangle ABC$ is 9 cm^2 , show that the area of $\triangle A'B'C'$ is 81 cm^2 .
9. A pentagon has an area of 20 cm^2 . If all the side lengths are doubled, show that the area of the enlarged pentagon is 80 cm^2 .
10. Two rectangles are similar. If the width of the first rectangle is twice of width of the other, prove that the ratio of their areas is $4 : 1$.
11. A cube has a surface area of 253.5 cm^2 . (Give answers correct to 1 decimal place where appropriate.)
 - a. Show that the side length of the cube is 6.5 cm .
 - b. Show that the volume of the cube is 274.625 cm^3 .
 - c. Determine the volume of a similar cube that has side lengths twice as long.
 - d. Determine the volume of a similar cube that has side lengths half as long.
 - e. Determine the surface area of a similar cube that has side lengths one third as long.
12. In the diagram shown, a light is shining through a hole, resulting in a circular bright spot with a radius of 5 cm on the screen.



The hole is 10 mm wide. If the light is 1 m behind the hole, show that the light is 10 m from the screen.

Problem solving

13. The areas of two similar trapeziums are 9 and 25. Determine the ratio of one pair of these trapeziums' corresponding side lengths.
14. Answer the following questions.
- Calculate the areas of squares with sides 2 cm, 5 cm, 10 cm and 20 cm.
 - State in words how the ratio of the areas of these squares is related to the ratio of their side lengths.
15. Two cones are similar. The ratio of these cones' volumes is 27 : 64. Determine the ratio of:
- the perpendicular heights of the cones
 - the areas of the bases of the cones.
16. Rectangle A has dimensions 5 by 4 units, rectangle B has the dimensions 4 by 3 units, and rectangle C has the dimensions 3 by 2.4 units.
- Determine which of these rectangles are similar. Explain your answer.
 - Evaluate the area scale factor for the similar rectangles that you have identified.
17. A balloon in the shape of a sphere has an initial volume of 840 cm^3 . Its volume is then increased to $430\,080 \text{ cm}^3$. Determine the increase in the radius of the balloon.
18. **WE18** The bottom half of an egg timer, which is shaped like two cones connected at their apexes, has sand poured into it as shown by the blue section of this diagram.



Using the measurements given, evaluate the ratio of the volume of sand in the bottom half of the egg timer to the volume of empty space that is left in the bottom half of the egg timer. You could also use technology to evaluate the answer.

LESSON

4.7 Review

4.7.1 Topic summary

Units of measurement

When converting units of **length**:

$\div 10$ $\div 100$ $\div 1000$
 millimetres (mm) centimetres (cm) metres (m) kilometres (km)
 $\times 10$ $\times 100$ $\times 1000$

When converting units of **area**:

$\div 10^2$ $\div 100^2$ $\div 1000^2$
 millimetres (mm²) centimetres (cm²) metres (m²) kilometres (km²)
 $\times 10^2$ $\times 100^2$ $\times 1000^2$

When converting units of **volume**:

$\div 10^3$ $\div 100^3$ $\div 1000^3$
 millimetres (mm³) centimetres (cm³) metres (m³) kilometres (km³)
 $\times 10^3$ $\times 100^3$ $\times 1000^3$

CONGRUENCE AND SIMILARITY

Congruent figures

- Congruent figures have exactly the same shape and size.
- The symbol for congruence is \cong .

Tests for congruent triangles

	SSS
	SAS
	ASA
	RHS

Ratio and scale factors

- Ratios compare quantities of the same type.
- Always simplify ratios. For example: $8 : 24 = 1 : 3$
- Scale factor = $\frac{\text{image length}}{\text{object length}}$.
- For a scale factor of n :
 - if $n > 1$ the image is larger than the object
 - if $n < 1$ the image is smaller than the object.

Similar figures

- Similar figures have the same shape, but different sizes.
- The symbol for similarity is \sim .

If the scale factor of similar figures' sides is n , then:

- the scale factor of their areas is n^2
- the scale factor of their volumes is n^3 .

Tests for similar triangles

	AAA
	SSS
	SAS
	RHS

• *Note:* Sides of similar triangles are not equal. They are proportional, or have the same scale factor.




Nets, polyhedra and Euler's formula

- The complete set of faces forming a solid is called its **net**.
- A 3-dimensional solid where each of the faces is a polygon is called a **polyhedron**.
- **Euler's formula** shows the relationship between the number of **edges (E)**, **faces (F)** and **vertices (V)** in any polyhedron: $F + V - 2 = E$

4.7.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
4.2	I can compare two quantities of the same type using ratios.			
	I can simplify ratios.			
	I can enlarge a figure using a scale factor.			
	I can calculate and use a scale factor.			
4.3	I can identify congruent figures.			
	I can show that two triangles are congruent using the appropriate congruency test.			
4.4	I can identify and produce nets of various solids.			
	I can define polyhedra and identify examples of Platonic solids.			
	I can verify Euler's formula for various polyhedra.			
4.5	I can identify similar figures.			
	I can determine the scale factor between similar figures.			
	I can show that two triangles are similar using the appropriate similarity test.			
4.6	I can convert between different units of length, area and volume.			
	I can, for a given scale factor of n^2 , determine the area of similar figures.			
	I can, for a given scale factor of n^3 , determine the volume of similar figures.			

4.7.3 Project

What's this object?

When using geometrical tools to construct shapes, we have to make sure that our measurements are precise. Even a small error in one single step of the measuring process can result in an incorrect shape. The object you will be making in this task is a very interesting one. This object is made by combining three congruent shapes.



The instructions for making the congruent shape that you will use to then make the final object are given below.


Part 1: Making the congruent shape

- Using a ruler, a protractor, a pencil and two compasses, draw a shape in your workbook by following the instructions below.
 - Measure a horizontal line (AB) that is 9.5 cm long. To make sure there is enough space for the whole of the final object, draw this first line close to the bottom of the page you are working on.
 - At point B, measure an angle of 120° above the line AB. Draw a line that follows this angle for 2 cm. Mark the end of this line as point C, creating the line BC.
 - At point C, measure an angle of 60° on the same side of line BC as point A. Draw a line that follows this angle for 7.5 cm. Mark the end of this line as point D, creating the line CD.
 - At point D, measure an angle of 60° above the line CD. Draw a line that follows this angle for 3.5 cm. Mark the end of this line as point E, creating the line DE.
 - At point E, measure an angle of 120° above the line DE. Draw a line that follows this angle for 2 cm. Mark the end of this line as point F, creating the line EF.
 - Join point F to point A.
Colour or shade this shape using any colour you wish.
- Determine the length of the line joining point A to point F.
- Describe what you notice about the size of the angles FAB and AFE.

Part 2: Making the final object

You have now constructed the shape that will be used three times to make the final object. To make this object, follow the instructions below.

- Trace the original shape onto a piece of tracing paper twice. Cut around the edges of both of the shapes you have just drawn. Label the shapes with the same letters you used to mark the points A–F when making the original shape.
 - Place line AF of the first traced copy of the original shape so that it is covering up line CD of the shape you first drew. Use the tracing paper to transfer this shape onto the original shape. Colour or shade the traced shape using a colour that is different to the colour you used for the first shape.
 - Place the line DC of your second traced copy so that it is covering up line FA of the shape you first drew. Use the tracing paper to transfer this shape onto the original shape. Colour or shade this third section with a third colour.
- Describe the object you have drawn.
 - Using the internet, your school library or other references, investigate other ‘impossible objects’ that can be drawn as 2-dimensional shapes. Recreate these shapes on a separate sheet of paper. Beside these shapes, briefly write some reasons for these shapes being referred to as ‘impossible’.



on Resources



eWorkbook

Topic 4 Workbook (worksheets, code puzzle and project) (ewbk-2004)



Interactivities

Crossword (int-2693)

Sudoku puzzle (int-3205)


Fluency

- Express each of the following ratios in simplest form.
 - $8 : 16$
 - $24 : 16$
 - $27 : 18$
- Express each of the following ratios in simplest form.
 - $56 : 80$
 - $8 : 20$
 - $49 : 35$
- MC** There are 9 girls and 17 boys in a Year 9 maths class. Select the ratio of boys to girls.
 - $9 : 17$
 - $17 : 9$
 - $17 : 26$
 - $9 : 26$
 - $26 : 9$
- Jan raised \$15 for a charity fundraising event, while her friend Lara raised \$25. Calculate the ratio of:
 - the amount Jan raised to the amount Lara raised
 - the amount Jan raised to the total amount raised by the pair
 - the amount the pair raised to the amount Lara raised.
- A pigeon breeder has 45 pigeons. These include 15 white, 21 speckled pigeons, and the rest grey. Calculate the following ratios.
 - White pigeons to grey pigeons.
 - Grey pigeons to speckled pigeons.
 - White pigeons to the total number of pigeons.
- Express each of the following ratios in simplest form.
 - $4\frac{1}{2} : 1$
 - $6\frac{1}{8} : 9$
 - $7\frac{1}{4} : 3\frac{1}{2}$
- Express each of the following ratios in simplest form.
 - $8.4 : 7.2$
 - $0.2 : 2.48$
 - $6.6 : 0.22$
- MC** Jack has completed 2.5 km of a 4.5 km race. Choose the ratio of the distance Jack has completed to the remaining distance he has left to run.
 - $9 : 5$
 - $4 : 5$
 - $4 : 9$
 - $5 : 4$
 - $5 : 9$
- MC** If $x : 16 = 5 : 4$, select the value of the pronumeral x .
 - 1
 - 4
 - 16
 - 20
 - 25
- MC** If $40 : 9 = 800 : y$, select the value of the pronumeral y .
 - 20
 - 40
 - 180
 - 450
 - 761
- Determine the value of the pronumeral in each of the following.
 - $a : 15 = 2 : 5$
 - $b : 20 = 5 : 8$
 - $9 : 10 = 12 : c$
- A polyhedron has a faces, $3a$ vertices and $(2a + 8)$ edges. Evaluate a .


13. $\triangle PQR \sim \triangle DEF$. Determine the length of the missing side in each of the following combinations.

- Calculate DF if $PQ = 10$ cm, $DE = 5$ cm and $PR = 6$ cm
- Calculate EF if $PQ = 4$ cm, $DE = 12$ cm and $QR = 5$ cm
- Calculate QR if $DE = 4$ cm, $PQ = 6$ cm and $EF = 8$ cm
- Calculate PQ if $DF = 5$ cm, $PR = 8$ cm and $DE = 6$ cm
- Calculate DE if $QR = 16$ cm, $EF = 6$ cm and $PQ = 12$ cm

14. Identify which of the following pairs of shapes are congruent.




15. Name the congruent triangles in these figures and determine the value of the pronumerals in each case.




16. Copy each of the following shapes and enlarge (or reduce) them by the given factor.

a.




Enlarge by a factor of 2.

b.



Reduce by a factor of 3.


c.




Enlarge by a factor of 4.

17. Determine the enlargement factors that have been used on the following shapes.


a.




b.




c.



d.











18. Evaluate the value of the pronumerals in the pair of congruent triangles shown.




19. The area of a family room is 16 m^2 and the length of the room is 6.4 m . Calculate the area of the room if it is drawn on a plan that uses a scale of $1 : 20$.

20. Each of the diagrams shown shows a pair of similar triangles. Calculate the value of x in each case.




Problem solving

21. The diagram shows a ramp made by Jinghua for her automotive class. The first post has a height of 0.25 m and is placed 2 m from the end of the ramp. If the second post is 1.5 m high, determine the distance it should be placed from the first post.



22. Consider the figure shown.


- Write a formal proof for a similarity relationship between two triangles in this figure. Give a reason for the similarity.
- Determine the values of the pronumerals.



23. *Slocum* is a yacht with a length of 12 m and a beam (width) of 2.5 m. A model of *Slocum* is constructed for a museum. If the length and width of the model are one fifth of the yacht's actual length and width, determine how the volume of the yacht and the volume of the model compare.
24. Before the start of a yacht race, judges must certify that all of the yachts have the same size of sail. Without removing the triangular sails from their masts, state the congruency rule that the official could most efficiently use to work out if the sails on each of the boats are exactly the same. Explain why this is the most appropriate rule.
25. Calculate the height of the top of the ladder in the photo shown by using similar triangles. Give your answer correct to 1 decimal place.




26. Poh is given a 1 m ruler and asked to estimate the height of a palm tree. She places the ruler vertically so that its shadow ends at exactly the same point as the shadow of the palm tree. The ruler's shadow is 2.5 m long and the palm tree's shadow is 12.5 m long.




Poh performed some calculations using similar triangles and calculated the height of the palm tree to be 4 m. Her friend Mikalya said that she thought Poh's calculations were incorrect, and that the answer should be 5 m.

- State the correct answer.
- Explain the error that was made by the person with the incorrect answer.

27. A flagpole casts a shadow 2 m long. If a 50 cm ruler is placed upright at the base of the flagpole it casts a shadow 20 cm long. Evaluate the height of the flagpole.



28. A student (S) uses a tape measure, backpack (B), water bottle (W), tree (T) and jacket (J) to help calculate the width of a river.



- a. Copy and complete the following:
- \angle ___ and \angle ___ are both ___.
 - \angle SBT and \angle ___ are equal because ___.
 - \angle STB and \angle ___ are equal because ___.
- b. Rewrite the following statement, selecting the correct alternatives from within each set of brackets.
 \angle STB and \angle JWB are (similar / congruent) (SSS / SAS / AAA / ASA / RHS)
- c. Copy and complete the following equation.

$$\frac{\overline{ST}}{\overline{JW}} = \frac{\square}{\square}$$

- d. Use your answer to part c to evaluate the width of the river.



To test your understanding and knowledge of this topic go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers




Topic 4 Congruence and similarity

4.1 Pre-test

- 25 : 39
- $x = 13.5$
- E
- C
- $y : z = 9 : 8$
- False
- D
- a. $\angle CED$ b. ASA
- A
- B, D
- $x = 2.3$ cm
- $b = 7.5$
- a. 3 : 5 b. 9 : 25
- 9 : 1
- B

4.2 Ratio and scale

- a. i. 240 : 7 ii. 7 : 240
 b. The track is $34\frac{2}{7}$ times as long as it is wide.
- a. i. 5 : 2 ii. 2 : 5
 b. The cliff is 2.5 times as high as the distance from the base of the cliff to the emu.
- a. 2 : 3 b. 1 : 7 c. 1 : 3 d. 2 : 5
- a. 4 : 15 b. 40 : 31 c. 8 : 25 d. 100 : 33
- a. 6 : 1 b. 13 : 15 c. 4 : 1 d. 5 : 1
- a. 1 : 2 b. 20 : 3 c. 1 : 6 d. 5 : 3
- a. 5 : 1 b. 11 : 5 c. 43 : 20 d. 1 : 200
- a. 3 : 4 b. 8 : 7 c. 2 : 3 d. 28 : 25
- a. 2 : 7 b. 10 : 3 c. 33 : 16 d. 23 : 15
- a. $a = 9$ b. $b = 24$ c. $c = 32$
- a. $e = 3\frac{3}{4}$ b. $f = 14\frac{2}{5}$ c. $g = 3\frac{3}{4}$
- a. $i = 29\frac{1}{3}$ b. $j = 8\frac{4}{7}$ c. $k = 10\frac{2}{5}$




- a. 3 b. 12 cm c. 3.5 cm
- a. 175 : 6428 b. 37
- 2.5 mL
- 1000 g
- a. 200 b. $6\text{ m} \times 6\text{ m}$ c. $5\text{ m} \times 5\text{ m}$
- a. Sample responses can be found in the worked solutions in the online resources.
 b. 1.69 g gold, 0.5 g copper, 0.06 g silver
- Sample responses can be found in the worked solutions in the online resources.
- $2k \times 3k \times 5k = 30k^3$
 $30k^3 = 21870$
 $k^3 = 729$
 $k = 9$
 Substituting k into the ratio $2k : 3k : 5k$, the dimensions are 18 cm, 27 cm and 45 cm.
- The profits aren't shared in a fair ratio. Tyler gets more profit than his share and Dylan gets less profit than his share. Only Ari gets the correct share of the profit.
- Sample responses can be found in the worked solutions in the online resources.

- 680 cm
- $y : z = 9 : 8$
- $P = 4, Q = 6$
- 49 students
- 125 students

4.3 Congruent figures

- b and c
- a and c
- a and b
- D
- $\triangle ABC$ and $\triangle PQR$, SAS
- $\triangle ABC$ and $\triangle LNM$, SSS
- $\triangle LMN$ and $\triangle PQR$, SAS
- $\triangle ABC$ and $\triangle PQR$, ASA
- $\triangle ABC$ and $\triangle LMN$, RHS
- $\triangle ABC$ and $\triangle DEF$, ASA
- $x = 3$ cm
- $x = 78^\circ$
- $x = 80^\circ, y = 30^\circ, z = 70^\circ$
- a. $x = 30^\circ, y = 7$
 b. $x = 40^\circ, y = 50^\circ, z = 50^\circ, n = 90^\circ, m = 90^\circ$
- a. 2 cm b. 8 cm c. 6 mm d. 7 mm
- Yes, because they are identical.
- No. The reason for congruence in this situation could alternatively be ASA.

18.



19. a. SSS


b. SSS

c. SAS


20. Sample responses can be found in the worked solutions in the online resources.

21. Because the angle is not between the two given sides, the general shape of the triangle is not set. This means that it is possible to draw many different shapes that fit these conditions.


22. This can be done by making an equilateral triangle and a regular tetrahedron.




23. This can be done by making a double regular tetrahedron.



24. Each piece shown in this solution is similar to the original shape.




25. 1 : 1




4.4 Nets, polyhedra and Euler's formula


1.




2. a.




b.




c.



3. Two possible nets are:



4. A possible net is:




5 to 7. Sample responses can be found in the worked solutions in the online resources.

8. a. $E = 12, V = 8, F = 6, 8 + 6 - 2 = 12$
 b. $E = 12, V = 6, F = 8, 6 + 8 - 2 = 12$
 9. a. $E = 18, V = 12, F = 8, 8 + 12 - 2 = 18$
 b. and c. $E = 12, V = 8, F = 6, 6 + 8 - 2 = 12$
 10. a, b. For each shape, $E = 9, V = 6$ and $F = 5$ where
 $5 + 6 - 2 = 9$.

11. Answers will vary. Sample response: A knowledge of 3D shapes reinforces key maths concepts such as volume relationships, geometrical models and shape recognition. Identify and organise visual information. Interpretations of data displays require an understanding of the representation of space.

12. No. Edges, vertices and faces can not be identified on a solid with curves.

13. No



14. Sample responses can be found in the worked solutions in the online resources.

15. a. Yes b. Yes c. No
 16. 20 triangular faces, 30 edges and 12 vertices
 $20 + 12 - 2 = 30$

4.5 Similar figures

1. a and c, AAA
 2. a and b, SSS
 3. a and b, SAS
 4. a and c, RHS
 5. a and c, SSS
 6. a. $\triangle ABC$ and $\triangle DEC$
 b. $\triangle PQR$ and $\triangle ABC$
 c. $\triangle PQR$ and $\triangle TSR$
 d. $\triangle ABC$ and $\triangle DEC$
 e. $\triangle ADB$ and $\triangle ADC$
 7. $\triangle EDC$
 8. a. $\frac{AB}{AD} = \frac{BC}{DE} = \frac{AC}{AE}$
 b. $y = 10$
 9. a. $h = 3.75, i = 7.5$
 b. $j = 2.4, k = 11.1$


10. $x = 12$
 11. $x = 20^\circ, y = 2\frac{1}{4}$
 12. $x = 6, y = 8$
 13. a. $x = 7.1$ b. $x = 3.1$ c. $x = 7.5, y = 7.7$
 14. Sample responses can be found in the worked solutions in the online resources.
 15. Sample responses can be found in the worked solutions in the online resources.
 16. Sample responses can be found in the worked solutions in the online resources.
 17. Sample responses can be found in the worked solutions in the online resources.
 18. a. True b. False c. False d. True
 19. a. 6.72 m b. 78 cm
 20. 3.75 m
 21. $a = 1$ cm
 22. a. $\triangle EDC$ and $\triangle EBA$
 b. $x = 3$ cm, $y \approx 3.29$ cm
 23. a. Sample response can be found in the worked solutions in the online resources.
 b. SAS
 c. 4 cm
 d. PR 6 cm, PS 5 cm and PQ 10 cm
 24. The triangles are similar (AAA). $l = \frac{y}{2}$.


4.6 Area and volume of similar figures

1. a. 900 cm^2 b. 67.5 cm^2 c. 855 mm^2
 d. 102 cm^2 e. 5 m^2 f. 25 mm^2
 2. $SA = 384 \text{ cm}^2, V = 288 \text{ cm}^3$
 3. a. 5400 cm^3 b. 337.5 cm^3 c. 1809 mm^3
 d. 100 cm^3 e. 2.5 m^3 f. 1 mm^3
 4. $50\,000 \text{ cm}^2$
 5. a. $250\,000 \text{ cm}^2$ b. 17.4 cm^2
 6. $8\,437\,500 \text{ cm}^3$
 7. a. $1\,200\,000 \text{ cm}^2$ b. 187.5 cm^2
 8. a. Sample responses can be found in the worked solutions in the online resources.
 b. 10 cm
 c. 15 cm
 d. Sample responses can be found in the worked solutions in the online resources.
 9. Sample responses can be found in the worked solutions in the online resources.
 10. Sample responses can be found in the worked solutions in the online resources.
 11. a. Sample responses can be found in the worked solutions in the online resources.
 b. Sample responses can be found in the worked solutions in the online resources.
 c. 2197 cm^3
 d. 34.3 cm^3
 e. 28.2 cm^2

12. Sample responses can be found in the worked solutions in the online resources.
13. 3 : 5
14. a. $4 \text{ cm}^2, 25 \text{ cm}^2, 100 \text{ cm}^2, 400 \text{ cm}^2$
 b. The ratio of the areas is equal to the square of the ratio of the side lengths.
15. a. 3 : 4 b. 9 : 16
16. a. A and C are similar rectangles by the ratio 5 : 3 or scale factor $\frac{5}{3}$.
 b. $\frac{25}{9}$
17. The new radius is 8 times the old radius.
18. The ratio is 7 : 1.

Project

1. 
2. 7.5 cm
3. $\angle \text{FAB} = 60^\circ, \angle \text{AFE} = 60^\circ$
4. The impossible triangle






5. Students are required to research and investigate other 'impossible' objects drawn as two-dimensional shapes. These need to be created on a separate sheet of paper with reasons as to why they are termed 'impossible'.

4.7 Review questions

1. a. 1 : 2 b. 3 : 2 c. 3 : 2
2. a. 7 : 10 b. 2 : 5 c. 7 : 5
3. B
4. a. 3 : 5 b. 3 : 8 c. 8 : 5
5. a. 5 : 3 b. 3 : 7 c. 1 : 3
6. a. 9 : 2 b. 49 : 72 c. 29 : 14
7. a. 7 : 6 b. 5 : 62 c. 30 : 1
8. D
9. D
10. C
11. a. $a = 6$ b. $b = 12\frac{1}{2}$ c. $c = 13\frac{1}{3}$
12. $a = 5$
13. a. 3 cm b. 15 cm c. 12 cm
 d. 9.6 cm e. 4.5 cm
14. a. Congruent b. Not congruent c. Congruent

15. a. $\triangle \text{ANP} \cong \triangle \text{DWR}, x = 3 \text{ cm}$
 b. $\triangle \text{TKB} \cong \triangle \text{KTS}$ or $\triangle \text{TKB} \cong \triangle \text{KST}$,
 $x = 75^\circ, y = 30^\circ, z = 75^\circ, p = 5 \text{ cm}, q = 5 \text{ cm}$



17. a. 3 b. 0.4 c. $\frac{3}{5}$ d. 3.5
18. $x = 12 \text{ cm}, y = 80^\circ, z = 40^\circ$
19. 400 cm^2
20. a. $x = 8$ b. $x = 4$ c. $x = 11\frac{1}{5}$ d. $x = 26\frac{2}{3}$
21. 10 m
22. a. $\triangle \text{ABC} \sim \triangle \text{QPC}, \text{AAA}$
 b. $y = 12, z = 3$
23. The model is $\frac{1}{125}$ the volume of the yacht.
24. The easiest way to check for congruency is to measure the parts of each sail that are easiest to reach. This would be the bottom of each sail and the angles at the bottom of each of those sails. In this situation ASA would be the most appropriate rule to use.

25. 2.1 m

26. a. 5 m

b. Mikalya had the correct answer. Poh used the distance of 10 m (from the ruler to the tree) in her calculations instead of 12.5 m (the whole length of the tree's shadow).

27. 5 m

28. a. i. $\angle TSB$ and $\angle WJB$ are both right angles.

ii. $\angle SBT$ and $\angle JBW$ are equal because they are vertically opposite.

iii. $\angle STB$ and $\angle JWB$ are equal because they are alternate.

b. $\angle SBT$ and $\angle BJW$ are similar (AAA).

c. $\frac{\overline{ST}}{\overline{JW}} = \frac{50}{5}$

d. 27 m

5 Pythagoras and trigonometry

LESSON SEQUENCE

5.1 Overview	254
5.2 Pythagoras' theorem	258
5.3 Applications of Pythagoras' theorem	267
5.4 Trigonometric ratios	274
5.5 Calculating unknown side lengths	282
5.6 Calculating unknown angles	291
5.7 Angles of elevation and depression	299
5.8 Review	305



LESSON

5.1 Overview

5.1.1 Why learn this?

Trigonometry is the branch of geometry that is concerned with triangles. The word ‘trigonometry’ was created in the sixteenth century from the Greek words *trigōnon* (‘triangle’) and *metron* (‘measure’), but the study of the geometry of triangles goes back to at least the sixth century BCE, when the ancient Greek philosopher Pythagoras of Samos developed his famous theorem.

Pythagoras was particularly interested in right-angled triangles and the relationships between their sides. Later, Greek mathematicians used Pythagoras’ theorem and the trigonometric ratios to calculate all kinds of distances, including Earth’s circumference. Trigonometry is still the primary tool used by surveyors and geographers today when working out distances between points on Earth’s surface.

None of the structures we build would be possible without our understanding of geometry and trigonometry. Engineers apply the principles of geometry and trigonometry regularly to make sure that buildings are strong, stable and capable of withstanding extreme conditions. Triangles are particularly useful to engineers and architects because they are the strongest shape. Any forces applied to a triangular frame will be distributed equally to all of its sides and joins. This fact has been known for thousands of years — triangular building frames were used as far back as the sixth century BCE.

A truss is an example of a structure that relies on the strength of triangles. Trusses are often used to hold up the roofs of houses and to keep bridges from falling down. Triangular frames can even be applied to curved shapes. Geodesic domes, like the one shown here, are rounded structures that are made up of many small triangular frames connected together. This use of triangular frames makes geodesic domes very strong, but also very light and easy to build.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept




Extra learning resources

Differentiated question sets


Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 5.1 Pre-test

1. Calculate the value of x in the diagram shown.




2. Calculate the value of a in the diagram shown, correct to 2 decimal places.



3. **MC** Identify which set of 3 numbers could represent the sides of a right-angled triangle.
A. 2, 3, 4 **B.** 5, 12, 13 **C.** 1, 2, 3 **D.** 5, 6, 7 **E.** 3, 5, 6
4. If a right-angled triangle has side lengths a , $a + b$ and $a - b$, where both a and b are greater than 0, state which one of the lengths is the hypotenuse.


5. **MC** Identify the correct rule linking the sides of the right-angled triangle shown.

- A.** $c^2 = (a + b)^2$
B. $c = a + b$
C. $c^2 = a^2 + b^2$
D. $c = \frac{b}{a}$
E. $c^2 = a^2 - b^2$




6. **MC** Consider the right-angled triangle shown. Identify the correct option for angle θ .

- A.** a is the adjacent side.
B. b is the opposite side.
C. a is the hypotenuse
D. b is the adjacent side.
E. c is the opposite side.




7. **MC** Identify the correct trigonometric ratio for the triangle shown.

- A. $\tan(\theta) = \frac{a}{c}$
- B. $\sin(\theta) = \frac{b}{a}$
- C. $\cos(\theta) = \frac{c}{b}$
- D. $\sin(\theta) = \frac{c}{b}$
- E. $\cos(\theta) = \frac{b}{c}$




8. Calculate the length of the unknown side of each of the following triangles, correct to 2 decimal places.

a.



b.



9. Evaluate, correct to 4 decimal places.


a. $\sin(62.5^\circ)$

b. $\cos(12.1^\circ)$


c. $\tan(74.9^\circ)$

10. **MC** Identify the lengths of the unknown sides in the triangle shown.

- A. $a = 6.1$, $b = 13.7$ and $c = 7.3$
- B. $a = 6.1$, $b = 13.7$ and $c = 5.1$
- C. $a = 5.1$, $b = 6.1$ and $c = 13.7$
- D. $a = 13.7$, $b = 6.1$ and $c = 5.1$
- E. $a = 13.7$, $b = 6.1$ and $c = 7.3$




11. Calculate the value of θ in the triangle shown, correct to 1 decimal place.




12. **MC** Identify the value of θ in the triangle shown, correct to 2 decimal places.

- A. 25.27°
- B. 28.16°
- C. 28.17°
- D. 61.83°
- E. 64.73°




13. **MC** Determine the correct value of the side length a .

- A. 7.62
- B. 10.04
- C. 14.34
- D. 20.48
- E. 29.25



14. A person who is 1.54 m tall stands 10 m from the foot of a tree and records the angle of elevation (using an inclinometer) to the top of the tree as 30° . Evaluate the height of the tree, in metres correct to 2 decimal places.



15. The angle of depression from a scuba diver who is floating on the water's surface to a shark swimming below them on the sea floor is 35.8° . The depth of the water is 35 m. Evaluate the horizontal distance from the scuba diver to the shark, in metres correct to 2 decimal places.

LESSON

5.2 Pythagoras' theorem

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- identify the hypotenuse in a right-angled triangle
- recall Pythagoras' theorem
- apply Pythagoras' theorem to calculate unknown side lengths.

5.2.1 Introducing Pythagoras' theorem

eles-4750

- A **right-angled triangle** is a triangle that contains a right angle (90°).
- The longest side of a right-angled triangle is always found opposite the right angle. This side is called the **hypotenuse**.
- Pythagoras (580–501 BCE) was a Greek mathematician who explored the relationship take between the lengths of the sides of right-angled triangles.
- The relationship he described is known as **Pythagoras' theorem**.
- If the lengths of two sides of a right-angled triangle are given, then Pythagoras' theorem lets us calculate the length of the unknown side.




Pythagoras' theorem

In any right-angled triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides. This rule is written as follows.

$$c^2 = a^2 + b^2$$

In this rule, a and b represent the two shorter sides of a right-angled triangle, while c represents the hypotenuse.




WORKED EXAMPLE 1 Assigning Pythagorean identities

For the right-angled triangles shown below:

i. state which side is the hypotenuse

a.




THINK

a. i. The hypotenuse is always opposite the right angle.


ii. write Pythagoras' theorem.

b.




WRITE

a. Side f is opposite the right angle. Therefore, side f is the hypotenuse.



- ii. If the triangle is labelled as usual with a , b and c , as shown in blue, Pythagoras' theorem can be written and then the letters can be replaced with their new values (names).




$$c = f; a = d; b = e$$

$$c^2 = a^2 + b^2$$


$$f^2 = d^2 + e^2$$

- b. i. The hypotenuse is opposite the right angle.



Side t is opposite the right angle.
Therefore, side t is the hypotenuse.

- ii. If the labels on the triangle are replaced with a , b and c , as shown in blue, Pythagoras' theorem can be written out, and then the letters a , b and c can be changed back to the original labels.



$$c = t; b = p; a = m$$

$$c^2 = a^2 + b^2$$

$$t^2 = m^2 + p^2$$


5.2.2 Calculating unknown side lengths

eles-4751

- Pythagoras' theorem can be used to calculate the unknown side length of a right-angled triangle if the other two side lengths are known.
- To calculate an unknown side length, substitute the known values into the equation $c^2 = a^2 + b^2$ and solve for the remaining pronumeral.
- Remember that the hypotenuse always corresponds to the pronumeral c .

WORKED EXAMPLE 2 Calculating the length of the hypotenuse


For the triangle shown, calculate the length of the hypotenuse, x , correct to 1 decimal place.



THINK

1. Copy the diagram and apply the labels a , b and c to the triangle, pairing them with the known values and the pronumeral. Remember to label the hypotenuse as c .

WRITE



2. Write out Pythagoras' theorem.

$$c^2 = a^2 + b^2$$

3. Substitute the values of a , b and c into this rule and simplify.


$$\begin{aligned}x^2 &= 7^2 + 11^2 \\ &= 49 + 121 \\ &= 170\end{aligned}$$

4. Calculate x by taking the square root of 170. Round the answer to 1 decimal place.

$$\begin{aligned}x &= \sqrt{170} \\ x &= 13.0\end{aligned}$$

WORKED EXAMPLE 3 Calculating the length of a shorter side

Calculate the length, as an exact value and correct to 1 decimal place, of the unmarked side of the triangle shown.



THINK


1. Copy the diagram and label the sides a , b and c . Remember to label the hypotenuse as c .

2. Write out Pythagoras' theorem for a shorter side.

3. Substitute the values of b and c into the equation and simplify.

4. Calculate the value of a by taking the square root of 132. Round the answer to 1 decimal place.

WRITE



$$a^2 = c^2 - b^2$$

$$\begin{aligned}a^2 &= 14^2 - 8^2 \\ &= 196 - 64 \\ &= 132\end{aligned}$$

$$\begin{aligned}a &= \sqrt{132} && \text{Exact answer.} \\ &= 11.5 \text{ cm} && \text{1 decimal place.}\end{aligned}$$

5.2.3 Applying Pythagoras' theorem

eles-4752

- If a problem involves a right-angled triangle, then Pythagoras' theorem can be applied to determine a solution.

Applying Pythagoras' theorem to a problem

1. Read the question carefully and draw a right-angled triangle to represent the problem.
2. Identify the known values and place them on the diagram in their correct positions.
3. Identify the variable that you will need to determine.
4. Substitute the known values into Pythagoras' theorem. Make sure that they are all using the same units of length. You may need to convert some units.
5. Solve for the unknown value.
6. Use the result to write the answer as a complete sentence.

WORKED EXAMPLE 4 Solving problems using Pythagoras' theorem

A wedge of cake has been cut so that it forms a right-angled triangle. The longest edge of the wedge is 12 cm long and its shortest edge is 84 mm long.


Calculate the length of the third side, in cm correct to 1 decimal place.



THINK

1. Draw a right-angled triangle.
Identify the longest side (this will always be the hypotenuse) and place its value on the diagram.
Identify the shortest side and place its value on the diagram.
Label the side with unknown length as x .
2. Identify the length of the hypotenuse and the two shorter sides.
Check that the units for all measurements are the same.
Convert 84 mm to centimetres by dividing by 10.
3. Substitute the values into the equation and simplify.
4. Solve the equation for x by:
subtracting 70.56 from both sides and taking the square root of both sides.
5. It is not possible for a triangle to have a side with a negative length, therefore the solution must be the positive value. Round the answer to 1 decimal place and include the appropriate units.
6. Write the answer as a sentence.

WRITE



$$\begin{aligned}c &= 12 \text{ cm} \\a &= 84 \text{ mm} = 8.4 \text{ cm} \\b &= x\end{aligned}$$

$$\begin{aligned}c^2 &= a^2 + b^2 \\12^2 &= (8.4)^2 + x^2 \\144 &= 70.56 + x^2 \\144 - 70.56 &= 70.56 - 70.56 + x^2 \\73.44 &= x^2 \\\sqrt{73.44} &= \sqrt{x^2} \\\pm 8.5697 &\approx x \\x &= 8.6 \text{ cm}\end{aligned}$$

The third side of the slice of cake is 8.6 cm long.

5.2.4 Pythagorean triads

eles-4753

- A **Pythagorean triad** is a group of any three whole numbers that satisfy Pythagoras' theorem. For example, $\{3, 4, 5\}$ is a Pythagorean triad because $3^2 = 9$, $4^2 = 16$ and $5^2 = 25$, and $9 + 16 = 25$.
- Pythagorean triads are useful when solving problems using Pythagoras' theorem.
- If two known side lengths in a triangle belong to a triad, the length of the third side can be stated without performing any calculations.
- Some well known Pythagorean triads are $\{3, 4, 5\}$, $\{5, 12, 13\}$, $\{8, 15, 17\}$ and $\{7, 24, 25\}$.

WORKED EXAMPLE 5 Identifying Pythagorean triads

Determine whether the following sets of numbers are Pythagorean triads.

a. {9, 10, 14}

b. {33, 65, 56}

THINK

a. 1. Pythagorean triads satisfy Pythagoras' theorem. Substitute the values into the equation $c^2 = a^2 + b^2$ and determine whether the equation is true. Remember that c is always the longest side.

2. Write your conclusion.

b. 1. Pythagorean triads satisfy Pythagoras' theorem. Substitute the values into the equation $c^2 = a^2 + b^2$ and determine whether the equation is true. Remember that c is always the longest side.

2. Write your conclusion.

WRITE

$$a. c^2 = a^2 + b^2$$

$$\begin{aligned} \text{LHS} &= c^2 \\ &= 14^2 \\ &= 196 \end{aligned}$$

$$\begin{aligned} \text{RHS} &= a^2 + b^2 \\ &= 9^2 + 10^2 \\ &= 81 + 100 \\ &= 181 \end{aligned}$$

Since $\text{LHS} \neq \text{RHS}$, the set {9, 10, 14} is not a Pythagorean triad.


$$b. c^2 = a^2 + b^2$$

$$\begin{aligned} \text{LHS} &= 65^2 \\ &= 4225 \end{aligned}$$

$$\begin{aligned} \text{RHS} &= 33^2 + 56^2 \\ &= 1089 + 3136 \\ &= 4225 \end{aligned}$$

Since $\text{LHS} = \text{RHS}$, the set {33, 65, 56} is a Pythagorean triad.

- If each term in a Pythagorean triad is multiplied by the same number, the result is also a triad.
For example, if we multiply each number in {5, 12, 13} by 2, the result {10, 24, 26} is also a triad.
- Builders and gardeners use multiples of the Pythagorean triad {3, 4, 5} to make sure that walls and floors are at right angles to each other.



Algorithm for Pythagorean Triads:



There are different ways to determine Pythagorean Triads, such as Fibonacci's Method and Sequences of mixed numbers to name a couple.

We will look at an algorithm using Dickson's Method of determining Pythagorean Triads.

Step	THINK	WRITE
1.	Select a number and call this r .	$r = 8$
2.	Calculate the value of $\frac{r^2}{2}$.	$\frac{r^2}{2} = \frac{8^2}{2} = \frac{64}{2} = 32$
3.	List all the pairs of factors of the $\frac{r^2}{2}$ value.	Factors of 32 are: (1, 32) (2, 16) (4, 8)
4.	Call the factors s and t .	Factor (1, 32) $s = 1$ and $t = 32$

5.	Use the following to calculate the Pythagorean Triads. $x = r + s$ $y = r + t$ $z = r + s + t$	$x = r + s$ $= 8 + 1$ $= 9$ $y = r + t$ $= 8 + 32$ $= 40$ $z = r + s + t$ $= 8 + 1 + 32$ $= 41$
6.	Write the Pythagorean Triad.	{9, 40, 41}
7.	Repeat for the other factor pairs.	
8.	Call the factors s and t .	Factor (2, 16) $s = 2$ and $t = 16$
9.	Use the following to calculate the Pythagorean Triads. $x = r + s$ $y = r + t$ $z = r + s + t$	$x = r + s$ $= 8 + 2$ $= 10$ $y = r + t$ $= 8 + 16$ $= 24$ $z = r + s + t$ $= 8 + 2 + 16$ $= 26$
10.	Write the Pythagorean Triad.	{10, 24, 26}
11.	Call the factors s and t .	Factor (4, 8) $s = 4$ and $t = 8$
12.	Use the following to calculate the Pythagorean Triads. $x = r + s$ $y = r + t$ $z = r + s + t$	$x = r + s$ $= 8 + 4$ $= 12$ $y = r + t$ $= 8 + 8$ $= 16$ $z = r + s + t$ $= 8 + 4 + 8$ $= 20$
13.	Write the Pythagorean Triad.	{12, 16, 20}

Resources

-  **eWorkbook** Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)
-  **Interactivities** Individual pathway interactivity: Pythagoras' theorem (int-4472)
 - Finding the hypotenuse (int-3844)
 - Finding the shorter side (int-3845)
 - Pythagorean triads (int-3848)

5.2 Quick quiz **on**

5.2 Exercise

Individual pathways

PRACTISE

1, 5, 6, 8, 12, 15

CONSOLIDATE

2, 7, 9, 13, 16





MASTER

3, 4, 10, 11, 14, 17




Fluency

1. **WE1** For the right-angled triangles shown below:





- i. state which side is the hypotenuse
- ii. write Pythagoras' theorem.




2. **WE2** For each of the following triangles, calculate the length of the hypotenuse, giving answers correct to 2 decimal places.



3. **WE3** Calculate the value of the pronumeral in each of the following triangles. Give your answers correct to 2 decimal places.



4. **WE4** Calculate the value of the missing side length. Give your answer correct to 1 decimal place.



5. **WE5** Determine whether the following sets of numbers are Pythagorean triads.

a. {2, 5, 6}

b. {7, 10, 12}

c. {18, 24, 30}

d. {72, 78, 30}

e. {13, 8, 15}

f. {50, 40, 30}

Understanding

6. In a right-angled triangle, the two shortest sides are 4.2 cm and 3.8 cm.

a. Draw a sketch of the triangle.

b. Calculate the length of the hypotenuse correct to 2 decimal places.

7. A right-angled triangle has a hypotenuse of 124 mm and another side of 8.5 cm.

a. Draw a diagram of the triangle.

b. Calculate the length of the third side of this triangle. Give your answer in millimetres, correct to 2 decimal places.

8. **MC** Identify which of the following sets is formed from the Pythagorean triad {21, 20, 29}.

A. {95, 100, 125}

B. {105, 145, 100}

C. {84, 80, 87}

D. {105, 80, 87}

E. {215, 205, 295}

9. **MC** Identify the length of the hypotenuse in the triangle shown.


A. 25 cm

B. 50 cm

C. 50 mm

D. 500 mm

E. 2500 mm



10. **MC** Identify the length of the third side in the triangle shown.


A. 48.75 cm

B. 0.698 m

C. 0.926 m

D. 92.6 cm

E. 69.8 mm



11. The lengths of two sides of a right-angled triangle are given below:

a. $a = 37$ cm, $c = 180$ cm; calculate b in cm.


b. $a = 856$ mm, $b = 1200$ mm; calculate c in cm.

c. $b = 4950$ m, $c = 5.6$ km; calculate a in km.

d. $a = 125\,600$ mm, $c = 450$ m; calculate b in m.

Calculate the value of the third side. Give your answer in the units specified (correct to 2 decimal places).

The diagram shown illustrates how each triangle is to be labelled.



Remember that c is always the hypotenuse.

Reasoning


12. A ladder that is 7 m long leans up against a vertical wall. The top of the ladder reaches 6.5 m up the wall. Determine how far the foot of the ladder is from the wall.

Give your answer correct to 2 decimal places.


13. A rectangular park measures 260 m by 480 m. Danny usually trains by running five circuits around the edge of the park. After heavy rain, two of the adjacent sides are too muddy to run along, so Danny runs along a triangular path that traces the other two sides and the diagonal. Danny does five circuits of this path for training.
Show that Danny has run about 970 metres less than his usual training session.
14. A water park has hired Sharnee to build part of a ramp for a new water slide. She builds a 12-metre-long ramp that rises to a height of 250 cm. To meet safety regulations, the ramp can only have a gradient between 0.1 and 0.25.
Show that the ramp Sharnee has built is within the regulations.

Problem solving


15. Spiridoula is trying to hang her newest painting on a hook in her hallway. She leans a ladder against the wall so that the distance between the foot of the ladder and the wall is 80 cm. The ladder is 1.2 m long.
- Draw a diagram showing the ladder leaning on the wall.
 - Calculate how far up the wall the ladder reaches. Give your answer correct to 2 decimal places.
 - Spiridoula climbs the ladder to check whether she can reach the hook from the step at the very top of the ladder. When she extends her arm, the distance from her feet to her fingertips is 1.7 m.
If the hook is 2.5 m above the floor, determine whether she can reach it from the top step of the ladder.
16. a. The smallest numbers of four Pythagorean triads are given below. Determine the middle number of each triad, and use this answer to determine the third number of each triad.
- i. 9 ii. 11 iii. 13 iv. 29
- b. Comment about the triads formed in part a.
17. We know that it is possible to determine the exact square root of some numbers, but not others. For example, we can determine $\sqrt{4}$ exactly, but not $\sqrt{3}$ or $\sqrt{5}$. Our calculators can calculate decimal approximations of these square roots, but because they cannot be found exactly they are called ‘irrational numbers’. However, there is a method of showing the exact location of irrational numbers on a number line. Using graph paper, draw a right-angled triangle with 2 equal sides of length 1 cm as shown below.



Using Pythagoras’ theorem, we know that the longest side of this triangle is $\sqrt{2}$ units. Place the compass point at zero and make an arc that will show the location of $\sqrt{2}$ on the number line.



Draw another right-angled triangle using the longest side of the first triangle as one side, and make the other side 1 cm in length.



The longest side of this triangle will have a length of $\sqrt{3}$ units. Draw an arc to determine the location of $\sqrt{3}$ on the number line.

Repeat these steps to draw triangles that will have sides of length $\sqrt{4}$, $\sqrt{5}$, $\sqrt{6}$ units, and so on.

LESSON

5.3 Applications of Pythagoras' theorem

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- divide composite shapes into simple shapes
- calculate unknown side lengths in composite shapes and irregular triangles
- use Pythagoras' theorem to solve problems in 3D contexts.

5.3.1 Using composite shapes to solve problems


eles-4754

- Complex or unusual shapes can be thought of as composites of simpler shapes with known properties.
- Dividing a shape into a series of these simpler shapes lets us use the known properties of those shapes to determine unknown values.
- For example, to calculate the value of x in the trapezium shown, a vertical line can be added to the trapezium to create two shapes: a right-angled triangle and a rectangle.
- Using Pythagoras' theorem, we can now calculate the length of x , which has become the hypotenuse of triangle ABE.



WORKED EXAMPLE 6 Calculating unknown side lengths in composite shapes


Calculate the value of x in the diagram shown, correct to 1 decimal place.



THINK

1. Add a horizontal line to divide the shape into 2 smaller shapes, one that is a right-angled triangle, as shown in orange. x is the hypotenuse of a right-angled triangle. To use Pythagoras' theorem, the length of the side shown in green must be known. This length can be calculated as the difference between the long and short vertical edges of the trapezium: $5.5 - 3.5 = 2$.

WRITE



2. Write out Pythagoras' theorem and substitute the values of a , b and c .
 $a = 2, b = 4.2, c = x$

$$c^2 = a^2 + b^2$$

$$x^2 = 2^2 + (4.2)^2$$

$$x^2 = 4 + 17.64$$

$$x^2 = 21.64$$


$$\sqrt{x^2} = \sqrt{21.64}$$

$$x \approx \pm 4.65188$$

3. It is not possible to have a side with a negative length. Therefore the solution is the positive value. Round the answer to 1 decimal place

$$x = 4.7$$

- To determine the value of y in the irregular triangle shown, the triangle can be split into two right-angled triangles: $\triangle ABD$ (pink) and $\triangle BDC$ (blue). There is now enough information to calculate the missing side length from $\triangle ABD$. This newly calculated length can then be used to determine the value of y .




WORKED EXAMPLE 7 Applying Pythagoras' theorem to non-right-angled triangles

- Calculate the perpendicular height of an isosceles triangle with equal sides that are both 15 mm long and a third side that is 18 mm long.
- Calculate the area of this triangle.

THINK

1. Draw the triangle and label all side lengths as described.
2. Draw an additional line to represent the height of the triangle (shown in pink) and label it appropriately. Because the triangle is an isosceles triangle, h bisects the base of 18 mm (shown in purple) to create two right-angled triangles.
3. Consider one of these right-angled triangles, which both have a height of h . The base length of each of these triangles is $\frac{18}{2} = 9$ mm.

WRITE



4. Substitute the values from this right-angled triangle into Pythagoras' theorem to calculate the value of h .

$$c^2 = a^2 + b^2$$

$$15^2 = h^2 + 9^2$$

$$225 = h^2 + 81$$

$$225 - 81 = h^2 + 81 - 81$$

$$144 = h^2$$

$$\sqrt{144} = \sqrt{h^2}$$

$$\pm 12 = h$$

5. It is not possible to have a side with a negative length. Therefore the solution is the positive value. Include the appropriate units in your answer.

$$h = 12 \text{ mm}$$

The perpendicular height of the triangle is 12 mm.

- b. 1. Write the formula for the area of a triangle. Using the answers from part a, the base is 18 mm and the height is 12 mm.

$$\text{b. } A_{\Delta} = \frac{1}{2}bh$$

$$= \frac{1}{2} \times 18 \times 12$$

$$= 108$$

2. Answer the question with appropriate units.


The area of the triangle is 108 mm^2 .



5.3.2 Pythagoras' theorem in 3D

eles-4755

- Pythagoras' theorem can be used to solve problems in three dimensions.
- 3D problems often involve multiple right-angled triangles.
- It is often helpful to redraw sections of a 3D diagram in two dimensions. This makes it easier to see any right angles.




WORKED EXAMPLE 8 Using Pythagoras' theorem in 3D problems

Calculate the maximum length of a metal rod that would fit into a rectangular crate with dimensions $1 \text{ m} \times 1.5 \text{ m} \times 0.5 \text{ m}$.


THINK

1. Draw a diagram of a rectangular box with a rod in it. Label the dimensions of this diagram.
2. Draw a right-angled triangle that has the metal rod as one of its sides (x , shown in pink). The lengths of x and y in this right-angled triangle are not known. Draw a second right-angled triangle (shown in green) to allow you to calculate the length of y .

WRITE



3. Calculate the length of y using Pythagoras' theorem.



$$c^2 = a^2 + b^2$$


$$y^2 = 1.5^2 + 1^2$$

$$y^2 = 3.25$$

$$y = \sqrt{3.25}$$

4. Draw the right-angled triangle that has the rod as its hypotenuse. Use Pythagoras' theorem to calculate the length of x .

z = height of the crate = 0.5 m



$$c^2 = a^2 + b^2$$

$$x^2 = (\sqrt{3.25})^2 + 0.5^2$$

$$x^2 = 3.25 + 0.25$$

$$x^2 = 3.5$$

$$x = \sqrt{3.5}$$

$$\approx 1.87$$

5. Answer the question with appropriate units.

The maximum length of the metal rod is 1.87 m (correct to 2 decimal places).

on Resources



eWorkbook Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)



Video eLesson Pythagoras' theorem in 3 dimensions (eles-1913)



Interactivities Individual pathway interactivity: Applications of Pythagoras' theorem (int-4475)
Composite shapes (int-3847)

5.3 Quick quiz **on**

5.3 Exercise

Individual pathways


PRACTISE
1, 2, 4, 8, 14, 18

CONSOLIDATE
3, 6, 7, 9, 13, 15, 16, 19



MASTER
5, 10, 11, 12, 17, 20

Fluency





1. **WE6** Calculate the length of the side x in the figure shown, correct to 2 decimal places.



2. Calculate the values of the pronumerals in each of the following photographs and diagrams. Give your answers correct to 2 decimal places.






3. For each of the following diagrams, calculate the lengths of the sides marked x and y . Give your answer correct to 2 decimal places (where necessary).



4. **WE7** The height of an isosceles triangle is 3.4 mm and its equal sides are twice as long.
- Sketch the triangle, showing all the information given.
 - Calculate the length of the third side, correct to 2 decimal places.
 - Calculate the area of the triangle, correct to 2 decimal places.


Understanding

5. **WE8** Calculate the length of the longest metal rod that could fit diagonally into each of the boxes shown below. Give your answers correct to 2 decimal places.



6. The side length of an equilateral triangle is 1 m. Calculate:
- the height of the triangle in metres, correct to 2 decimal places
 - the area of the triangle in m^2 , correct to 3 decimal places.
7. **MC** Identify the longest length of an object that fits into a box with dimensions of $42 \text{ cm} \times 60 \text{ cm} \times 13 \text{ cm}$.
- A. 74.5 cm B. 60 cm C. 73.2 cm D. 5533 cm E. 74 cm
8. Priya wants to pack an umbrella into her suitcase, which measures $89 \text{ cm} \times 21 \text{ cm} \times 44 \text{ cm}$.
- State whether her 1 m umbrella will fit in.
 - Give the length of the longest umbrella that Priya could fit in their suitcase, correct to 2 decimal places.
9. Giang is packing her lunch into her lunchbox, which measures $15 \text{ cm} \times 5 \text{ cm} \times 8 \text{ cm}$.
- State whether her 22 cm straw will fit in.
 - Calculate the length of the longest object that will fit into Giang's lunchbox, correct to 2 decimal places.
10. A cylindrical pipe is 2.4 m long. It has an internal diameter of 30 cm. Calculate the size of largest object that could fit inside the pipe. Give your answer correct to the nearest cm.

11. A classroom contains 20 identical desks shaped like the one shown.




- Calculate the width of each desktop (labelled w). Give your answer correct to 2 decimal places.
- Calculate the area of each desktop, correct to 2 decimal places.
- If the top surface of each desk needs to be painted, determine the total area of the desktops that need to be painted. Give your answer correct to 2 decimal places.
- Each desktop needs to be given 2 coats of paint. The paint being used is sold in 1-litre tins that cost \$29.95 each. Each of these tins holds enough paint to cover an area of 12 m^2 . Using your answer to part c, calculate how much it will cost to paint all 20 desks.

12. A utility knife has a blade shaped like a right-angled trapezium. Its sloping edge is 2 cm long and its parallel sides are 32 mm and 48 mm long.
Calculate the width of the blade, then use your answer to calculate the blade's area.
13. A cylindrical pencil holder has an internal height of 16 cm. If its diameter is 8.5 cm and its width is 2 mm, calculate the length of the longest pen that could fit inside the holder. Give your answer correct to the nearest mm.

Reasoning

14. Consider the figure shown.

- State whether you can calculate the length of AC without first calculating the lengths of AD and DC.
- Calculate the lengths of the sides in part a. Give your answer correct to 2 decimal places.
- Explain whether the triangle ABC is right-angled.




15. Explain in your own words how a 2D right-angled triangle can be seen in a 3D figure.

16. Katie goes on a hike. She walks 2.5 km north, then 3.1 km east. She then walks 1 km north and 2 km west. Finally, she walks in a straight line back to her starting point. Show that Katie has walked a total distance of 12.27 km.

17. Consider the following diagram:


- Show that the distance AB is 18.44 metres.
- Show that the angle $\angle ACB$ is not a right angle.




Problem solving

18. The diagram shown illustrates a cross-section of the roof of Gabriel's house.


- Calculate the height of the roof, h , to the nearest millimetre.
- The roof's longer supports, labelled L , are 5200 mm long. Show that the shorter supports, labelled s , are 2193 mm shorter than the longer supports.



19. A flagpole is attached to the ground by two wires, as illustrated in the diagram shown. Use the information in the diagram to determine the length of the lower wire, x .
Give your answer correct to 1 decimal place.



20. Evaluate the values of the pronumerals w , x , y and z in the diagram shown.



LESSON

5.4 Trigonometric ratios

LEARNING INTENTIONS



At the end of this lesson you should be able to:

- identify the hypotenuse, opposite and adjacent sides of a right-angled triangle with respect to a given angle
- use the ratios of a triangle's sides to calculate the sine, cosine and tangent of an angle.

5.4.1 What is trigonometry?

eles-4756

- The word **trigonometry** is derived from the Greek words *trigonon* ('triangle') and *metron* ('measurement'). The literal translation of the word is 'to measure a triangle'.
- Trigonometry deals with the relationship between the sides and the angles of a triangle.
- As we have already seen, the longest side of a right-angled triangle is called the hypotenuse. It is always located opposite the right angle.
- In order to name the remaining two sides of a triangle, another angle, called the 'given angle', θ , must be labelled on the triangle. θ is pronounced 'theta'; it is the eighth letter of the Greek alphabet.
- The side that is across from the given angle is called the **opposite side** (or **altitude**). You can see that the opposite side does not touch the given angle at all.
- The remaining side is called the **adjacent side** (or **base**).
- Note that the given angle always sits between the hypotenuse and the adjacent side.
- When the hypotenuse equals one, $\sin(\theta) = \text{Opposite (or altitude)}$ and $\cos(\theta) = \text{Adjacent (or base)}$.




WORKED EXAMPLE 9 Labelling the sides of a right-angled triangle

Label the sides of the right-angled triangle shown using the words 'hypotenuse', 'adjacent' and 'opposite'.




THINK


1. Identify the hypotenuse. Remember that the hypotenuse always lies opposite the right angle.

WRITE

2. Look at the position of the given angle. The given angle always sits between the hypotenuse and the adjacent side; therefore, the adjacent side is the bottom side of this triangle.



3. The opposite side does not touch the given angle; therefore, it is the vertical side on the right of this triangle.



5.4.2 Trigonometric ratios


eles-4757

- Trigonometry is based upon the ratios between pairs of side lengths. Each of these ratios is given a special name.
- In any right-angled triangle:

$$\mathbf{sine}(\theta) = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\mathbf{cosine}(\theta) = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\mathbf{tangent}(\theta) = \frac{\text{opposite}}{\text{adjacent}}$$



- These rules are often abbreviated as shown in the following.

Trigonometric ratios

$$\sin(\theta) = \frac{O}{H} \quad \cos(\theta) = \frac{A}{H} \quad \tan(\theta) = \frac{O}{A}$$

- The following mnemonic can be used to help remember the trigonometric ratios.


SOH – CAH – TOA

- In this mnemonic:

- **SOH** refers to $\sin(\theta) = \frac{\text{Opposite}}{\text{Hypotenuse}}$


- **CAH** refers to $\cos(\theta) = \frac{\text{Adjacent}}{\text{Hypotenuse}}$

- **TOA** refers to $\tan(\theta) = \frac{\text{Opposite}}{\text{Adjacent}}$.



WORKED EXAMPLE 10 Identifying trigonometric ratios


For the triangle shown, write the equations for the sine, cosine and tangent ratios of the given angle (θ).



THINK

1. Label the sides of the triangle.

WRITE



2. Write the trigonometric ratios.


$$\sin(\theta) = \frac{O}{H}, \quad \cos(\theta) = \frac{A}{H}, \quad \tan(\theta) = \frac{O}{A}$$

3. Substitute the values of A, O and H into each formula.

$$\sin(\theta) = \frac{5}{13}, \quad \cos(\theta) = \frac{12}{13}, \quad \tan(\theta) = \frac{5}{12}$$

WORKED EXAMPLE 11 Relating sides to trigonometric ratios

For each of the following triangles, write the trigonometric ratio that relates the two given sides and the given angle.




THINK

a. 1. Label the given sides.


2. We are given O and H. These are used in SOH. Write the ratio.
3. Substitute the values of the pronumerals into the ratio.
4. Simplify the fraction.

b. 1. Label the given sides.

2. We are given A and O. These are used in TOA. Write the ratio.
3. Substitute the values of the angle and the pronumerals into the ratio.




WRITE



$$\sin(\theta) = \frac{O}{H}$$

$$\sin(b) = \frac{6}{15}$$

$$\sin(b) = \frac{2}{5}$$



$$\tan(\theta) = \frac{O}{A}$$

$$\tan(50^\circ) = \frac{x}{18}$$

on Resources



eWorkbook Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)



Interactivities Individual pathway interactivity: Trigonometric ratios (int-4498)
 Triangles (int-3843)
 Trigonometric ratios (int-2577)

5.4 Quick quiz **on**

5.4 Exercise

Individual pathways

PRACTISE

1, 3, 5, 8, 11, 13, 16, 19

CONSOLIDATE






2, 6, 9, 14, 17, 20

MASTER




4, 7, 10, 12, 15, 18, 21

Fluency


WE9 For questions 1 and 2, label the sides of the following right-angled triangles using the words ‘hypotenuse’, ‘adjacent’ and ‘opposite’.



3. For each of the following right-angled triangles, label the given angle and, where appropriate, the hypotenuse, the adjacent side and the opposite side.




4. **MC** Identify which option correctly names the sides and angle of the triangle shown.




- A. $\angle B = \theta$, AB = adjacent side, AC = hypotenuse, BC = opposite side
- B. $\angle C = \theta$, AC = opposite side, BC = hypotenuse, AC = adjacent side
- C. $\angle A = \theta$, AB = opposite side, BC = hypotenuse, AC = adjacent side
- D. $\angle C = \theta$, AB = opposite side, AC = hypotenuse, BC = adjacent side
- E. $\angle C = \theta$, BC = adjacent side, AB = hypotenuse, AC = opposite side

WE10 For questions 5–10, based on the given angle in the triangles shown, write the expressions for the ratios of sine, cosine and tangent.


5.




6.




7.




8.



9.




10.




11. WE11 Write the trigonometric ratio that relates the two given sides and the given angle in each of the following triangles.


a.




b.







c.



d.



12. Write the trigonometric ratio that relates the two given sides and the given angle in each of the following triangles.



Understanding

13. **MC** Identify the correct trigonometric ratio for the triangle shown.


A. $\tan(\gamma) = \frac{a}{c}$

B. $\sin(\gamma) = \frac{c}{a}$

C. $\cos(\gamma) = \frac{c}{b}$

D. $\sin(\gamma) = \frac{c}{b}$

E. $\tan(\gamma) = \frac{a}{b}$



14. **MC** Identify the incorrect trigonometric ratio for the triangle shown.


A. $\sin(\alpha) = \frac{b}{c}$

B. $\sin(\alpha) = \frac{a}{c}$

C. $\cos(\alpha) = \frac{a}{c}$

D. $\tan(\alpha) = \frac{b}{a}$

E. $\cos(\alpha) = \frac{b}{c}$



15. **MC** A right-angled triangle contains angles α and β as well as its right angle. Identify which of the following statements is always correct.

A. $\sin(\alpha) = \sin(\beta)$

B. $\tan(\alpha) = \tan(\beta)$

C. $\cos(\alpha) = 1 - \sin(\beta)$


D. $\tan(\alpha) = \frac{1}{\tan(\beta)}$

E. $\cos(\alpha) = 1 + \sin(\beta)$

Reasoning

16. If a right-angled triangle has side lengths m , $(m + n)$ and $(m - n)$, explain which of the lengths is the hypotenuse.

17. Given the triangle shown:




a. explain why $a = b$

b. determine the value of $\tan(45^\circ)$.

18. Using a protractor and ruler, carefully measure and draw a right-angled triangle with a base 10 cm long and an angle of 60° , as shown in the diagram. Measure the length of the other two sides of this triangle and mark these lengths on the diagram as well. Give the lengths of these sides to the nearest millimetre. Use your measurements to calculate the following three ratios, correct to 2 decimal places.

$$\frac{\text{opposite}}{\text{adjacent}} = \quad , \quad \frac{\text{opposite}}{\text{hypotenuse}} = \quad , \quad \frac{\text{adjacent}}{\text{hypotenuse}} =$$



Draw another triangle, similar to the first. Make the base length any length you want, but make sure all angles are equal to the angles in the first triangle. Once this is done, measure the length of the remaining two sides. Calculate the following three ratios correct to 2 decimal places.


$$\frac{\text{opposite}}{\text{adjacent}} = \quad , \quad \frac{\text{opposite}}{\text{hypotenuse}} = \quad , \quad \frac{\text{adjacent}}{\text{hypotenuse}} =$$

Comment on the conclusions you can draw from these two sets of ratios.

Problem solving


19. A ladder leans on a wall as shown. In relation to the angle given, identify the part of the image that represents:

- the adjacent side
- the hypotenuse
- the opposite side.



20. Consider the right-angled triangle shown.

- Label each of the sides using the letters O, A and H with respect to the 41° angle.
- Determine the value of each of the following trigonometric ratios, correct to 2 decimal places.
 - $\sin(41^\circ)$
 - $\cos(41^\circ)$
 - $\tan(41^\circ)$
- Calculate the value of the unknown angle, α .
- Determine the value of each of the following trigonometric ratios, correct to 2 decimal places. (*Hint:* Start by re-labelling the sides of the triangle with respect to angle α .)
 - $\sin(\alpha)$
 - $\cos(\alpha)$
 - $\tan(\alpha)$
- Comment about the relationship between $\sin(41^\circ)$ and $\cos(\alpha)$.
- Comment about the relationship between $\sin(\alpha)$ and $\cos(41^\circ)$.
- Make a general statement about the two angles.



21. In relation to right-angled triangles, investigate the following.

- As the acute angle increases in size, determine what happens to the ratio of the length of the opposite side to the length of the hypotenuse in any right-angled triangle.
- As the acute angle increases in size, determine what happens to the ratio of the length of the adjacent side to the length of the hypotenuse. Determine what happens to the ratio of the length of the opposite side to the length of the adjacent side.
- Evaluate the largest possible value for the following.
 - $\sin(\theta)$
 - $\cos(\theta)$
 - $\tan(\theta)$

LESSON

5.5 Calculating unknown side lengths

LEARNING INTENTIONS



At the end of this lesson you should be able to:

- use a calculator to calculate the trigonometric ratios for a given angle
- determine the lengths of unknown sides in a right-angled triangle given an angle and one known side length.

5.5.1 Values of trigonometric ratios

eles-4758

- The values of the trigonometric ratios are always the same for a given angle.
For example, if a right-angled triangle has a given angle of 30° , then the ratio of the opposite side to the hypotenuse will always be equal to $\sin(30^\circ) = 0.5$.
- The ratio of the adjacent side to the hypotenuse will always be equal to $\cos(30^\circ) \approx 0.87$. This relationship is demonstrated in the table.

	$\begin{aligned}\sin(30^\circ) &= \frac{O}{H} \\ &= \frac{5}{10} \\ &= 0.5\end{aligned}$	$\begin{aligned}\cos(30^\circ) &= \frac{A}{H} \\ &= \frac{8.66}{10} \\ &= 0.87\end{aligned}$	$\begin{aligned}\tan(30^\circ) &= \frac{O}{A} \\ &= \frac{5}{8.66} \\ &= 0.58\end{aligned}$
	$\begin{aligned}\sin(30^\circ) &= \frac{O}{H} \\ &= \frac{1}{2} \\ &= 0.5\end{aligned}$	$\begin{aligned}\cos(30^\circ) &= \frac{A}{H} \\ &= \frac{1.74}{2} \\ &= 0.87\end{aligned}$	$\begin{aligned}\tan(30^\circ) &= \frac{O}{A} \\ &= \frac{1}{1.74} \\ &= 0.58\end{aligned}$

WORKED EXAMPLE 12 Evaluating trigonometric ratios using a calculator

Evaluate each of the following. Give your answers correct to 4 decimal places.

a. $\sin(53^\circ)$

b. $\cos(31^\circ)$

c. $\tan(79^\circ)$

THINK

- a. 1. Set the calculator to degree mode. Make the calculation and then write out the first 5 decimal places.
2. Round the answer to 4 decimal places.
- b. 1. Set the calculator to degree mode. Make the calculation and then write out the first 5 decimal places.
2. Round the answer to 4 decimal places.
- c. 1. Set the calculator to degree mode. Make the calculation and then write out the first 5 decimal places.
2. Round the answer to 4 places.

WRITE

a. $\sin(53^\circ) = 0.79863$

≈ 0.7986

b. $\cos(31^\circ) = 0.85716$

≈ 0.8572

c. $\tan(79^\circ) = 5.14455$

≈ 5.1446

TI | THINK

a-c. 1. Ensure your calculator is set to Degree and Approximate modes. To do this, press:

- HOME
 - 5: settings
 - 2: Document settings
- In the Display Digits, select Fix 4, tab to Angle and select Degree, then tab to Calculation Mode and select Approximate.



Tab to OK and press ENTER.

2. On a Calculator page, complete the entry lines as:

sin(53)
cos(31)
tan(79)

Press CTRL ENTER after each entry to get a decimal approximation.

DISPLAY/WRITE



- a.** $\sin(53^\circ) \approx 0.7986$
- b.** $\cos(31^\circ) \approx 0.8572$
- c.** $\tan(79^\circ) \approx 5.1446$


CASIO | THINK

a-c. Ensure the calculator is in Degree and Decimal modes. On the Main screen, complete the entry lines as:

sin(53)
cos(31)
tan(79)

Press EXE after each entry.

DISPLAY/WRITE



- a.** $\sin(53^\circ) \approx 0.7986$
- b.** $\cos(31^\circ) \approx 0.8572$
- c.** $\tan(79^\circ) \approx 5.1446$


5.5.2 Determining side lengths using trigonometric ratios

eles-4759

- If a given angle and any side length of a right-angled triangle are known, it is possible to calculate the lengths of the other sides using trigonometry.

WORKED EXAMPLE 13 Calculating an unknown side length using the tangent ratio


Use the appropriate trigonometric ratio to calculate the length of the unknown side in the triangle shown. Give your answer correct to 2 decimal places.



THINK

1. Label the given sides.
2. These sides can be used in TOA. Write the ratio.
3. Substitute the values of θ , O and A into the ratio.

WRITE



$$\tan(\theta) = \frac{O}{A}$$

$$\tan(58^\circ) = \frac{x}{16.2}$$

4. Solve the equation for x .

$$16.2 \times \tan(58^\circ) = x$$

5. Calculate the value of x to 3 decimal places, then round the answer to 2 decimal places.

$$x = 16.2 \tan(58^\circ)$$

$$x = 25.925$$

$$x \approx 25.93 \text{ m}$$


TI | THINK

Ensure your calculator is set to Degree mode. In a new problem, on a Calculator page, complete the entry line as:

$$\text{solve} \left(\tan(58) = \frac{x}{16.2}, x \right)$$

Then press ENTER.

DISPLAY/WRITE



$x = 25.93 \text{ m}$ (correct to 2 decimal places)


CASIO | THINK

Ensure your calculator is set to Degree mode. On the Main screen, complete the entry line as:

$$\text{solve} \left(\tan(58) = \frac{x}{16.2}, x \right)$$

Then press EXE.


DISPLAY/WRITE



$x = 25.93 \text{ m}$ (correct to 2 decimal places)

WORKED EXAMPLE 14 Calculating an unknown side length using the cosine ratio


Use the appropriate trigonometric ratios to calculate the length of the side marked m in the triangle shown. Give your answer correct to 2 decimal places.



THINK

1. Label the given sides.
2. These sides can be used in CAH. Write the ratio.
3. Substitute the values of θ , A and H into the ratio.
4. Solve the equation for m .
5. Calculate the value of m to 3 decimal places, then round the answer to 2 decimal places.

WRITE



$$\cos(\theta) = \frac{A}{H}$$

$$\cos(22^\circ) = \frac{17.4}{m}$$

$$m \cos(22^\circ) = 17.4$$


$$m = \frac{17.4}{\cos(22^\circ)}$$

$$m = 18.766$$

$$m \approx 18.77 \text{ cm}$$

WORKED EXAMPLE 15 Solving worded problems using trigonometric ratios

Bachar set out on a bushwalking expedition. Using a compass, he set off on a course $N 70^\circ E$ (or $070^\circ T$) and travelled a distance of 5 km from his base camp.




- a. Calculate how far east Bachar travelled.
- b. Calculate how far north Bachar travelled from the base camp.
Give answers in km correct to 2 decimal places.

THINK

1. Label the eastern distance x .
Label the northern distance y .
Label the sides of the triangle as hypotenuse, opposite and adjacent.
 2. To calculate the value of x , use the sides of the triangle: $x = O$, $H = 5$.
These are used in SOH. Write the ratio.
 3. Substitute the values of the angle and the pronumerals into the sine ratio.
 4. Make x the subject of the equation.
 5. Solve for x using a calculator. Write out the answer to 3 decimal places.
 6. Round the answer to 2 decimal places.
 7. Write the answer in sentence form.
1. To calculate the value of y , use the hypotenuse and the adjacent side: $y = A$, $H = 5$.
These are used in CAH. Write the ratio.

WRITE

a.



$$\sin(\theta) = \frac{O}{H}$$

$$\sin(70^\circ) = \frac{x}{5}$$

$$x = 5 \sin(70^\circ)$$

$$= 4.698$$

$$\approx 4.70 \text{ km}$$

Bachar has travelled 4.70 km east of the base camp.

b. $\cos(\theta) = \frac{A}{H}$

2. Substitute the values of the angle and the pronumerals into the cosine ratio.
3. Make y the subject of the equation.
4. Solve for y using a calculator. Write out the answer to 3 decimal places.
5. Round the answer to 2 decimal places.
6. Write the answer in sentence form.

$$\cos(70^\circ) = \frac{y}{5}$$

$$y = 5 \cos(70^\circ)$$

$$= 1.710$$

$$\approx 1.71 \text{ km}$$

Bachar has travelled 1.71 km north of the base camp.

on Resources



eWorkbook Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)



Interactivity Individual pathway interactivity: Calculating unknown side lengths (int-4499)

Exercise 5.5 Calculating unknown side lengths

learn on

5.5 Quick quiz

on

5.5 Exercise

Individual pathways

PRACTISE

1, 4, 6, 9, 12, 13, 16, 19, 22

CONSOLIDATE

2, 5, 7, 10, 14, 17, 20, 23

MASTER

3, 8, 11, 15, 18, 21, 24, 25

Fluency

1. a. **WE12** Calculate the following. Give your answers correct to 4 decimal places.
 - i. $\sin(55^\circ)$
 - ii. $\sin(11.6^\circ)$
- b. Complete the table shown. Use your calculator to calculate each value of $\sin(\theta)$. Give your answers correct to 2 decimal places.

θ	0°	15°	30°	45°	60°	75°	90°
$\sin(\theta)$							

- c. Summarise the trend in these values.
2. a. Calculate the following. Give your answers correct to 4 decimal places.
 - i. $\cos(38^\circ)$
 - ii. $\cos(53.71^\circ)$
- b. Complete the table shown. Use your calculator to calculate each value of $\cos(\theta)$. Give your answers correct to 2 decimal places.

θ	0°	15°	30°	45°	60°	75°	90°
$\cos(\theta)$							

- c. Summarise the trend in these values.

3. a. Calculate the following. Give your answers correct to 4 decimal places.

i. $\tan(18^\circ)$

ii. $\tan(51.9^\circ)$

b. Complete the table shown. Use your calculator to calculate each value of $\tan(\theta)$. Give your answers correct to 2 decimal places.

θ	0°	15°	30°	45°	60°	75°	90°
$\tan(\theta)$							


c. Determine the value of $\tan(89^\circ)$ and $\tan(89.9^\circ)$.

d. Comment on these results.


For questions 4–10, use the appropriate trigonometric ratios to calculate the length of the unknown side in each of the triangles shown. Give your answers correct to 2 decimal places.

4. WE13


a.




b.




c.




5. a.



b.




c.




6. WE14


a.




b.




c.




7. a.




b.




c.




8. a.









b.






c.






11. Calculate the lengths of the unknown sides in the triangles shown. Give your answers correct to 2 decimal places.



Understanding


12. **MC** Identify the value of x in the triangle shown, correct to 2 decimal places.

- A. 59.65
- B. 23.31
- C. 64.80
- D. 27.51
- E. 27.15




13. **MC** Identify the value of x in the triangle shown, correct to 2 decimal places.

- A. 99.24 mm
- B. 92.55 mm
- C. 185.55 mm
- D. 198.97 mm
- E. 92.55 mm




14. **MC** Identify y in the triangle shown, correct to 2 decimal places.

- A. 47.19
- B. 7.94
- C. 1.37
- D. 0.23
- E. 6.33




15. **MC** Identify the value of y in the triangle shown, correct to 2 decimal places.

- A. 0.76 km
- B. 1.79 km
- C. 3.83 km
- D. 3.47 km
- E. 3.47 m



16. **WE15** A ship that was supposed to travel due north veers off course and travels N 80° E (or 080° T) for a distance of 280 km instead. The ship's path is shown in the diagram.


- Calculate the distance east that the ship has travelled, in km correct to 2 decimal places.
- Determine the distance north that the ship has travelled, in km correct to 2 decimal places.




17. A rescue helicopter spots a missing surfer drifting out to sea on their damaged board. The helicopter descends vertically to a height of 19 m above sea level and drops down an emergency rope, which the surfer grabs onto.

Due to the high winds, the rope swings at an angle of 27° to the vertical, as shown in the diagram.

Calculate the length of the rope, in metres correct to 2 decimal places.




18. Walking along the coastline, Michelle (M) looks up at an angle of 55° and sees her friend Hella (H) at the lookout point on top of the cliff. If Michelle is 200 m from the cliff's base, determine the height of the cliff, in metres correct to 2 decimal places (Assume Michelle and Hella are the same height.)



Reasoning

- Using a diagram, explain why $\sin(70^\circ) = \cos(20^\circ)$ and $\cos(70^\circ) = \sin(20^\circ)$. Generally speaking, explain which cosine $\sin(\theta)$ will be equal to.
- One method for determining the distance across a body of water is illustrated in the diagram shown.



The required distance is AB. A surveyor moves 50 m at a right angle to point C, then uses a tool called a transit to measure the angle θ ($\angle ACB$).


- If $\theta = 12.3^\circ$, show that the distance from A to B is 10.90 m.
 - Show that a value of $\theta = 63.44^\circ$ gives a distance from A to B of 100 m.
 - Determine a rule that can be used to calculate the distance from A to C.
21. Explain why $\cos(0^\circ) = 1$ and $\sin(0^\circ) = 0$, but $\cos(90^\circ) = 0$ and $\sin(90^\circ) = 1$

Problem solving

22. Calculate the value of the pronumeral in each of the triangles shown in the following photos.




23. A tile in the shape of a parallelogram has the measurements shown. Determine the tile's width, w , to the nearest mm.




24. A pole is supported by two wires as shown. If the length of the lower wire is 4.3 m, evaluate the following to 1 decimal place.

- a. The length of the top wire
- b. The height of the pole



25. The frame of a kite is built from six wooden rods as shown. Evaluate the total length of wood used to make the frame of the kite. Give your answer to the nearest metre.



LESSON

5.6 Calculating unknown angles

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- use a calculator to identify an angle from a given trigonometric ratio
- determine the given angle in a right-angled triangle when given two side lengths.

5.6.1 Inverse trigonometric ratios

eles-4760

- When we are given an angle, the sine, cosine and tangent functions let us determine the ratio of side lengths, but what if we only know the side lengths and we want to determine the angle in a right-angled triangle?
- The inverse trigonometric functions allow us to calculate angles using sine, cosine or tangent ratios.
- We have seen that $\sin(30^\circ) = 0.5$. This means that 30° is the inverse sine of 0.5. This is written as $\sin^{-1}(0.5) = 30^\circ$.

Inverse trigonometric ratios

Sine function	Inverse sine function
$\sin(30^\circ) = 0.5$	$\sin^{-1}(0.5) = 30^\circ$

- When reading the expression $\sin^{-1}(x)$ out loud, we say, ‘the inverse sine of x ’.
- When reading the expression $\cos^{-1}(x)$ out loud, we say, ‘the inverse cosine of x ’.
- When reading the expression $\tan^{-1}(x)$ out loud, we say, ‘the inverse tangent of x ’.
- You can calculate inverse trigonometric ratios using the SIN^{-1} , COS^{-1} and TAN^{-1} buttons on your calculator.

Digital technology

1. Use your calculator to determine $\sin(30^\circ)$, then determine the inverse sine (\sin^{-1}) of the answer. Choose another angle and do the same thing.
2. Now determine $\cos(30^\circ)$ and then determine the inverse cosine (\cos^{-1}) of the answer. Choose another angle and do the same thing.
3. Lastly, determine $\tan(45^\circ)$ and then determine the inverse tangent (\tan^{-1}) of the answer. Choose another angle and do the same thing.

The fact that \sin and \sin^{-1} cancel each other out is useful when solving equations like the following.

Consider: $\sin(\theta) = 0.3$.

Take the inverse sine of both sides

$$\sin^{-1}(\sin(\theta)) = \sin^{-1}(0.3)$$

$$\theta = \sin^{-1}(0.3)$$

Consider: $\sin^{-1}(x) = 15^\circ$.

Take the sine of both sides.

$$\begin{aligned}\sin(\sin^{-1}(x)) &= \sin(15^\circ) \\ x &= \sin(15^\circ)\end{aligned}$$

Similarly, $\cos(\theta) = 0.522$ means that

$$\begin{aligned}\theta &= \cos^{-1}(0.522) \\ \text{and } \tan(\theta) &= 1.25 \text{ means that} \\ \theta &= \tan^{-1}(1.25).\end{aligned}$$

WORKED EXAMPLE 16 Evaluating inverse cosine values

Evaluate $\cos^{-1}(0.3678)$, correct to the nearest degree.

THINK

1. Set your calculator to degree mode and make the calculation.
2. Round the answer to the nearest whole number and include the degree symbol in your answer.

WRITE

$$\begin{aligned}\cos^{-1}(0.3678) &= 68.4 \\ &\approx 68^\circ\end{aligned}$$

TI | THINK

Ensure your calculator is set to Degree mode. In a new problem, on a Calculator page, complete the entry line as: solve $\cos^{-1}(0.3678)$ Then press ENTER.

WRITE/DRAW




$$\cos^{-1}(0.3678) = 68^\circ \text{ (to the nearest degree)}$$

CASIO | THINK

Ensure your calculator is set to Degree mode. On the Main screen, complete the entry line as: $\cos^{-1}(0.3678)$ Then press EXE.

WRITE/DRAW



$$\cos^{-1}(0.3678) = 68^\circ \text{ (to the nearest degree)}$$

WORKED EXAMPLE 17 Determining angles using inverse trigonometric ratios

Determine the size of angle θ in each of the following. Give answers correct to the nearest degree.

a. $\sin(\theta) = 0.6543$

b. $\tan(\theta) = 1.745$

THINK

- a. 1. θ is the inverse sine of 0.6543.
2. Make the calculation using your calculator's inverse sine function and record the answer.
3. Round the answer to the nearest degree and include the degree symbol in your answer.

WRITE

$$\begin{aligned}\text{a. } \sin(\theta) &= 0.6543 \\ \theta &= \sin^{-1}(0.6543) \\ &= 40.8 \\ &\approx 41^\circ\end{aligned}$$

- b. 1. θ is the inverse tangent of 1.745.
2. Make the calculation using your calculator's inverse tangent function and record the answer.
3. Round the answer to the nearest degree and include the degree symbol in your answer.

$$\begin{aligned} \text{b. } \tan(\theta) &= 1.745 \\ \theta &= \tan^{-1}(1.745) \\ &= 60.18 \\ &\approx 60^\circ \end{aligned}$$




5.6.2 Determining the angle when two sides are known

eles-4761

- If the lengths of any two sides of a right-angled triangle are known, it is possible to determine an angle using inverse sine, inverse cosine or inverse tangent.

WORKED EXAMPLE 18 Determining an unknown angle when two sides are known


Determine the value of θ in the triangle shown. Give your answer correct to the nearest degree.



THINK

1. Label the given sides. These are used in CAH.
Write out the ratio.
2. Substitute the given values into the cosine ratio.
3. θ is the inverse cosine of $\frac{12}{63}$.
4. Evaluate.
5. Round the answer to the nearest degree.

WRITE



$$\cos(\theta) = \frac{A}{H}$$

$$\cos(\theta) = \frac{12}{63}$$


$$\theta = \cos^{-1}\left(\frac{12}{63}\right)$$

$$= 79.0$$

$$\approx 79^\circ$$


5.6.3 Relating the tangent of the angle to the gradient

- $\tan(\theta) = \frac{O}{A}$



- The gradient is given as:

$$\text{Gradient } (m) = \frac{\text{Rise}}{\text{Run}}$$



- Comparing the two right-angled triangles, we can see:
Opposite (Altitude) = Rise
and
Adjacent (Base) = Run
- Therefore, we can say that:


$$\tan(\theta) = \frac{O}{A} = \frac{\text{Rise}}{\text{Run}} = m$$

Therefore, $\tan(\theta) = m$
Where m is the gradient.

WORKED EXAMPLE 19 Solving word problems using the inverse tangent

Roberta goes water skiing on the Hawkesbury River. She is going to try out a new ramp. The ramp rises 1.5 m above the water level and has a length of 6.4 m.


Determine the magnitude (size) of the angle that the ramp makes with the water's surface. Give your answer to the nearest degree. Hence determine the gradient of the ramp to 2 decimal places.



THINK

1. Draw a simple diagram, showing the known lengths and the angle to be found.

WRITE



2. Label the given sides. These are used in TOA.
Write out the ratio.

$$\tan(\theta) = \frac{O}{A}$$

3. Substitute the values of the pronumerals into the tangent ratio.

$$\tan(\theta) = \frac{1.5}{6.4}$$

4. θ is the inverse inverse tangent of $\frac{1.5}{6.4}$.

$$\theta = \tan^{-1}\left(\frac{1.5}{6.4}\right)$$

5. Evaluate.

$$= 13.19$$

6. Round the answer to the nearest degree.

$$\approx 13^\circ$$

7. Write the answer in words.

The ramp makes an angle of 13° with the water's surface.

8. Since $m = \tan(\theta)$ and $\theta = 13.19^\circ$.
Substitute the angle into the equation to determine the gradient.

$$\begin{aligned} m &= \tan(\theta) \\ &= \tan(13.19^\circ) \\ &= 0.23 \end{aligned}$$

9. Write the answer.

The gradient of the ramp is 0.23.

on Resources



eWorkbook Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)



Interactivities Individual pathway interactivity: Calculating unknown angles (int-4500)
Finding the angle when two sides are known (int-6046)

Exercise 5.6 Calculating unknown angles

learn on

5.6 Quick quiz

5.6 Exercise

Individual pathways

■ PRACTISE

1, 4, 7, 10, 11, 14, 18, 21

■ CONSOLIDATE

2, 5, 8, 12, 15, 17, 19, 22

■ MASTER

3, 6, 9, 13, 16, 20, 23, 24

Fluency

- WE16** Calculate each of the following, rounding your answers to the nearest degree.
a. $\sin^{-1}(0.6294)$ b. $\cos^{-1}(0.3110)$ c. $\tan^{-1}(0.7409)$
- Calculate each of the following, rounding your answers to the nearest degree.
a. $\tan^{-1}(1.3061)$ b. $\sin^{-1}(0.9357)$ c. $\cos^{-1}(0.3275)$
- Calculate each of the following, rounding your answers to the nearest degree.
a. $\cos^{-1}(0.1928)$ b. $\tan^{-1}(4.1966)$ c. $\sin^{-1}(0.2554)$
- WE17** Determine the size of the angle in each of the following, rounding your answers to the nearest degree.
a. $\sin(\theta) = 0.3214$ b. $\sin(\theta) = 0.6752$ c. $\sin(\beta) = 0.8235$ d. $\cos(\beta) = 0.9351$




5. Determine the size of the angle in each of the following, rounding your answers to the nearest degree.

- a. $\cos(\alpha) = 0.6529$ b. $\cos(\alpha) = 0.1722$ c. $\tan(\theta) = 0.7065$ d. $\tan(\alpha) = 1$




6. Determine the size of the angle in each of the following, rounding your answers to the nearest degree.

- a. $\tan(b) = 0.876$ b. $\sin(c) = 0.3936$ c. $\cos(\theta) = 0.5241$ d. $\tan(\alpha) = 5.6214$




7. **WE18** Determine the value of θ in each of the following triangles, rounding your answers to the nearest degree.



8. Determine the value of θ in each of the following triangles, rounding your answers to the nearest degree.



9. Determine the value of θ in each of the following triangles, rounding your answers to the nearest degree.



10. **MC** If $\cos(\theta) = 0.8752$, identify the value of θ , correct to 2 decimal places.


- A. 61.07° B. 41.19° C. 25.84° D. 28.93° E. 29.93°

11. **MC** If $\sin(\theta) = 0.5530$, identify the value of θ , correct to 2 decimal places.

- A. 56.43° B. 33.57° C. 28.94° D. 36.87° E. 33.67°


12. **MC** Identify the value of θ in the triangle shown, correct to 2 decimal places.

- A. 41.30°
B. 28.55°
C. 48.70°
D. 61.45°
E. 60.45°




13. **MC** Identify the value of θ in the triangle shown, correct to 2 decimal places.

- A. 42.10°
B. 64.63°
C. 25.37°
D. 47.90°
E. 22.37°




Understanding

14. A piece of fabric measuring 2.54 m by 1.5 m has been printed with parallel diagonal stripes. Determine the angle each diagonal makes with the length of the fabric.
Give your answer in degrees correct to 2 decimal places.



15. **WE19** Danny Dingo is perched on top of a 20-metre-high cliff. They are watching Erwin Emu, who is feeding on a bush that stands 8 m away from the base of the cliff. Danny has purchased a flying contraption that they hope will help them capture Erwin.
Calculate the angle to the cliff that Danny should follow downwards to catch their prey.
Give your answer in degrees correct to 2 decimal places.



16. a. Complete the table shown.

x	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
$y = \cos^{-1}(x)$	90°					60°					0°

- b. Plot the table from part a on graph paper. Alternatively you can use a spreadsheet program or a suitable calculator.
17. A zipline runs from an observation platform 400 m above sea level to a landing platform on the ground 1200 metres away.
Calculate the angle that the zipline makes with the ground at the landing platform and hence determine the gradient of the zipline.
Give your answers correct to 2 decimal places.

Reasoning


18. Safety guidelines for wheelchair access ramps used to state that the gradient of a ramp had to be in the ratio 1 : 20.
- Using this ratio, show that the angle that any ramp needs to make with the horizontal is closest to 3° .
 - New regulations have changed the guidelines about the ratio of the gradient of a ramp. Now the angle the ramp makes with the horizontal must be closest to 6° .
Explain why, using this new angle size, the new ratio could be 1 : 9.5.
19. Jayani and Lee are camping with their friends Awer and Susie. Both couples have tents that are 2 m high. The top of each 2 m tent pole has to be tied with a piece of rope so that the pole stays upright. To make sure this rope doesn't trip anyone, Jayani and Lee decide that the angle between the rope and the ground should be 60° .
- Determine the length of the rope that Jayani and Lee need to run from the top of their tent pole to the ground. Give your answer in metres correct to 2 decimal places.
 - Awer and Susie set up their tent further into the camping ground. They want to secure their tent pole using a piece of rope that they know is somewhere between 2 and 3 metres long.
 - Explain why Awer and Susie's rope will have to be longer than 2 m.
 - Show that the minimum angle Awer and Susie's rope will make with the ground will be 41.8° .

20. Use the formulas $\sin(\theta) = \frac{O}{H}$ and $\cos(\theta) = \frac{A}{H}$ to prove that $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$.


Problem solving

21. Calculate the value of the pronumeral in each of the following, in degrees correct to 2 decimal places.

a.



b.



c.



22. A family is building a patio extension for their house. One section of the new patio will have a gable roof. A similar structure is shown, with the planned post heights and the span of the roof given.

To allow more light in, the family wants the peak (the highest point) of the roof to be at least 5 m above ground level.


According to building regulations, the slope of the roof (the angle that the sloping edge makes with the horizontal) must be 22° .

- Use trigonometry to calculate whether the roof would be high enough if the angle was 22° .
- Using trigonometry, determine the size of the obtuse angle formed at the peak of the roof.



23. The height of a square-based prism is twice its base length. Calculate the angle the diagonal of the prism makes with the diagonal of the base.

24. A series of seven shapes are marked out as shown inside a square with a side length of 10 cm. If the dots marked on the diagram represent the midpoints of the square's sides, determine the dimensions of each of the seven smaller shapes.



LESSON

5.7 Angles of elevation and depression

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- use angles of depression and elevation to determine heights and distances
- use trigonometric ratios to determine angles of depression and elevation.


5.7.1 Angles of elevation and depression

eles-4762


- When looking up towards an object, the **angle of elevation** is the angle between the horizontal line and the line of vision.



- When looking down at an object, the **angle of depression** is the angle between the horizontal line and the line of vision.



- Angles of elevation and depression are always measured from horizontal lines.
- For any two objects, A and B, the angle of elevation of B, as seen from A, is equal to the angle of depression of A, as seen from B.




WORKED EXAMPLE 20 Determining height using angle of elevation

At a point 10 m from the base of a tree, the angle of elevation of the top of the tree is 38° . Calculate the height of the tree. Give your answer to the nearest centimetre.

THINK

1. Draw a simple diagram to represent the situation.
The angle of elevation is 38° from the horizontal.

WRITE



2. Label the given sides of the triangle. These sides are used in TOA. Write out the ratio.
3. Multiply both sides by 10.
4. Calculate the answer, correct to 4 decimal places.
5. Convert metres into centimetres by multiplying the height of the tree by 100.
6. Write the answer in words to the nearest centimetre.

$$\tan(38^\circ) = \frac{h}{10}$$

$$10 \tan(38^\circ) = h$$


$$h = 7.8129 \text{ m}$$

$$\approx 7.8129 \times 100 = 781.29 \text{ cm}$$

The tree is 781 cm tall.

WORKED EXAMPLE 21 Calculating an angle of depression


A 30-metre-tall lighthouse stands on top of a cliff that is 180 m high. Determine the angle of depression (θ) of a ship (to the nearest degree) from the top of the lighthouse if the ship is 3700 m from the bottom of the cliff.



THINK

1. Draw a simple diagram to represent the situation.
The height of the triangle is $180 + 30 = 210$ m. Draw a horizontal line from the top of the triangle and mark the angle of depression, θ . Mark the alternate angle as well.

WRITE



2. Label the given sides of the triangle. These sides are used in TOA. Write out the ratio.

$$\tan(\theta) = \frac{O}{A}$$

3. Substitute the given values into the ratio.

$$\tan(\theta) = \frac{210}{3700}$$

4. θ is the inverse tangent of $\frac{210}{3700}$.

$$\theta = \tan^{-1} \left(\frac{210}{3700} \right)$$

5. Calculate the answer.

$$= 3.24$$

6. Round the answer to the nearest degree.

$$\approx 3^\circ$$

7. Write the answer in words.

The angle of depression of the ship from the top of the lighthouse is 3° .

on Resources



eWorkbook

Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)



Interactivities

Individual pathway interactivity: Angles of elevation and depression (int-4501)

Finding the angle of elevation and angle of depression (int-6047)

Exercise 5.7 Angles of elevation and depression

learn on

5.7 Quick quiz

on

5.7 Exercise

Individual pathways

PRACTISE

1, 4, 6, 9, 12

CONSOLIDATE


2, 5, 7, 10, 13

MASTER


3, 8, 11, 14, 15, 16

Fluency

1. **WE20** A lifesaver standing on their tower 3 m above the ground spots a swimmer experiencing difficulty. The angle of depression of the swimmer from the lifesaver is 12° . Calculate how far the swimmer is from the lifesaver's tower. Give your answer in metres correct to 2 decimal places.



2. From the top of a 50-metre-high lookout, the angle of depression of a camp site that is level with the base of the lookout is 37° . Calculate how far the camp site is from the base of the lookout. Give your answer in metres correct to 2 decimal places.




3. Building specifications require the angle of elevation of any ramp constructed for public use to be less than 3° .

A new shopping centre is constructing its access ramps with a ratio of 7 m horizontal length to 1 m vertical height.

Calculate the angle of elevation of these new ramps, in degrees correct to 2 decimal places.

Determine whether the new ramps meet the specifications required by ramps intended for public use.

Hence, calculate the gradient of the ramp to 2 decimal places.



Understanding


4. From a point on the ground 60 m away from a tree, the angle of elevation of the top of the tree is 35° .

- Draw a labelled diagram to represent this situation.
- Calculate the height of the tree to the nearest metre.


5. Miriam wants to take a video of her daughter Alexandra's first attempts at crawling. When Alexandra lies on the floor and looks up at her mother, the angle of elevation is 17° .

If Alexandra is 5.2 m away from her mother, calculate how tall Miriam is.


Give your answer in metres correct to 1 decimal place.







6. **WE21** Hien, who is 1.95 m tall, measures the length of the shadow he casts along the ground as 0.98 m. Determine the angle of depression of the sun's rays to the nearest degree.



7. A ladder is 3.8 m tall and leaning against a wall.
- Determine the angle the ladder makes with the ground if it reaches 2.1 m up the wall. Give your answer to the nearest degree.
 - Determine how far the foot of the ladder is from the wall. Give your answer to the nearest metre.




8. **MC** A lighthouse is 78 metres tall. The angle of elevation to the top of the lighthouse from point B, which is level with the base of the lighthouse, is 60° . Select the correct diagram for this information.



E. None of these


Reasoning

9. Con and John are practising shots at goal. Con is 3.6 m away from the goal and John is 4.2 m away, as shown in the diagram. If the height of the goal post is 2.44 m, determine the maximum angle of elevation that each player can kick the ball in order to score a goal. Give your answers to the nearest degree.



10. Lifesaver Sami is sitting in a tower 10 m from the water's edge and 4 m above sea level. They spot some dolphins playing near a marker at sea directly in front of them. The marker the dolphins are swimming near is 20 m from the water's edge.
- Draw a diagram to represent this information.
 - Show that the angle of depression of Sami's view of the dolphins, correct to 1 decimal place, is 7.6° .
 - As the dolphins swim towards Sami, determine whether the angle of depression would increase or decrease. Justify your answer in terms of the tangent ratios.

11. A pair of office buildings are 100 m and 75 m high. From the top of the north side of the taller building, the angle of depression to the top of the south side of the shorter building is 20° , as shown. Show that the horizontal distance between the north side of the taller building and the south side of the shorter building is closest to 69 m.



Problem solving

12. From a rescue helicopter 80 m above the ocean, the angles of depression of two shipwreck survivors are 40° and 60° respectively. The two survivors and the helicopter are in line with each other.


- Draw a labelled diagram to represent the situation.
- Calculate the distance between the two survivors. Give your answer to the nearest metre.

13. Rouka was hiking in the mountains when she spotted an eagle sitting up in a tree. The angle of elevation of her view of the eagle was 35° . She then walked 20 m towards the tree. From her new position, her angle of elevation was 50° . The distance between the eagle and the ground was 35.5 m.


- Draw a labelled diagram to represent this information.
- If Rouka's eyes are located 9 cm below the very top of her head, calculate how tall she is. Give your answer in metres, correct to the nearest centimetre.

14. A lookout in a lighthouse tower can see two ships approaching the coast. Their angles of depression are 25° and 30° . If the ships are 100 m apart, show that the height of the lighthouse, to the nearest metre, is 242 metres.

15. As shown in the diagram, at a certain distance from an office building, the angle of elevation to the top of the building is 60° . From a distance 12 m further back, the angle of elevation to the top of the building is 45° . Show that the building is 28.4 m high.



16. A gum tree stands in a courtyard in the middle of a group of office buildings. A group of three Year 9 students, Jackie, Pho and Theo, measure the angle of elevation from three different positions. They are unable to measure the distance to the base of the tree because of the steel tree guard around the base. This diagram shows the angles of elevation and the distances measured.




- Show that $x = \frac{15 \tan(\alpha)}{\tan(\beta) - \tan(\alpha)}$, where x is the distance, in metres, from the base of the tree to Pho's position.
- The students estimate the tree to be 15 m taller than them. Pho measured the angle of elevation to be 72° . If these measurements are correct, calculate Jackie's angle of elevation, correct to the nearest degree.
- Theo did some calculations and determined that the tree was only about 10.4 m taller than them. Jackie claims that Theo's calculation of 10.4 m is incorrect.
 - Decide if Jackie's claim is correct. Show how Theo calculated a height of 10.4 m.
 - If the height of the tree was actually 15 metres above the height of the students, determine the horizontal distance Theo should have used in his calculations. Write your answer to the nearest centimetre.

LESSON

5.8 Review

5.8.1 Topic summary


Trigonometric ratios	Inverse trigonometric ratios
<ul style="list-style-type: none"> For a given angle, θ, in a right-angled triangle, the trigonometric ratios sine, cosine and tangent are constant. $\sin(\theta) = \frac{\text{opposite (O)}}{\text{hypotenuse (H)}}$ $\cos(\theta) = \frac{\text{adjacent (A)}}{\text{hypotenuse (H)}}$ $\tan(\theta) = \frac{\text{opposite (O)}}{\text{adjacent (A)}}$ To remember these ratios, use the mnemonic SOH CAH TOA (Sine Opposite Hypotenuse, Cos Adjacent Hypotenuse, Tan Opposite Adjacent). 	<ul style="list-style-type: none"> These ratios can be used to determine the size of a given angle when given two side lengths. $\sin^{-1}\left(\frac{O}{H}\right) = \theta$ $\cos^{-1}\left(\frac{A}{H}\right) = \theta$ $\tan^{-1}\left(\frac{O}{A}\right) = \theta$

PYTHAGORAS AND TRIGONOMETRY

Pythagoras' theorem


Pythagoras' theorem states the relationship between the lengths of the sides of right-angled triangles. Mathematically it is written as:

- $a^2 + b^2 = c^2$, where c is the length of the hypotenuse (the longest side) and a and b are the lengths of the other sides.
- This rule can be applied to any right-angled triangle.
- The hypotenuse is always located opposite the right angle.
- A Pythagorean triad is a group of any three whole numbers that satisfy Pythagoras' theorem.
e.g. $\{3, 4, 5\}$ is a Pythagorean triad because $3^2 = 9$, $4^2 = 16$ and $5^2 = 25$, and $9 + 16 = 25$.



Angles of elevation and depression




- When looking upwards at an object, the **angle of elevation** is the angle between the horizontal and the upward line of sight.
- When looking downwards at an object, the **angle of depression** is the angle between the horizontal and the downward line of sight.
- The angle of elevation when looking from A to B is equal to the angle of depression when looking from B to A.



5.8.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
5.2	I can identify the hypotenuse in a right-angled triangle.			
	I can recall Pythagoras' theorem.			
	I can apply Pythagoras' theorem to calculate unknown side lengths.			
5.3	I can divide composite shapes into simple shapes.			
	I can calculate unknown side lengths in composite shapes and irregular triangles.			
	I can use Pythagoras' theorem to solve problems in 3D contexts.			
5.4	I can identify the hypotenuse, opposite and adjacent sides of a right-angled triangle with respect to a given angle.			
	I can use the ratios of a triangle's sides to calculate the sine, cosine and tangent of an angle.			
5.5	I can use a calculator to calculate the trigonometric ratios for a given angle.			
	I can determine the lengths of unknown sides in a right-angled triangle given an angle and one known side length.			
5.6	I can use a calculator to identify an angle from a given trigonometric ratio.			
	I can determine the given angle in a right-angled triangle when given two side lengths.			
5.7	I can use angles of depression and elevation to determine heights and distances.			
	I can use trigonometric ratios to determine angles of depression and elevation.			


5.8.3 Project

The Great Pyramid of Giza

The Great Pyramid of Giza, in Egypt, was built over 4500 years ago. It was constructed using approximately 2 300 000 rectangular granite blocks. This construction took over 20 years to complete. When it was finished, one side of its square base was 230 m long and its vertical height was 146.5 m.



1. Each side of the Great Pyramid of Giza has a triangular face. Use Pythagoras' theorem and the dimensions provided to calculate the height of each triangular face. Give your answer correct to 2 decimal places.
2. Special finishing blocks were added to the ends of each row of stones that were used to make the pyramid. These finishing blocks gave each triangular face a smooth, flat finish. Calculate the area of each face of the pyramid.




3. The edges of the pyramid join two of its faces and run all the way from the ground to the pyramid's tip. Use Pythagoras' theorem to calculate the length of these edges.

Wall braces

Builders use braces to strengthen wall frames. These braces typically run between the top and bottom horizontal sections of the wall frame.



Building industry standards require that the acute angle that a brace makes with the horizontal sections needs to be somewhere between 37° and 53° . Sometimes more than one brace may be required if the frame is particularly long, as shown in the diagram.





1. Cut out some thin strips of cardboard and arrange them in the shape of a rectangle to represent a wall frame. Place pins at the corners of the rectangle to hold the strips together. Notice that the frame can easily be moved out of shape. Attach a brace to this frame according to the building industry standards discussed. Write a brief comment to describe what effect this brace has on the frame.
2. Investigate what happens to the length of the brace as the acute angle it creates with the base is increased from 37° to 53° .
3. Use your findings from question 2 to calculate the angle that requires the shortest brace and the angle that requires the longest brace.

Most modern houses are constructed with a ceiling height (the height of the walls from floor to ceiling) of 2.4 m. Use this information to help with your calculations for the following questions.

4. Assume you are working with a section of wall that is 3.5 m long. Calculate the length of the longest possible brace. Draw a diagram and show your working to support your answer.
5. Calculate the minimum wall length for which two braces are required. Draw a diagram and show your working to support your answer.
6. Some older houses have ceilings that are over 2.4. m high. Answer the questions in 4 and 5 for a house with a ceiling that is 3 m high. Draw diagrams and show your workings to support your answers.
7. Take the measurements of a wall with no windows at your school or at home. Draw a scale drawing of the frame of this wall and show where the brace or braces for this wall might lie. Calculate the length and angle of each brace.

on Resources


-  **eWorkbook** Topic 5 Workbook (worksheets, code puzzle and project) (ewbk-2005)
-  **Interactivities** Crossword (int-0703)
Sudoku puzzle (int-3206)

Exercise 5.8 Review questions

Fluency


1. **MC** Identify the length of the third side in the triangle shown.

- A. 34.71 m B. 2.96 m C. 5.89 m
 D. 1722 cm E. 58.9 cm




2. **MC** Select the most accurate measure for the length of the third side in the triangle shown.

- A. 4.83 m. B. 23.3 cm. C. 3.94 m.
 D. 4826 mm. E. 4.83 mm.



3. **MC** Select the value of x in this figure.

- A. 5.4 B. 7.5 C. 10.1
 D. 10.3 E. 10.5




4. **MC** Identify which of the following is not a Pythagorean triad.

- A. 3, 4, 5 B. 6, 8, 10 C. 5, 12, 13
 D. 2, 3, 4 E. 12, 16, 20

5. **MC** Identify which of the following correctly names the sides and angle of the triangle shown.


- A. $\angle C = \theta$, AB = adjacent side, AC = hypotenuse, BC = opposite side
 B. $\angle C = \theta$, AB = opposite side, BC = hypotenuse, AC = adjacent side
 C. $\angle A = \theta$, AB = opposite side, AC = hypotenuse, BC = adjacent side
 D. $\angle A = \theta$, AB = adjacent side, AC = hypotenuse, BC = opposite side
 E. $\angle A = \theta$, AB = adjacent side, BC = hypotenuse, AC = opposite side



6. **MC** Select which of the following statements is correct.

- A. $\sin(60^\circ) = \cos(60^\circ)$ B. $\cos(25^\circ) = \cos(65^\circ)$ C. $\cos(60^\circ) = \sin(30^\circ)$
 D. $\sin(70^\circ) = \cos(70^\circ)$ E. $\cos(30^\circ) = \sin(60^\circ)$

7. **MC** Identify the value of x in the triangle shown, correct to 2 decimal places.



- A. 26.49 B. 10.04 C. 12.85
 D. 20.70 E. 10.06

8. **MC** Identify which of the following could be used to calculate the value of x in the triangle shown.


A. $x = \frac{172.1}{\cos(29^\circ)}$

B. $x = \frac{172.1}{\sin(29^\circ)}$

C. $x = 172.1 \times \sin(29^\circ)$

D. $x = 172.1 \times \cos(29^\circ)$

E. $x = 172.1 \times \tan(29^\circ)$



9. **MC** Identify which of the following could be used to calculate the value of x in the triangle shown.


A. $x = \frac{115.3}{\sin(23^\circ)}$

B. $x = \frac{115.3}{\cos(67^\circ)}$

C. $x = 115.3 \times \sin(67^\circ)$

D. $x = \frac{115.3}{\cos(23^\circ)}$

E. $x = 115.3 \times \cos(67^\circ)$



10. **MC** Identify which of the following could be used to calculate the value of x in the triangle shown.


A. $x = \frac{28.74}{\cos(17^\circ)}$

B. $x = 28.74 \times \sin(17^\circ)$

C. $x = 28.74 \times \cos(17^\circ)$

D. $x = 28.74 \times \cos(73^\circ)$

E. $x = 28.74 \times \sin(73^\circ)$



11. **MC** Select which of the following is closest to the value of $\tan^{-1}(1.8931)$.

A. 62°


B. 0.0331°

C. 1.08°

D. 69°

E. 63°

12. **MC** Select the value of θ in the triangle shown, correct to 2 decimal places.



A. 40.89°

B. 60°


C. 35.27°

D. 30°


E. 0.5°

13. Calculate x , correct to 2 decimal places.


a.



b.




c.




14. Calculate x , correct to 2 decimal places.


a.



b.



c.




15. The top of a kitchen table measures 160 cm by 90 cm. A beetle walks diagonally across the table from one corner to the other. Calculate how far it walks. Give your answer in cm correct to 2 decimal places.

16. A broom leans against a wall. The broom is 1.5 m long and reaches 1.2 m up the wall. Calculate how far the bottom of the broom is from the base of the wall. Write your answer in metres correct to 1 decimal place.




17. Calculate the unknown values in the figures shown. Give your answers correct to 2 decimal places.


a.



b.



18. A beach athletics event involves three swimming legs and a run along the beach back to the start of the first swimming leg. Some of the distances of the race course are shown. Calculate the total distance covered in the race, correct to the nearest metre.






19. True or False? The following are Pythagorean triads.

a. 15, 36, 39

b. 50, 51, 10




c. 50, 48, 14

20. Label the unlabelled sides of the following right-angled triangles using the symbol θ and the words 'hypotenuse', 'adjacent' and 'opposite' where appropriate.



21. Write a trigonometric ratio, in fraction form, that connects the lengths of the given sides and the size of the given angle in the following triangles.

Use your ratio to then calculate the size of the angle, correct to the nearest degree.









22. Use a calculator to evaluate the following trigonometric ratios, correct to 4 decimal places.

a. $\sin(54^\circ)$

b. $\cos(39^\circ)$

c. $\tan(12^\circ)$

23. Calculate the values of the pronumerals in each of the following triangles. Give your answers correct to 2 decimal places.



24. Evaluate each of the following, correct to the nearest degree.

a. $\sin^{-1}(0.1572)$

b. $\cos^{-1}(0.8361)$

c. $\tan^{-1}(0.5237)$

25. Calculate the size of the angle in each of the following. Give your answers correct to the nearest degree.

a. $\sin(\theta) = 0.5321$

b. $\cos(\theta) = 0.7071$

c. $\tan(\theta) = 0.8235$

d. $\cos(\alpha) = 0.3729$

e. $\tan(\alpha) = 0.5774$

f. $\sin(\beta) = 0.8660$

g. $\cos(\beta) = 0.5050$

h. $\tan(\beta) = 8.3791$

26. A tree is 6.7 m tall. At a certain time of the day it casts a shadow that is 1.87 m long. Determine the angle of depression of the rays of the sun at that time.

Write your answer in degrees rounded to 2 decimal places.


Problem solving

27. A 10-metre-high flagpole stands in the corner of a rectangular park that measures 240 m by 150 m.

a. Determine the following, in metres correct to 2 decimal places.


- The length of the diagonal of the park
- The distance from A to the top of the pole
- The distance from B to the top of the pole

b. A bird flies from the top of the pole to the middle of the park. Determine far the bird has to fly to reach this point, in metres correct to 2 decimal places.




28. Calculate the perimeters of the shapes shown, round your answers to 2 decimal places.


a.



b.



29. Evaluate the length of the shortest distance between points B and D inside the figure shown. Write your answer in metres correct to 2 decimal places.



30. A pair of towers stand 30 m apart. From the top of tower A, the angle of depression of the base of tower B is 60° and the angle of depression of the top of tower B is 30° .

Rounding your answer to the nearest metre, calculate the height of tower B.


31. Calculate the angles of a triangle whose sides can be described using the Pythagorean triad 3, 4, 5. Round your answers to the nearest degree.

32. A stack of chairs is 2 m tall. The stack needs to fit through a doorway that is 1.8 m high. The maximum angle that the stack of chairs can be safely tilted is 25° to the vertical.

Based on this information, determine if it is safe to try to move the stack of chairs through the doorway.

33. Evaluate the side length of the largest square that can be drawn within a circle that has a radius of r . Give your answer as a surd.

34. A Biltmore stick is used by foresters to measure the height and diameter of a tree. The diagram below shows how a Biltmore stick was used to measure the height of a tree.



A Biltmore stick is usually used by standing 20.12 m from the base of a tree. If the Biltmore stick reads 35.24 m to merchantable height (the height to the top of the trunk of the tree) and 1.80 m to the stump height (the height from eye level down to the base of the tree stump), as per the diagram.

- Determine the angle of elevation to the merchantable height, in degrees correct to 2 decimal places.
- If the average diameter of the tree is 0.8 m, approximate the surface area of bark that could be taken off the tree to be used by First Nation artists to produce paintings.
Write your answer in square metres correct to 2 decimal places.



To test your understanding and knowledge of this topic go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers


Topic 5 Pythagoras and trigonometry

5.1 Pre-test


- 5 cm
- 30.96 m
- B
- $a + b$
- C
- D
- D
- a. 3.47 cm b. 5.14 mm
- a. 0.8870 b. 0.9778 c. 3.7062
- E
- 23.4°
- A
- D
- 7.31 m
- 48.53 m

5.2 Pythagoras' theorem

- a. i. r
ii. $r^2 = p^2 + s^2$
- i. x
ii. $x = y^2 + z^2$
- i. k
ii. $k^2 = m^2 + w^2$
- i. FU
ii. $(FU)^2 = (VU)^2 + (VF)^2$
- a. 7.86 b. 33.27 c. 980.95
- a. $x = 12.49$ b. $p = 11.76$ cm
c. $f = 5.14$ m d. $c = 97.08$ mm
- 10.2 cm
- a. No b. No
c. Yes d. Yes
e. No f. Yes



b. 5.66 cm



b. 90.28 mm

- B
- C
- B

- a. 176.16 cm b. 147.40 cm
c. 2.62 km d. 432.12 m
- 2.60 m

13. The length of the diagonal can be calculated using Pythagoras' theorem.

$$c^2 = a^2 + b^2$$

$$c^2 = 260^2 + 480^2$$

$$c^2 = 298\,000$$

$$c = \sqrt{298\,000} = 546 \text{ m}$$

The length of the triangular circuit is

$$260 \text{ m} + 480 \text{ m} + 546 \text{ m} = 1286 \text{ m.}$$

$$5 \text{ laps of this circuit is } 5 \times 1286 = 6430 \text{ m.}$$


5 laps of the regular circuit is

$$5 \times (260 + 260 + 480 + 480) = 7400 \text{ m.}$$

$$7400 \text{ m} - 6430 \text{ m} = 970 \text{ m.}$$

14. The horizontal distance is 11.74 m, so the gradient is 0.21, which is within the limits.

15. a.




- b. 89.44 cm
c. Yes, she can reach the hook from the top step.

- a. i. {9, 40, 41}
ii. {11, 60, 61}
iii. {13, 84, 85}
iv. {29, 420, 421}

b. The middle number and the large number are one number apart.


17. The following figure shows the first three triangles, with the values of $\sqrt{2}$, $\sqrt{3}$ and $\sqrt{4}$.



5.3 Applications of Pythagoras' theorem

- 12.08 cm
- a. $k = 16.40$ m b. $x = 6.78$ cm c. $g = 4.10$ km
- a. $x = 4$, $y = 9.17$ b. $x = 6.93$, $y = 5.80$
c. $x = 13$, $y = 15.20$ d. $x = 2.52$, $y = 4.32$

4. a.



b. 11.78 mm

c. 20.02 mm²

5. a. 30.48 cm

b. 2.61 cm

c. 47.27 cm

6. a. 0.87 m

b. 0.433 m²

7. E

8. a. Yes.

b. The longest umbrella that can fit in the suitcase is an umbrella that is 101.48 cm long.

9. a. No.

b. The longest straw that can fit in the lunchbox is a straw that is 17.72 cm long.

10. 242 cm

11. a. $w = 0.47$ m

b. 0.64 m²

c. 12.79 m²

d. \$89.85

12. Width = 12 mm, area = 480 mm²

13. 179 mm

14. a. No, AC is calculated by summing the lengths of AD and DC, so these lengths have to be calculated first.

b. AD = 4.47 cm, DC = 9.17 cm, AC = 13.64 cm

c. The triangle ABC is not right-angled because $(AB)^2 + (BC)^2 \neq (AC)^2$.

15. Even though a problem may be represented in 3D, right-angled triangles in 2D can often be found within the problem. This can be done by drawing a cross-section of the shape or by looking at individual faces of the shape.

16. Sample responses can be found in the worked solutions in the online resources.

17. a, b. Sample responses can be found in the worked solutions in the online resources.

18. a. 2606 mm




b. Sample responses can be found in the worked solutions in the online resources.

19. 16.7 m




20. $w = 3.536$ m, $x = 7.071$ cm, $y = 15.909$ cm, $z = 3.536$ cm

5.4 Trigonometric ratios

1. a.



2. a.



3. a. DE = hypotenuse DF = opposite $\angle E = \theta$

b. GH = hypotenuse IH = adjacent $\angle H = \theta$

c. JL = hypotenuse KL = opposite $\angle J = \theta$

4. D

5. $\sin(\theta) = \frac{4}{5}$, $\cos(\theta) = \frac{3}{5}$, $\tan(\theta) = \frac{4}{3}$

6. $\sin(\alpha) = \frac{i}{g}$, $\cos(\alpha) = \frac{h}{g}$, $\tan(\alpha) = \frac{i}{h}$

7. $\sin(\beta) = 0.8$, $\cos(\beta) = 0.6$, $\tan(\beta) = 1.3$

8. $\sin(\gamma) = \frac{24}{25}$, $\cos(\gamma) = \frac{7}{25}$, $\tan(\gamma) = \frac{24}{7}$

9. $\sin(\beta) = \frac{b}{c}$, $\cos(\beta) = \frac{a}{c}$, $\tan(\beta) = \frac{b}{a}$

10. $\sin(\gamma) = \frac{v}{u}$, $\cos(\gamma) = \frac{t}{u}$, $\tan(\gamma) = \frac{v}{t}$

11. a. $\sin(\theta) = \frac{12}{15} = \frac{4}{5}$

b. $\cos(\theta) = \frac{25}{30} = \frac{5}{6}$

c. $\tan(\theta) = \frac{4}{5}$

d. $\sin(35^\circ) = \frac{17}{t}$

12. a. $\sin(\alpha) = \frac{14.3}{17.5}$

b. $\sin(15^\circ) = \frac{7}{x}$

c. $\tan(\theta) = \frac{20}{31}$

d. $\cos(\alpha) = \frac{3.1}{9.8}$

13. D

14. B

15. D

16. Provided n is a positive value, $(m + n)$ would be the hypotenuse, because it has a greater value than both m and $(m - n)$.

17. a. Sample responses can be found in the worked solutions in the online resources.

b. 1


18. The ratios remain constant when the angle is unchanged. The size of the triangle has no effect.

19. a. Ground

b. Ladder

c. Brick wall

20. a.



b. i. $\sin(41^\circ) = 0.65$

ii. $\cos(41^\circ) = 0.76$

iii. $\tan(41^\circ) = 0.86$

c. $\alpha = 49^\circ$

d. i. $\sin(49^\circ) = 0.76$

ii. $\cos(49^\circ) = 0.65$

iii. $\tan(49^\circ) = 1.16$

e. They are equal.

f. They are equal.

g. The sine of an angle is equal to the cosine of its complement.

21. a. The ratio of the length of the opposite side to the length of the hypotenuse will increase.

b. The ratio of the length of the adjacent side will decrease, and the ratio of the opposite side to the adjacent will increase.

c. i. 1

ii. 1

iii. ∞

5.5 Calculating unknown side lengths

1. a. i. 0.8192
ii. 0.2011

θ	0°	15°	30°	45°	60°	75°	90°
$\sin(\theta)$	0	0.26	0.50	0.71	0.87	0.97	1.00

- c. As θ increases, so does $\sin(\theta)$, starting at 0 and increasing to 1.

2. a. i. 0.7880
ii. 0.5919

θ	0°	15°	30°	45°	60°	75°	90°
$\cos(\theta)$	1.00	0.97	0.87	0.71	0.50	0.26	0

- c. As θ increases, $\cos(\theta)$ decreases, starting at 1 and decreasing to 0.

3. a. i. 0.3249
ii. 1.2753

θ	0°	15°	30°	45°	60°	75°	90°
$\tan(\theta)$	0	0.27	0.58	1.00	1.73	3.73	Undefined

- c. $\tan(89^\circ) = 57.29$, $\tan(89.9^\circ) = 572.96$

- d. As θ increases, $\tan(\theta)$ increases, starting at 0 and becoming very large. There is no value for $\tan(90^\circ)$.

4. a. 13.02 m b. 7.04 m c. 27.64 mm
5. a. 2.79 cm b. 6.27 m c. 14.16 m
6. a. 2.95 cm b. 25.99 cm c. 184.73 cm
7. a. 14.06 km b. 8.43 km c. 31.04 m
8. a. 26.96 mm b. 60.09 cm c. 0.84 km
9. a. 0.94 km b. 5.59 m c. 41.67 m
10. a. 54.73 m b. 106.46 cm c. 298.54 mm
11. a. $a = 17.95$, $b = 55.92$
b. $a = 15.59$, $b = 9.00$, $c = 10.73$
c. $a = 12.96$, $b = 28.24$, $c = 15.28$

12. D

13. B

14. A

15. D

16. a. 275.75 km b. 48.62 km

17. 21.32 m

18. 285.63 m

19. Sample responses can be found in the worked solutions in the online resources.

20. a. Sample responses can be found in the worked solutions in the online resources.

b. Sample responses can be found in the worked solutions in the online resources.

c. $AC = \frac{AB}{\tan(\theta)}$

21. Sample responses can be found in the worked solutions in the online resources.

22. a. $x = 12.87$ m b. $h = 3.01$ m c. $x = 2.60$ m

23. $w = 41$ mm

24. a. 5.9 m b. 5.2 m

25. 4 m

5.6 Calculating unknown angles

1. a. 39° b. 72° c. 37°
2. a. 53° b. 69° c. 71°
3. a. 79° b. 77° c. 15°
4. a. 19° b. 42° c. 55° d. 21°
5. a. 49° b. 80° c. 35° d. 45°
6. a. 41° b. 23° c. 58° d. 80°
7. a. 47° b. 45° c. 24°
8. a. 43° b. 45° c. 18°
9. a. 26° b. 12° c. 76°

10. D

11. B

12. D


13. C

14. 30.56°

15. 21.80°

16. a.

x	$y = \cos^{-1}(x)$
0.0	90°
0.1	84°
0.2	78°
0.3	73°
0.4	66°
0.5	60°
0.6	53°
0.7	46°
0.8	37°
0.9	26°
1.0	0°




17. 18.43° and $m = 0.33$.


18. a. Sample responses can be found in the worked solutions in the online resources.


b. Sample responses can be found in the worked solutions in the online resources.


19. a. 2.31 m
 b. Sample responses can be found in the worked solutions in the online resources.
20. Sample responses can be found in the worked solutions in the online resources.
21. a. $\theta = 26.01^\circ$ b. $\theta = 32.01^\circ$ c. $x = 22.62^\circ$
22. a. The roof would not be high enough.
 b. 136°
23. 54.74°
24. Large square dimensions: length = 5 cm, width = 5 cm
 Large triangle dimensions: base = 5 cm, height = 5 cm
 Small square dimensions: length = $\frac{5\sqrt{2}}{2}$ cm,
 width $\frac{5\sqrt{2}}{2}$ cm
 Small triangle dimensions: base = $\frac{5\sqrt{2}}{2}$ cm, height $\frac{5\sqrt{2}}{2}$ cm
 Parallelogram dimensions: height = 5 cm, length $5\sqrt{2}$ cm


5.7 Angles of elevation and depression

1. 14.11 m
 2. 66.35 m
 3. The new ramps have an angle of inclination of 8.13° .
 This does not meet the required specifications. $m = 0.14$.
4. a.  b. 42 m



5. 1.6 m
 6. 63°
 7. a. 34° b. 3 m
 8. B
 9. Con: 34° , John: 30°
10. a. 



- b. Sample responses can be found in the worked solutions in the online resources.
 c. Sample responses can be found in the worked solutions in the online resources.
11. Sample responses can be found in the worked solutions in the online resources.
12. a.  b. 49 m
13. a. Sample responses can be found in the worked solutions in the online resources.
 b. 1.64 m
14. Sample responses can be found in the worked solutions in the online resources.
15. Sample responses can be found in the worked solutions in the online resources.
16. a. Sample responses can be found in the worked solutions in the online resources.
 b. 37°
 c. i. Yes
 ii. 17.26 m

Project

The Great Pyramid of Giza

1. 186.25 m
 2. 21 418.75 m²
 3. 218.89 m


Wall braces

1. Sample responses can be found in the worked solutions in the online resources.
 2. Sample responses can be found in the worked solutions in the online resources.
 3. 53° requires the shortest brace and 37° requires the longest brace.
 4. 4.24 m
 5. 3.62 m
 6. 4.61 m, 4.52 m
 7. Sample responses can be found in the worked solutions in the online resources.


5.8 Review questions

1. C
 2. D
 3. D
 4. D
 5. D
 6. C
 7. B
 8. B
 9. D
 10. C
 11. A
 12. D
 13. a. 11.06 m
 b. 12.40 cm
 c. 429.70 cm or 4.30 m
 14. a. 113.06 cm
 b. 83.46 mm
 c. 55.50 mm or 5.55 cm
 15. 183.58 cm
 16. 0.9 m
 17. a. $x = 12.69$
 b. $l = 11.53, k = 10.34$
 18. 593 m
 19. a. True b. False c. True


20. a.



b.



c.



21. a. $\cos(\theta) = \frac{6}{7}$

$\theta = 31^\circ$

b. $\tan(\beta) = \frac{12}{5}$

$\beta = 67^\circ$

c. $\sin(\gamma) = \frac{11}{13}$

$\gamma = 58^\circ$

22. a. 0.8090 b. 0.7771 c. 0.2126

23. a. 7.76 b. 36.00 c. 2.56 m
d. 19.03 e. 6.79 km f. 394.29 mm

24. a. 9° b. 33° c. 28°

25. a. 32° b. 45° c. 39°
d. 68° e. 30° f. 60°
g. 60° h. 83°

26. 74.41°

27. a. i. 283.02 m
ii. 240.21 m
iii. 150.33 m

b. 141.86 m

28. a. 64.81 cm b. 84.06 m

29. 59.24 m

30. 35 m

31. 90° , 53° and 37°

32. No. The stack of chairs must be tilted by 25.84° to fit through the doorway, which is more than the safe angle of 25° .

33. $\frac{2r}{\sqrt{2}} = \sqrt{2}r$

34. a. $\theta = 60.28^\circ$ b. 93.09 m^2

6 Linear and non-linear graphs

LESSON SEQUENCE

6.1 Overview	322
6.2 Plotting linear graphs	325
6.3 Features of linear graphs	331
6.4 The equation of a straight line	339
6.5 Sketching linear graphs	348
6.6 Technology and linear graphs	356
6.7 Practical applications of linear graphs	360
6.8 Midpoint of a line segment and distance between two points	365
6.9 Non-linear relations	371
6.10 Review	385



LESSON

6.1 Overview

6.1.1 Why learn this?

In many aspects of life, one quantity depends on another quantity, and the relationship between them can be described by equations. These equations can then be used in mathematical modelling to gain an understanding of the situation — we can draw graphs and use them to analyse, interpret and explain the relationship between the variables, and to make predictions about the future. Scientists, engineers, health professionals and financial analysts all rely heavily on mathematical equations to model real-life situations and solve problems.

Linear graphs are used extensively to represent trends, for example in the stock market or when considering the population growth of various countries. Some of the most common curves we see each day are in the arches of bridges. The arch shape is a parabolic function. Architects and structural engineers use arches extensively in buildings and other structures because its shape is very strong and stable. In space, satellites orbiting Earth follow an elliptical path, while the orbits of planets are almost circular.

Because of the many uses of linear and non-linear graphs, it is important to understand the basic concepts that you will study in this topic, such as gradient, how to calculate the distance between two points, and how to sketch linear and non-linear relationships.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities



Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 6.1 Pre-test

- State the location of the point $(-3, 0)$.
- Using the table shown, match each point in the left-hand column with the line in the right-hand column that passes through that point.

Point	Line
a. $(3, -1)$	A. $y = 3 - 2x$
b. $(0, 4)$	B. $2y = x$
c. $(2, 1)$	C. $x + y = 2$
d. $(-1, 5)$	D. $y = x + 4$

- MC** Select the rule that corresponds to the table of values shown.

x	-3	-2	-1	-0	1
y	2	1	0	-1	-2


- A. $y = -x + 1$ B. $y = x + 1$ C. $y = -x - 1$ D. $y = x - 1$ E. $y = -x$
- Calculate the gradient of the line passing through the points $\left(-3, \frac{1}{2}\right)$ and $\left(5, -\frac{7}{2}\right)$.
 - MC** Select the gradient of the line $3x + 4y = 12$.
 A. $\frac{4}{3}$ B. 3 C. 4 D. $-\frac{3}{4}$ E. -3
 - Match the gradients and y -intercepts to the rules given by $y = mx + c$.

Gradient and y -intercept	Rule
a. $m = \frac{1}{2}, c = 3$	A. $2y - 6x = 4$
b. $m = \frac{1}{4}, c = 0$	B. $y = \frac{1}{2}x + 3$
c. $m = 4, c = -1$	C. $4y = x$
d. $m = 3, c = 2$	D. $y = 4x - 1$

- Determine the rule for a straight line that passes through the origin and point $(2.4, -0.6)$.
 - Determine the rule for a straight line that has an x -intercept = -20 and y -intercept = 400 .
- Determine the value of a so that the point $M\left(-\frac{1}{2}, -1\frac{1}{2}\right)$ is the midpoint of the segment joining points $A(3, 2)$ and $B(-4, a)$.
- Calculate the exact distance between the points $(2, 5)$ and $(-3, 7)$.

10. **MC** Select the equation of the linear graph shown.


- A. $y = 2x - 6$
- B. $y = 2x - 8$
- C. $y = -2x - 6$
- D. $y = -2x - 4$
- E. $y = -2x - 8$



11. A student plays the game Blue Blobs. She has the following blobs on her screen, as shown in the diagram.

The student types in an equation for a line that will pass through four blobs.

- a. Determine the equation of the straight line that passes through four blobs.
- b. The student then types in the equation $x = -1$. State the coordinate of the blob that both lines would hit.



12. **MC** Select the vertical asymptote for the equation $y = \frac{5}{x}$.

- A. $x = 0$
- B. $x = 5$
- C. $y = 0$
- D. $y = 5$
- E. $y = x$

13. **MC** Select the centre and radius for the graph $(x + 1)^2 + (y - 2)^2 = 9$.


- A. $(1, -2), r = 9$
- B. $(1, -2), r = 3$
- C. $(-1, 2), r = 9$
- D. $(-1, 2), r = 3$
- E. $(1, -2), r = 81$

14. **MC** For the graph $y = -x^2 + 4$, select the correct turning point of the parabola.

- A. a maximum at $(0, -4)$
- B. a minimum at $(4, 0)$
- C. a maximum at $(-4, 0)$
- D. a minimum at $(0, -4)$
- E. a maximum at $(0, 4)$

15. **MC** Select the correct equation for the graph shown.

- A. $y = x^2 - 4$
- B. $y = -(x - 2)^2 - 4$
- C. $y = (x + 2)^2 + 4$
- D. $y = -(x + 2)^2$
- E. $y = -x^2 + 2$



LESSON

6.2 Plotting linear graphs

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- plot points on the Cartesian plane
- plot a linear graph on the Cartesian plane.

6.2.1 Plotting linear graphs on the Cartesian plane

eles-4779

- The **Cartesian plane** is divided into four regions, known as quadrants, by the x - and y -axes, as shown.
- Every point in the plane is described exactly by a pair of coordinates (x, y) , where x is the horizontal location and y is the vertical location of the point. For example, the point P $(4, 2)$ is marked on the diagram.



Plotting linear graphs from a rule

- A straight-line graph is called a **linear graph**.
- The rule for a linear graph can always be written in the form $y = mx + c$.
For example, $y = 4x - 5$ or $y = x + 2$.
- A minimum of two points are needed to plot a linear graph.

Plotting a linear graph from a rule on a Cartesian plane

1. Draw a table of values for the x -values.
2. Substitute the x -values into the rule to find the y -values.
3. Plot the points on the Cartesian plane.
4. Join the points with a straight line and label the graph.

WORKED EXAMPLE 1 Plotting a graph from a rule

Plot the graph specified by the rule $y = x + 2$ for the x -values $-3, -2, -1, 0, 1, 2, 3$.


THINK

1. Draw a table and write in the required x -values.
2. Substitute each x -value into the rule $y = x + 2$ to obtain the corresponding y -value.
When $x = -3$, $y = -3 + 2 = -1$.
When $x = -2$, $y = -2 + 2 = 0$ etc.
Write the y -values into the table.

WRITE/DRAW

x	-3	-2	-1	0	1	2	3
y							
x	-3	-2	-1	0	1	2	3
y	-1	0	1	2	3	4	5


- Plot the points from the table.
- Join the points with a straight line and label the graph with its equation: $y = x + 2$.




TI | THINK

- In a new document, on a Lists & Spreadsheet page, label column A as x -values and column B as y -values. Enter the x -values from -3 to 3 into column A. Then in cell B1, complete the entry line as: $=a1 + 2$. Press ENTER. Hold down SHIFT and the down arrow key to fill down the y -values.

WRITE




- Open a Data & Statistics page. Press TAB to locate the label of the horizontal axis and select the variable x -values. Press TAB again to locate the label of the vertical axis and select the variable y -values. The points will be plotted. To change the colour of the points, select any point and press:
 - CTRL
 - MENU
 - 3: Colour.
 Choose a colour from the palette.




CASIO | THINK

- On the Spreadsheet screen, enter the x -values into column A. In cell B1, complete the entry line as: $=A1 + 2$. Press EXE.


WRITE




- Highlight the required cells in column B. Press:
 - Edit
 - Fill
 - Fill Range




3. To join the points with a line, press:
- MENU
 - 4: Analyze
 - 4: Plot Function.
- Complete the entry line as:
 $f1(x) = x + 2$



3. Highlight both columns and then press Graph. Select the scatterplot.



4. To determine the equation, press:
- Calc
 - Regression
 - Linear Regression
- The equation is
 $y = x + 2$. The line will appear as continuous.



WORKED EXAMPLE 2 Drawing linear graphs by plotting two points


Plot two points and use them to draw the linear graph $y = 2x - 1$.

THINK

1. Choose any two x -values, for example $x = -2$ and $x = 3$.
2. Calculate y by substituting each x -value into $y = 2x - 1$.
 $y = 2 \times (-2) - 1 = -5$
 $x = 3: y = 2 \times 3 - 1 = 5$
3. Plot the points $(-2, -5)$ and $(3, 5)$.
4. Draw a line through the points and label the line.

WRITE/DRAW


x	-2	3
y	-5	5



6.2.2 Points on a line

eles-4781

- Consider the line that has the rule $y = 2x + 3$ as shown in the graph.
If $x = 1$, then $y = 2(1) + 3 = 5$ so the point $(1, 5)$ lies on the line $y = 2x + 3$.
- The points $(1, -3)$, $(1, 9)$, $(1, 12)$... are not on the line, but lie above or below it.
- The point $(-6, 4)$ is in quadrant 2.
If $x = -6$: $y = 2(-6) + 3 = -9$
 $y \neq 4$
This shows that the point does not lie on the line.



WORKED EXAMPLE 3 Determining whether a point lies on a line

Determine whether the point $(2, 4)$ lies on the line given by:

a. $y = 3x - 2$

b. $x + y = 5$

THINK

a. 1. Substitute $x = 2$ into the equation $y = 3x - 2$ and solve for y .

2. When $x = 2$, $y = 4$, so the point $(2, 4)$ lies on the line. Write the answer.

b. 1. Substitute $x = 2$ into the equation $x + y = 5$ and solve for y .

2. The point $(2, 3)$ lies on the line, but the point $(2, 4)$ does not. Write the answer.

WRITE

a. $y = 3x - 2$
 $x = 2: y = 3(2) - 2$
 $= 6 - 2$
 $= 4$

The point $(2, 4)$ lies on the line $y = 3x - 2$.

b. $x + y = 5$
 $x = 2: 2 + y = 5$
 $y = 3$

The point $(2, 4)$ does not lie on the line $x + y = 5$.

DISCUSSION

In linear equations, what does the coefficient of x determine?

on Resources

- eWorkbook** Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)
- Interactivities** Individual pathway interactivity: Plotting linear graphs (int-4502)
Plotting linear graphs (int-3834)

Individual pathways

PRACTISE

1, 2, 4, 9, 12

CONSOLIDATE

3, 5, 8, 11, 13

MASTER

6, 7, 10, 14

Fluency

- MC** a. The point with coordinates $(-2, 3)$ is:

A. in quadrant 1 B. in quadrant 2 C. in quadrant 3
 D. in quadrant 4 E. on the x -axis

b. The point with coordinates $(-1, -5)$ is:

A. in the first quadrant B. in the second quadrant C. in the third quadrant
 D. in the fourth quadrant E. on the y -axis

c. The point with coordinates $(0, -2)$ is:

A. in the third quadrant B. in the fourth quadrant C. on the x -axis
 D. on the y -axis E. in the second quadrant
- WE1** For each of the following rules:

i. complete the table ii. plot the linear graph.

x	-3	-2	-1	0	1	2	3
y							

- a. $y = x$ b. $y = 2x + 2$ c. $y = 3x - 1$ d. $y = -2x$

Understanding

- WE2** By first plotting two points, draw the linear graph given by each of the following.

a. $y = -x$ b. $y = \frac{1}{2}x + 4$ c. $y = -2x + 3$ d. $y = x - 3$
- WE3** Determine whether these points lie on the graph of $y = 2x - 5$.

a. $(3, 1)$ b. $(-1, 3)$ c. $(0, 5)$ d. $(5, 5)$
- Determine whether the given point lies on the given line.


a. $y = -x - 7, (1, -8)$ b. $y = 3x + 5, (0, 5)$ c. $y = x + 6, (-1, 5)$
 d. $y = 5 - x, (8, 3)$ e. $y = -2x + 11, (5, -1)$ f. $y = x - 4, (-4, 0)$
- MC** Select the line that passes through the point $(2, -1)$.

A. $y = -2x + 5$ B. $y = 2x - 1$ C. $y = -2x + 1$ D. $x + y = 1$ E. $x + y = -1$
- Match each point with a line passing through that point.

Point	Line
a. $(1, 1)$	A. $x + y = 4$
b. $(1, 3)$	B. $2x - y = 1$
c. $(1, 6)$	C. $y = 3x - 7$
d. $(1, -4)$	D. $y = 7 - x$

Reasoning

8. The line through $(1, 3)$ and $(0, 4)$ passes through every quadrant except one. State the quadrant through which this line does not pass. Explain your answer.
9. Answer the following questions.
 - a. Determine which quadrant(s) the line $y = x + 1$ passes through.
 - b. Show that the point $(1, 3)$ does not lie on the line $y = x + 1$.
10. Explain the process of how to check whether a point lies on a given line.
11. Using the coordinates $(-1, -3)$, $(0, -1)$ and $(2, 3)$, show that a rule for the linear graph is $y = 2x - 1$.




Problem solving

12. Consider the pattern of squares on the grid shown. Determine the coordinates of the centre of the 20th square.
13. It is known that the mass of a certain kind of genetically modified tomato increases linearly over time. The following results were recorded.

Time, t (weeks)	1	4	6	9	16
Mass, m (grams)	6	21	31	46	81



- a. Plot the above points on a Cartesian plane.
 - b. Determine the rule connecting mass with time.
 - c. Show that the mass of one of these tomatoes is 101 grams after 20 weeks.
14. As a particular chemical reaction proceeds, the temperature increases at a constant rate. The graph represents the same chemical reaction with and without stirring. Interpret the graph and explain how stirring affects the reaction.



LESSON

6.3 Features of linear graphs

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the gradient of a line or line segment
- calculate the y-intercept of a linear graph.


6.3.1 The gradient (m)

eles-4782

- The **gradient** of a line is a measure of the steepness of its slope.
- The symbol for the gradient is m .
- The gradient of an interval AB is defined as the rise (distance up) divided by the run (distance across).

Gradient formula

$$m = \frac{\text{rise}}{\text{run}}$$




- If the line is sloping upwards (from left to right), the gradient is positive.
- If the line is sloping downwards, the gradient is negative.

Positive gradient (sloping upwards)	Negative gradient (sloping downwards)
<p>3 6 $m = 2$</p>	<p>2 1 $m = -2$</p>
<p>2 1 $m = \frac{1}{2}$</p>	<p>1 2 $m = -\frac{1}{2}$</p>

Determining the gradient of a line passing through two points

- Suppose a line passes through the points $(1, 4)$ and $(3, 8)$, as shown in the graph.
- By completing a right-angled triangle, it can be seen that:
 - the rise (difference in y -values): $8 - 4 = 4$
 - the run (difference in x -values): $3 - 1 = 2$
 - to determine the gradient:

$$\begin{aligned}m &= \frac{8 - 4}{3 - 1} \\ &= \frac{4}{2} \\ &= 2\end{aligned}$$



Gradient of a line

In general, if a line passes through the points (x_1, y_1) and (x_2, y_2) , then:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

WORKED EXAMPLE 4 Determining the gradient of a line passing through two points

Calculate the gradient of the line passing through the points $(-2, 5)$ and $(1, 14)$.

THINK

1. Let the two points be (x_1, y_1) and (x_2, y_2) .
2. Write the formula for gradient.
3. Substitute the coordinates of the given points into the formula and evaluate.

WRITE

$$(-2, 5) = (x_1, y_1), (1, 14) = (x_2, y_2)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \frac{14 - 5}{1 - (-2)}$$

$$= \frac{9}{1 + 2}$$

$$= \frac{9}{3}$$

$$= 3$$

4. Write the answer.

The gradient of the line passing through $(-2, 5)$ and $(1, 14)$ is 3.

Note: If you were to switch the order of the points and let $(x_1, y_1) = (1, 14)$ and $(x_2, y_2) = (-2, 5)$, then the gradient could be calculated as shown and it can be seen that the result is the same.

$$\begin{aligned} m &= \frac{y_2 - y_1}{x_2 - x_1} \\ &= \frac{5 - 14}{-2 - 1} \\ &= \frac{-9}{-3} \\ &= 3 \end{aligned}$$

Rates of change

- A rate of change measures the change in the y -values with respect to the change in the x -values.
- The rate of change is equivalent to the gradient of the graph.

The gradient of horizontal and vertical lines

- The gradient of a horizontal line is zero, since it does not have any upwards or downwards slope.





$$m = \frac{\text{rise}}{\text{run}} = \frac{0}{\text{run}} = 0$$

- The gradient of a vertical line is undefined, as the run is zero.

$$m = \frac{\text{rise}}{\text{run}} = \frac{\text{rise}}{0} = \text{undefined}$$

WORKED EXAMPLE 5 Calculating the gradient of a straight line

Calculate the gradients of the lines shown.



THINK

a. 1. Write down two points that lie on the line.

2. Calculate the gradient by finding the ratio $\frac{\text{rise}}{\text{run}}$.

b. 1. Write down two points that lie on the line.

2. Calculate the gradient.

c. 1. Write down two points that lie on the line.

2. There is no rise between the two points.

3. Calculate the gradient.

Note: The gradient of a horizontal line is always zero. The line has no slope.

d. 1. Write down two points that lie on the line.

2. The vertical distance between the selected points is 13 units. There is no run between the two points.

3. Calculate the gradient.

Note: The gradient of a vertical line is always undefined.

WRITE

a. Let $(x_1, y_1) = (0, -2)$ and $(x_2, y_2) = (10, 13)$.

$$\text{Rise} = y_2 - y_1 = 13 - (-2) = 15$$

$$\text{Run} = x_2 - x_1 = 10 - 0 = 10$$

$$m = \frac{\text{rise}}{\text{run}}$$

$$= \frac{15}{10}$$

$$= \frac{3}{2} \text{ or } 1.5$$

b. Let $(x_1, y_1) = (0, 6)$ and $(x_2, y_2) = (10, -24)$.

$$\text{Rise} = y_2 - y_1 = -24 - 6 = -30$$

$$\text{Run} = x_2 - x_1 = 10 - 0 = 10$$

$$m = \frac{\text{rise}}{\text{run}}$$

$$= \frac{-30}{10}$$

$$= -3$$

c. Let $(x_1, y_1) = (5, -6)$ and $(x_2, y_2) = (10, -6)$.

$$\text{Rise} = y_2 - y_1$$

$$= -6 - (-6) = 0$$

$$\text{Run} = x_2 - x_1$$

$$= 10 - 5 = 5$$

$$m = \frac{\text{rise}}{\text{run}}$$

$$= \frac{0}{5}$$

$$= 0$$

d. Let $(x_1, y_1) = (7, 10)$ and $(x_2, y_2) = (7, -3)$.

$$\text{Rise} = y_2 - y_1 = -3 - 10 = 13$$

$$\text{Run} = x_2 - x_1 = 7 - 7 = 0$$

$$m = \frac{\text{rise}}{\text{run}}$$

$$= \frac{13}{0}$$


$$= \text{undefined}$$



6.3.2 The y-intercept

eles-4783

- The point where a line cuts the y-axis is called the **y-intercept**.
- In the graph shown the line cuts the y-axis at $y = 2$, so the y-intercept is $(0, 2)$.
- The y-intercept of any line is found by substituting 0 for x and calculating the y -value.



WORKED EXAMPLE 6 Determining the y-intercepts of lines

Determine the y-intercepts of the lines whose linear rules are given, and state the coordinates of the y-intercept.

a. $y = -4x + 7$

b. $5y + 2x = 10$

c. $y = 2x$

d. $y = -8$

THINK

a. 1. To calculate the y-intercept, substitute $x = 0$ into the equation.

2. Solve for y .

3. Write the coordinates of the y-intercept.

b. 1. To calculate the y-intercept, substitute $x = 0$ into the equation.

2. Solve for y .

3. Write the coordinates of the y-intercept.

c. 1. To calculate the y-intercept, substitute $x = 0$ into the equation.

2. Write the coordinates of the y-intercept.

d. The value of y is -8 regardless of the x -value.

WRITE

a. $y = -4x + 7$
 $y = -4(0) + 7$

$$y = 7$$

y-intercept: $(0, 7)$

b. $5y + 2x = 10$
 $5y + 2(0) = 10$

$$5y = 10$$

$$y = 2$$

y-intercept: $(0, 2)$

c. $y = 2x$
 $y = 2(0)$
 $y = 0$




y-intercept: $(0, 0)$

d. y-intercept: $(0, -8)$

DISCUSSION

Why is the y-intercept of a graph found by substituting $x = 0$ into the equation?

on Resources

-  **eWorkbook** Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)
-  **Video eLesson** Gradient (eles-1889)
-  **Interactivities** Individual pathway interactivity: Determining linear rules (int-4506)
The gradient (int-3836)
Linear graphs (int-6484)

Exercise 6.3 Features of linear graphs

learn on

6.3 Quick quiz

6.3 Exercise

Individual pathways

PRACTISE

1, 3, 5, 8, 11, 14, 16

CONSOLIDATE

2, 4, 6, 9, 12, 15, 17, 18

MASTER

7, 10, 13, 19, 20

Fluency

1. **WE4** Calculate the gradients of the lines passing through the following pairs of points.

- a. (2, 10) and (4, 22)
- b. (1, -2) and (3, -10)
- c. (-3, 0) and (7, 0)
- d. (-4, -7) and (1, -1)
- e. (0, 4) and (4, -4.8)
- f. (-2, 122) and (1, -13)

2. Calculate the gradients of the lines passing through the following pairs of points.

- a. (2, 3) and (17, 3)
- b. (-2, 2) and (2, 2.4)
- c. (1, -5) and (5, -15.4)
- d. (-12, -7) and (8.4, -7)
- e. (-2, -17.7) and (0, 0.3)
- f. (-3, 3.4) and (5, 2.6)




3. **WE6** Determine the y-intercepts of the lines whose rules are given below.

- a. $y = 5x + 23$
- b. $y = 54 - 3x$
- c. $y = 3(x - 2)$
- d. $y = 70 - 2x$




4. Determine the y-intercepts of the lines whose rules are given below.

- a. $y = \frac{1}{2}(x + 2)$
- b. $y = \frac{x}{2} + 5.2$
- c. $y = 100 - x$
- d. $y = 100$




5. **WE5** Calculate the gradients of the lines shown.



6. Calculate the gradients of the lines shown.



7. Calculate the gradients of the lines shown.



Understanding

8. **MC** Select which of the following statements about linear graphs is false.

- A. A gradient of zero means the graph is a horizontal line.
- B. A gradient can be any real number.
- C. A linear graph can have two y -intercepts.
- D. In the form $y = mx + c$, the y -intercept equals c .
- E. An undefined gradient means the graph is a vertical line.

9. Identify the y -intercept of the line $y = mx + c$.

10. Determine the coordinates of the y -intercepts of the lines with the following rules.

- a. $y = -6x - 10$
- b. $3y + 3x = -12$
- c. $7x - 5y + 15 = 0$
- d. $y = 7$
- e. $x = 9$

Reasoning


11. Explain why the gradient of a vertical line is undefined.

12. Explain why the gradient of a horizontal line is zero.


13. Show the gradient of the line passing through the points (a, b) and (c, d) is $\frac{d-b}{c-a}$.

Problem solving


14. When using the gradient to draw a line, does it matter if you rise before you run or run before you rise? Explain.
15. Consider the graph shown.
- Determine a general formula for the gradient m in terms of x , y and c .
 - Transpose your formula to make y the subject.
Discuss what you notice about this equation.



16. Calculate the gradient, correct to 2 decimal places, of the playground equipment in the diagrams given below.



17. Safety standards for escalators outline that the gradient of an escalator must not exceed 0.58. Determine if an escalator with a height of 18 metres that has a horizontal distance of 30 metres meets the required safety standards.
18. The maximum gradient for wheelchair ramps is $\frac{1}{14}$. If a ramp is being built across a horizontal distance of 4.2 metres determine the maximum height of the ramp.
19. The price per kilogram for three different types of meat is illustrated in the graph shown.




- Calculate the gradient (using units) for each graph.
- Determine the cost of 1 kg of each type of meat.
- Evaluate the cost of purchasing:
 - 1 kg of lamb
 - 0.5 kg of chicken
 - 2 kg of beef.
- Calculate the total cost of the order in part c.

e. Complete the table below to confirm your answer from part d.

Meat type	Cost per kilogram (\$/kg)	Weight required (kg)	Cost = \$/kg × kg
Lamb		1	
Chicken		0.5	
Beef		2	
Total cost			

20. Three right-angled triangles have been superimposed on the graph shown.

- Use each of these to determine the gradient of the line.
- Does it matter which points are chosen to determine the gradient of a line? Explain your answer.
- Describe the shape of the graph.



LESSON

6.4 The equation of a straight line

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- determine the equation of a straight line given the gradient and the y-intercept
- determine the equation of a straight line given the gradient and one point
- determine the equation of a straight line given two points
- determine the equation of a straight line from a graph.

6.4.1 The general equation of a straight line

eles-4784

- The general equation of a straight line is

$$y = mx + c$$

where m is the gradient and c is the y-intercept of the line.

- Once the gradient and y-intercept are known the equation of the straight line can be stated.

WORKED EXAMPLE 7 Determining the equation from the gradient and y-intercept

Determine the equation of the line with a gradient of -2 and y-intercept of 3 .

THINK

1. Write the equation of a straight line.
2. Substitute the values $m = -2$, $c = 3$ to write the equation.

WRITE

$$y = mx + c$$
$$m = -2, c = 3$$
$$y = -2x + 3$$

6.4.2 Determining the equation of a straight line

eles-4785

- If the gradient (m) and any single point on a straight line are known, then the y-intercept can be calculated algebraically.

WORKED EXAMPLE 8 Determining the equation of a line from the gradient and a point

Determine the equation of a straight line that goes through the point $(1, -3)$, if its gradient is -2 .

THINK

1. Write the equation of a straight line.
2. Substitute the value $m = -2$.
3. Since the line passes through the point $(1, -3)$, substitute $x = 1$ and $y = -3$ into $y = -2x + c$ to calculate the value of c .
4. Solve for c .
5. Write the rule.

WRITE

$$y = mx + c$$
$$y = -2x + c$$
$$\text{When } x = 1, y = -3.$$
$$-3 = -2 \times 1 + c$$
$$-3 = -2 + c$$
$$-3 + 2 = c$$
$$c = -1$$
$$y = -2x - 1$$

- If two points on a straight line are known, then the gradient (m) can be calculated using the formula $m = \frac{y_2 - y_1}{x_2 - x_1}$, or $m = \frac{\text{rise}}{\text{run}}$.
- Using the gradient and one of the points, the equation can then be found using the method in Worked example 8.

WORKED EXAMPLE 9 Determining the equation of a line from two points

Determine the equation of the straight line passing through the points $(-1, 6)$ and $(3, -2)$.

THINK

1. Write the equation of a straight line.
2. Write the formula for calculating the gradient, m .
3. Let $(x_1, y_1) = (-1, 6)$ and $(x_2, y_2) = (3, -2)$. Substitute the values into the formula and determine the value of m .

WRITE

$$y = mx + c$$
$$m = \frac{y_2 - y_1}{x_2 - x_1}$$
$$m = \frac{-2 - 6}{3 - (-1)}$$
$$= \frac{-2 - 6}{3 + 1}$$
$$= \frac{-8}{4}$$
$$= -2$$

- Substitute the value of m into the general equation $y = mx + c$.
- Select either of the two points, say $(3, -2)$, and substitute into $y = -2x + c$.
- Solve for c .
- Write the rule using the values $c = 4, m = -2$.

$$y = -2x + c$$

Point $(3, -2)$:

$$-2 = -2(3) + c$$

$$-2 = -6 + c$$

$$c = 4$$

$$y = -2x + 4$$

TI | THINK

In a new problem, on a Calculator page, complete the entry lines as:

$$m := \frac{y_2 - y_1}{x_2 - x_1} | x_1 = -1$$

and $y_1 = 6$ and $x_2 = 3$ and $y_2 = -2$

$$y = -2x + c$$


solve $(y = -2x + c, c)$

$(|x = 3)$ and $y = -2$

$$y = -2x + c | c = 4$$

Press ENTER after each entry.

WRITE




The equation is $y = 4 - 2x$.

CASIO | THINK


- On the Spreadsheet screen, complete the entries as:

	A	B
1	-1	6
2	3	-2


WRITE



- Highlight the data in both columns. Press Graph and select the scatterplot. The points will be plotted.



- To get the equation of the line, press the Line icon. The equation will appear above the graph, and the points will be joined by a line.




The equation is $y = -2x + 4$.

- The equation of a straight line can be determined from a graph by observing the y-intercept and calculating the gradient.

WORKED EXAMPLE 10 Determining the equation of a line from its graph

Determine the equation of the linear graph shown.



THINK

1. Read the important information from the graph. The y-intercept is 4 and the x-intercept is 2. Write the coordinates of each point.
2. Calculate the gradient using the formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.
Let $(x_1, y_1) = (2, 0)$ and $(x_2, y_2) = (0, 4)$.




3. From the graph, the y-intercept is 4.
4. Write the equation.

WRITE

The graph passes through $(0, 4)$ and $(2, 0)$.

$$\begin{aligned}
 m &= \frac{4 - 0}{0 - 2} \\
 &= \frac{4}{-2} \\
 &= -2 \\
 y &= -2x + c \\
 c &= 4 \\
 y &= -2x + 4
 \end{aligned}$$

on Resources

-  **eWorkbook** Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)
-  **Video eLesson** The equation of a straight line (eles-2313)
-  **Interactivity** Individual pathway interactivity: The equation of a straight line (int-4503)

Individual pathways

PRACTISE

1, 3, 5, 10, 15, 16, 19, 22, 25

CONSOLIDATE

2, 4, 6, 8, 11, 13, 17, 20, 23, 26, 27

MASTER

7, 9, 12, 14, 18, 21, 24, 28, 29

Fluency

1. **WE7** Determine the equations of the straight lines with the gradients and y-intercepts given.
 - a. Gradient = 4, y-intercept = 2
 - b. Gradient = -4, y-intercept = 1
 - c. Gradient = 4, y-intercept = 8
2. Determine the equations of the straight lines with the gradients and y-intercepts given.
 - a. Gradient = 6, y-intercept = 7
 - b. Gradient = -2.5, y-intercept = 6
 - c. Gradient = 45, y-intercept = 135
3. Determine the equation for each straight line passing through the origin and with the gradient given.
 - a. Gradient = -2
 - b. Gradient = 4
 - c. Gradient = 10.5
4. Determine the equation for each straight line passing through the origin and with the gradient.
 - a. Gradient = -20
 - b. Gradient = 1.07
 - c. Gradient = 32
5. **WE8** Determine the equation of the straight lines with:
 - a. gradient = 1, point = (3, 5)
 - b. gradient = -1, point = (3, 5)
 - c. gradient = -4, point = (-3, 4)
 - d. gradient = 2, point = (5, -3).
6. Determine the equation of the straight lines with:
 - a. gradient = -5, point = (13, 5)
 - b. gradient = 2, point = (10, -3)
 - c. gradient = -6, point = (2, -1)
 - d. gradient = -1, point = (-2, 0.5).
7. Determine the equation of the straight lines with:
 - a. gradient = 6, point = (-6, -6)
 - b. gradient = -3.5, point = (3, 5)
 - c. gradient = 1.2, point = (2.4, -1.2)
 - d. gradient = 0.2, point = (1.3, -1.5).

8. Determine the equations of the straight lines with:

- a. gradient = -4 , x -intercept = -6
- b. gradient = 2 , x -intercept = 3
- c. gradient = -2 , x -intercept = 2 .

9. Determine the equation of the straight lines with:

- a. gradient = 5 , x -intercept = -7
- b. gradient = 1.5 , x -intercept = 2.5
- c. gradient = 0.4 , x -intercept = 2.4 .

10. **WE9** Determine the equation for each straight line passing through the given points.

- a. $(-6, 11)$ and $(6, 23)$
- b. $(1, 2)$ and $(-5, 8)$
- c. $(4, 11)$ and $(6, 11)$
- d. $(3, 6.5)$ and $(6.5, 10)$

11. Determine the equation for each straight line passing through the given points.

- a. $(1.5, 2)$ and $(6, -2.5)$
- b. $(-7, 3)$ and $(2, 4)$
- c. $(25, -60)$ and $(10, 30)$
- d. $(5, 100)$ and $(25, 500)$

12. Determine the equation for each straight lines passing through the given points.

- a. $(1, 3)$ and $(3, 1)$
- b. $(2, 5)$ and $(-2, 6)$
- c. $(9, -2)$ and $(2, -4)$
- d. $(1, 4)$ and $(-0.5, 3)$

13. Determine the rules for the linear graphs that have the following x - and y -intercepts.

- a. x -intercept = -3 , y -intercept = 3
- b. x -intercept = 4 , y -intercept = 5
- c. x -intercept = 1 , y -intercept = 6
- d. x -intercept = -40 , y -intercept = 35



14. Determine the rules for the linear graphs that have the following x - and y -intercepts.

- a. x -intercept = -8 , y -intercept = 8
- b. x -intercept = 3 , y -intercept = 6
- c. x -intercept = -7 , y -intercept = -3
- d. x -intercept = -200 , y -intercept = 50



15. Determine the rule for each straight line passing through the origin and:

- a. the point $(4, 7)$
- b. the point $(5, 5)$
- c. the point $(-4, 8)$
- d. the point $(-1.2, 3.6)$
- e. the point $(-22, 48)$
- f. the point $(-105, 35)$.



16. **WE10** Determine the equation of the line shown on each of the following graphs.



17. Determine the equation of the line shown on each of the following graphs.



18. Determine the equation of the line shown on each of the following graphs.



Understanding

19. **MC** a. The gradient of the straight line that passes through (3, 5) and (5, 3) is:

- A. -2 B. -1 C. 0 D. 1 E. 2

b. A straight line with an x -intercept of 10 and a y -intercept of 20 has a gradient of:

- A. -2 B. -1 C. -0.5 D. 0 E. 0.5


c. The rule $2y - 3x = 20$ has an x -intercept at:

- A. $-\frac{3}{2}$ B. $-\frac{2}{3}$ C. 0
 D. $\frac{2}{3}$ E. None of these

20. Given that the x -intercept of a straight line graph is $(-5, 0)$ and the y -intercept is $(0, -12)$:


- a. determine the equation of the straight line
 b. calculate the value of y when $x = 19.3$.

21. a. Determine the equation of the straight line shown in the graph.
Use the fact that when $x = 5, y = 7$.
b. Determine the x -intercept.




Reasoning

22. The graph shows the carbon dioxide (CO_2) concentration in the atmosphere, measured in parts per million (ppm).
- If the trend follows a linear pattern, determine the equation for the line.
 - Explain why c cannot be read directly from the graph.
 - Infer the CO_2 concentration predicted for 2020. State the assumption made when determining this value.
23. Show that the equation for the line that passes through the point $(3, 6)$ parallel to the line through the points $(0, -7)$ and $(4, -15)$ is $y = -2x + 12$.




24. a. Determine the equations for line A and line B as shown in this graph.
b. Write the point of intersection between line A and line B and mark it on the Cartesian plane.
c. Show that the equation of the line that is perpendicular to line B and passes through the point $(-4, 6)$ is $y = x + 10$.




Problem solving


25. The graph shown describes the mass in kilograms of metric cups of water. Write a rule to describe the mass of water (m) relative to the number of cups (c).



26. Harpinder plays the game *Space Galaxy* on her phone. The stars and spaceships are displayed on her screen as shown.
- Copy the diagram. On the diagram, draw a straight line that will hit three stars.
 - Determine the equation of the straight line that will hit three stars.
 - Harpinder types in the equation $y = \frac{1}{2}x + \frac{1}{2}$ and manages to hit two stars. Draw the straight line on your diagram.
 - If Harpinder types in the equation from part **b** and the equation from part **c**, determine the coordinate of the star that both lines will hit.
 - If she types $y = 2$, state how many stars will she hit.
 - Give another equation of a straight line that will hit two stars.



27. The temperature of water in a kettle is 15°C before the temperature increases at a constant rate for 20 seconds to reach boiling point (100°C). Adel argues that $T = 5t + 15$ describes the water temperature, citing the starting temperature of 15°C and that to reach 100°C in 20 seconds an increase of 5°C for every second is required.
- Explain why Adel's equation is incorrect and devise another equation that correctly describes the temperature of the water.
28. A father wants to administer a children's liquid painkiller to his child. The recommended dosage is a range, $7.5 - 9$ mL for an average weight of $12 - 14$ kg. The child weighs 12.8 kg. The father uses a linear relationship to calculate an exact dosage.
- Evaluate the dosage that the father calculates.
29. The graph shown displays the wages earned in three different workplaces.



- Identify the set allowance for each workplace.
- Determine the hourly rates for each workplace.
- Using your answers from parts **a** and **b**, determine linear equations that describe the wages at each workplace.
- Match each working lifestyle below to the most appropriate workplace.
 - Working lifestyle 1: Earn the most money possible while working at most 4 hours in a day.
 - Working lifestyle 2: Earn the most money possible while working an 8-hour day.
- If a person works an average of 8 hours a day, determine the advantage of workplace C.
- Calculate how much money is earned at each workplace for a:
 - 2-hour day
 - 6-hour day.

LESSON

6.5 Sketching linear graphs

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- sketch linear graphs using the x - and y -intercept method
- sketch linear graphs using the gradient–intercept method
- sketch vertical and horizontal lines.

6.5.1 The x - and y -intercept method

eles-4786

- To sketch a linear graph all you need is two points on the line. Once you have found two points, the line can be drawn through those points.
- One method to sketch a linear graph is to find both the **x -intercept** and the **y -intercept**.
- To determine the intercepts of a graph:
 x -intercept: let $y = 0$ and solve for x
 y -intercept: let $x = 0$ and solve for y .
- If both intercepts are zero (at the origin), another point is needed to sketch the line.



WORKED EXAMPLE 11 Sketching graphs using the x - and y -intercept method

Using the x - and y -intercept method, sketch the graphs of:

a. $2y + 3x = 6$

b. $y = \frac{4}{5}x + 5$

c. $y = 2x$.

THINK

1. Write the rule.
2. To calculate the y -intercept, let $x = 0$.
Write the coordinates of the y -intercept.
3. To calculate the x -intercept, let $y = 0$.
Write the coordinates of the x -intercept.

WRITE/DRAW

a. $2y + 3x = 6$


$$\begin{aligned}x = 0: \quad 2y + 3 \times 0 &= 6 \\ 2y &= 6 \\ y &= 3\end{aligned}$$

y -intercept: $(0, 3)$

$$\begin{aligned}y = 0: \quad 2 \times 0 + 3x &= 6 \\ 3x &= 6 \\ x &= 2\end{aligned}$$

x -intercept: $(2, 0)$

4. Plot and label the x - and y -intercepts on a set of axes and rule a straight line through them. Label the graph.



- b. 1. Write the rule.
2. The rule is in the form $y = mx + c$, so the y -intercept is the value of c .
3. To calculate the x -intercept, let $y = 0$. Write the coordinates of the x -intercept.

$$b. y = \frac{4}{5}x + 5$$

$$c = 5$$

$$y\text{-intercept: } (0, 5)$$

$$y = 0: \quad y = \frac{4}{5}x + 5$$


$$0 = \frac{4}{5}x + 5$$

$$-5 = \frac{4}{5}x$$

$$x = -\frac{25}{4} \left(= -6\frac{1}{4} \right)$$

$$x\text{-intercept: } \left(-\frac{25}{4}, 0 \right)$$

4. Plot and label the intercepts on a set of axes and rule a straight line through them. Label the graph.



- c. 1. Write the rule.
2. To calculate the y -intercept, let $x = 0$.
Write the coordinates of the y -intercept.
3. The x - and y -intercepts are the same point, $(0, 0)$,
so one more point is required.
Choose any value for x , such as $x = 3$.
Substitute and write the coordinates of the point.
4. Plot the points, then rule and label the graph.


c. $y = 2x$

$$x = 0: \quad y = 2 \times 0 \\ = 0$$

y -intercept: $(0, 0)$

$$x = 3: \quad y = 2 \times 3 \\ = 6$$

Another point: $(3, 6)$



6.5.2 The gradient–intercept method

eles-4787

- To use this method, the gradient and the y -intercept must be known.
- The line is drawn by plotting the y -intercept, then drawing a line with the correct gradient through that point.

A line interval of gradient

$3 \left(= \frac{3}{1} \right)$ can be drawn with a

rise of 3 and a run of 1.



A line interval of gradient

$-2 \left(= \frac{-2}{1} \right)$ can be drawn with


a rise downwards of 2 and a run of 1.



A line interval of gradient

$\frac{3}{5}$ can be shown with a rise of

3 and a run of 5.



WORKED EXAMPLE 12 Sketching graphs using the gradient–intercept method

Using the gradient–intercept method, sketch the graphs of:

a. $y = \frac{3}{4}x + 2$


b. $4x + 2y = 3$.

THINK

- a. 1. From the equation, the y-intercept is 2. Plot the point (0, 2).
2. From the equation, the gradient is $\frac{3}{4}$, so $\frac{\text{rise}}{\text{run}} = \frac{3}{4}$.
From (0, 2), run 4 units and rise 3 units. Mark the point P (4, 5).
3. Draw a line through (0, 2) and P (4, 5). Label the graph.

WRITE

a.



- b. 1. Write the rule in gradient–intercept form:

$$y = mx + c.$$

From the equation, $m = -2$, $c = \frac{3}{2}$.

Plot the point $\left(0, \frac{3}{2}\right)$.

- b. $4x + 2y = 3$

$$2y = 3 - 4x$$


$$y = \frac{3}{2} - 2x$$

$$y = -2x + \frac{3}{2}$$

2. The gradient is -2 , so $\frac{\text{rise}}{\text{run}} = \frac{-2}{1}$.

From $\left(0, \frac{3}{2}\right)$, run 1 units and rise -2 units (i.e. go down 2 units). Mark the point P $\left(1, -\frac{1}{2}\right)$.

3. Draw a line through $\left(0, \frac{3}{2}\right)$ and P $\left(1, -\frac{1}{2}\right)$. Label the graph.



6.5.3 Horizontal and vertical lines


eles-4788

Horizontal lines ($y = c$)

- Horizontal lines are expressed in the form $y = c$, where c is the y -intercept.
- In horizontal lines the y -value remains the same regardless of the x -value.
- This can be seen by looking at a table of values, like the one shown.

x	-2	0	2	4
y	c	c	c	c

- Plotting these points gives a horizontal line, as shown in the graph.
- Horizontal lines have a gradient of 0. They do not rise or fall.




Vertical lines ($x = a$)

- Vertical lines are expressed in the form $x = a$, where a is the x -intercept.
- In vertical lines the x -value remains the same regardless of the y -value.
- This can be seen by looking at a table of values, like the one shown.

x	a	a	a	a
y	-2	0	2	4

- Plotting these points gives a vertical line, as shown in the graph.
- The run of the graph is 0, so using the formula $m = \frac{\text{rise}}{\text{run}}$ involves dividing by zero, which cannot be done. The gradient is said to be **undefined**.



WORKED EXAMPLE 13 Sketching horizontal and vertical lines

On a pair of axes, sketch the graphs of the following and label the point of intersection of the two lines.


a. $x = -3$

b. $y = 4$


THINK

1. The line $x = -3$ is in the form $x = a$.
This is a vertical line.
2. Rule the vertical line where $x = -3$.
Label the graph.

WRITE/DRAW



- b. 1. The line $y = 4$ is in the form $y = c$. This is a horizontal line.
2. Rule the horizontal line where $y = 4$. Label the graph.
3. The lines intersect at $(-3, 4)$.



on Resources



eWorkbook

Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)



Video eLessons

Sketching linear graphs (eles-1919)

Sketching linear graphs using the gradient–intercept method (eles-1920)



Interactivities

Individual pathway interactivity: Sketching linear graphs (int-4504)

The intercept method (int-3840)

The gradient–intercept method (int-3839)

Vertical and horizontal lines (int-6049)

Exercise 6.5 Sketching linear graphs

learn on

6.5 Quick quiz

on

6.5 Exercise

Individual pathways

PRACTISE

1, 3, 5, 7, 9, 11, 15, 20

CONSOLIDATE

4, 8, 12, 13, 16, 17, 21

MASTER

2, 6, 10, 14, 18, 19, 22

Fluency

- WE11** Using the x - and y -intercept method, sketch the graphs of:

a. $5y - 4x = 20$ b. $y = x + 2$ c. $y = -3x + 6$.
- Using the x - and y -intercept method, sketch the graphs of:

a. $3y + 4x = -12$ b. $x - y = 5$ c. $2y + 7x - 8 = 0$.
- WE12** Using the gradient–intercept method, sketch the graphs of:

a. $y = x - 7$ b. $y = 2x + 2$ c. $y = -2x + 2$.
- Using the gradient–intercept method, sketch the graphs of:

a. $y = \frac{1}{2}x - 1$ b. $y = 4 - x$ c. $y = -x - 10$.
- WE13** On a pair of axes, sketch the graphs of the following.

a. $y = 4$ b. $y = -3$

6. On a pair of axes, sketch the graphs of the following.

a. $y = -12.5$

b. $y = \frac{4}{5}$

7. Sketch the graphs of the following.

a. $x = 2$

b. $x = -6$

8. Sketch the graphs of the following.

a. $x = -2.5$

b. $x = \frac{3}{4}$

9. Sketch the graphs of the following.

a. $y = 3x$

b. $y = -2x$

10. Sketch the graphs of the following.

a. $y = \frac{3}{4}x$

b. $y = -\frac{1}{3}x$

Understanding

11. **MC** Select which of the following statements about the rule $y = 4$ is false.

- A. The gradient $m = 0$.
- B. The y -intercept is at $(0, 4)$.
- C. The graph is parallel to the x -axis.
- D. The point $(4, 2)$ lies on this graph.
- E. There is no x -intercept for this graph.

12. **MC** Select which of the following statements is not true about the rule $y = -\frac{3}{5}x$.

- A. The graph passes through the origin.
- B. The gradient $m = -\frac{3}{5}$.
- C. The x -intercept is at $x = 0$.
- D. The graph can be sketched using the x - and the y -intercept method.
- E. The y -intercept is at $y = 0$.

13. $2x + 5y = 20$ is a linear equation in the form $ax + by = c$.

- a. Rearrange this equation into the form $y = mx + c$.
- b. State the gradient.
- c. State the x - and y -intercepts.
- d. Sketch this straight line.

14. A straight line has an x -intercept of -3 and a y -intercept of 5 .

- a. State the gradient.
- b. Draw the graph.
- c. Write the equation in the form:
 - i. $y = mx + c$
 - ii. $ax + by = c$.

Reasoning


15. Consider the relationship $4x - 3y = 24$.

- a. Rewrite this relationship, making y the subject.
- b. Show that the x - and y -intercepts are $(6, 0)$ and $(0, -8)$ respectively.
- c. Sketch a graph of this relationship.

16. Consider the relationship $ax + by = c$.
- Rewrite the relationship, making y the subject.
 - If a , b and c are positive integer values, explain how the gradient is negative.
17. Josie accidentally spilled a drink on her work. Part of her calculations were smudged. The line $y = \frac{1}{2}x + \frac{3}{4}$ was written in the form $ax + 4y = 3$. Show that the value of $a = -2$.
18. Explain why the descriptions 'right 3 up 2', 'right 6 up 4', 'left 3 down 2', 'right $\frac{3}{2}$ up 1' and 'left 1 down $\frac{2}{3}$ ', all describe the same gradient for a straight line.
19. A straight line passes through the points (3, 5) and (6, 11).
- Determine the slope of the line.
 - Determine the equation of the line.
 - State the coordinates of another point that lies on the same line.

Problem solving

20. Consider the following graph.



- Match the descriptions given below with their corresponding line.
 - Straight line with a y -intercept of (0, 1) and a positive gradient.
 - Straight line with a gradient of $1\frac{1}{2}$.
 - Straight line with a gradient of -1 .
 - Write a description for the unmatched graph.
21. a. Sketch the linear equation $y = -\frac{5}{7}x - \frac{3}{4}$:
- using the y -intercept and the gradient
 - using the x - and y -intercepts
 - using two other points.
- b. Compare and contrast the methods and generate a list of advantages and disadvantages for each method used in part a. Explain which method you think is best. Give your reasons.
22. Consider these two linear graphs.

$$y - ax = b \text{ and } y - cx = d.$$

Show that if these two graphs intersect where both x and y are positive, then $a > c$ when $d > b$.

LESSON

6.6 Technology and linear graphs

LEARNING INTENTION


At the end of this lesson you should be able to:

- produce linear graphs using digital technologies.

6.6.1 Graphing with technology


eles-4789

- There are many digital technologies that can be used to graph linear relationships.
- The Desmos Graphing Calculator is a free graphing tool that can be found online.
- Other commonly used digital technologies include Microsoft Excel and graphing calculators.
- Digital technologies can help identify important features and patterns in graphs.



Sketching a linear graph using technology


- Depending on the choice of digital technology used, the steps involved to produce a linear graph may vary slightly.
- Most graphing calculators have an entry (or input) line to type in the equation of the line you wish to sketch.
- When using the Desmos Graphing Calculator you can simply type $y = x + 1$ into the input box to produce its graph.
- When using technologies such as a Texas Instruments or CASIO graphing calculator you will need to open a Graphs page first. The entry line for these types of technologies has a template that needs to be followed carefully. Both of these graphing calculators begin with $f1(x) =$ or $y1 =$, which is effectively saying $y =$.
- Thus, to draw the graph of $y = x + 1$, you would simply enter ' $x + 1$ '.
- The screen shows the graph of $y = x + 1$ sketched on a TI-Nspire CAS calculator.



6.6.2 Graphing parallel lines

eles-4790


- Lines with the same gradient are called **parallel lines**. For a pair of parallel lines, $m_1 = m_2$.
- Digital technologies can be used to help visualise this concept. For example, $y = 3x + 1$, $y = 3x - 4$ and $y = 3x$ are all parallel lines, because $m = 3$.
- Select the digital technology of your choice and sketch these three parallel lines on the same set of axes.



6.6.3 Graphing perpendicular lines

eles-4791

- Lines that meet at right angles are called **perpendicular lines**.
- The product of the gradients of two perpendicular lines is equal to -1 .
- For a pair of perpendicular lines, $m_1 \times m_2 = -1$.
- Digital technologies can be used to help visualise this concept. For example, $y = 2x + 1$ and $y = -\frac{1}{2}x + 6$ are perpendicular, because $2 \times -\frac{1}{2} = -1$.
- Select a digital technology of your choice and sketch these two lines on the same set of axes.



Resources



eWorkbook Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)



Interactivities Individual pathway interactivity: Technology and linear graphs (int-4505)
Parallel lines (int-3841)

Exercise 6.6 Technology and linear graphs

learn **on**

6.6 Quick quiz **on**

6.6 Exercise

Individual pathways

PRACTISE

1, 3, 4, 11, 15, 16

CONSOLIDATE

2, 5, 6, 7, 12, 14, 17

MASTER

8, 9, 10, 13, 18

Understanding

Use technology wherever possible to answer the following questions.

1. Sketch the following graphs on the same Cartesian plane.

a. $y = x$

b. $y = 2x$

c. $y = 3x$

- Describe what happens to the steepness of the graph (the gradient of the line) as the coefficient of x increases in value.
- Identify where each graph cuts the x -axis (its x -intercept).
- Identify where each graph cuts the y -axis (its y -intercept).

2. Sketch the following graphs on the same Cartesian plane.

a. $y = x$

b. $y = \frac{1}{2}x$

c. $y = \frac{1}{3}x$

- Describe what happens to the steepness of the graph (the gradient of the line) as the coefficient of x decreases in value.
- Identify where each graph cuts the x -axis (its x -intercept).
- Identify where each graph cuts the y -axis (its y -intercept).

3. Sketch the following graphs on the same Cartesian plane.
- a. $y = -x$ b. $y = -2x$ c. $y = -3x$
- Describe what happens to the steepness of the graph as the magnitude of the coefficient of x decreases in value (becomes more negative).
 - Identify where each graph cuts the x -axis (its x -intercept).
 - Identify where each graph cuts the y -axis (its y -intercept).
4. Identify the correct word or words from the options given and rewrite the following sentences using the correct option.
- If the coefficient of x is (**positive/negative**), then the graph will have an upward slope to the right. That is, the gradient of the graph is (**positive/negative**).
 - If the coefficient of x is negative, then the graph will have a (**downward/upward**) slope to the right. That is, the gradient of the graph is (**positive/negative**).
 - The bigger the magnitude of the coefficient of x (more positive or more negative), the (**bigger/smaller**) the steepness of the graph.
 - If there is no constant term in the equation, the graph (**will/will not**) pass through the origin.
5. Sketch the following graphs on the same Cartesian plane and answer the following questions for each graph.
- Is the coefficient of x the same for each graph? If so, state the coefficient.
 - State whether the steepness (gradient) of each graph differs.
 - Identify where each graph cuts the x -axis (its x -intercept).
 - Identify where each graph cuts the y -axis (its y -intercept).
- i. $y = x$ ii. $y = x + 2$ iii. $y = x - 2$
6. Sketch the following graphs on the same Cartesian plane and answer the following questions for each graph.
- Is the coefficient of x the same for each graph? If so, state the coefficient.
 - State whether the steepness (gradient) of each graph differs.
 - Identify where each graph cuts the x -axis (its x -intercept).
 - Identify where each graph cuts the y -axis (its y -intercept).
- i. $y = -x$ ii. $y = -x + 2$ iii. $y = -x - 2$
7. Identify the correct word or words from the options given and rewrite the following sentences using the correct option.
- For a given set of linear graphs, if the coefficient of x is (**the same/different**), the graphs will be parallel.
 - The constant term in the equation is the (**y -intercept/ x -intercept**) or where the graph cuts the (**y -axis/ x -axis**).
 - The (**y -intercept/ x -intercept**) can be found by substituting $x = 0$ into the equation.
 - The (**y -intercept/ x -intercept**) can be found by substituting $y = 0$ into the equation.
8. On the same Cartesian plane, sketch the following graphs.
- a. $y = x + 5$ b. $y = -x + 5$
c. $y = 3x + 5$ d. $y = -\frac{2}{5}x + 5$
- Is the coefficient of x the same for each graph? If so, state the coefficient.
 - State whether the steepness (gradient) of each graph differ.
 - Write down the gradient of each linear graph.
 - Identify where each graph cuts the x -axis (its x -intercept).
 - Identify where each graph cuts the y -axis (its y -intercept).

9. Identify the correct word or words from the options given and rewrite the following sentences using the correct option.
- One of the general forms of the equation of a linear graph is $y = mx + c$, where m is the (**steepness/x-coordinate**) of the graph. We call the steepness of the graph the *gradient*.
 - The value of c is the (**x-coordinate/y-coordinate**) where the graph cuts the (**x-axis/y-axis**).
 - All linear graphs with the (**same/different**) gradient are (**parallel/perpendicular**).
 - All linear graphs that have the same y-intercept pass through (**the same/different**) point on the y-axis.
10. For each of the following lines, identify the gradient and the y-intercept.
- $y = 2x$
 - $y = x + 1$
 - $y = -3x + 5$
 - $y = \frac{2}{3}x - 7$

Reasoning

11. Using a CAS calculator, graph the lines $y = 3(x - 1) + 5$, $y = 2(x - 1) + 5$ and $y = -\frac{1}{2}(x - 1) + 5$. Describe what they have in common and how they differ from each other.
12. Show on a CAS calculator how $y = \frac{1}{2}x - \frac{5}{3}$ can be written as $6y = 3x - 10$.
13. A phone company charges \$2.20 for international calls of 1 minute or less and \$0.55 for each additional minute. Using a CAS calculator, graph the cost for calls that last for whole numbers of minutes. Explain all the important values needed to sketch the graph.
14. Shirly walks dogs after school for extra pocket money. She finds that she can use the equation $P = -15 + 10N$ to calculate her profit (in dollars) each week.
- Explain the real-world meaning of the numbers -15 and 10 and the variable N .
 - What is the minimum number of dogs that Shirly must walk in order to earn a profit?
 - Using a CAS calculator, sketch the equation.
15. Graph $y = 0.2x + 3.71$ on a CAS calculator. Explain how to use the calculator to find an approximate value when $x = 70.3$.

Problem solving

16. Plot the points $(6, 3.5)$ and $(-1, -10.5)$ using a CAS calculator and:
- determine the equation of the line
 - sketch the graph, showing x - and y -intercepts
 - calculate the value of y when $x = 8$
 - calculate the value of x when $y = 12$.
17. A school investigating the price of a site licence for their computer network found that it would cost \$1750 for 30 computers and \$2500 for 60 computers.
- Using a CAS calculator, find a linear equation that represents the cost of a site licence in terms of the number of computers in the school.
 - Determine the y -intercept of the linear equation and explain how it relates to the cost of a site licence.
 - Calculate the cost for 200 computers.
 - Evaluate how many computers you could connect for \$3000.
18. Dylan starts his exercise routine by jogging to the gym, which burns 325 calories. He then pedals a stationary bike burning 3.8 calories a minute.
- Graph this data using a CAS calculator.
 - Evaluate how many calories Dylan has burnt after 15 minutes of pedalling.
 - Evaluate how long it took Dylan to burn a total of 450 calories.



LESSON

6.7 Practical applications of linear graphs

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- determine the linear rule from a table of values
- determine the linear rule that models a real-life problem
- interpret linear rules in real-life problems.


6.7.1 Determining a linear rule from a table of values

eles-4792

- If two **variables** are linked by a linear rule, then as one variable increases, the other increases, or decreases, at a steady rate.
- The table shown gives an example of a linear relationship.

x	0	1	2	3
y	5	8	11	14


- Each time x increases by 1, y increases by 3.
- The linear rule connecting x and y is $y = 3x + 5$.



- Consider the relationship depicted in the table shown.

x	0	1	2	3
y	7	5	3	1

- Each time x increases by 1, y decreases by 2.
- The linear rule in this case is $y = -2x + 7$.



WORKED EXAMPLE 14 Determining a rule from a table of values

Determine the rule connecting x and y in each of the following tables.

a.

x	0	1	2	3
y	-3	2	7	12

b.

x	3	4	5	6
y	12	11	10	9

THINK

1. y increases at a steady rate, so this is a linear relationship.
Write the rule.
2. To calculate m : y increases by 5 each time x increases by 1.
Write the value of the gradient.
3. To determine c : From the table, when $x = 0$, $y = -3$.
Write the value of the y -intercept.
4. Write the rule.

WRITE

a. $y = mx + c$

$$m = 5$$

$$c = -3$$

$$y = 5x - 3$$

- b. 1. y decreases at a steady rate, so this is a linear relationship.
Write the rule.
2. To calculate m : y decreases by 1 each time x increases by 1.
Write the value of the gradient.
3. To determine c : From the table, when $x = 3$, $y = 12$.
To calculate the y -intercept, substitute the x - and y -values of one of the points, and solve for c .
4. Write the rule.

$$b. y = mx + c$$

$$m = -1$$

$$(3, 12): \quad y = -x + c$$

$$12 = -(3) + c$$

$$12 = -3 + c$$

$$c = 15$$

$$y = -x + 15$$



eles-4793

6.7.2 Modelling linear relationships

- Relationships between real-life variables are often modelled by a mathematical equation. In other words, an equation or formula is used to link the two variables.
For example:
 - $A = l^2$ represents the relationship between the area and the side length of a square
 - $C = \pi d$ represents the relationship between the circumference and the diameter of a circle.
- If one variable changes at a constant rate compared to the other, then the two variables have a linear relationship.

WORKED EXAMPLE 15 Determining the rule to model a relationship in a worded question

An online bookstore sells a certain textbook for \$21 and charges \$10 for delivery, regardless of the number of books being delivered.

- Determine the rule connecting the cost (\$ C) with the number of copies of the textbook delivered (n).
- Use the rule to calculate the cost of delivering 35 copies of the textbook.
- Calculate how many copies of the textbook can be delivered for \$1000.

THINK

- Set up a table.
Cost for 1 copy = $21 + 10$
Cost for 2 copies = $2(21) + 10$
= 52
- The cost rises steadily, so there is a linear relationship. Write the rule.
- To calculate the gradient, use the formula
 $m = \frac{y_2 - y_1}{x_2 - x_1}$ with the points (1, 31) and (2, 52).
- To determine the value of c , substitute $C = 31$ and $n = 1$.
- Write the rule.

WRITE

a.

n	1	2	3
C	31	52	73

$$C = mn + c$$

$$m = \frac{52 - 31}{2 - 1}$$

$$= 21$$

$$C = 21n + c$$

$$(1, 31): 31 = 21(1) + c$$

$$c = 10$$

$$C = 21n + 10$$

b. 1. Substitute $n = 35$ and calculate the value of C .

2. Write the answer.

c. 1. Substitute $C = 1000$ and calculate the value of n .

2. You cannot buy 47.14 books, so round down.
Write the answer.

$$\begin{aligned} \text{b. } C &= 21(35) + 10 \\ &= 735 + 10 \\ &= 745 \end{aligned}$$

The cost including delivery for 35 copies is \$745.

$$\begin{aligned} \text{c. } 1000 &= 21n + 10 \\ 21n &= 990 \\ n &= 47.14 \end{aligned}$$

For \$1000, 47 copies of the textbook can be bought and delivered.

- In Worked example 15, compare the rule $C = 21n + 10$ with the original question. It is clear that the $21n$ refers to the cost of the textbooks (a variable amount, depending on the number of copies) and that 10 refers to the fixed (constant) delivery charge.
- In this case C is called the **response variable** or the **dependent variable**, because it depends on the number of copies of the textbooks (n).
- The variable n is called the **explanatory variable** or the **independent variable** because it is the variable that may explain the changes in the response variable.
- When graphing numerical data, the response variable is plotted on the vertical axis and the explanatory variable is plotted on the horizontal axis.

on Resources



eWorkbook Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)



Interactivities Individual pathway interactivity: Practical applications of linear graphs (int-4507)
Dependent and independent variables (int-6050)

Exercise 6.7 Practical applications of linear graphs

learn on

6.7 Quick quiz

6.7 Exercise

Individual pathways

PRACTISE

1, 3, 5, 6, 11, 16

CONSOLIDATE

2, 7, 9, 12, 13, 17

MASTER

4, 8, 10, 14, 15, 18

Fluency

1. **WE14** Determine the linear rule linking the variables in each of the following tables.

a.

x	0	1	2	3
y	-5	1	7	13

b.

x	0	1	2	3
y	8	5	2	-1

2. Determine the linear rule linking the variables in each of the following tables.

a.

x	0	1	2	3
y	4	6	8	10

b.

x	0	1	2	3
y	1.1	2.0	2.9	3.8

3. Determine the linear rule linking the variables in each of the following tables.

a.

x	2	3	4	5
y	7	10	13	16

b.

x	5	6	7	8
y	12	11	10	9

4. Determine the linear rule linking the variables in each of the following tables.

a.

t	3	4	5	6
v	18	15	12	9

b.

d	1	2	3	4
C	11	14	17	20

5. **MC** Sasha and Fiame hire a car. They are charged a fixed fee of \$150 for hiring the car and then \$25 per day. They hire the car for d days.

Select which one of the following rules describes the number of days the car is hired and the total cost, C , they would be charged for that number of days charged.

A. $C = 25d$

B. $C = 150d$

C. $C = 175d$

D. $C = 25d + 150$

E. $C = 150d + 25$

Understanding

6. **WE15** Fady's bank balance has increased in a linear manner since he started his part-time job. After 20 weeks of work his bank balance was at \$560 and after 21 weeks of work it was at \$585.

- Determine the rule that relates the size of Fady's bank balance, A , and the time (in weeks) worked, t .
- Use the rule to calculate the amount in Fady's account after 200 weeks.
- Use the rule to identify the initial amount in Fady's account.



7. The cost of making a shoe increases as the size of the shoe increases. It costs \$5.30 to make a size 6 shoe, and \$6.40 to make a size 8 shoe. Assuming that a linear relationship exists:

- determine the rule relating cost (C) to shoe size (s)
- calculate much it costs to produce a size 12 shoe.

8. The number of books in a library (N) increases steadily with time (t). After 10 years there are 7200 publications in the library, and after 12 years there are 8000 publications.

- Determine the rule predicting the number of books in the library.
- Calculate how many books were there after 5.5 years.
- Calculate how many books there will be after 25 years.

9. A skyscraper can be built at a rate of 4.5 storeys per month.

- Calculate how many storeys will be built after 6 months.
- Calculate how many storeys will be built after 24 months.


10. The Nguyens' water tank sprang a leak and has been losing water at a steady rate. Four days after the leak occurred, the tank contained 552 L of water, and ten days later it held only 312 L.

- Determine the rule linking the amount of water in the tank (w) and the number of days (t) since the leak occurred.
- Calculate the amount of water that was initially in the tank.
- If water loss continues at the same rate, determine when the tank will be empty.



Reasoning

11. The pressure inside a boiler increases steadily as the temperature increases. For each 1°C increase in temperature, the pressure increases by 10 units, and at a temperature of 100°C the pressure is 1200 units. If the maximum pressure allowed is 2000 units, show that the temperature cannot exceed 180°C .
12. After 11 pm a taxi company charges a \$3.50 flag fall plus \$2.57 for each kilometre travelled.
 - a. Determine the linear rule connecting the cost, C , and the distance travelled, d .
 - b. Calculate how much an 11.5 km trip will cost.
 - c. If you have \$22 in your pocket, calculate how far you can afford to travel, correct to 1 decimal place.
13. A certain kind of eucalyptus tree grows at a linear rate for its first 2 years of growth. If the growth rate is 5 cm per month, show that the tree will be 1.07 m tall after 21.4 months.
14. A software company claims that its staff can fix 22 bugs each month. They are working on a project to fix a program that started out with 164 bugs.
 - a. Determine the linear rule connecting the number of bugs left, N , and the time in months, t , from the beginning of the project.
 - b. Calculate how many bugs will be left after 2 months.
 - c. Determine how long it will be until there are only 54 bugs left.
 - d. Determine how long it will take to eliminate all of the bugs. Justify your answer.
15. Michael produces and sells prints of his art at a local gallery. For each print run his profit (P) is given by the equation $P = 200n - 800$, where n is the number of prints sold.
 - a. Sketch the graph of this rule.
 - b. Identify the y -intercept. Determine what it represents in this situation.
 - c. Identify the x -intercept. Determine what it represents in this example.
 - d. Identify the gradient of the graph. Determine what this means in this situation.



Problem solving

16. The cost of a taxi ride is \$3.50 flag fall plus \$2.14 for each kilometre travelled.
 - a. Determine the linear rule connecting the cost, C , and the distance travelled, d .
 - b. Calculate how much an 11.5 km trip will cost.
 - c. Calculate how much a 23.1 km trip will cost.
17. Theo is going on holiday to Japan. One yen (¥) buys 0.0127 Australian dollars (A\$).
 - a. Write an equation that converts Australian dollars to Japanese yen (¥), where A represents amount of Australian dollars and Y represents amount of yen.
 - b. Using the equation from part a, calculate how many yen Theo will receive in exchange if they have A\$2500.
 - c. There is a commission to be paid on exchanging currency. Theo needs to pay 2.8% for each Australian dollar they exchange into yen. Write down an equation that calculates the total amount of yen Theo will receive. Write your equation in terms of Y_T (total amount of yen) and $\text{\$}A$ (Australian dollars).
18. Burchill and Cody need to make a journey to the other branch of their store across town. The traffic is very busy at this time of the day, so Burchill catches a train that travels halfway, then walks the rest of the way. Cody travels by bike the whole way. The bike path travels along the train line and then follows the road to the other branch of their store. Cody's bike travels twice as fast as Burchill's walking speed, and the train travels 4 times faster than Cody's bike. Evaluate who arrives at the destination first.

LESSON

6.8 Midpoint of a line segment and distance between two points

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- calculate the midpoint of a line segment
- calculate the distance between two points.



6.8.1 Calculating the midpoint of a line segment

eles-4794

- The x - and y -coordinates of the **midpoint** are halfway between those of the end points of a line segment.
- The coordinates of the midpoint, M , of a line can be found by averaging the x - and y -coordinates of the end points.



The midpoint of a line segment

The midpoint, M , of the line segment joining the points (x_1, y_1) and (x_2, y_2) is:

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

WORKED EXAMPLE 16 Calculating the midpoint of a line segment

Calculate the midpoint of the line segment joining $(5, 9)$ and $(-3, 11)$.

THINK

1. Average the x -values: $\frac{x_1 + x_2}{2}$.

2. Average the y -values: $\frac{y_1 + y_2}{2}$.

3. Write the answer.

WRITE

$$x = \frac{5 - 3}{2}$$

$$= \frac{2}{2}$$

$$= 1$$

$$y = \frac{9 + 11}{2}$$

$$= \frac{20}{2}$$

$$= 10$$

The midpoint is $(1, 10)$.

TI | THINK

- On a Graphs page, set an appropriate window as shown and press:
 - MENU
 - 8: Geometry
 - Points & Lines
 - 3: Point by Coordinates

Type the coordinates of one point on the keypad, pressing ENTER after each number.

Repeat the above step to plot the second point.

- To determine the midpoint, press:

- MENU
- 8: Geometry
- 4: Construction
- 5: Midpoint.

Move the cursor to one of the points and press ENTER.

Move the cursor to the other point and press ENTER. This will place a dot on the screen representing the midpoint of the two plotted points.



To display the midpoint's coordinates, press:

- MENU
- 1: Actions
- 8: Coordinates and Equations

Move the cursor to the midpoint and press ENTER twice.

- State the coordinates of the midpoint.

WRITE



The coordinates of the midpoint are (1, 10).

CASIO | THINK

On a Geometry screen, select a pair of axes with scale and grid.

This is achieved by continuous clicking on the graph icon.

To get the desired window, select:

- View
- Zoom out

Using the Line segment tool, place the pointer at (−3, 11) and hold to place the second point at (5, 9). The line segment will be automatically labelled.

To locate the midpoint:



- Click on the line segment.
- Click again for the point C to be placed in the centre or midpoint.

To determine the coordinates of the midpoint C:

- Click on the point C.
- Use the measuring tool at the top right to display the coordinates.

State the coordinates of the midpoint.

WRITE



The coordinates of the midpoint are (1, 10).

WORKED EXAMPLE 17 Determining the coordinates of a point given the midpoint and another point of an interval

M(7, 2) is the midpoint of the line segment AB. If the coordinates of A are (1, −4), determine the coordinates of B.

THINK

- Let B have the coordinates (x, y).
- The midpoint is (7, 2), so the average of the x-values is 7. Solve for x.

WRITE

A(1, −4), B(x, y), M(7, 2)

$$\frac{1+x}{2} = 7$$

$$1+x = 14$$

$$x = 13$$

3. The average of the y -values is 2.
Solve for y .

$$\frac{-4 + y}{2} = 2$$

$$\begin{aligned} -4 + y &= 4 \\ y &= 8 \end{aligned}$$

4. Write the answer.

The coordinates of point B are (13, 8).



6.8.2 Calculating the distance between two points


eles-4795

- The distance between two points on the Cartesian plane is calculated using Pythagoras' theorem applied to a right-angled triangle.

Distance between two points

The distance, d , of the line segment joining the points (x_1, y_1) and (x_2, y_2) is:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$



WORKED EXAMPLE 18 Calculating the distance between two points

Calculate the distance between the points $(-1, 3)$ and $(4, 5)$:


a. in exact form

b. correct to 3 decimal places.

THINK

- a. 1. Draw a diagram showing the right-angled triangle (optional).

WRITE/DRAW



2. Write the formula for the distance between two points.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

3. Let $(x_1, y_1) = (-1, 3)$ and $(x_2, y_2) = (4, 5)$. Substitute the x - and y -values into the equation.

$$d = \sqrt{(4 - (-1))^2 + (5 - 3)^2}$$

4. Simplify.

$$d = \sqrt{5^2 + 2^2}$$

$$d = \sqrt{25 + 4}$$

$$d = \sqrt{29}$$

5. Write the answer.

The exact distance between points $(-1, 3)$ and $(4, 5)$ is $\sqrt{29}$.


- b. 1. Write $\sqrt{29}$ as a decimal to 5 decimal places. b. $\sqrt{29} = 5.38516$
 2. Round the answer correct to 3 decimal places. $d \approx 5.385$

TI | THINK

a-b. In a new problem, on a Calculator page, complete the entry lines as:
 $x1 := -1$
 $y1 := 3$
 $x2 := 4$
 $y2 := 5$
 $d := \sqrt{(x2 - x1)^2 + (y2 - y1)^2}$
 Press ENTER after each entry.
 Press CTRL ENTER to get a decimal approximation.

WRITE

a-b.




The distance is $\sqrt{29} \approx 5.385$ to 3 decimal places.

CASIO | THINK

a. Make sure the calculator is set to Standard mode. On the Main screen, complete the entry line as:
 $\sqrt{(4 - (-1))^2 + (5 - 3)^2}$
 Press EXE.

WRITE



b. Convert the answer to a decimal by tapping Standard and changing it to Decimal. Then press EXE.

- a. $\sqrt{29}$
 b. 5.385

on Resources

- eWorkbook** Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)
- Interactivities** Individual pathway interactivity: Midpoint of a line segment and distance between two points (int-4508)
 Distance between two points (int-6051)
 Midpoints (int-6052)

Exercise 6.8 Midpoint of a line segment and distance between two points

6.8 Quick quiz **on**

6.8 Exercise

Individual pathways

PRACTISE

1, 4, 8, 11, 13, 16, 20

CONSOLIDATE

2, 5, 9, 12, 14, 17, 21

MASTER



3, 6, 7, 10, 15, 18, 19, 22

Fluency



- WE16** Calculate the midpoint of the line segment joining each of the following pairs of points.
 - (1, 3) and (3, 5)
 - (6, 4) and (4, -2)
 - (2, 3) and (12, 1)
- Calculate the midpoint of the line segment joining each of the following pairs of points.
 - (6, 3) and (10, 15)
 - (4, 2) and (-4, 8)
 - (0, -5) and (-2, 9)
- Calculate the midpoint of the line segment joining each of the following pairs of points.
 - (8, 2) and (-18, -6)
 - (-3, -5) and (7, 11)
 - (-8, -3) and (8, 27)
- Calculate the midpoint of the segment joining each of the following pairs of points.
 - (7, -2) and (-4, 13)
 - (0, 22) and (-6, -29)
 - (-15, 8) and (-4, 11)
 - (-3, 40) and (0, -27)
- WE17** Determine the value of a in each series of points so that the point M is the midpoint of the line segment joining points A and B .
 - $A(-2, a), B(-6, 5), M(-4, 5)$
 - $A(a, 0), B(7, 3), M\left(8, \frac{3}{2}\right)$
 - $A(3, 3), B(4, a), M\left(3\frac{1}{2}, -6\frac{1}{2}\right)$
 - $A(-4, 4), B(a, 0), M(-2, 2)$
- M is the midpoint of the line interval AB . Determine the coordinates of B if:
 - $A = (0, 0)$ and $M = (2, 3)$
 - $A = (2, 3)$ and $M = (0, 0)$
 - $A = (-3, 2)$ and $M = (4, 2)$
 - $A = (3, -1)$ and $M = (-2, -2)$.
- Determine the equation of a line that has a gradient of 5 and passes through the midpoint of the line segment joining $(-1, -7)$ and $(3, 3)$.
- WE18** Calculate the distance between each of the following pairs of points.
 - (4, 5) and (1, 1)
 - (7, 14) and (15, 8)
 - (2, 4) and (2, 3)
- Calculate the distance between each of the following pairs of points.
 - (12, 8) and (10, 8)
 - (14, 9) and (2, 14)
 - (5, -13) and (-3, -7)
- Calculate the distance between each of the following pairs of points.
 - (-14, -9) and (-10, -6)
 - (0, 1) and (-15, 9)
 - (-4, -8) and (1, 4)
- Calculate the distance between the following pairs of points, correct to 3 decimal places.
 - (-14, 10) and (-8, 14)
 - (6, -7) and (13, 6)
 - (-11, 1) and (2, 2)
- Calculate the distance between the following pairs of points, correct to 3 decimal places.
 - (9, 0) and (5, -8)
 - (2, -7) and (-2, 12)
 - (9, 4) and (-10, 0)

Understanding


13. Calculate the perimeter of each figure shown, giving your answers correct to 3 decimal places.



14. Calculate the perimeter of each triangle shown, giving your answers correct to 3 decimal places.



15. Two hikers are about to hike from A to B (shown on the map). Calculate the straight-line distance from A to B.



Reasoning

16. Show that the distance between the points A (2, 2) and B (6, -1) is 5.
17. The point M (-2, -4) is the midpoint of the interval AB. Show that the point B is (-9, -2), given A is (5, -6).
18. Show that the point B (6, -10) is equidistant from the points A (15, 3) and C (-7, -1).
19. Answer the following questions.
 - a. Plot the following points on a Cartesian plane: A (-1, -4), B (2, 3), C (-3, 8) and D (4, -5).
 - b. Show that the midpoint of the interval AC is (-2, 2).
 - c. Calculate the exact distance between the points A and C.
 - d. If B is the midpoint of an interval CM, determine the coordinates of point M.
 - e. Show that the gradient of the line segment AB is $\frac{7}{3}$.
 - f. Determine the equation of the line that passes through points B and D.

Problem solving

20. Explain what type of triangle $\triangle ABC$ is if it has vertices A (-4, 1), B (2, 3) and C (0, -3).
21. Calculate the gradient of the line through the points (-1, 3) and (3 + 4t, 5 + 2t).
22. A map of a town drawn on a Cartesian plane shows the main street extending from (-4, 5) to (0, -7). There are five streetlights positioned in the street. There is one streetlight at either end, and three streetlights spaced evenly down the street.
Give the position of the five lights in the street.

LESSON

6.9 Non-linear relations

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- recognise and sketch a basic parabola from its equation
 - sketch a parabola that has been translated horizontally or vertically
 - recognise and use digital technology to sketch hyperbolas, cubics and exponentials
 - recognise and sketch a circle from its equation.
- There are many examples of non-linear relationships in mathematics. Some of them are the parabola, the hyperbola and the circle.
- Note:* This lesson is an introduction to non-linear relations — parabolas will be discussed in more depth in topic 12.

6.9.1 The parabola

eles-4796

- The **parabola** is a curve that is often found in nature. It is also commonly used in architecture and engineering.
- The simplest parabola has the equation $y = x^2$.
- The key features of the parabola $y = x^2$ are:
 - it has an **axis of symmetry**, which is the y -axis
 - the graph is concave up (it opens upwards)
 - it has a minimum **turning point**, or **vertex**, at $(0, 0)$.
- All parabolas have the same basic shape; however, they can be wider or narrower depending on their equation, or can be translated vertically or horizontally or reflected.
- To draw a parabola we can create a table of values and plot points, or use calculator technology.



WORKED EXAMPLE 19 Graphing a parabola

Plot the graph of $y = x^2$ for values of x from -3 to 3 . State the equation of the axis of symmetry and the coordinates of the turning point.


THINK

1. Write the equation.
2. Produce a table of values using x -values from -3 to 3 .
3. Draw a set of clearly labelled axes. Plot the points and join them with a smooth curve. The scale on the y -axis would be from -2 to 10 and from -4 to 4 on the x -axis.

WRITE/DRAW

$$y = x^2$$

x	-3	-2	-1	0	1	2	3
y	9	4	1	0	1	4	9



- Write the equation of the line that divides the parabola exactly in half.
- Write the coordinates of the turning point.


The equation of the axis of symmetry is $x = 0$.

The turning point is $(0, 0)$.

TI | THINK


- Open a Graphs page, and press:
 - MENU
 - 4: Window/Zoom
 - 1: Window Settings
 Complete the entry lines as shown. Press OK.

WRITE



- To sketch the graph, complete the entry line as:


$$f1(x) = x^2$$
 Press ENTER. The graph will be displayed.



CASIO | THINK


Open a Graph & Table screen and select the Window icon. Complete the entry lines as shown. Press OK.

WRITE



To sketch the graph, complete the entry line as:


$$y1 = x^2$$
 Press the Graph icon. The graph will be displayed.



- The parabola $y = x^2$ can be transformed into other parabolas using transformations, such as:
 - vertical translations
 - horizontal translations
 - reflections.
- Other parabolas can be sketched using their key features, such as the axis of symmetry and vertex.

Vertical translations

- Parabolas can be **translated** vertically by adding a number to or subtracting a number from the right-hand side of the equation.
- When a number is added to the equation the parabola is translated vertically upwards.
- When a number is subtracted from the equation the parabola is translated vertically downwards.
- The graph shown displays the parabolas $y = x^2$, $y = x^2 + 2$ and $y = x^2 - 1$ on the same set of axes.
- $y = x^2 + 2$ has been translated 2 units upwards. The turning point is now $(0, 2)$.
- $y = x^2 - 1$ has been translated 1 unit downwards. The turning point is now $(0, -1)$.



WORKED EXAMPLE 20 Stating vertical translations of parabolas

State the vertical translation (when compared with the graph of $y = x^2$) and the coordinates of the turning point for the graphs of each of the following equations.

a. $y = x^2 + 5$

b. $y = x^2 - 4$

THINK

- a. 1. Write the equation.
2. $+5$ means the graph is translated upwards 5 units.
3. Translate the turning point of $y = x^2$, which is $(0, 0)$.
The x -coordinate of the turning point remains 0 and the y -coordinate has 5 added to it.
- b. 1. Write the equation.
2. -4 means the graph is translated downwards 4 units.
3. Translate the turning point of $y = x^2$, which is $(0, 0)$.
The x -coordinate of the turning point remains 0 and the y -coordinate has 4 subtracted from it.

WRITE

a. $y = x^2 + 5$


Vertical translation of 5 units up
The turning point becomes $(0, 5)$.

b. $y = x^2 - 4$

Vertical translation of 4 units down
The turning point becomes $(0, -4)$.

Horizontal translations

- Parabolas can be translated horizontally by placing x in brackets and adding a number to it or subtracting a number from it.
- When a number is added to x , the parabola is translated to the left.
- When a number is subtracted from x , the parabola is translated to the right.
- The graphs shown displays the parabolas $y = x^2$, $y = (x + 1)^2$ and $y = (x - 2)^2$ on the same set of axes.
- $y = (x + 1)^2$ has been translated 1 unit left. The turning point is now $(-1, 0)$.
- $y = (x - 2)^2$ has been translated 2 units right. The turning point is now $(2, 0)$.



WORKED EXAMPLE 21 Stating horizontal translations of parabolas

State the horizontal translation (when compared to the graph of $y = x^2$) and the coordinates of the turning point for the graphs of each of the following equations.

a. $y = (x - 3)^2$

b. $y = (x + 2)^2$

THINK

- a. 1. Write the equation.
2. -3 means the graph is translated to the right 3 units.
3. Translate the turning point of $y = x^2$, which is $(0, 0)$.
The y -coordinate of the turning point remains 0, and the x -coordinate has 3 added to it.

WRITE

a. $y = (x - 3)^2$


Horizontal translation of 3 units right
The turning point becomes $(3, 0)$.

- b. 1. Write the equation.
 2. +2 means the graph is translated to the left 2 units.
 3. Translate the turning point of $y = x^2$, which is $(0, 0)$.
 The y -coordinate of the turning point remains 0, and the x -coordinate has 2 subtracted from it.

- b. $y = (x + 2)^2$
 Horizontal translation of 2 units left
 The turning point becomes $(-2, 0)$.

Reflection

- Compare the graph of $y = -x^2$ with that of $y = x^2$.
 - In each case the axis of symmetry is the line $x = 0$ and the turning point is $(0, 0)$.
 - The only difference between the equations is the $-$ sign in $y = -x^2$.
 - The difference between the graphs is that $y = x^2$ 'sits' on the x -axis and $y = -x^2$ 'hangs' from the x -axis (one is a reflection, or mirror image, of the other).
 - $y = x^2$ has a minimum turning point and $y = -x^2$ has a maximum turning point.
- Any quadratic graph for which x^2 is positive has a U shape and is said to be concave up.
- Conversely, if x^2 is negative the graph has a \cap shape and is said to be concave down.



6.9.2 The hyperbola

eles-4797

- A hyperbola is a non-linear graph for which the equation is $y = \frac{1}{x}$ (or $xy = 1$).

WORKED EXAMPLE 22 Graphing a hyperbola

Complete the table of values below and use it to plot the graph of $y = \frac{1}{x}$.

x	-3	-2	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	2	3
y								


THINK

- Substitute each x -value into the function $y = \frac{1}{x}$ to obtain the corresponding y -value.

WRITE/DRAW

x	-3	-2	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	2	3
y	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	-2	undefined	2	1	$\frac{1}{2}$	$\frac{1}{3}$

2. Draw a set of axes and plot the point from the table. Join them with a smooth curve.



- The graph in Worked example 22 has several key features.
 - There are no values when $x = 0$ or $y = 0$.
 - The curve gets closer to the axes but never touches the x - or y -axes. We say the axes are horizontal and vertical **asymptotes**.
 - The hyperbola has two separate branches. It cannot be drawn without lifting your pen from the paper. We say it is a discontinuous graph.
 - All hyperbolas have the same basic shape but they can be wider or narrower depending on the equation.

6.9.3 The cubic

- A cubic is a non-linear graph for which the equation is $y = x^3$

WORKED EXAMPLE 23 Graphing a cubic

Complete the table of values below and use it to plot $y = x^3$.


x	-3	-2	-1	0	1	2	3
y							

THINK

- Substitute each x -value into the equation $y = x^3$ to obtain the corresponding y -value.
- Draw a set of axes and plot the points from the table. Join them with a smooth curve.


WRITE/DRAW

x	-3	-2	-1	0	1	2	3
y	-27	-8	-1	0	1	8	27



TI | THINK

In a new problem, on a Graphs page, complete the function entry line as:
 $f1(x) = x^3$
 Then press ENTER.

DISPLAY/WRITE**CASIO | THINK**

On the Main screen complete the entry line as:
 $y = x^3$ and then press the Graph icon to bring up the graphing screen.
 Highlight the equation and drag it down to the graphing screen.

DISPLAY/WRITE

- A key feature of the graph in Worked example 23 is the **point of inflection**.
 - The point of inflection is where the graph changes from concave up to concave down.
 - It appears as a horizontal, or flat part, of the graph.
 - The point of inflection of the graph in Worked example 23, $y = x^3$, is at the origin $(0, 0)$.
- Cubics can undergo transformations in the same way as parabolas and hyperbolas.

**6.9.4 The exponential**

eles-3690

- Exponentials are non-linear graphs of the form $y = a^x$, where a is a real number not equal to 1.

WORKED EXAMPLE 24 Graphing an exponential

Complete the table of values below and use it to plot $y = 2^x$.

x	-3	-2	-1	0	1	2	3
y							


THINK

1. Substitute each x -value into the equation $y = 2^x$ to obtain the corresponding y -value.

WRITE/DRAW

x	-3	-2	-1	0	1	2	3
y	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8


2. Draw a set of axes and plot the points from the table. Join them with a smooth curve.



TI | THINK

In a new problem, on a Graphs page, complete the function entry line as:
 $f1(x) = 2^x$
 Then press ENTER.


WRITE



CASIO|THINK

On the Main screen complete the entry line as:
 $y = 2^x$ and then press the Graph icon to bring up the graphing screen. Highlight the equation and drag it down to the graphing screen.


WRITE



- The graph in Worked example 24 has several key features.
 - The value of y is always greater than zero.
 - The y -intercept is 1.
 - As x increases, y becomes very large.
 - As x decreases, y gets closer to but never touches zero, which is the x -axis. The x -axis is therefore called an **asymptote**.
- Exponentials can undergo transformations.

Comparing exponential graphs

- The diagram shows the graphs of $y = 2^x$ and $y = \left(\frac{1}{2}\right)^x$.




- The graphs both pass through the point $(0, 1)$.
- $y = 0$ is an asymptote for both graphs.
- The graph of $y = 2^x$ is reflected across the y -axis to produce the graph of $y = \left(\frac{1}{2}\right)^x$.

6.9.5 The circle

eles-4798


- A circle is the path traced out by a point at a constant distance (the radius) from a fixed point (the centre).
- Consider the circle shown. Its centre is at the origin and its radius is r .



- Let $P(x, y)$ be a point on the circle.
- By Pythagoras' theorem, $x^2 + y^2 = r^2$.
- This relationship is true for all points on the circle.
- The equation of a circle with centre $(0, 0)$ and radius r is: $x^2 + y^2 = r^2$.
- If the circle is translated h units to the right, parallel to the x -axis, and k units upwards, parallel to the y -axis, then the equation of a circle with centre (h, k) and radius r is: $(x - h)^2 + (y - k)^2 = r^2$.

Equation of a circle with centre (h, k) and radius r

$$(x - h)^2 + (y - k)^2 = r^2$$



WORKED EXAMPLE 25 Graphing a circle centered at the origin

Sketch the graph of $x^2 + y^2 = 49$, stating the centre and radius.

THINK

1. Write the equation.
2. State the coordinates of the centre.
3. Calculate the length of the radius by taking the square root of both sides. (Ignore the negative results.)
4. Sketch the graph.

WRITE/DRAW


$$x^2 + y^2 = 49$$

$$\text{Centre} = (0, 0)$$

$$r^2 = 49$$

$$r = 7$$

Radius = 7 units




TI | THINK


In a new problem, on a Graphs page, press:

- MENU
- 3: Graph Entry/Edit
- 2: Equation
- 3: Circle
- 1: Centre form.

WRITE




Complete the function entry line as shown. Press TAB to move between the fields. Press ENTER.




CASIO | THINK

On the Main screen, complete the entry line as:
 $x^2 + y^2 = 49$


WRITE




Press the Graph icon to bring up the graphing screen.



The graph of the circle is shown.



Highlight the equation and drag it down to the graphing screen. Press the Resize icon to look at the graph on the full screen.



To determine the centre of the circle, press:


- MENU
- 6: Analyze Graph
- 9: Analyze Conics
- 1: Centre.

Click on the circle and press ENTER. The coordinates of the centre are displayed.

To find the radius of the circle, press:

- MENU
- 6: Analyze Graph
- 9: Analyze Conics
- 7: Radius.

Click on the circle and press ENTER. The radius of the circle is displayed.



The circle $x^2 + y^2 = 49$ has its centre at $(0, 0)$ and has a radius of 7.

WORKED EXAMPLE 26 Graphing a circle

Sketch the graph of $(x - 2)^2 + (y + 3)^2 = 16$, clearly showing the centre and radius.

THINK

1. Express the equation in general form.
2. State the coordinates of the centre.
3. State the length of the radius.

WRITE

$$(x - h)^2 + (y - k)^2 = r^2$$

$$(x - 2)^2 + (y + 3)^2 = 4^2$$


Centre $(2, -3)$

$$r^2 = 16$$

$$r = 4$$

Radius = 4 units

4. Sketch the graph.



Resources



eWorkbook Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)



Interactivities Individual pathway interactivity: Non-linear relations (parabolas, hyperbolas, circles) (int-4509)
 Equations of circles (int-6053)
 Horizontal translations of parabolas (int-6054)
 Vertical translations of parabolas (int-6055)

Exercise 6.9 Non-linear relations

learn **on**

6.9 Quick quiz



6.9 Exercise

Individual pathways

PRACTISE

1, 2, 4, 8, 11, 15, 17, 21, 24

CONSOLIDATE

3, 5, 9, 12, 14, 16, 20, 23

MASTER

6, 7, 10, 13, 18, 19, 22, 25

Fluency

- WE20** State the vertical translation and the coordinates of the turning point for the graph of each of the following equations.

a. $y = x^2 + 3$ b. $y = x^2 - 1$ c. $y = x^2 - 7$ d. $y = x^2 + \frac{1}{4}$
- WE21** State the horizontal translation and the coordinates of the turning point for the graph of each of the following equations.

a. $y = (x - 1)^2$ b. $y = (x - 2)^2$ c. $y = (x + 10)^2$ d. $y = (x + 4)^2$
- For each of the following graphs, give the coordinates of the turning point and state whether it is a maximum or a minimum.

a. $y = -x^2 + 1$ b. $y = x^2 - 3$ c. $y = -(x + 2)^2$
- WE22** Complete the table of values shown and use it to plot the graph of $y = \frac{10}{x}$.

x	-5	-4	-3	-2	-1	0	1	2	3	4	5
y											

5. a. Plot the graph of each of the following hyperbolas.

b. Write the equation of each asymptote.

i. $y = \frac{5}{x}$

ii. $y = \frac{20}{x}$

iii. $y = \frac{100}{x}$

6. On the same set of axes, draw the graphs of $y = \frac{2}{x}$, $y = \frac{3}{x}$ and $y = \frac{4}{x}$.

7. Use your answer to question 6 to describe the effect of increasing the value of k on the graph of $y = \frac{k}{x}$.

8. **WE23** Complete the table of values shown and use it to plot the graph of $y = 2x^3$

x	-3	-2	-1	0	1	2	3
y							

9. Use digital tools to sketch the graph of each of the following cubics on the same axes, noting the point of inflection: $y = \frac{1}{2}x^3$, $y = 3x^3$, $y = 7x^3$

10. Use your graphs in question 9 to describe the effect of increasing the value of k on the graph of $y = kx^3$.

11. **WE24** Complete the table of values shown and use it to plot the graph of $y = 4^x$

x	-3	-2	-1	0	1	2	3
y							

12. Use digital tools to sketch the graph of each of the following exponentials on the same axes:
 $y = 3^x$, $y = 5^x$, $y = 10^x$.

13. Use your graphs in question 12 to describe the effect of increasing the value of a on the graph of $y = a^x$.

14. Use digital tools to sketch the graph of $y = \sqrt{x}$.

15. **WE25** Sketch the graphs of the following, stating the centre and radius of each.

a. $x^2 + y^2 = 49$

b. $x^2 + y^2 = 4^2$

c. $x^2 + y^2 = 36$

d. $x^2 + y^2 = 81$

e. $x^2 + y^2 = 25$

f. $x^2 + y^2 = 100$

16. **WE26** Sketch the graphs of the following, clearly showing the centre and the radius.

a. $(x - 1)^2 + (y - 2)^2 = 5^2$

b. $(x + 2)^2 + (y + 3)^2 = 6^2$

c. $(x + 3)^2 + (y - 1)^2 = 49$


d. $(x - 4)^2 + (y + 5)^2 = 64$

e. $x^2 + (y + 3)^2 = 4$


f. $(x - 5)^2 + y^2 = 100$

17. **MC** Select which of the following is the graph of $(x - 2)^2 + (y + 5)^2 = 4$.


A.




B.




C.



D.



E.



Reasoning

18. Show that $y = x^2$ is the same as $y = (-x)^2$.
19. Explain the similarities and differences between $y = \frac{1}{x}$ and $y = -\frac{1}{x}$.
20. Show that the turning point for $y = (x + 3)^2 - 1$ is $(-3, -1)$. Explain whether this is a maximum or minimum turning point.
21. Show the equation of the circle that has centre $(2, 1)$ and passes through the point $(6, 1)$ is $(x - 2)^2 + (y - 1)^2 = 16$.
22. Identify the point on the circle $x^2 + y^2 - 12x - 4y = 9$ that has the lowest y -value.

Problem solving


23. At a kindergarten sports day, Edmond, Huot and Nalini ran a 20-metre race. Each child ran according to the following equations, where d is the distance, in metres, from the starting line, and t is the time, in seconds.

Edmond: $d = 2.4 + 0.5t$

Huot: $d = 0.1t(t - 5)$

Nalini: $d = 0.2t(t - 5)(t - 9)$

The following sketch shows their running paths.



- Describe Nalini's race style.
 - Determine how long it took Huot to start running.
 - At the start of the race, determine how far Edmond was from the starting line.
 - If the winner of the 20-metre race won the race in 10.66 seconds, determine whether Huot or Nalini won the race. Justify your answer using calculations.
24. A circle with centre $(0, 2)$ passes through the point $(3, 6)$. Determine the coordinates of the points where the circle crosses the y -axis.
25. Consider the relationship $\frac{1}{y} = \frac{1}{x} + 1$. It appears that both variables are being raised to the power of 1. The relationship can be transposed as shown.
- Generate a table of values for x versus y using the transposed relationship.
 - Plot a graph to show the points contained in the table of values.
 - Use your plot to confirm whether this relationship is linear.

$$\begin{aligned} \frac{1}{y} &= \frac{1}{x} + 1 \\ \frac{1}{y} &= \frac{1}{x} + \frac{x}{x} \\ \frac{1}{y} &= \frac{1+x}{x} \\ y &= \frac{x}{1+x} \end{aligned}$$

LESSON

6.10 Review

6.10.1 Topic summary


LINEAR AND NON-LINEAR GRAPHS

Gradient and axis intercepts

- The gradient of a straight line is given the label m . It can be determined by the following formula:

$$m = \frac{\text{rise}}{\text{run}} = \frac{y_2 - y_1}{x_2 - x_1}$$

- If the line is sloping upwards (from left to right), the gradient is positive.
- If the line is sloping downwards, the gradient is negative.
- The y -intercept of a line can be determined by letting $x = 0$, then solving for y .
- The x -intercept of a line can be determined by letting $y = 0$, then solving for x .



Equation of a straight line

- The equation of a straight line is $y = mx + c$ where:
 - m is the gradient
 - c is the y -intercept.

Sketching linear graphs

- To sketch a linear graph, all you need are two points that the line passes through.
- The x - and y -intercept method involves finding both axis intercepts, then drawing the line through them.
- The gradient and y -intercept method involves plotting the y -intercept and then one other point (usually $x = 1$) using the gradient.

Horizontal and vertical lines

- Horizontal lines are of the form $y = c$. They have a gradient of zero ($m = 0$).
- Vertical lines are of the form $x = a$. They have an undefined gradient.


Midpoint and distance

- The midpoint of two points (x_1, y_1) and (x_2, y_2) is

$$M = \left(\frac{x_2 + x_1}{2}, \frac{y_2 + y_1}{2} \right)$$


- The distance between two points, (x_1, y_1) and (x_2, y_2) is

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$




Non-linear graphs


- A basic **parabola** has the equation $y = x^2$.
- It has a concave upward shape and a turning point at the origin.




- A **hyperbola** has the equation $y = \frac{1}{x}$.
- It consists of two curved arcs that do not touch.



- A **circle** of radius r , when centred at the origin, has the equation $x^2 + y^2 = r^2$.



- A **circle** of radius r , when centred at the point (h, k) , has the equation $(x - h)^2 + (y - k)^2 = r^2$.



6.10.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
6.2	I can plot points on the Cartesian plane.			
	I can plot a linear graph on the Cartesian plane.			
6.3	I can calculate the gradient of a line or line segment.			
	I can calculate the y -intercept of a linear graph.			
6.4	I can determine the equation of a straight line given the gradient and the y -intercept.			
	I can determine the equation of a straight line given the gradient and a single point.			
	I can determine the equation of a straight line given two points.			
	I can determine the equation of a straight line from a graph.			
6.5	I can sketch linear graphs using the x - and y -intercept method.			
	I can sketch linear graphs using the gradient–intercept method.			
	I can sketch vertical and horizontal lines.			
6.6	I can produce linear graphs using digital technologies.			
6.7	I can determine the linear rule from a table of values.			
	I can determine the linear rule that models a real-life problem.			
	I can interpret linear rules in real-life problems.			
6.8	I can calculate the midpoint of a line segment.			
	I can calculate the distance between two points.			
6.9	I can recognise and sketch a basic parabola from its equation.			
	I can sketch a parabola that has been translated horizontally or vertically.			
	I can recognise and use digital technology to sketch hyperbolas, cubics and exponentials.			
	I can recognise and sketch a circle from its equation.			

6.10.3 Project

Path of a billiard ball

The path of a billiard ball can be mapped using mathematics. A billiard table can be represented by a rectangle, a ball by a point and its path by line segments. In this investigation, we will look at the trajectory of a single ball, unobstructed by other balls.

Note: A billiard table has a pocket at each of its corners and one in the middle of each of its long sides.

Consider the path of a ball that is hit on its side, from the lower left-hand corner of the table, so that it travels at a 45° angle from the corner of the table. Assume that the ball continues to move, rebounding from the sides and stopping only when it comes to a pocket.

Diagrams of the table drawn on grid paper are shown. Each grid square has a side length of 0.25 metres.












Diagram A shows the trajectory of a ball that has been hit on its side at a 45° angle, from the lower left-hand corner of a table that is 2.5 m long and 1.25 m wide.

Note that, because the ball has been hit on its side at a 45° angle, it travels diagonally through each square in its path, from corner to corner. Diagram B shows the trajectory of a ball on a 2.5 m by 1.25 m table.

1. In Diagram B, how many times does the ball rebound off the sides before going into a pocket?
2. A series of eight tables of different sizes are drawn on grid paper in the following diagrams. For each table, determine the trajectory of a ball hit at 45° on its side, from the lower left-hand corner of the table. Draw the path each ball travels until it reaches a pocket.





- Determine for which tables the ball will travel through the simplest path. What is special about the shape of these tables?
- Determine for which table the ball will travel through the most complicated path. What is special about this path? Draw another table (and path of the ball) with the same feature.
- Determine for which tables the ball will travel through a path that does not cross itself. Draw another table (and path of the ball) with the same feature.
- Consider the variety of table shapes. Will a ball hit on its side from the lower left-hand corner of a table at 45° always end up in a pocket (assuming it does not run out of energy)?

Simplify matters a little and consider a billiard table with no pockets in the middle of the long sides. Use a systematic way to look for patterns for tables whose dimensions are related in a special way.

- Draw a series of billiard tables of length 3 m. Increase the width of these tables from an initial value of 0.25 m in increments of 0.25 m. Investigate the final destination (the pocket the ball lands in) of a ball hit from the lower left-hand corner. Complete the table.

	Length of table (m)	Width of table (m)	Destination pocket
a.	3	0.25	
b.	3	0.5	
c.	3	0.75	
d.	3	1	
e.	3	1.25	
f.	3	1.5	
g.	3	1.75	
h.	3	2	

- How can you predict (without drawing a diagram) the destination pocket of a ball hit from the lower left-hand corner of a table that is 3 m long? Provide an illustration to verify your prediction.



eWorkbook Topic 6 Workbook (worksheets, code puzzle and project) (ewbk-2006)



Interactivities Crossword (int-2688)
Sudoku puzzle (int-3207)

Exercise 6.10 Review questions

Fluency

- MC** For the rule $y = 3x - 1$, select the value of y when $x = 2$.

A. -1 B. 1 C. 2 D. 5 E. 7
- MC** Identify the gradient of the linear rule $y = 4 - 6x$.

A. 6 B. -6 C. 4 D. -4 E. -2
- MC** The graph with the rule $2y - x + 6 = 0$ has an x -intercept of:

A. -2 B. 0 C. 2 D. 6 E. 7
- MC** The graph with the rule $2y - x + 6 = 0$ has a y -intercept of:

A. -6 B. -3 C. 0 D. 3 E. 6
- MC** Consider a linear graph that goes through the points $(6, -1)$ and $(0, 5)$. The gradient of this line is:

A. 5 B. -5 C. 1 D. -1 E. 3
- MC** A straight line passes through the points $(2, 1)$ and $(5, 4)$. Its rule is:

A. $y = x - 1$ B. $y = x + 1$ C. $y = 2x$ D. $4y = 5x$ E. $y = 3x$
- MC** The rule for a line whose gradient is -4 and y -intercept $= 8$ is:

A. $y = -4x + 32$ B. $y = -4x + 8$ C. $y = 4x - 32$ D. $y = 4x - 8$ E. $-4y = x + 8$
- MC** Identify which of the following linear rules will *not* intersect with the straight line defined by $y = 3x$.

A. $y = 3x + 2$ B. $y = -3x + 1$ C. $y = -3x + 2$ D. $y = -2x + 1$ E. $y = 3$
- MC** If $y = 2x + 1$, select from the following the point that could not be on the line.

A. $(3, 7)$ B. $(-3, -5)$ C. $(0, 1)$ D. $(-3, 0)$ E. $(5, 11)$
- MC** The solution to $y = 3x + 1$ and $y = -3x + 1$ is:

A. $(0, 1)$ B. $(1, 0)$ C. $\left(0, -\frac{1}{3}\right)$ D. $\left(-\frac{1}{3}, 0\right)$ E. $(3, 1)$

11. Write down the gradient and the y -intercept of the following linear graphs.

a. $y = 8x - 3$

b. $y = 5 - 9x$

c. $2x + y - 6 = 0$

d. $4x - 2y = 0$

e. $y = \frac{2x - 1}{3}$

12. **MC** Determine which of the following lines are parallel to the line with the equation $y = 6 - x$. There could be more than one correct answer.

A. $y + x = 4$

B. $y = 13 - x$

C. $2y - 2x = 1$

D. $x + 2y - 4 = 0$

E. $2x - 3y - 18 = 0$

13. **MC** Determine which of the following rules will yield a linear graph. There could be more than one correct answer.

A. $3y = -5x - \frac{1}{2}$




B. $y = 3x^2 - 2\frac{1}{2}$

C. $3x + 4y + 6 = 0$

D. $x + y - 2xy = 0$

E. $2y = 4^2x + 9^2$

14. Determine the gradient of the lines shown.



15. Calculate the gradient of the line passing through the following pairs of points.

a. $(2, -3)$ and $(4, 1)$

b. $(0, -5)$ and $(4, 0)$.

16. For each of the following rules, state the gradient and the y -intercept.

a. $y = -3x + 7$

b. $2y - 3x = 6$

c. $y = -\frac{2}{5}x$

d. $y = 4$

17. For the following rules, use the gradient–intercept method to sketch linear graphs.

a. $y = -x + 5$

b. $y = 4x - 2.5$

c. $y = \frac{2}{3}x - 1$



d. $y = 3 - \frac{5}{4}x$

18. For the following rules, use the x - and y -intercept method to sketch linear graphs.
- a. $y = -6x + 25$ b. $y = 20x + 45$ c. $2y + x = -5$ d. $4y + x - 2.5 = 0$

19. For the following rules, use an appropriate method to sketch linear graphs.

- a. $y = -3x$ b. $y = \frac{1}{4}x$ c. $y = -2$ d. $x = 3$

20. Determine the rule of the lines shown.



21. Determine the linear rules given the following pieces of information.

- a. Gradient = 2, y -intercept = -7 b. Gradient = 2, x -intercept = 7
 c. Gradient = 2, passing through (7, 9) d. Gradient = -5 , passing through the origin
 e. y -intercept = -2 , passing through (1, -3) f. Passing through (1, 5) and (5, -6)
 g. x -intercept = 3, y -intercept = -3 h. y -intercept = 5, passing through (-4 , 13)

22. Determine the midpoint of the line interval joining the points (-2 , 3) and (4, -1).

23. Calculate the distance between the points (1, 1) and (4, 5).

24. Sketch each of the following, comparing it with the graph of $y = x^2$.

- a. $y = x^2 - 3$ b. $y = (x + 2)^2$

25. a. Plot the graph of $y = \frac{4}{x}$.

- b. Identify what type of graph this is.

26. Sketch the circle with the equation $x^2 + y^2 = 16$. Clearly show the centre and radius.

Problem solving

27. Louise owes her friend Sula \$400 and agrees to pay her back \$15 per week.
- a. State a linear rule that demonstrates this debt reduction schedule and sketch the graph.
 b. Calculate how many weeks it takes Louise to repay the debt.
 c. Calculate how much she owes after 15 weeks.
 d. Evaluate how many repayments Louise needs to make to reduce her debt to \$85.

28. A bushwalker is 40 km from their base camp when they decide to head back.

If they are able to walk 3.5 km each hour:

- determine the linear rule that describes this situation and sketch its graph
- calculate how long, correct to 1 decimal place, it will take them to reach base camp
- calculate how far they will have walked in 6.5 hours.



29. Udaya is writing test questions. She has already written 25 questions and can write a further 5 questions per hour.
- Represent this information as a linear equation where t hours is the time spent writing test questions and n is the number of questions written.
 - Predict the total number of questions written after a further 8 hours assuming the same linear rule.
 - Calculate how long, to the nearest minute, it will take Udaya to have written 53 questions.

30. Catherine earns a daily rate of \$200 for working in her mother's store. She receives \$5 for each necklace that she sells.
- Write an equation to show how much money (m) Catherine earned for the day after selling (n) necklaces.
 - Graph the equation that you created in part **a**, showing the two intercepts.
 - Explain which part of the line applies to her earnings.
 - Explain which part of the line does not apply to her earnings.



31. Calculate the gradient of the line passing through the points $(2, 3)$ and $(6 + 4t, 5 + 2t)$. Write your answer in simplest form.
32. Determine the point on the line $y = 2x + 7$ that is also 5 units above the x -axis.
33. An experiment was conducted to collect data for two variables p and t . The data are presented in the table shown.

p	$-\frac{1}{2}$		$\frac{1}{2}$	3.6
t	$2\frac{1}{4}$	1.75		-5.95

It is known that the relationship between p and t is a linear one. Determine the two values missing from the table.

34. The distance from the origin to the y -intercept of a linear graph is three times the distance from the origin to the x -intercept. The area of the triangle formed by the line and the axes is 3.375 units^2 . The line has a negative gradient and a negative y -intercept. Determine the equation of the line.



To test your understanding and knowledge of this topic go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 6 Linear and non-linear graphs

6.1 Pre-test

1. On the x -axis (because $y = 0$).

2. a. C b. D
c. B d. A

3. C

4. -0.5

5. D

6. a. B b. C
c. D d. A

7. a. $y = -\frac{1}{4}x$ b. $y = 20x + 400$

8. $a = -5$

9. $\sqrt{29}$

10. D

11. a. $y = -x + 2$ b. $(-1, 3)$

12. A

13. D

14. E

15. D


6.2 Plotting linear graphs

1. a. B b. C c. D

2. a. i. $y = x$

x	-3	-2	-1	0	1	2	3
y	-3	-2	-1	0	1	2	3


ii. $y = x$



b. i. $y = 2x + 2$

x	-3	-2	-1	0	1	2	3
y	-4	-2	0	2	4	6	8


ii. $y = 2x + 2$



c. i. $y = 3x - 1$

x	-3	-2	-1	0	1	2	3
y	-10	-7	-4	-1	2	5	8


ii. $y = 3x - 1$




d. i. $y = -2x$

x	-3	-2	-1	0	1	2	3
y	6	4	2	0	-2	-4	-6


ii. $y = -2x$




3. a. Sample points shown; $y = -x$




- b. Sample points shown; $y = \frac{1}{2}x + 4$



- c. Sample points shown; $y = -2x + 3$



- d. Sample points shown; $y = x - 3$



4. a. Yes b. No c. No d. Yes
 5. a. Yes b. Yes c. Yes
 d. No e. No f. No
 6. D
 7. a. B b. A c. D d. C

8. By plotting the points on the Cartesian plane and joining them to make a line it can be seen that this line does not pass through the third quadrant.

9. a. The first, second and third quadrants.


b. Answers will vary. Sample response: Substitute point (1, 3) into line equation. $3 \neq 1 + 1$.

10. Answers will vary. Sample response: Substitute the point into the line equation. If the LHS equals the RHS of the equation, then the point lies on the line. Alternatively, graph the line and check the point on the Cartesian plane.

11. Answers will vary. Sample response: Plot the points on the Cartesian plane and draw a line through the points. Find the equation of the line. Check all points lie on the line by substituting into the line equation.

12. (59, 21)

13. a.



- b. $m = 5t + 1$

c. Sample responses can be found in the worked solutions in the online resources.

14. Stirring increases the rate of reaction.

6.3 Features of linear graphs

1. a. 6 b. -4 c. 0
 d. $\frac{6}{5}$ e. -2.2 f. -45
 2. a. 0 b. 0.1 c. -2.6
 d. 0 e. 9 f. -0.1
 3. a. 23 b. 54 c. -6 d. 70
 4. a. 1 b. 5.2 c. 100 d. 100
 5. a. 4 b. 1 c. 0
 6. a. 20 b. 400 c. -5.2
 7. a. 0 b. -5 c. -300
 8. C
 9. y-intercept = c
 10. a. (0, -10) b. (0, -4) c. (0, 3)
 d. (0, 7) e. No y-intercept
 11. There is a rise and no run; $\frac{\text{rise}}{0} = \text{undefined}$.
 12. There is no rise and a run; $\frac{0}{\text{run}} = 0$
 13. Sample responses can be found in the worked solutions in the online resources.

14. It does not matter if you rise before you run or run before you rise, as long as you take into account whether the rise or run is negative.

15. a. $m = \frac{y-c}{x}$ b. $y = mx + c$

16. a. 0.28 b. 0.64

17. $0.6 > 0.58$, therefore the safety standards have not been met.

18. 0.3 metres

19. a. Lamb: $m = \$15/\text{kg}$, Chicken: $m = \$10/\text{kg}$,
Beef: $m = \$7.50/\text{kg}$

b. Lamb: \$15, Chicken: \$10, Beef: \$7.50

c. i. \$15 ii. \$5 iii. \$15

d. \$35

Meat type	Cost per kilogram (\$/kg)	Weight required (kg)	Cost = \$/kg \times kg
Lamb	15	1	15
Chicken	10	0.5	5
Beef	7.50	2	15
Total cost			35

20. a. $m = -\frac{3}{2}$

b. It does not matter which points are chosen to determine the gradient of the graph because the gradient will always remain the same.

c. Straight line with a y -intercept of $(0, -2)$ and a slope of $-\frac{3}{2}$

6.4 The equation of a straight line

1. a. $y = 4x + 2$ b. $y = -4x + 1$ c. $y = 4x + 8$
 2. a. $y = 6x + 7$ b. $y = -2.5x + 6$ c. $y = 45x + 135$
 3. a. $y = -2x$ b. $y = 4x$ c. $y = 10.5x$
 4. a. $y = -20x$ b. $y = 1.07x$ c. $y = 32x$
 5. a. $y = x + 2$ b. $y = -x + 8$
 c. $y = -4x - 8$ d. $y = 2x - 13$
 6. a. $y = -5x + 70$ b. $y = 2x - 23$
 c. $y = -6x + 11$ d. $y = -x - 1.5$
 7. a. $y = 6x + 30$ b. $y = -3.5x + 15.5$
 c. $y = 1.2x - 4.08$ d. $y = 0.2x - 1.76$
 8. a. $y = -4x - 24$ b. $y = 2x - 6$
 c. $y = -2x + 4$
 9. a. $y = 5x + 35$ b. $y = 1.5x - 3.75$
 c. $y = 0.4x - 0.96$
 10. a. $y = x + 17$ b. $y = -x + 3$
 c. $y = 11$ d. $y = x + 3.5$
 11. a. $y = -x + 3.5$
 b. $y = \frac{1}{9}x + \frac{34}{9}$ (or $9y = x + 34$)
 c. $y = -6x + 90$

d. $y = 20x$

12. a. $y = -x + 4$

b. $y = -0.25x + 5.5$

c. $y = \frac{2}{7}x - \frac{32}{7}$ (or $7y = 2x - 32$)

d. $y = \frac{2}{3}x + \frac{10}{3}$ (or $3y = 2x + 10$)

13. a. $y = x + 3$

b. $y = \frac{-5}{4}x + 5$

c. $y = -6x + 6$

d. $y = \frac{7}{8}x + 35$

14. a. $y = x + 8$

b. $y = -2x + 6$

c. $y = \frac{-3}{7}x - 3$

d. $y = \frac{1}{4}x + 50$

15. a. $y = \frac{7}{4}x$

b. $y = x$

c. $y = -2x$

d. $y = -3x$

e. $y = \frac{-24}{11}x$

f. $y = \frac{-x}{3}$

16. a. $y = 2x + 4$

b. $y = 5x$

17. a. $y = -x + 3$

b. $y = \frac{1}{5}x$

18. a. $y = x + 2$

b. $y = -2x + 3$

19. a. B

b. A

c. E

20. a. $y = -2.4x - 12$

b. $y = -58.32$

21. a. $y = -0.6x + 10$

b. $(16\frac{2}{3}, 0)$

22. a. The equation is $y = \frac{5}{4}x - 2135$

b. Values of c cannot be read directly from the graph because the graph doesn't contain the origin, and since c is found at $x = 0$, we need to use the equation.

c. In 2020 the concentration of CO_2 will be 390 ppm. The assumption is that the concentration of CO_2 will continue to follow this linear pattern.

23. $m = \frac{-15 + 7}{4 - 0} = -2$

$y = -2x + c$

$6 = -6 + c$

$c = 12$

The equation is $y = -2x + 12$.

24. a. Line A: $y = x - 1$, Line B: $y = -x + 8$

b. $(4.5, 3.5)$

c. Sample responses can be found in the worked solutions in the online resources. Possible methods include graphing on a Cartesian plane or using algebra.

Line B: $y = -x + 8, m = -1, m_{\perp} = 1$


$y = x + c$

$6 = -4 + c$

$c = 10$

25. $m = \frac{1}{4}C$

26. a, c



b. $y = -2x + 3$ c. (1, 1) d. 1 star

e. Sample responses can be found in the worked solutions in the online resources. Possible responses could include the following.

$y = -x - 1, x = 2$

27. Sample responses can be found in the worked solutions in the online resources. Responses should include the correct equation: $T = 4.25t + 15$.

28. 8.1 mL

29. a. A: \$0, B: \$100, C: \$200

b. A: \$80, B: \$50 C: \$33.33

c. A: $w = 80h$, B: $w = 50h + 100$, C: $w = 33.3h + 200$

d.i. C ii. A


e. The advantage of location C is that it has the highest minimum pay, so you would be guaranteed at least \$200 per day, regardless of how many hours you work.

f. i. A: \$ 160, B: \$ 200, C: \$ 266.66


ii. A: \$ 480, B: \$ 400, C: \$ 400

6.5 Sketching linear graphs


1. a. $5y - 4x = 20$




b. $y = x + 2$




c. $y = -3x + 6$




2. a. $3y + 4x = -12$





















b. $x - y = 5$



c. $2y + 7x - 8 = 0$








11. D

12. D


13. a. $y = -\frac{2x}{5} + 4$

b. $-\frac{2}{5}$

c. The x-intercept is 10, the y-intercept is 4.



14. a. $m = \frac{\text{rise}}{\text{run}} = \frac{5}{3}$




c. i. $y = \frac{5x}{3} + 5$

ii. $-5x + 3y = 15$

15. a. $y = \frac{4}{3}x - 8$

b. $y = 0, 4x = 24, x = 6$
 $x = 0, -3y = 24, y = -8$



16. a. $y = \frac{-ax + c}{b}$

b. Sample response: The gradient is $\frac{-a}{b}$ substituting positive values always results in a negative gradient.

17. Sample response:

$$4y = \frac{4}{2}x + 3$$

$$4y = 2x + 3$$

$$-2x + 4y = 3$$



18. Sample response: All descriptions use the idea that a gradient is equal to $\frac{\text{rise}}{\text{run}}$, which equals $\frac{2}{3}$ in all of these cases.

19. a. The gradient is 2. b. The equation of the line is $y = 2x - 1$.

c. One point is (1, 1).

20. a. i. C ii. B iii. A

b. Straight line y-intercept of (0, 2) with a slope of $-\frac{1}{4}$



iii. Sample responses can be found in the worked solutions in the online resources.

b. Sample responses can be found in the worked solutions in the online resources.

22. Sample responses can be found in the worked solutions in the online resources.

6.6 Technology and linear graphs

1. i. As the size of the coefficient increases, the steepness of the graph increases.

ii. Each graph cuts the x-axis at (0, 0).

iii. Each graph cuts the y-axis at (0, 0).

2. i. As the size of the coefficient decreases, the steepness of the graph decreases.

ii. Each graph cuts the x-axis at (0, 0)

iii. Each graph cuts the y-axis at (0, 0)

3. i. As the magnitude of the coefficient decreases, the steepness of the graph increases.

ii. Each graph cuts the x-axis at (0, 0).

iii. Each graph cuts the y-axis at (0, 0).

4. a. positive, positive b. downward, negative

c. bigger d. will

5. a. Yes, 1

b. No, the lines are parallel.

c. i. (0, 0), $x = 0$

ii. (-2, 0), $x = -2$

iii. (2, 0), $x = 2$

d. i. (0, 0), $y = 0$

ii. (0, 2), $y = 2$

iii. (0, -2), $y = -2$

6. a. Yes, -1

b. No, the lines are parallel.

c. i. (0, 0), $x = 0$

ii. (2, 0), $x = 2$

iii. (-2, 0), $x = -2$

d. i. (0, 0), $y = 0$

ii. (0, 2), $y = 2$

iii. (0, -2), $y = -2$

7. a. same

c. y-intercept

b. y-intercept, y-axis

d. x-intercept

8. i. No

ii. Yes

iii. a. 1

b. -1

c. 3

d. $-\frac{2}{5}$

iv. a. $(-5, 0)$, i.e. $x = -5$

b. $(5, 0)$, i.e. $x = 5$

c. $(-\frac{5}{3}, 0)$, i.e. $x = -\frac{5}{3}$

d. $(\frac{25}{2}, 0)$, i.e. $x = \frac{25}{2}$

v. a. $(0, 5)$, i.e. $y = 5$

b. $(0, 5)$, i.e. $y = 5$

c. $(0, 5)$, i.e. $y = 5$

d. $(0, 5)$, i.e. $y = 5$

9. a. steepness b. y-coordinate, y-axis

c. same, parallel d. the same

10. a. Gradient = 2, y-intercept = $(0, 0)$, $y = 0$

b. Gradient = 1, y-intercept = $(0, 1)$, $y = 1$

c. Gradient = -3, y-intercept = $(0, 5)$, $y = 5$

d. Gradient = $\frac{2}{3}$, y-intercept = $(0, -7)$, $y = -7$


11. Sample response: They all have in common the point $(1, 5)$. Their gradients are all different.

12. Sample responses can be found in the worked solutions in the online resources. Use solve on your CAS calculator.

13. Answers should show the y-intercept $(0, 2.2)$, point $(1, 2.75)$ and gradient 0.55.


14. a. N is the number of dogs walked; $-\$15$ is Shirley's starting cost and out-of-pocket expense before she walks a dog, and she earns $\$10$ for every dog walked.

b. She needs to walk at least two dogs a week before she can make a profit.



15. Sample responses can be found in the worked solutions in the online resources.

16. a. $y = 2x - 8.5$



c. $y = 7.5$

d. $x = 10.25$

17. a. $C = 25n + 1000$

b. $(0, 1000)$. This is the initial cost of the site licence.

c. $\$6000$

d. 80

18. a. Sample response: Possible methods include plotting the points $(0, 325)$ and $(1, 328.8)$ or using the equation $c = 3.8m + 325$.

b. 382 calories

c. 32.9 minutes

6.7 Practical applications of linear graphs

1. a. $y = 6x - 5$ b. $y = -3x + 8$

2. a. $y = 2x + 4$ b. $y = 0.9x + 1.1$

3. a. $y = 3x + 1$ b. $y = -x + 17$

4. a. $v = -3t + 27$ b. $C = 3d + 8$

5. D

6. a. $A = 25t + 60$ b. $\$5060$

c. $\$60$

7. a. $C = 0.55s + 2.0$ b. $\$8.60$

8. a. $N = 400t + 3200$ b. 5400

c. 13 200

9. a. 27 b. 108

10. a. $w = -40t + 712$ b. 712 L

c. 18 days

11. Sample responses can be found in the worked solutions in the online resources.
12. a. $C = 2.57d + 3.50$
 b. \$33.06
 c. 7.20 km
13. $5 \times 21.4 = 107 \text{ cm} = 1.07 \text{ m}$
14. a. $N = -22t + 164$ b. 120
 c. 5 d. 7.5 months
15. a. See the figure at the foot of the page.*
 b. $(0, -800)$, $y = -800$. Fixed costs are \$800.
 c. $(4, 0)$, $x = 4$, this is the break-even amount
 d. 200, this is the sale price per print that contributes to profit
16. a. $C = 2.14d + 3.50$
 b. \$28.11
 c. \$52.93
17. a. $Y = 78.7A$ b. ¥196 750 c. $Y_T = 76.5A$
18. Cody arrives first.

5. a. 5 b. 9 c. -16 d. 0
6. a. $(4, 6)$ b. $(-2, -3)$
 c. $(11, 2)$ d. $(-7, -3)$
7. $y = 5x - 7$
8. a. 5 b. 10 c. 1
9. a. 2 b. 13 c. 10
10. a. 5 b. 17 c. 13
11. a. 7.211 b. 14.765 c. 13.038
12. a. 8.944 b. 19.416 c. 19.416
13. a. 24.472 b. 25.464
14. a. 17.788 b. 25.763
15. 21.024 km
16. $d = 5$

$$17. -2 = \frac{x+5}{2} \Rightarrow x = -9$$

$$-4 = \frac{-6+y}{2} \Rightarrow y = -2$$

The required point is $(-9, -2)$.

$$18. d_{AB} = \sqrt[5]{10}$$


$$d_{BC} = \sqrt[5]{10}$$


Therefore, B is equidistant from A and C.

6.8 Midpoint of a line segment and distance between two points

1. a. $(2, 4)$ b. $(5, 1)$ c. $(7, 2)$
 2. a. $(8, 9)$ b. $(0, 5)$ c. $(-1, 2)$
 3. a. $(-5, -2)$ b. $(2, 3)$ c. $(0, 12)$
 4. a. $\left(1\frac{1}{2}, 5\frac{1}{2}\right)$ b. $\left(-3, -3\frac{1}{2}\right)$
 c. $\left(-9\frac{1}{2}, 9\frac{1}{2}\right)$ d. $\left(-1\frac{1}{2}, 6\frac{1}{2}\right)$

*15. a.





b. $M = (-2, 2)$

c. $2\sqrt{37}$

d. $(7, -2)$

e. $\frac{3 - (-4)}{2 - (-1)} = \frac{7}{3}$

f. $y = -4x + 11$

20. Sample responses can be found in the worked solutions in the online resources.

$d_{AB} = 2\sqrt{10}$

$d_{BC} = 2\sqrt{10}$

$d_{AC} = 4\sqrt{2}$

Side length AB is equal to side length BC but not equal to side length AC. Therefore, $\triangle ABC$ is an isosceles triangle.

21. $\frac{1}{2}$

22. $(-4, 5), (-3, 2), (-2, -1), (-1, -4), (0, -7)$

6.9 Non-linear relations

1. a. Vertical 3 up, TP $(0, 3)$

b. Vertical 1 down, TP $(0, -1)$

c. Vertical 7 down, TP $(0, -7)$

d. Vertical $\frac{1}{4}$ up, TP $(0, \frac{1}{4})$

2. a. Horizontal 1 right, $(1, 0)$


b. Horizontal 2 right, $(2, 0)$

c. Horizontal 10 left, $(-10, 0)$


d. Horizontal 4 left, $(-4, 0)$

3. a. $(0, 1)$, max b. $(0, -3)$, min c. $(-2, 0)$, max


4. See the table at the foot of the page.*




5. a. i.



ii.



iii.




b. i. $x = 0, y = 0$

ii. $x = 0, y = 0$


iii. $x = 0, y = 0$

*4.



x	-5	-4	-3	-2	-1	0	1	2	3	4	5
y	-2	-2.5	-3.3	-5	-10	Undefined	10	5	3.3	2.5	2




7. It increases the y -values by a factor of k and hence dilates the curve by a factor of k . As k increases, the graph is further from the origin.




9. The point of inflection for all three graphs is at the origin $(0,0)$
TI




10. It increases the y -values by a factor of k so as the value of k increases the arms of the cubic move closer to the y -axis.



12. TI




CASIO




13. As the value of a increases the values of y increase more and more rapidly resulting in the graph moving closer to the y -axis.


14. TI



CASIO




15. a.




Centre (0, 0), radius 7


b.






Centre (0, 0), radius 4



Centre (0, 0), radius 6




Centre (0, 0), radius 5




17. D




18. Sample responses can be found in the worked solutions in the online resources. Possible methods include using a table of values or graphing both equations on a Cartesian plane.
19. Similarity: they both have asymptotes of $x = 0$ and $y = 0$.
Difference: one is a reflection of the other about the y -axis.
20. Sample responses can be found in the worked solutions in the online resources. Use algebra or graph the equation. The turning point is a minimum.
21. $r = 4$
The centre is (2, 1). The equation is $(x - 2)^2 + (y - 1)^2 = 16$
Sample responses can be found in the worked solutions in the online resources.



Centre (0, 0), radius 9




Centre (0, 0), radius 10



22. (6, -5)


23. a. Nalini started well and raced ahead of Huot and Edmond. She then turned around and went back to the starting line. She stayed at the starting line for an amount of time before turning around and sprinting to the finishing line, passing Huot and Edmond.
- b. From the graph, Huot took 5 seconds to start running.
- c. From the graph, at $t = 0$, Edmond is 2.4 metres from the starting line.
- d. Nalini won the race.



At (0, 7) and (0, -3)

25. a.

x	-3	-2	-1	0	1	2	3
y	$\frac{3}{2}$	2	Undef.	0	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{3}{4}$



c. This relationship is not linear.

Project





- 6
- Sample responses can be found in the worked solutions in the online resources.
- a and h
- e
- a, b, c and h
- Yes









	Length of table (m)	Width of table (m)	Destination pocket
a	3	0.25	Far left
b	3	0.5	Far left
c	3	0.75	Far left
d	3	1	Far right
e	3	1.25	Far left
f	3	1.5	Far left
g	3	1.75	Far right
h	3	2	Close right

8. Sample responses can be found in the worked solutions in the online resources.

6.10 Review questions


- D
- B
- D
- B
- D
- A
- B
- A
- D
- A
- a. Gradient = 8, y-intercept -3
b. Gradient = -9, y-intercept 5
c. Gradient = -2, y-intercept 6
d. Gradient = 2, y-intercept 0
e. Gradient = $\frac{2}{3}$, y-intercept $-\frac{1}{3}$
- A, B
- A, C, E
- a. 2 b. $-\frac{2}{5}$ c. 0
- a. 2 b. $\frac{5}{4}$
- a. $m = -3, c = 7$
b. $m = \frac{3}{2}, c = 3$
c. $m = -\frac{2}{5}, c = 0$
d. $m = 0, c = 4$






20. a. $y = 2x$
 b. $y = 2 - \frac{2}{7}x$
21. a. $y = 2x - 7$
 b. $y = 2x - 14$
 c. $y = 2x - 5$
 d. $y = -5x$
 e. $y = -x - 2$
 f. $y = -2.75x + 7.75$
 g. $y = x - 3$
 h. $y = -2x + 5$
22. (1,1)
23. 5 units


24. a. Vertical translation 3 units down, TP = (0, -3)



- b. Horizontal translation 2 units to the left; TP = (2, 0)




25. a.




- b. Hyperbola

- 26.




27. a. $y = 400 - 15x$



- b. 26.7 or 27 weeks
 c. \$175
 d. 21 repayments


28. a. $y = 40 - 3.5x$



- b. 11.4 hours
 c. 22.75 km
 29. a. $n = 5t + 25$ b. 65 c. 336 minutes

30. a. $m = 5n + 200$
- b. See the figure at the foot of the page.*
- c. Everything to the right of the vertical axis and including the vertical axis applies to her earnings because she can only sell 0 or more necklaces.
- d. Everything to the left of the vertical axis, because she cannot sell a negative number of necklaces and because she cannot make less than her base salary of \$200.
31. $\frac{1}{2}$
32. $(-1, 5)$
33. $p = -0.25, t = 0.25$
34. $y = -3x - 4.5$

*30. b.




Semester review 1

The learnON platform is a powerful tool that enables students to complete revision independently and allows teachers to set mixed and spaced practice with ease.

Student self-study

Review the **Course Content** to determine which topics and subtopics you studied throughout the year. Notice the green bubbles showing which elements were covered.



Review your results in **My Reports** and highlight the areas where you may need additional practice.





Use these and other tools to help identify areas of strengths and weakness and target those areas for improvement.


Teachers

It is possible to set questions that span multiple topics. These assignments can be given to individual students, to groups or to the whole class in a few easy steps.

Go to **Menu** and select **Assignments** and then **Create Assignment**. You can select questions from one or many topics simply by ticking the boxes as shown below.



Once your selections are made, you can assign to your whole class or subsets of your class, with individualised start and finish times. You can also share with other teachers.



More instructions and helpful hints are available at www.jacplus.com.au.

7 Proportion and rates

LESSON SEQUENCE

7.1 Overview	414
7.2 Direct linear proportion	416
7.3 Proportionality	421
7.4 Introduction to rates	426
7.5 Constant and variable rates	429
7.6 Review	436



LESSON


7.1 Overview

7.1.1 Why learn this?

Proportion and rates are often used to compare quantities. Kilometres per hour, price per kilogram, dollars per litre and pay per hour are all examples of rates that are used in everyday life. Supermarkets give the prices of items as a rate, such as dollars per kilogram, so customers can compare different-sized packages of the same item and make decisions about which is the better value. Chefs use proportions when combining different ingredients from a recipe.

In geometry, the circumference of a circle is proportional to its diameter. We say, simply, that the circumference, C , is pi (π) times its diameter, d ; that is, $C = \pi d$. Calculations of π date back to the time of Archimedes, but the symbol became widely used in the 1700s. The Golden Ratio, approximately 1.618, is a proportion that is found in geometry, art and architecture and has been made famous in the illustrations of Leonardo da Vinci (1452–1519).

In the construction industry, concrete is made in different proportions of gravel, sand, cement and water depending on the particular application. Exchange rates allow business people and the general population to convert Australian dollars to an equivalent amount in an overseas currency. Artists, architects and designers use the concept of proportion in many spheres of their work.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 7.1 Pre-test

1. State whether this statement is True or False.

The expression $y \propto x$ means y is directly proportional to x .

2. **MC** Identify which of the following relationships is directly proportional.

A.

a	0	1	2	3
b	0	2	4	6

B.

a	0	1	2	3
b	0	1	4	9

C.

a	1	2	3	4
b	4	3	2	1

D.

a	0	1	2	3
b	1	3	5	7

E.

a	1	2	3	4
b	2	3	4	5

3. Mobile phone calls are charged at \$1 per minute. State if the cost of a phone bill and the number of 1-minute time periods are directly proportional.

4. State whether this statement is True or False.

The perimeter of a square and its side length have a relationship that is directly proportional.

5. **MC** If $a \propto b$ and $a = 3$ when $b = 9$, select the rule that links a and b .

A. $a = \frac{1}{3}b$ B. $a = 3b$ C. $a = \frac{3}{b}$ D. $a = 9b$ E. $3a = 9b$

6. The quantities x and y are related to each other by direct variation. If k is the constant of proportionality, complete the proportion statement $x = \underline{\hspace{2cm}}$.

7. The voltage, v , that powers a mobile phone is directly proportional to the current, I amps.

A mobile phone with consistent resistance uses a current of two amps when powered by a voltage of 5.1 volts.

a. Determine the consistent resistance or constant of proportionality, k .

b. Calculate the voltage required to operate a mobile phone that would use a current of 1.2 amps.

8. State whether this statement is True or False.

The statement ' x varies directly with y ' can be written as $y \propto \frac{1}{x}$ or $y = \frac{k}{x}$.

9. **MC** If y varies with the square of x and $y = 100$ when $x = 2$, then the constant of variation is:

A. 400 B. 200 C. 100 D. 50 E. 25

10. **MC** If b is directly proportional to c^2 , select the constant of variation if $b = 72$ when $c = 12$.

A. 0.5 B. 2 C. 6 D. 36 E. 144

11. **MC** y is directly proportional to x .
If x is tripled, select what y is.
- A. Doubled B. Tripled C. Reduced to one-third
D. Reduced to one-ninth E. Halved
12. **MC** On average, Noah kicks 2.25 goals per game of soccer. In a 16-game season, the number of goals Noah would be expected to kick is:
- A. 32 B. 36 C. 30 D. 38 E. 34
13. Tea bags in a supermarket can be bought for \$1.45 per pack of 10 or for \$3.85 per pack of 25. Determine the cheaper way of buying tea bags.
14. Given that $y \propto x$ and $\frac{y}{x} = 4$, determine the rule linking y and x .
15. Determine your average speed if you drive 320 km in five hours.

LESSON

7.2 Direct linear proportion

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- understand the concept of direct linear proportion
- recognise and write direct linear proportion statements.

7.2.1 Direct linear proportion

eles-4818


- Proportion is described by two equal ratios.
- In **direct linear proportion**, the two variables change at the same rate.
- Suppose that ice-cream costs \$3 and that you are to buy some for your friends.
- There is a relationship between the cost of the ice-cream (C) and the number of ice-creams that you buy (n).
- The relationship between the two variables (number of ice-creams and cost of the ice-cream) can be illustrated in a table or a graph.

n	0	1	2	3	4
C (\$)	0	3	6	9	12

- This relationship is called direct linear proportion as it shows the following characteristics:
 - As n increases, so does C .
 - The graph of the relationship is a straight line passing through the origin as $n = 0$ and $C = 0$.



- We say that ‘ C is directly proportional to n ’ or ‘ C varies directly as n ’.



Expressing direct linear proportion

If the y -values increase (or decrease) directly with the x -values, then this relationship can be expressed as:

$$y \propto x$$

where \propto is the symbol for proportionality — this means that y is directly proportional to x .

WORKED EXAMPLE 1 Checking for direct linear proportion

For each of the following pairs of variables, state whether direct linear proportion exists.

- The height of a stack of photocopy paper (h) and the number of sheets (n) in the stack.
- Your Maths mark (m) and the number of hours of Maths homework you have completed (n).

THINK

- When n increases, so does h .
When $n = 0$, $h = 0$.
If graphed, the relationship would be linear.
- As n increases, so does m .
When $n = 0$, I may get a low mark but it is not necessarily zero, so $m \neq 0$.



WRITE

- $h \propto n$
- m is not directly proportional to n .

WORKED EXAMPLE 2 Checking for direct linear proportion from a graph

For each of the following, determine whether direct linear proportion exists between the variables.


a.

t	0	1	2	3	4
y	0	1	3	7	15

b.

n	1	2	3	4
c	5	10	15	20

c.



THINK

- a. From the table, when t increases, so does y .
When $t = 0$, $y = 0$. The t -values increase by a constant amount but the y -values do not, so the relationship is not linear.
When t is doubled, y is not.
- b. From the table, as n increases, so does c .
Extending the pattern gives $n = 0$, $c = 0$. The n -values and C -values increase by constant amounts, so the relationship is linear.
- c. When x increases, so does y .
When $x = 0$, $y = 0$.
The graph is not a straight line.

WRITE

- a. y is not directly proportional to t .
- b. $C \propto n$
- c. y is not directly proportional to x .


Note: Not all direct proportion relationships are linear. Non-linear direct proportion will be studied in later years.

DISCUSSION

How do you know when two quantities are directly proportional?
Discuss the impact on image editing if proportion is not maintained.



 **eWorkbook** Topic 7 Workbook (worksheets, code puzzle and project) (ewbk-2007)

 **Interactivity** Individual pathway interactivity: Direct linear proportion (int-4510)
Direct linear proportion (int-2767)

Exercise 7.2 Direct linear proportion

learn**on**

7.2 Quick quiz **on**

7.2 Exercise

Individual pathways

PRACTISE

1, 3, 6, 9, 13

CONSOLIDATE

2, 5, 7, 10, 11, 14

MASTER

4, 8, 12, 15, 16

Fluency

1. **WE1** For each of the following pairs of variables, state whether direct linear proportion exists. If it does not exist, give a reason why.

- The distance (d) travelled in a car travelling at 60 km/h and the time taken (t).
- The speed of a swimmer (s) and the time the swimmer takes to complete one lap of the pool (t).
- The cost of a bus ticket (c) and the distance travelled (d).



2. For each of the following pairs of variables, state whether direct linear proportion exists. If it does not exist, give a reason why.

- The perimeter (p) of a square and the side length (l).
- The area of a square (A) and the side length (l).
- The total cost (C) and the number of boxes of pencils purchased (n).
- The length of a line in centimetres (c) and the length in millimetres (m).

3. For each of the following pairs of variables, state whether direct linear proportion exists. If it does not exist, give a reason why.

- The weight of an object in kilograms (k) and in pounds (p).
- The distance (d) travelled in a taxi and the cost (c).
- A person's height (h) and their age (a).
- The cost of an item in Australian dollars (A) and the cost of an item in Euros (E).

4. State whether the following statements are True or False.

There is a direct linear relationship between:

- the total cost, \$ C , of purchasing netballs and the number, n , purchased
- the circumference of a circle and its diameter
- the area of a semicircle and its radius.

5. **MC** For the table shown, if $y \propto x$, then the values of a and b would be:

x	1	2	4	8
y	3	6	a	b

- A.** $a = 8, b = 16$
D. $a = 12, b = 18$

- B.** $a = 8, b = 24$
E. $a = 12, b = 24$

- C.** $a = 9, b = 12$

Understanding

6. **WE2** For each of the following, determine whether direct linear proportion exists between the variables. If it does not, explain why.

a.

x	0	1	2	3	4
y	0	1	3	8	15

b.

a	0	1	2	3	6
M	0	8	16	24	48





c.

t	0	1	2	4	8
d	0	3	6	9	12

d.

n	0	1	2	3	4
C	10	20	30	40	50

7. For each of the following, determine whether direct proportion exists between the variables. If it does not, explain why.



8. List five pairs of real-life variables that exhibit direct linear proportion.

Reasoning

9. Explain which point must always exist in a table of values if the two variables exhibit direct linear proportionality.
10. If direct linear proportion exists between two variables m and n , fill out the table and explain your reasoning.

m	0	2	5
n	0		20

11. If the variables x and y in the following table are directly proportional, find the values of a , b and c . Explain your reasoning.

x	2	4	9	a
y	8	b	c	128

12. If $y \propto x$, explain what happens to:

a. y if x is doubled

b. y if x is halved

c. x if y is tripled.

Problem solving

13. Mobile phone calls are charged at 17 cents per 30 seconds.

a. Does direct linear proportion exist between the cost of a phone bill and the number of 30-second time periods? Justify your answer.

b. If a call went for 7.5 minutes, determine how much the call would cost.

14. An electrician charges \$55 for every 30 minutes or part of 30 minutes for his labour on a building site.

a. Does direct linear proportion exist between the cost of hiring the electrician and the time spent at the building site? Justify your answer.

b. If the electrician worked for 8.5 hours, determine how much this would cost.

15. A one-litre can of paint covers five square metres of wall.

a. Does a direct linear proportion exist between the number of litres purchased and the area of the walls to be painted? Justify your answer.

b. Evaluate the number of cans needed to paint a wall 5 metres long and 2.9 metres high with two coats of paint.

16. Bruce is building a pergola and needs to buy treated pine timber. He wants 4.2-metre and 5.4-metre lengths of timber. If a 4.2-metre length costs \$23.10 and a 5.4-metre length costs \$29.16, determine if direct linear proportion exists between the cost of the timber and the length of the timber per metre.

LESSON

7.3 Proportionality

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the constant of proportionality
- express the constant of proportionality as a ratio of the two quantities.



7.3.1 Constant of proportionality

eles-4819


- Proportionality relationships ($y \propto x$) are shown by the graph with four straight lines:

$$y = \frac{1}{2}x$$

$$y = x$$

$$y = 2x$$

$$y = 3x$$



- For the equation $y = kx$:
 - k is a constant, called the **constant of proportionality**
 - y is directly proportional to x , or $y \propto x$
 - all the graphs start at $(0, 0)$
 - y is called the *dependent variable* and is normally placed on the vertical axis (y -axis)
 - x is called the *independent variable* and is normally placed on the horizontal axis (x -axis).

WORKED EXAMPLE 3 Calculating the constant of proportionality

Given that $y \propto x$ and $y = 12$ when $x = 3$, calculate the constant of proportionality and state the rule linking y and x .

THINK

1. $y \propto x$, so write the linear rule.
2. Substitute $y = 12$ and $x = 3$ into $y = kx$.
3. Calculate the constant of proportionality by solving for k .
4. Write the rule.

WRITE

$$y = kx$$

$$12 = 3k$$

$$k = 4$$

$$y = 4x$$

WORKED EXAMPLE 4 Determining the direct proportion relationship

The weight (W) of \$1 coins in a bag is directly proportional to the number of coins (n). Twenty coins weigh 180 g.

- a. Determine the relationship between W and n .
- b. Calculate how much 57 coins weigh.
- c. Determine how many coins weigh 252 g.



THINK

Summarise the information given in a table.

- a. 1. Substitute $n = 20$ and $W = 180$ into $W = kn$.
2. Solve for k .
3. Write the relationship between W and n .
- b. 1. State the rule.
2. Substitute 57 for n to find W .

3. Write the answer in a sentence.
- c. 1. State the rule.
2. Substitute 252 for W .

3. Solve for n .

4. Write the answer in a sentence.

WRITE

$$W = kn$$

n	20	57	
W	180		252

- a. $180 = 20k$
 $k = 9$
 $W = 9n$
- b. $W = 9n$
 $W = 9 \times 57$
 $= 513$
Fifty-seven coins weigh 513 g.
- c. $W = 9n$
 $252 = 9n$
 $n = \frac{252}{9}$
 $= 28$
Twenty-eight coins weigh 252 g.

7.3.2 Ratio

eles-4820

- If $y \propto x$, then $y = kx$, where k is constant.
Transposing this formula gives $\frac{y}{x} = k$.
- The constant of proportionality, k , is the ratio of any pair of values (x, y) .
- For example, this table shows that $v \propto t$.

t	1	2	3	4
v	5	10	15	20

It is clear that $\frac{20}{4} = \frac{15}{3} = \frac{10}{2} = \frac{5}{1} = 5$, therefore, $k = 5$ and $v = 5t$.



WORKED EXAMPLE 5 Applying the constant of proportionality as a ratio

Sharon works part-time and is paid at a fixed rate per hour. If she earns \$135 for 6 hours' work, calculate how much she will earn for 11 hours.

THINK

- Sharon's payment is directly proportional to the number of hours worked. Write the rule.
- Summarise the information given. The value of x needs to be calculated.
- Since $p = kn$, $\frac{P}{n}$ is constant.
Solve for x .
- Write the answer in a sentence.

WRITE

$$P = kn$$

n	6	11
P	135	x

$$\frac{135}{6} = \frac{x}{11}$$

$$x = \frac{135 \times 11}{6}$$

$$x = 247.5$$

Sharon earns \$247.50 for 11 hours of work.

Resources



eWorkbook

Topic 7 Workbook (worksheets, code puzzle and project) (ewbk-2007)



Interactivities

Individual pathway interactivity: Proportionality (int-4511)

Constants of proportionality (int-6057)

7.3 Quick quiz **on**

7.3 Exercise

Individual pathways

PRACTISE

1, 2, 6, 7, 12, 15

CONSOLIDATE

3, 4, 8, 9, 13, 16

MASTER

5, 10, 11, 14, 17, 18

Fluency

- WE3** If a is directly proportional to b , and $a = 30$ when $b = 5$, calculate the constant of proportionality and state the rule linking a and b .
- If $a \propto b$, and $a = 2.5$ when $b = 5$, determine the rule linking a and b .
- If $C \propto t$, and $C = 100$ when $t = 8$, determine the rule linking C and t .
- If $v \propto t$, and $t = 20$ when $v = 10$, determine the rule linking v and t .
- If $F \propto a$, and $a = 40$ when $F = 100$, determine the rule linking a and F .

Understanding

- WE4** Springs are often used to weigh objects, because the extension of a spring (E) is directly proportional to the weight (W) of the object hanging from the spring. A 4-kg load stretches a spring by 2.5 cm.
 - Determine the relationship between E and W .
 - Calculate the load that will stretch the spring by 12 cm.
 - Determine how far 7 kg will extend the spring.
- Han finds that 40 shelled almonds weigh 52 g.
 - Determine the relationship between the weight (W) and the number of almonds (n).
 - Calculate how many almonds there would be in a 500 g bag.
 - Calculate how much 250 almonds would weigh.
- Petra knows that her bicycle wheel turns 40 times when she travels 100 m.
 - Determine the relationship between the distance travelled (d) and the number of turns of the wheel (n).
 - Calculate how far she goes if her wheel turns 807 times.
 - Determine how many times her wheel turns if she travels 5 km.
- Fiona, who operates a plant nursery, uses large quantities of potting mix. Last week she used 96 kg of potting mix to place 800 seedlings in medium-sized pots.
 - Determine the relationship between the mass of potting mix (M) and the number of seedlings (n).
 - Calculate how many seedlings she can pot with her remaining 54 kg of potting mix.
 - Determine how much potting mix she will need to pot 3000 more seedlings.
- WE5** Tamara is paid at a fixed rate per hour. If she earns \$136 for five hours of work, calculate how much she will earn for eight hours of work.
- It costs \$158 to buy 40 bags of birdseed. Calculate how much 55 bags will cost.



Reasoning

12. If 2.5 L of lawn fertiliser will cover an area of 150 m^2 , determine how much fertiliser is needed to cover an area of 800 m^2 at the same rate. Justify your answer.
13. Paul paid \$68.13 for 45 L of fuel. At the same rate, determine how much he would pay for 70 L. Justify your answer.
14. Rose gold is an alloy of gold and copper that is used to make high-quality musical instruments. If it takes 45 g of gold to produce 60 g of rose gold, evaluate how much gold would be needed to make 500 g of rose gold. Justify your answer.



Problem solving

15. If Noah takes a group of friends to the movies for his birthday and it would cost \$62.50 for five tickets, determine how much it would cost if there were 12 people (including Noah) in the group.
16. Sharyn enjoys quality chocolate, so she makes a trip to her favourite chocolate shop. She is able to select her favourite chocolates for \$7.50 per 150 grams. Since Sharyn loves her chocolate, she decides to purchase 675 grams. Evaluate how much she spent.
17. Anthony drives to Mildura, covering an average of 75 kilometers in 45 minutes. He has to travel 610 kilometres to get to Mildura.



- a. Determine the relationship between the distance he travelled in kilometres, D , and his driving time in hours, T .
 - b. Evaluate how long it will take Anthony to complete his trip.
 - c. If he stops at Bendigo, 220 kilometres from his starting position, determine how long it will take him to reach Bendigo.
 - d. Determine how much longer it will take him to arrive at Mildura after leaving Bendigo.
18. The volume of a bird's egg can be determined by the formula $V = kl^3$, where V is the volume in cm^3 , l is the length of the egg in cm and k is a constant. A typical ostrich egg is 15 cm long and has a volume of 7425 cm^3 . Evaluate the volume of a chicken egg that is 5 cm long.

LESSON

7.4 Introduction to rates

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the rate connecting two quantities
- apply rates to worded problems.

7.4.1 Rates

eles-4821

- The word **rate** occurs commonly in news reports and conversation.
'Home ownership rates are falling.'
'People work at different rates.'
'What is the current rate of inflation?'
'Do you offer a student rate?'
'The crime rate seems to be increasing.'
- Rate is a word often used when referring to a specific ratio.
- In general, to calculate a rate (or specific ratio), one quantity is divided by another.
- State the units for the rate, such as grams/litre (g/L) or kilometres/hour (km/h).



7.4.2 Average speed

eles-4822

- If you travel 160 km in 4 hours, your average speed is 40 km per hour.
- Speed is an example of a rate, and its unit of measurement contains a formula, such as kilometres per hour.
- The word 'per' can be replaced with 'divided by', so speed = $\frac{\text{distance (in km)}}{\text{time (in hours)}}$.
- It is important to note the units involved.
For example, an athlete's speed is often measured in metres per second (m/s) rather than km/h.

WORKED EXAMPLE 6 Calculating rates from statements

Calculate the rates suggested by these statements.

- Eight litres of fuel costs \$12.56.
- A cricket team scored 152 runs in 20 overs.
- A car travels 210 kilometres in 3 hours.

THINK

1. The rate suggested is 'dollars per litre'.
2. Substitute the values for the price per 8 litres and the number of litres.
3. Write the rate.

WRITE

$$\begin{aligned} \text{a. Rate} &= \frac{\text{number of dollars}}{\text{number of litres}} \\ &= \frac{12.56}{8} \\ &= \$1.57 \text{ per litre} \end{aligned}$$

b. 1. The rate suggested is 'runs per over'.

2. Substitute the values for the number of runs and the number of overs.

3. Write the rate.

$$\text{b. Rate} = \frac{\text{number of runs}}{\text{number of overs}}$$

$$= \frac{152}{20}$$

$$= 7.6 \text{ runs per over}$$

c. 1. The rate suggested is 'kilometres per hour', or the speed.

2. Substitute the values for the distance and the time.

3. Write the rate.

$$\text{c. Speed} = \frac{\text{distance}}{\text{time}}$$

$$= \frac{210}{3}$$

$$= 70 \text{ km per hour}$$

WORKED EXAMPLE 7 Calculating rates

The concentration of a solution is measured in g/L (grams per litre). Calculate the concentration of the solution when 10 g of salt is dissolved in 750 mL of water.

THINK

1. Concentration is measured in g/L, which means that concentration = number of grams (mass) ÷ number of litres (volume).
2. Substitute the values of mass and volume.
3. Simplify.
4. Write the rate in a sentence.

WRITE

$$\text{Concentration} = \frac{\text{mass (g)}}{\text{volume (L)}}$$

$$= \frac{10}{0.75}$$

$$= 13.3 \text{ g/L}$$

The concentration of the solution is 13.3 g/L

on Resources



eWorkbook Topic 7 Workbook (worksheets, code puzzle and project) (ewbk-2007)



Interactivities Individual pathway interactivity: Introduction to rates (int-4513)
Speed (int-6457)

7.4 Quick quiz

7.4 Exercise

Individual pathways

PRACTISE

1, 3, 6, 9

CONSOLIDATE

2, 5, 7, 10

MASTER

4, 8, 11

Fluency

- Hayden drove from Hay to Bee. He covered a total distance of 96 km and took 1.5 hours for the trip. Calculate Hayden's average speed for the journey.
- Calculate the average speed in km/h for each of the following.
 - 90 km in 45 min.
 - 5500 km in 3 h 15 min (correct to 1 decimal place).
- Calculate the rates suggested by these statements, giving your answers to 2 decimal places where necessary.
 - It costs \$736 for 8 theatre tickets.
 - Penelope decorated 72 small cakes in 3 hours.
 - Usain Bolt has a 100-metre world sprint record of 9.58 seconds.
 - It takes 30 hours to fill the swimming pool to a depth of 90 cm.
 - Peter received \$260 for 15 hours' work.
 - Yan received \$300 for assembling 6 air conditioners.



Understanding

- A metal bolt of volume 25 cm^3 has a mass of 100 g. Calculate its density (mass per unit of volume).
- One hundred and twenty grams of sugar is dissolved in 200 mL of water. Calculate the concentration of this solution in g/L. Explain your answer.

Reasoning

- In a race between a tortoise and a hare, the hare ran at 72 km per hour while the giant tortoise moved at 240 cm per minute. Compare and explain the difference in the speed of the two animals.
- A school had 300 students in 2013 and 450 students in 2015. Determine the average rate of growth in the number of students per year. Show your working.
- The average speed of a car is determined by the distance of the journey and the time the journey takes. Explain the two ways in which the speed can be increased.



Problem solving

9. Mt Feathertop is Victoria's second-highest peak. To walk to the top involves an increase in height of 1500 m over a horizontal distance of 10 km (10 000 m). Determine the average gradient of the track.
10. In the 2000 Sydney Olympics, Cathy Freeman won gold in the 400-metre race. Her time was 49.11 seconds. In the 2000 Beijing Olympics, Usain Bolt set a new world record for the men's 100-metre race. His time was 9.69 seconds. Evaluate the average speed of the winner for each race in kilometres per hour.
11. Beaches are sometimes unfit for swimming if heavy rain has washed pollution into the water. A beach is declared unsafe for swimming if the concentration of bacteria is more than 5000 organisms per litre. A sample of 20 millilitres was tested and found to contain 55 organisms. Evaluate the concentration in the sample (in organisms/litre) and state whether or not the beach should be closed.

LESSON

7.5 Constant and variable rates

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- interpret a graph showing constant rates
- draw a graph showing constant rates
- interpret a graph showing variable rates.



7.5.1 Constant rates


eles-4823

- A **constant rate** of change means the quantity always changes by the same amount in a given time.
- Consider a car travelling along a highway at a constant speed (rate) of 90 km/h. After one hour it will have travelled 90 km, as shown in the table.

Time (h)	0	1	2	3
Distance (km)	0	90	180	270

Diagram illustrating constant rates with arrows above the table showing +1 for each hour and below showing +90 for each hour.



- The distance–time graph is also shown.



- The gradient of the graph is $\frac{270-0}{3-0} = 90$.
- The equation of the graph is $d = 90t$, and the gradient is equal to the speed, or rate of progress.
- A constant rate of change produces a linear (straight-line) graph.

WORKED EXAMPLE 8 Describing constant rates from a graph

Each diagram shown illustrates the distance travelled by a car over time. Describe the journey, including the speed of the car.



THINK

- a. There are three distinct sections.
1. In the first two hours, the car travels 80 km: $\frac{80}{2} = 40$ km/h.
 2. In the next hour, the car does not move.
 3. In the fourth hour, the car travels 40 km.

- b. There are two distinct sections.
1. In the first two hours, the car travels 80 km: $\frac{80}{2} = 40$ km/h.
 2. In the next two hours, the car travels 20 km: $\frac{20}{2} = 10$ km/h.

WRITE

- a.
- The car travels at a speed of 40 km/h for 2 hours.
- The car stops for 1 hour.
- The car then travels for 1 hour at 40 km/h.

- b.
- The car travels at a speed of 40 km/h for 2 hours.
- The car then travels at 10 km/h for a further 2 hours.

WORKED EXAMPLE 9 Drawing a distance–time graph to illustrate constant rates

A cyclist travels for 1 hour at a constant speed of 10 km/h, then stops for a 30-minute break before riding a further 6 km for half an hour at a constant speed.

Draw a distance–time graph to illustrate the following journey.


THINK

There are three phases to the journey.

The graph starts at $(0, 0)$.

1. In the first hour, the cyclist travels 10 km. Draw a line segment from $(0, 0)$ to $(1, 10)$.
2. For the next half-hour, the cyclist is stationary, so draw a horizontal line segment from $(1, 10)$ to $(1.5, 10)$.
3. In the next half-hour, the cyclist travels 6 km. Draw a line segment from $(1.5, 10)$ to $(2, 16)$.

WRITE/DRAW






7.5.2 Variable rates

eles-4824

- A variable rate means a quantity does not change by the same amount in a given time.
- In reality, a car tends not to travel at constant speed. It starts from rest and gradually picks up speed.
- Since $\text{speed} = \frac{\text{distance}}{\text{time}}$, the


distance–time graph will:

- be flat, or horizontal, when the car is not moving (is stationary)
- be fairly flat when the car is moving slowly
- be steeper and steeper as the speed of the car increases.
- This is demonstrated in the distance–time graph shown.
- A variable rate of change produces a non-linear graph.



WORKED EXAMPLE 10 Describing graphs with variable rates

The diagram shown illustrates the distance travelled by a car over time. Describe what is happening, in terms of speed, at each of the marked points.



THINK

1. At point A on the graph, the gradient is small but becoming steeper.
2. At point B on the graph, the gradient is at its steepest and is not changing.
3. At point C, the graph is becoming flatter.
4. At point D, the graph is horizontal.

WRITE

At point A the car is travelling slowly but accelerating.



At point B the car is at its greatest speed.

At point C the car is slowing down.

At point D the car is stationary.

DISCUSSION

How can you tell the difference between constant and variable rates?

-  **eWorkbook** Topic 7 Workbook (worksheets, code puzzle and project) (ewbk-2007)
-  **Interactivities** Individual pathway interactivity: Constant and variable rates (int-4514)
 Constant rates (int-6060)
 Variable rates (int-6061)

Exercise 7.5 Constant and variable rates

7.5 Quick quiz **on**

7.5 Exercise

Individual pathways

PRACTISE

1, 3, 7, 10

CONSOLIDATE




2, 4, 8, 11

MASTER

5, 6, 9, 12, 13



Fluency

1. **WE8** Each diagram shown illustrates the distance travelled by a car over time. Describe the journey, including the speed of the car.



2. Two friends take part in a 24-kilometre mini-marathon. They run at constant speed. Ali takes 2 hours and Beth takes 3 hours to complete the journey.
- a. On the same diagram, draw a distance–time graph for each runner.
 - b. i. Calculate the equation for each graph.
 ii. State the difference between the two graphs.
3. **WE9** Draw a distance–time graph to illustrate each of the following journeys.
- a. A cyclist rides at 40 km/h for 30 minutes, stops for a 30-minute break, and then travels another 30 km at a speed of 15 km/h.
 - b. Zelko jogs at a speed of 10 km/h for one hour, and then at half the speed for another hour.

4. **WE10** The diagrams shown illustrate the distances travelled by two cars over time.



- Describe what is happening in terms of speed at each of the marked points.
- For each diagram, state at which point:
 - the speed is the greatest
 - the speed is the lowest
 - the car is stationary.

Understanding

5. The table below shows the distance travelled, D , as a person runs for R minutes.

R (minutes)	10	20	50
D (km)	2	4	10

- Calculate the rate in km/minute between the time 10 minutes and 20 minutes.
 - Determine the rate in km/minute between the distance 4 km and 10 km.
 - Explain whether the person's speed is constant.
6. The table below shows the water used, W , after the start of a shower, where T is the time after the shower was started.

T (minutes)	0	1	2	4
W (litres)	0	20	30	100

- Calculate the rate of water usage in L/minute for the four stages of the shower.
- Explain whether the rate of water usage was constant.




Reasoning

7. Margaret and Brian left Brisbane airport at 9:00 am. They travelled separately but on the same road and in the same direction.

Their journeys are represented by the travel graph below. Show your working.

- Determine the distance from the airport at which their paths crossed.
- Determine how far apart they were at 1:00 pm.
- Determine for how long each person stopped on the way.
- Determine the total time spent driving and the total distance for each person.
- Evaluate the average speed while driving for each person.



8. Hannah rode her bike along the bay one morning. She left home at 7:30 am and covered 12 km in the first hour. She felt tired and rested for half an hour. After resting she completed another 8 km in the next hour to reach her destination. Show your working.

- Determine how long Hannah took for the entire journey.
- Calculate the total distance for which she actually rode her bike.
- Draw a travel graph for Hannah's journey.

9. Rebecca and Joanne set off at the same time to jog 12 kilometres. Joanne ran the entire journey at a constant speed and finished at the same time as Rebecca.

Rebecca set out at 12 km/h, stopping after 30 minutes to let Joanne catch up, then she ran at a steady rate to complete the distance in 2 hours.

- Show the progress of the two runners on a distance–time graph.
- Determine how long Rebecca waited for Joanna to catch up. Show your working.




Problem solving


10. An internet service provider charges \$30 per month plus \$0.10 per megabyte downloaded. The table of monthly cost versus download amount is shown.

Download (MB)	0	100	200	300	400	500
Cost (\$)	30	40	50	60	70	80

- Determine how much the cost increases by when the download amount increases from:
 - 0 to 100 MB
 - 100 to 200 MB
 - 200 to 300 MB.
 - Explain whether the cost is increasing at a constant rate.
11. Use the graph showing the volume of water in a rainwater tank to answer the following questions.
- Determine the day(s) during which the rate of change was positive.
 - Determine the day(s) during which the rate of change was negative.
 - Determine the day(s) during which the rate of change was zero.
 - Determine the day on which the volume of water increased at the fastest rate.
 - Determine the day on which the volume of water decreased at the fastest rate.




12. The graph shows the number of soft-drink cans in a vending machine at the end of each day.



- Determine how much the number of cans changed by in the first day.
 - Determine how much the number of cans changed by in the fifth day.
 - Explain whether the number of soft-drink cans is changing at a constant or variable rate.
13. The table and graph show Melbourne's average daily maximum temperature over the year.

	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Mean maximum (°C)	25.8	25.8	23.8	20.3	16.7	14.0	13.4	14.9	17.2	19.6	21.9	24.2



- Determine the average maximum temperature in:
 - February
 - June.
- Calculate the change in temperature from:
 - January to August
 - November to December.
- Explain whether the temperature is changing at a constant rate.


LESSON

7.6 Review

7.6.1 Topic summary

Direct linear proportion


- If y is directly proportional to x , then:
 - y varies with the value of x
 - $y \propto x$
 - as x increases (or decreases), y increases (or decreases)
 - $x = 0$ and $y = 0$ results in a straight line.



- Not all direct proportion relationships are linear.

Constant of proportionality

- If y is directly proportional to x , then:
 - $y = kx$
 - k is a constant called the **constant of proportionality** — this is the ratio of any pair of values (x, y)
 - the gradient of the straight line is k
 - all graphs start at $(0, 0)$.
- y is called the dependent variable and is normally placed on the vertical axis.
- x is called the independent variable and is normally placed on the horizontal axis.




PROPORTION AND RATES

Rates

- A rate compares two quantities that have different units.
- Rates are special ratios.
- A rate is calculated by dividing one quantity with another.

Variable rates

- A variable rate means a quantity does not change by the same amount in a given time.
- Variable rates produce a non-linear graph.




Average speed

- Average speed is an example of a rate; it is calculated by dividing total distance by total time.
- Unit: metres per second (m/s) or kilometres per hour (km/h)
- Average speed = $\frac{\text{distance in kilometres}}{\text{time in hours}}$
e.g. if the distance travelled is 200 km in 4 hours, then
average speed = $\frac{200 \text{ km}}{4 \text{ hours}} = 50 \text{ km/h}$.

Constant rates




- A constant rate means a quantity always changes by the same amount in a given time.
- Constant rates produce a linear (straight line) graph.



7.6.2 Success criteria

Tick a column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
7.2	I can understand the concept of direct linear proportion.			
	I can recognise and write direct linear proportion statements.			
7.3	I can determine the constant of proportionality.			
	I can express the constant of proportionality as a ratio of the two quantities.			
7.4	I can calculate the rate connecting two quantities.			
	I can apply rates to a worded problem.			
7.5	I can interpret a graph showing constant rates.			
	I can draw a graph showing constant rates.			
	I can interpret a graph showing variable rates.			

7.6.3 Project

Fastest speeds

To calculate the average speed of a journey, we can use the speed formula:

$$\text{speed} = \frac{\text{distance}}{\text{time}}$$

Speed is a *rate*, because it compares two quantities of different units. Speed can be expressed in units such as km/h, km/min and m/s. The units of the quantities substituted into the numerator and denominator determine the final units of speed. In order to compare the speeds of different events, it is useful to convert them to the same units.

In 2008, the summer and winter Olympics were held. At the summer Olympics, sprinter Usain Bolt ran 100 m in 9.69 seconds. At the winter Olympics, cross-country skier Petter Northug covered 50 km in 2 hours and 5 minutes, and speed skater Mika Poutala covered 0.5 km in 34.86 seconds. Which competitor was the fastest? In order to answer this question, we need to determine the speed of each competitor. Since the information is quoted in a variety of units, we need to decide on a common unit for speed.



1. Calculate the speed of each athlete in m/s, correct to one decimal place.
2. Determine which athlete was the fastest.



Consider the speed of objects in the world around you. This task requires you to order the following objects from fastest to slowest, assuming that each object is travelling at its fastest possible speed:

electric car, submarine, diesel train, car ferry, skateboard, bowled cricket ball, windsurfer, served tennis ball, solar-powered car, motorcycle, aircraft carrier, Jaguar car, helicopter, airliner, rocket-powered car, the Concorde supersonic airliner.

3. From your personal understanding and experience, order the above list from fastest to slowest.

We can make a more informed judgement if we have facts available about the movement of each of these objects. Consider the following facts:

- When travelling at its fastest speed, the Concorde could cover a distance of about 7000 km in 3 hours.
 - The fastest airliner can cover a distance of 5174 km in 4 hours and a helicopter 600 km in 1.5 hours.
 - It would take 9.5 hours for a rocket-powered car to travel 9652 km, 3.25 hours for an electric car to travel 1280 km and 1.5 hours for a Jaguar car to travel 525 km.
 - On a sunny day, a solar-powered car can travel 39 km in half an hour.
 - In 15 minutes, a car ferry can travel 26.75 km, an aircraft carrier 14 km and a submarine 18.5 km.
 - It takes the fastest diesel train 0.42 hours to travel 100 km and a motorcyclist just 12 minutes to travel the same distance.
 - In perfect weather conditions, a skateboarder can travel 30 km in 20 minutes and a windsurfer can travel 42 km in 30 minutes.
 - A bowled cricket ball can travel 20 m in about 0.45 seconds, while a served tennis ball can travel 25 m in about 0.4 seconds.
4. Using the information above, decide on a common unit of speed and determine the speed of each of the objects.
 5. Order the above objects from fastest to slowest.
 6. Compare the order from question 5 with the list you made in question 3.
 7. Conduct an experiment with your classmates to record the times that objects take to cover a certain distance. For example, record the time it takes to run 100 m or throw a ball 20 m. Compare the speeds with the speeds of the objects calculated during this investigation.



eWorkbook Topic 7 Workbook (worksheets, code puzzle and project) (ewbk-2007)



Interactivities Crossword (int-2690)
Sudoku puzzle (int-3208)

Exercise 7.6 Review questions

Fluency


- MC** y is directly proportional to x and $y = 450$ when $x = 15$. The rule relating x and y is:
A. $y = 0.033x$ **B.** $y = 30x$ **C.** $y = 60x$ **D.** $y = 6750x$ **E.** $y = 0.3x$
- MC** If $y \propto x$ and $y = 10$ when $x = 50$, the constant of proportionality is:
A. 10 **B.** 5 **C.** 1 **D.** 0.2 **E.** 0.5
- MC** If $y \propto x$ and $y = 10$ when $x = 50$, the value of x when $y = 12$ is:
A. 6 **B.** 60 **C.** 40 **D.** 2.4 **E.** 24
- In the table below, if $y \propto x$, state the rule for the relationship between x and y .

x	0	1	2	3
y	0	4	8	12

- In the table below, if $q \propto p$, state the rule for the relationship between p and q .


p	0	3	6	9
q	0	1	2	3

- In the graph below, y varies directly as x . State the rule for the relationship between x and y .




- MC** A speed of 60 km/h is equivalent to approximately:
A. 17 m/min **B.** 1 km/s **C.** 1 m/s
D. 17 m/s **E.** 60 m/s

8. **MC** A metal part has a density of 37 g/mm^3 . If its volume is 6 mm^3 , it has a mass of:
 A. 6.1 g B. 6.1 kg C. 222 kg
 D. 222 g E. 22.2 g
9. **MC** Identify which of the following is *not* a rate.
 A. 50 km/h B. 70 beats/min C. 40 kg
 D. Gradient E. Slope
10. **MC** Calculate how long it takes to travel 240 km at an average speed of 60 km/h.
 A. 0.25 hours B. 240 minutes C. 180 hours
 D. 14 400 hours E. 4 minutes
11. **MC** A plane takes two hours to travel 1600 km. Calculate how long it takes to travel 1000 km.
 A. 150 minutes B. 48 minutes C. 2 hours
 D. 75 minutes E. 85 minutes
12. **MC** The graph shows Sandy's bank balance. Determine the length of time (in months) that Sandy has been saving.



- A. 2 months B. 5 months C. 10 months D. 12 months E. 8 months

13. True or False: The following graph shows that $y \propto x$. Give your reason.



14. If $w \propto v$, and $w = 7.5$ when $v = 5$, calculate k , the constant of proportionality.

15. The wages earned by a worker for different numbers of hours are shown in the table.

Time (hours)	1	2	3	4
Wages (\$)	25	50	75	100

Determine the rule relating wages, \$ w , to the hours worked, t , if $w \propto t$. Give your answer in hours.

16. Calculate the missing quantities in the table.

Mass	Volume	Density
500 g	20 cm ³	
1500 g		50 g/cm ³
	120 cm ³	17 g/cm ³

17. Answer the following questions.


- A 2000-litre water tank takes two days to fill. Express this rate in litres per hour correct to 2 decimal places.
- Determine how far you have to travel vertically to travel 600 metres horizontally if the gradient of the track is 0.3.
- Determine how long it takes for a 60-watt (60 joules/second) light globe to use 100 kilojoules.
- Evaluate the cost of 2.3 m³ of sand at \$40 per m³.

Problem solving

- Lisa drove to the city from her school. She covered a distance of 180 km in two hours.
 - Calculate Lisa's average speed.
 - Lisa travelled back at an average speed of 60 km/h. Determine the travel time.
- Seventy grams of ammonium sulfate crystals are dissolved in 0.5 L of water.
 - Calculate the concentration of the solution in g/mL.
 - Another 500 mL of water is added. Determine the concentration of the solution now.
- A skyscraper can be built at a rate of 4.5 storeys per month.
 - Determine how many storeys will be built in 6 months.
 - Determine how many storeys will be built in 24 months.
- A certain kind of eucalyptus tree grows at a linear rate for its first two years of growth. If the growth rate is 5 cm per month, determine how long it will take to grow to a height of 1.07 m.
- The pressure inside a boiler increases as the temperature increases. For each 1°C, the pressure increases by 10 units. At a temperature of 100°C, the pressure is 600 units. If the boiler can withstand a pressure of 2000 units, determine the temperature at which this pressure is reached.
- Hector has a part-time job as a waiter at a local cafe and is paid \$8.50 per hour. Complete the table of values relating the amount of money Hector receives to the number of hours he works.

Number of hours	0	2	4	6	8	10
Pay						

24. Karina left home at 9:00 am. She spent some time at a friend's house, then travelled to the airport to pick up her sister. She then travelled straight back home. Her journey is shown by the travel graph.



- Determine how far Karina's house is from her friend's house.
 - Determine how much time Karina spent at her friend's place.
 - Calculate how far the airport is from Karina's house.
 - Determine how much time Karina spent at the airport.
 - Calculate how much time Karina took to drive home.
 - Calculate the average speed of Karina's journey:
 - from her home to her friend's place
 - from her friend's place to the airport
 - from the airport to her home.
25. A commercial aircraft covers a distance of 1700 km in 2 hours 5 minutes. Evaluate the distance in metres that the aircraft travels each second.
26. A fun park charges a \$10 entry fee and an additional \$3 per ride. Complete the following table of values relating the total cost to the number of rides.

Rides	0	2	4	6	8	10
Cost						



Use the following information to answer questions 27–30.

To help compare speed in different units, a length conversion chart is useful. Time is measured in the same units throughout the world, so a time conversion chart is not necessary.

Length conversion chart

Imperial unit	Conversion factor	Metric unit
Inches (in)	25.4	mm
Feet (ft)	30.5	cm
Yards (yd)	0.915	m
Miles (mi)	1.61	km

To convert an imperial length unit into its equivalent metric unit, multiply by the conversion factor. Divide by the conversion factor when converting from metric to imperial units.

27. a. Use the table to complete each of the following.

i. $12 \text{ in} = \underline{\hspace{1cm}} \text{ mm}$

ii. $3 \text{ ft} = \underline{\hspace{1cm}} \text{ mm}$

iii. $1 \text{ m} = \underline{\hspace{1cm}} \text{ yd}$

iv. $1 \text{ km} = \underline{\hspace{1cm}} \text{ mi}$

b. Determine which is the faster: a car travelling at 100 km/h or a car travelling at 100 miles/h.

28. The speed limit of 60 km/h in Australia would be equivalent to a speed limit of $\underline{\hspace{1cm}}$ miles/h in USA. Give your answer correct to three decimal places.

29. A launched rocket covers a distance of 17 miles in 10 seconds. Calculate its speed in km/h.

30. Here are some record speeds for moving objects.

Object	Speed
Motorcycle	149 m/sec
Train	302 miles/h
Human skiing	244 km/h
Bullet from 38-calibre revolver	4000 ft/sec

Convert these speeds to the same unit. Then place them in order from fastest to slowest.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 7 Proportion and rates

7.1 Pre-test

1. True
2. A
3. Yes
4. True
5. A
6. $x = \frac{k}{y}$
7. a. $k = 2.55$ b. $v = 3.06$ amps
8. False
9. E
10. A
11. B
12. B
13. \$1.45 for a pack of 10
14. $y = 4x$
15. 64 km/h

7.2 Direct linear proportion

1. a. Yes
b. No; as speed increases, time decreases.
c. No; doubling distance doesn't double the cost.
2. a. Yes
b. No; doubling the side length doesn't double the area.
c. Yes
d. Yes
3. a. Yes
b. No; doubling distance doesn't double the cost (due to the initial fee).
c. No; doubling age doesn't double height.
d. Yes
4. a. True: as the number, n , increases, so does the cost, \$ C .
b. True: $C = \pi D$, as C increases with D . π is a constant number.
c. False: $A = \frac{1}{2}\pi r^2$, so it is not linear.
5. E
6. a. No; as the value of x increases, the value of y does not increase by a constant amount.
b. Yes; as the value of a increases, the value of M increases by a constant amount. For every increase in a by 1, the value of M increases by 8.
c. No; as the value of x increases, the value of y does not increase by a constant amount.
d. No; when $n = 0$, C does not equal 0.
7. a. Yes
b. No; when $t = 0$, d does not.
c. No; as x doubles, y does not.
d. No; when $d = 0$, w does not.
8. Sample responses can be found in the worked solutions in the online resources.

9. The point (0, 0) must always exist in a table of values of two variables exhibiting direct proportionality.
10. The missing value is 8. The relationship between n and m can be calculated by the values 5 and 20 in the table ($n = 4m$).
11. $a = 32, b = 16, c = 36$
12. a. y is doubled.
b. y is halved.
c. x is tripled.
13. a. Yes, direct proportion does exist.
b. \$2.55
14. a. Yes b. \$935
15. a. Yes, direct linear proportion does exist.
b. 6 cans of paint
16. Direct proportion does not exist. The price per metre for the 4.2-metre length is \$5.50, and the price per metre for the 5.4-metre length is \$5.40.

7.3 Proportionality

1. $k = 6, a = 6b$
2. $a = 0.5b$
3. $C = 12.5t$
4. $v = 0.5t$
5. $F = 2.5a$
6. a. $E = 0.625W$ b. 19.2 kg c. 4.375 cm
7. a. $W = 1.3n$
b. ≈ 385 almonds
c. 325 g
8. a. $d = 2.5n$ b. 2017.5 m c. 2000 turns
9. a. $M = 0.12n$ b. 450 seedlings c. 360 kg
10. \$217.60
11. \$217.25
12. 13.33 L
13. \$105.98
14. 375 g
15. \$150
16. \$33.75
17. a. $D = 100T$ c. 2 hours 12 minutes
b. 6 hours 6 minutes d. 3 hours 54 minutes
18. 275 cm^3

7.4 Introduction to rates

1. 64 km/h
2. a. 120 km/h
b. 1692.3 km/h
3. a. \$92 per ticket b. 24 cakes per hour
c. 10.44 m/s d. 3 cm/h
e. \$17.33/h f. \$50 per air conditioner
4. 4 g/cm^3
5. 600 g/L
6. The hare runs at 120 000 cm per minute; the tortoise runs at 0.144 km per hour. The hare runs 500 times faster than the tortoise.
7. 75 students/year

8. Either the distance of the journey increases and the time remains constant, or the distance of the journey remains constant and the time decreases.

9. 0.15


10. Cathy Freeman: 29.32 km/h; Usain Bolt: 37.15 km/h

11. 2750 organisms/litre. The beach should not be closed.

7.5 Constant and variable rates


1. a. 1 hour at 60 km/h, then 1 hour at 120 km/h
- b. 1 hour at 50 km/h, a 30-minute stop, 30 minutes at 100 km/h, a 1-hour stop, then 1 hour at 100 km/h
- c. 1 hour at 20 km/h, 1 hour at 100 km/h, 1 hour at 40 km/h

2. a. d (km)




- b. i. A: $d = 12t$; B: $d = 8t$
- ii. A has a steeper gradient than B.

3. a. d (km)



b. d (km)



4. a. A. The car is moving with steady speed.
- B. The car is momentarily stationary.
- C. The speed is increasing.
- D. The car is slowing down.
- E. The car is moving at a slow steady speed.

F. The speed is increasing.

G. The car is moving at a faster steady speed.

b. i. A, G ii. B, E iii. B

5. a. 0.2 km/min

b. 0.2 km/min

c. Yes, both rates are the same.

6. a. 20, 10, 35, 0 L/min

b. No

7. a. 150 km, 200 km and 250 km

b. 50 km

c. Both stop for 1 hour.


d. Brian — 300 km, 3.5 h; Margaret — 300 km, 4 h

e. Brian — 85.7 km/h; Margaret — 75 km/h


8. a. 2.5 hours

b. 20 km

c.



9. a. d (km)



b. 30 minutes

10. a. i. \$10

 ii. \$10

 iii. \$10

b. Yes

11. a. 1, 3, 5, 10

b. 4, 6, 7, 9

c. 2, 8

d. 10

e. 4

12. a. 30

b. 30

c. Variable

13. a. i. 25.8 °C

 ii. 14.0 °C

b. i. 10.9 °C

 ii. 2.3 °C

c. Sample responses can be found in the worked solutions in the online resources.

Project

- Usain Bolt = 10.3 m/s
Petter Northug = 6.7 m/s
Mika Poutala = 14.3 m/s
- Mika Poutala
- Answers will vary. Sample response: the Concorde supersonic airliner, rocket-powered car, Jaguar motorcar, helicopter, airliner, motorcycle, electric car, diesel train, served tennis ball, car ferry, skateboard, bowled cricket ball, windsurfer, submarine, aircraft carrier, solar-powered car.
- Concorde: 2333.3 km/h
Airliner: 1293.5 km/h
Helicopter: 400 km/h
Rocket-powered car: 1016 km/h
Electric car: 393.8 km/h
Jaguar car: 350 km/h
Solar-powered car: 78 km/h
Car ferry: 107 km/h
Aircraft carrier: 56 km/h
Submarine: 74 km/h
Diesel train: 238.1 km/h
Motorcyclist: 500 km/h
Skateboarder: 90 km/h
Windsurfer: 84 km/h
Bowled cricket ball: 160 km/h
Served tennis ball: 225 km/h
- Concorde, airliner, rocket-powered car, motorcyclist, helicopter, electric car, Jaguar car, diesel train, served tennis ball, bowled cricket ball, oar ferry, skateboarder, windsurfer, solar-powered car, submarine, aircraft carrier
- Answers will vary. Concorde was the fastest and the submarine was the slowest.
- Answers will vary. Students could measure the speed of different objects by measuring the distance covered in certain time.

7.6 Review questions

- B
- D
- B
- $y = 4x$
- $q = \frac{1}{3}p$
- $y = 2.5x$
- D
- D
- C
- B
- D
- C
- False. The graph starts at (0, 0) and increases, but it is not linear. Therefore, y does not vary directly with x .
- 1.5
- $w = 25t$

Mass	Volume	Density
500 g	20 cm ³	25 g/cm ³
1500 g	30 cm ³	50 g/cm ³
2040 g	120 cm ³	17 g/cm ³

- a. 41.67 L/h
c. 28 minutes
- b. 180 m
d. \$92
- a. 90 km/h
- a. 0.14 g/mL
- a. 27
- 21.4 months
- 240 °C

Number of hours	0	2	4	6	8	10
Pay	\$0	\$17	\$34	\$51	\$68	\$85

- a. 20 km
b. 15 min
c. 40 km
d. 30 min
e. 30 min
f. i. 80 km/h
ii. 40 km/h
iii. 80 km/h
- $226 \frac{2}{3}$ m/sec
- | Rides | 0 | 2 | 4 | 6 | 8 | 10 |
|-------|------|------|------|------|------|------|
| Cost | \$10 | \$16 | \$22 | \$28 | \$34 | \$40 |
- a. i. 304.8 mm
iii. 1.09 yd
b. The one travelling at 100 miles/h
- 37.267 miles/h
- 9853.2 km/h
- The conversions may vary, but the order from fastest to slowest is:
Bullet from 38-calibre revolver
Motorcycle
Train
Human skiing

8 Financial mathematics (extending)

LESSON SEQUENCE

8.1 Overview	448
8.2 Salaries and wages	451
8.3 Penalty rates	455
8.4 Simple interest	461
8.5 Compound interest	469
8.6 Review	478



LESSON

8.1 Overview

8.1.1 Why learn this?

There are many famous sayings about money. You have probably heard that ‘money doesn’t grow on trees’, ‘money can’t buy happiness’ and ‘a fool and his money are soon parted’. Most of us have to be conscious of where our money comes from and where it goes to. Understanding the basic principles of finance is very helpful for managing everyday life. Some of you will already be earning money from a part-time or casual job. Do you know what you spend your money on, or where it goes? Do you have your own savings account? Many of you may already be saving for a new games console, your first car or your dream holiday.

In this topic you will investigate different kinds of employment as well as different investment options for saving your money. Every branch of industry and business, whether large or small, international or domestic, will have to pay their employees either an annual salary or a wage based on an hourly rate. Financial investments are another way for businesses and individuals to make money — it is important to understand how investments work to be able to decide whether an investment is a good idea or not.

A sound knowledge of financial mathematics is essential in a range of careers, including financial consultancy, accountancy, business management and pay administration.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities



Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 8.1 Pre-test

- Kira has an annual salary of \$85 450.70 (without tax). Calculate how much she is paid:
 - weekly
 - fortnightly
 - monthly.
- Ishmael has a casual job at a department store. He is paid \$11.50 per hour. He gets double time on a Sunday. Calculate his wage for a week in which he worked from 4 pm to 8 pm on Thursday and 12 pm to 6 pm on Sunday.
- Jose is a tutor. He receives \$975 for a week in which he works 30 hours. Calculate his hourly rate of pay.
- MC** This table shows a timesheet for Yumi, who works in an electronics store.

Day	Pay rate	Start time	Finish time
Monday	Normal	9 am	1 pm
Tuesday	Normal	9 am	5 pm
Wednesday	Normal	11 am	7 pm
Thursday	Normal	2 pm	7 pm
Friday	Normal	4 pm	8 pm
Saturday	Time-and-a-half	9 am	2 pm
Sunday	Double	12 pm	6 pm

If Yumi's normal hourly rate is \$15.30, the amount she earned for the week is:

- A.** \$612.00 **B.** \$734.00 **C.** \$818.55 **D.** \$742.05 **E.** \$902.70
- Anita was offered a job with a salary of \$65 000 per year, or \$32.50 per hour. Assuming that she works a 40-hour week, compare the two rates to decide which is the better pay.
 - Ali earns \$20.40 an hour. He needs to earn a minimum of \$710 each week for his living expenses. Calculate the minimum number of hours he has to work each week. Give your answer to the nearest hour.
 - Mawluda is a casual worker in a café. She is paid time-and-a-half for working on Saturday and double time for working on Sunday. Her normal pay rate is \$18.75 an hour. Over the weekend, she worked 8 hours on Saturday and 5 hours on Sunday. Calculate how much Mawluda earned over the weekend.



8. A bank has a special offer on a savings account of 3% p.a. simple interest. Peter opens an account with \$800 and leaves his money there for 5 months.
- Calculate how much interest he earned over the 5 months.
 - Calculate the balance of Peter's account after the 5 months.
9. **MC** If an investment of \$40 000 pays 8% simple interest per year, then the value of the investment at the end of 3 years is:
- A. \$3 200 B. \$9 600 C. \$43 200 D. \$49 600 E. \$50 390

10. The Liang family wants to buy a family car at a cost of \$22 700. They pay a deposit of \$4500 and borrow the balance at an interest rate of 11.5% p.a. The loan will be paid off with 48 equal monthly payments.
- Calculate how much interest they will pay.
 - Calculate the total cost of the car, including the interest paid.
 - Calculate the amount of each repayment.



11. **MC** If \$10 000 is invested at a simple interest rate of 6% per annum, select the correct formula for calculating the interest earned for 6 months.
- A. $I = 10\,000 \times \frac{6}{100} \times 6$ B. $I = 10\,000 \times 0.6 \times 6$ C. $I = 10\,000 \times \frac{6}{100} \times \frac{1}{2}$
- D. $I = 10\,000 \times (1.6)^6$ E. $I = 10\,000 \times (1.06)^6$
12. If the simple interest charged on a loan of \$6400 over 30 months was \$1450, calculate the percentage rate of interest charged. Give your answer to the nearest whole percentage.
13. **MC** The amount of interest earned if \$112 000 is invested for 4 years at 6.8% p.a. interest compounded annually is:
- A. \$23 947.14 B. \$30 464.00 C. \$33 714.59 D. \$142 464.00 E. \$145 714.59
14. **MC** The total value of \$3500 invested for 18 months at 12% p.a. compounding monthly can be calculated using:
- A. $A = 3500(1.12)^{18}$ B. $A = 3500(1.12)^{1.5}$ C. $A = 3500(1.01)^{18}$
- D. $A = 3500(1.01)^{1.5}$ E. $A = 3500 \times \frac{12}{100} \times 18.$
15. Alecia invests \$5000 at 15% p.a. compounded annually and Cooper invests \$5000 at a 15% p.a. flat rate. Compare the two investments over a period of 10 years to determine how much Alecia's investment will be worth compared to Cooper's.

LESSON

8.2 Salaries and wages

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- understand the difference between a salary and a wage
- calculate weekly, fortnightly or monthly pays from a given annual salary
- calculate the amount earned from the hours worked and the hourly rate of pay.

8.2.1 Employees' salaries and wages

eles-4836

- Employees may be paid for their work in a variety of ways — most receive either a **wage** or a **salary**.



Key points

- **Normal working hours in Australia are 38 hours per week.**
- **There are 52 weeks in a year.**
- **There are 26 fortnights in a year (this value is slightly different for a leap year).**
- **There are 12 months in a year.**

WORKED EXAMPLE 1 Calculating pay from annual salary

Susan has an annual salary of \$63 048.92. Calculate how much she is paid:

- a. weekly b. fortnightly c. monthly.

THINK

- a. 1. Annual means per year, so divide the salary by 52 because there are 52 weeks in a year.
2. Write the answer in a sentence.
- b. 1. There are 26 fortnights in a year, so divide the salary by 26.
2. Write the answer in a sentence.
- c. 1. There are 12 months in a year, so divide the salary by 12.
2. Write the answer in a sentence.

WRITE

- a. Weekly salary = $63\,048.92 \div 52$
 ≈ 1212.48
Susan's weekly salary is \$1212.48.
- b. Fortnightly salary = $63\,048.92 \div 26$
 ≈ 2424.96
Susan's fortnightly salary is \$2424.96.
- c. Monthly salary = $63\,048.92 \div 12$
 ≈ 5254.08
Susan's monthly salary is \$5254.08.

WORKED EXAMPLE 2 Calculating wage given hourly rate of pay

Frisco has casual work at a fast-food store. He is paid \$12.27 per hour Monday to Saturday and \$24.54 per hour on Sunday.

Calculate his wage for a week in which he worked from 5:00 pm to 10:00 pm on Friday and from 6:00 pm to 9:00 pm on Sunday.

THINK

1. Work out the number of hours Frisco worked each day.
He worked 5 hours on Friday and 3 hours on Sunday.
2. Calculate the total amount earned by adding the wages earned on Friday and Sunday.
3. Write the answer in a sentence.

WRITE

- Friday: $5 \times 12.27 = 61.35$
Sunday: $3 \times 24.54 = 73.62$
 $61.35 + 73.62 = 134.97$
Frisco's wage was \$134.97.

Resources



eWorkbook

Topic 8 Workbook (worksheets, code puzzle and project) (ewbk-2008)



Interactivities

Individual pathway interactivity: Salaries and wages (int-4520)
Salaries (int-6067)

Individual pathways

PRACTISE

1, 3, 7, 10, 13, 14, 17

CONSOLIDATE

2, 4, 8, 11, 15, 18

MASTER

5, 6, 9, 12, 16, 19, 20

Fluency

- WE1** Johann has an annual salary of \$57 482. Calculate how much he is paid:
 - weekly
 - fortnightly
 - monthly.
- Nigara earns \$62 300 per annum. Calculate how much she earns:
 - weekly
 - fortnightly
 - monthly.
- Calculate the annual salary of workers with the following weekly incomes.
 - \$368
 - \$892.50
 - \$1320.85
- Calculate how much is earned per annum by people who are paid fortnightly salaries of:
 - \$995
 - \$1622.46
 - \$3865.31.
- Compare the following salaries, and determine which of each pair is the higher.
 - \$3890 per month or \$45 700 per annum.
 - \$3200.58 per fortnight or \$6700 per month.
- Calculate the hourly rate for these workers.
 - Rahni earns \$98.75 for working 5 hours.
 - Francisco is paid \$54.75 for working $4\frac{1}{2}$ hours.
 - Nhan earns \$977.74 for working a 38-hour week.
 - Jessica works $7\frac{1}{2}$ hours a day for 5 days to earn \$1464.75.

Understanding

- Henry is a second-year apprentice motor mechanic. He receives an award wage of \$12.08 per hour. Jenny, a fourth-year apprentice, earns \$17.65 per hour.
 - Calculate how much Henry earns in a 38-hour week.
 - Determine how much more Jenny earns in the same period of time.



- WE2** Juan has casual work for which he is paid \$13.17 per hour Monday to Saturday and \$26.34 per hour on Sundays. Calculate his total pay for a week in which he worked from 11 am to 5 pm on Thursday and from 2 pm to 7 pm on Sunday.

9. Mimi worked the following hours in one week.

Wednesday	5 pm to 9 pm
Thursday	6 pm to 9 pm
Friday	7 pm to 11 pm

If her pay is \$21.79 per hour up to 9 pm and \$32.69 per hour after that, calculate her total pay for that week.

10. Decide who earns more money each week: Rhonda, who receives \$38.55 an hour for 38 hours of work, or Rob, who receives \$41.87 an hour for 36 hours of work.
11. Glenn is a chef. He receives \$1076.92 for a week in which he works 35 hours. Calculate his hourly rate of pay.
12. Zack and Kaylah work in different department stores. Zack is paid \$981.77 per week. Kaylah is paid \$26.36 per hour. Calculate how many hours Kaylah must work to earn more money than Zack.
13. Calculate what pay each of the following salary earners will receive for each of the periods specified.
- Annual salary \$83 500, paid each week.
 - Annual salary \$72 509, paid each fortnight.
 - Annual salary \$57 200, paid each week.
 - Annual salary \$105 240, paid each month.



Reasoning

14. **MC** When Jack was successful in getting a job as a trainee journalist, he was offered the following choice of four salary packages. Decide which salary package Jack should choose, and justify your answer.
- \$456 per week
 - \$915 per fortnight
 - \$1980 per calendar month
 - \$23 700 per year
 - \$356 per week



15. Julie is considering two job offers for work as a receptionist. Job A pays \$878.56 for a 38-hour working week. Job B pays \$812.16 for a 36-hour working week. Determine which job has the higher hourly rate of pay. Justify your answer.
16. In his job as a bookkeeper, Minh works 38 hours per week and is paid \$32.26 per hour. Michelle, who works 38 hours per week in a similar job, is paid a salary of \$55 280 per year. Decide who has the higher-paying job. Justify your answer.

Problem solving

17. A lawyer is offered a choice between two jobs: one with a salary of \$74 000 per year and a second that pays \$40 per hour. Assuming that the lawyer will work 80 hours every fortnight, decide which job pays the most. Justify your answer.
18. Over the last four weeks Shahni has worked 35, 36, 34 and 41 hours. If she earns \$24.45 per hour, calculate how much she earned for each of the two fortnights.

19. Jackson works a 40-hour week (8 hours a day, Monday–Friday) and earns \$62 000 per annum.
- Calculate Jackson’s hourly rate.
 - If he works on average an extra half an hour per day from Monday to Friday and then another 4 hours over the weekend (for the same annual salary), explain how his hourly rate is affected.
 - If Jackson was earning the hourly rate and was being paid for every hour worked, calculate his potential earnings for the year.
20. Mark saves \$10 per week. Phil saves 5 cents in the first week, 10 cents the second week, and doubles the amount each week from that time on. Determine how many weeks it will take for Phil’s savings to be worth more than Mark’s.

LESSON

8.3 Penalty rates

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate total wages when overtime or penalty rates are included
- calculate hours paid when time-and-a-half or double time is involved.



eles-4837

8.3.1 Overtime and penalty rates

- **Overtime** is paid when a wage earner works more than their regular hours each week. These additional payments are often referred to as penalty rates.
- Penalty rates are usually paid for working on weekends, public holidays or at night.
- The extra hours are paid at a higher hourly rate, normally calculated at either time-and-a-half or double time.

Calculating overtime

The **overtime** hourly rate is usually a multiple of the regular hourly rate.

Some examples of overtime rates include:

- $1.5 \times$ regular hourly wage (time-and-a-half)
- $2 \times$ regular hourly wage (double time)
- $2.5 \times$ regular hourly wage (double time-and-a-half).



WORKED EXAMPLE 3 Calculating wages with overtime

Ursula works in hospitality and earns \$23.30 per hour. Last week she received the normal rate for 30 hours of work as well as time-and-a-half for 3 hours of overtime and double time for 5 hours of work on Sunday. Calculate her total pay.



THINK

1. Calculate Ursula's normal pay for 30 hours.
2. Calculate Ursula's pay for 3 hours at time-and-a-half ($1.5 \times$ regular hourly wage)
3. Calculate Ursula's pay for 5 hours at double time ($2 \times$ regular hourly wage).
4. Calculate the total amount by adding normal pay and overtime pay.
5. Write the answer in a sentence.

WRITE

$$\text{Normal pay: } 30 \times 23.30 = 699.00$$

$$\text{Overtime: } 3 \times 1.5 \times 23.30 = 104.85$$

$$\text{Sunday: } 5 \times 2 \times 23.30 = 233.00$$

$$\begin{aligned} \text{Total} &= 699.00 + 104.85 + 233.00 \\ &= 1036.85 \end{aligned}$$

Ursula's total pay was \$1036.85.

8.3.2 Time sheets and pay slips

eles-4398

- Employers often use records to monitor the number of working hours of their employees.



WORKED EXAMPLE 4 Calculating pay slips showing overtime

Fiona works in a department store, and in the week before Christmas she works overtime. Her time sheet is shown. Fill in the details on her pay slip.

	Start	Finish	Normal hours	Over time 1.5	Pay slip for: Fiona Lee	Week ending December 21
M	9:00	15:00	6		Normal hours	
T	9:00	17:00	8		Normal rate	\$17.95
W	9:00	17:00	8			
T	9:00	19:00	8	2	Overtime hours	
F	9:00	19:00	8	2	Overtime rate	
S					Total wage	

THINK

- Calculate the number of normal hours worked by adding the hours worked on each day of the week.
- Calculate the number of overtime hours worked by adding the overtime hours worked on Thursday and Friday.
- Calculate the overtime rate ($1.5 \times$ regular hourly wage).
- Calculate the total pay by multiplying the number of normal hours by the normal rate and adding the overtime amount (calculated by multiplying the number of overtime hours by the overtime rate).
- Fill in the amounts on the pay slip.

WRITE

$$\text{Normal hours: } 6 + 8 + 8 + 8 + 8 = 38$$

$$\text{Overtime hours: } 2 + 2 = 4$$



$$\begin{aligned} \text{Overtime rate} &= 1.5 \times 17.95 \\ &= 26.93 \end{aligned}$$

$$\begin{aligned} \text{Total pay} &= 38 \times 17.95 + 4 \times 26.93 \\ &= 789.82 \end{aligned}$$

Pay slip for: Fiona Lee	Week ending December 21
Normal hours	38
Normal rate	\$17.95
Overtime hours	4
Overtime rate	\$26.93
Total wage	\$789.82

COLLABORATIVE TASK: Comparing pay rates and conditions

Working in small groups, collect several job advertisements from a mixture of print and digital sources. Compare the pay rates and conditions for the different positions and present a report to the class.

-  **eWorkbook** Topic 8 Workbook (worksheets, code puzzle and project) (ewbk-2008)
-  **Interactivities** Individual pathway interactivity: Penalty rates (int-4521)
Special rates (int-6068)

Exercise 8.3 Penalty rates

8.3 Quick quiz **on**

8.3 Exercise

Individual pathways

PRACTISE
1, 2, 6, 9, 12, 15

CONSOLIDATE
3, 4, 7, 10, 13, 16

MASTER
5, 8, 11, 14, 17

Fluency

1. Calculate the following penalty rates:
 - a. time-and-a-half when the hourly rate is \$15.96
 - b. double time when the hourly rate is \$23.90
 - c. double time-and-a-half when the hourly rate is \$17.40.
2. Calculate the following total weekly wages:
 - a. 38 hours at \$22.10 per hour, plus 2 hours at time-and-a-half
 - b. 40 hours at \$17.85 per hour, plus 3 hours at time-and-a-half
 - c. 37 hours at \$18.32 per hour, plus 3 hours at time-and-a-half and 2 hours at double time.
3. Julio is paid \$956.08 for a regular 38-hour week. Calculate:
 - a. his hourly rate of pay
 - b. how much he is paid for 3 hours of overtime at time-and-a-half rates
 - c. his wage for a week in which he works 41 hours.
4. **WE3** Geoff is a waiter in a cafe and works 8 hours most days. Calculate what he earns for 8 hours work on the following days:
 - a. a Monday, when he receives his standard rate of \$21.30 per hour
 - b. a Sunday, when he is paid double time
 - c. a public holiday, when he is paid double time-and-a-half.
5. Albert is paid \$870.58 for a 38-hour week. Determine his total wage for a week in which he works 5 extra hours on a public holiday with a double-time-and-a-half penalty rate.



Understanding

6. Jeleesa (aged 16) works at a supermarket on Thursday nights and weekends. The award rate for a 16-year-old is \$7.55 per hour. Calculate what she would earn for working:
 - a. 4 hours on Thursday night
 - b. 6 hours on Saturday
 - c. 4 hours on Sunday at double time
 - d. the total of the 3 days described in parts a, b and c.

7. Jacob works in a pizza shop and is paid \$13.17 per hour.
- Jacob is paid double time-and-a-half for public holiday work. Calculate what he earns per hour on public holidays. Give your answer to the nearest cent.
 - Calculate Jacob's pay for working 6 hours on a public holiday.



8. If Bronte earns \$7.80 on normal time, calculate how much she receives per hour:
- at time-and-a-half
 - at double time
 - at double time-and-a-half.
9. **WE4** a. Complete the time sheet shown. Calculate the number of hours Susan worked this week.

Day	Pay rate	Start time	Finish time	Hours worked
Monday	Normal	9:00 am	5:00 pm	
Tuesday	Normal	9:00 am	5:00 pm	
Wednesday	Normal	9:00 am	5:00 pm	
Thursday	Normal	9:00 am	5:00 pm	
Friday	Normal	9:00 am	3:00 pm	

- b. Complete Susan's pay slip for this week.

Pay slip for: Susan Jones	Week ending 17 August
Normal hours	
Normal pay rate	\$25.60
Overtime hours	0
Overtime pay rate	\$38.40
Total pay	

10. a. Manu works in a department store. His time sheet is shown. Complete the table.

Day	Pay rate	Start time	Finish time	Hours worked
Monday	Normal	9:00 am	5:00 pm	
Tuesday	Normal	9:00 am	5:00 pm	
Wednesday	Normal	—		
Thursday	Normal	1:00 pm	9:00 pm	
Friday	Normal	—		
Saturday	Time-and-a-half	8:00 am	12:00 pm	

- b. Complete Manu's pay slip for this week.

Pay slip for: Manu Taumata	Week ending 21 December
Normal hours	
Normal pay rate	\$10.90
Overtime hours	
Overtime pay rate (time-and-a-half)	
Total pay	

11. a. Eleanor does shift work. Copy and complete their time sheet.

Day	Pay rate	Start time	Finish time	Hours worked
Monday	Normal	7:00 am	3:00 pm	
Tuesday	Normal	7:00 am	3:00 pm	
Wednesday				
Thursday				
Friday	Normal	11:00 pm	7:00 am	
Saturday	Time-and-a-half	11:00 pm	7:00 am	
Sunday	Double time	11:00 pm	7:00 am	

- b. Copy and complete Eleanor's pay slip for the week.

Pay slip for: Eleanor Rigby	Week ending 15 September
Normal hours	
Normal pay rate	\$16.80
Time-and-a-half hours	
Time-and-a-half pay rate	
Double time hours	
Double time pay rate	
Gross pay	



Reasoning

12. Calculate the following total weekly wages:
- 38 hours at \$18.40 per hour, plus 2 hours at time-and-a-half
 - 32 hours at \$23.70 per hour plus 6 hours on a Sunday at double time.
13. Ruby earns \$979.64 for her normal 38-hour week, but last week she also worked 6 hours of overtime at time-and-a-half rates.
- Calculate how much extra she earned and give a possible reason for her getting time-and-a-half rates.
 - Calculate Ruby's total wage.
14. A standard working week is 38 hours and a worker puts in 3 hours overtime at time-and-a-half, plus 2 hours at double time. Calculate the amount of standard work hours equivalent to the total time they have worked.



Problem solving

15. Joshua's basic wage is \$22 per hour. His overtime during the week is paid at time-and-a-half. Over the weekend he is paid double time. Calculate Joshua's gross wage in a week when he works his basic 40 hours, together with 1 hour overtime on Monday, 2 hours overtime on Wednesday and 4 hours overtime on Saturday.
16. Lin works 32 hours per week at \$22 per hour and is paid overtime for any time worked over the 32 hours per week. In one week Lin worked 42 hours and was paid \$814. Overtime is paid at 1.5 times the standard wage. Determine whether Lin was paid the correct amount. If not, then provide the correct amount.

17. The table shows the pay sheet for a small company. If a person works up to 36 hours, the regular pay is \$14.50 per hour.
 For hours over 36 and up to 40, the overtime is time-and-a-half.
 For hours over 40, the overtime is double time.
 Complete the table.

	Hours worked	Regular pay	Overtime pay	Total pay
a.	32			
b.	38.5			
c.	40.5			
d.	47.2			

LESSON

8.4 Simple interest

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the simple interest on a loan or an investment
- apply the simple interest formula to determine the time, the rate or the principal.



8.4.1 Principal and interest

eles-4399

- When you put money in a financial institution such as a bank or credit union, the amount of money you start with is called the **principal**.
- **Interest** is the fee charged for the use of someone else's money.
- Investors are people who place money in a financial institution — they receive interest from the financial institution in return for the use of their money.
- Borrowers pay interest to financial institutions in return for being given money as a loan.
- The amount of interest is determined by the interest rate.
- Interest rates are quoted as a percentage over a given time period, usually a year.
- The total amount of money, which combines the principal (the initial amount) and the interest earned, is known as the value of the investment.

Value of an investment or loan

$$A = P + I$$

where:

- **A** = the total amount of money at the end of the investment or loan period
- **P** = the principal (the initial amount invested or borrowed)
- **I** = the amount of interest earned.



WORKED EXAMPLE 5 Calculating the total amount

Sam borrowed \$450 000 from a bank for his business. After 30 years Sam had repaid his loan, with a total of \$500 000 paid to the bank.

Calculate how much interest Sam paid the bank.

THINK

1. Write the known quantities: principal (P) and the total amount (A).
2. Write the formula to calculate interest paid.
3. Substitute the given values into the formula.
4. Rearrange the formula to calculate the value of interest.
5. Write the answer in a sentence.

WRITE

$$P = \$450\,000$$

$$A = \$500\,000$$

$$A = P + I$$

$$500\,000 = 450\,000 + I$$

$$\begin{aligned} I &= 500\,000 - 450\,000 \\ &= 50\,000 \end{aligned}$$

Sam paid \$50 000 in interest to the bank.



eles-4400

8.4.2 Simple interest

- **Simple interest** is the interest paid based on the principal of an investment.
- The principal remains constant and does not change from one period to the next.
- Since the amount of interest paid for each time period is based on the principal, the amount of interest paid is constant.
- Simple interest is usually used for short-term assets or loans.
 - For example, \$500 is placed in an account that earns 10% p.a. simple interest every year.
 - This means \$50 is earned each year (10% of $\$500 = \50).
 - After 4 years the total interest earned is $4 \times 50 = \$200$, so the value of the investment after 4 years is $500 + 200 = \$700$.



Formula for simple interest

$$I = Pin$$

where:

- I = the amount of interest earned or paid
- P = the principal
- i = the interest rate as a percentage per annum (yearly), written as a decimal (e.g. 2% p.a. is equal to 0.02 p.a.)
- n = the duration of the investment in years.

WORKED EXAMPLE 6 Calculating simple interest

Calculate the amount of simple interest earned over 4 years on an investment of \$35 000 that returns 2.1% p.a.

THINK

1. Write the simple interest formula.
2. State the known values of the variables.
3. Substitute the given values into the formula and evaluate the amount of interest.
4. Write the answer in a sentence.

WRITE

$$I = Pin$$

$$P = \$35\,000$$

$$i = 0.021$$

$$n = 4$$

$$I = 35\,000 \times 0.021 \times 4$$

$$I = 2940$$

The amount of simple interest over 4 years is \$2940.

- Note that the time, n , is in years. Months need to be expressed as a fraction of a year. For example, 27 months = 2 years and 3 months = 2.25 years.

WORKED EXAMPLE 7 Calculating the time of an investment

Karine invests \$2000 at a simple interest rate of 4% p.a. Calculate how long the money needs to be invested to earn \$120 in interest.

THINK

1. Write the simple interest formula.
2. State the known values of the variables.
3. Substitute the given values into the formula and rearrange to calculate the value of the time.
4. Change the years into years and months.
5. Write the answer in a sentence.

WRITE

$$I = Pin$$

$$P = \$2000$$

$$i = 0.04$$

$$n = \$120$$

$$120 = 2000 \times 0.04 \times n$$

$$120 = 80n$$

$$\frac{120}{80} = \frac{80n}{80}$$

$$n = 1.5$$

$$n = 1 \text{ year and 6 months}$$

Karine will need to invest the \$2000 for 1 year and 6 months

WORKED EXAMPLE 8 Calculating the total amount of a loan

Zac borrows \$3 000 for 2 years at 9% p.a. simple interest.

- Calculate how much interest Zac is charged over the term of the loan.
- Calculate the total amount Zac must repay.

THINK

- Write the simple interest formula.
 - State the known values of the variables.
 - Substitute the given values into the simple interest formula and evaluate the value of interest.
 - Write the answer in a sentence.
- State the relationship for the value of a loan.
 - Substitute the values for P and I and calculate the total amount.
 - Write the answer in a sentence.

WRITE

- $I = Pin$
 $P = \$3000$
 $i = 0.09$
 $n = 2$
 $I = 3000 \times 0.09 \times 2$
 $I = 540$
Zac is charged \$540 in interest.
- $A = P + I$
 $A = 3000 + 540$
 $A = 3540$
To repay the loan, Zac must pay \$3540 in total.

WORKED EXAMPLE 9 Calculating simple interest on an item

The Carlon-Tozer family needs to buy a new refrigerator at a cost of \$1679. They will pay a deposit of \$200 and borrow the balance at an interest rate of 19.5% p.a.

The loan will be paid off in 24 equal monthly payments.

- Calculate how much money the Carlon-Tozers need to borrow.
- Calculate how much interest they will pay.
- Determine the total cost of the refrigerator.
- Calculate the amount of each payment.

THINK

- Subtract the deposit from the cost to calculate the amount still owing.
 - Write the answer in a sentence.
- State the known values of the variables and write the simple formula.
 - Substitute the values into the formula to calculate interest (I), by multiplying the values.
 - Write the answer in a sentence.

WRITE

- $1679 - 200 = 1479$
They must borrow \$1479.
- $P = 1479$
 $i = 0.195$
 $n = 2$
 $I = Pin$
 $= 1479 \times 0.195 \times 2$
 $= 576.81$
The interest will be \$576.81.



- | | |
|---|---|
| <p>c. 1. Add the interest to the initial cost to calculate the total cost of the refrigerator.</p> <p>2. Write the answer in a sentence.</p> | <p>c. $1679.00 + 576.81 = 2255.81$</p> <p>The total cost of the refrigerator will be \$2255.81.</p> |
| <p>d. 1. Subtract the deposit from the total cost to calculate the amount to be repaid.</p> <p>2. Divide the total payment into 24 equal payments to calculate each repayment.</p> <p>3. Write the answer in a sentence.</p> | <p>d. $2255.81 - 200 = 2055.81$.</p> <p>$2055.81 \div 24 = 85.66$</p> <p>Each payment will be \$85.66.</p> |

Technology and simple interest

- Digital technologies such as Microsoft Excel can be helpful in understanding the effects of changing P , i and n . Alternatively, you can investigate these effects using the interactivities in the Resources tab for this lesson.



COLLABORATIVE TASK: Interesting time periods

1. In groups, complete the table below for the pair of values $P = \$100$ and $R = 6\%$, with interest calculated and paid:
- each year
 - every 6 months
 - every 4 months
 - every 3 months
 - every 2 months
 - every month.

Time period	2 years	3 years	5 years	7 years	10 years	15 years
Value of the investment						

2. As a class, discuss the effect of increasing the frequency of payments (for example, by paying interest monthly) at the same time as decreasing the interest rate (for example, by dividing it by 12). Would these changes make a difference to the final investment value? Why?

Resources

-  **eWorkbook** Topic 8 Workbook (worksheets, code puzzle and project) (ewbk-2008)
-  **Interactivities** Individual pathway interactivity: Simple interest (int-4526)
Simple interest (int-6074)

8.4 Quick quiz **on**

8.4 Exercise

Individual pathways

PRACTISE

1, 4, 6, 7, 10, 13, 18, 22

CONSOLIDATE

2, 5, 8, 11, 14, 17, 19, 20, 23, 24

MASTER

3, 9, 12, 15, 16, 21, 25, 26

Fluency

- WE5** Alecia invests \$400. After 3 years, she withdraws the total value of her investment, which is \$496. Calculate the simple interest she earned per year.
- WE6** Calculate the amount of simple interest earned on an investment of \$4500:
 - returning 3.5% p.a. for 6 years
 - returning 2.75% p.a. for 4 years
 - returning 7.5% p.a. for 2 years
 - returning 2.5% p.a. for 8 years.
- Calculate the amount of simple interest earned on an investment of:
 - \$2000, returning 4.5% p.a. for 3 years and 6 months
 - \$18 200, returning 3.6% p.a. for 6 months
 - \$460, returning 2.15% p.a. for 2 years and 3 months
 - \$6700, returning 3.2% p.a. for 7 years.
- WE7** Calculate how long it will take an investment of \$12 000 to earn the following:
 - \$1260 interest at a simple interest rate of 3.5% p.a.
 - \$2520 interest at a simple interest rate of 4.2% p.a.
 - \$405 interest at a simple interest rate of 2.25% p.a.
 - \$1485 interest at a simple interest rate of 4.5% p.a.
- Calculate the simple interest rate per annum if:
 - \$3000 earns \$270 in 3 years
 - \$480 earns \$16.20 in 9 months
 - \$5500 earns \$660 in 2.5 years
 - \$2750 earns \$748 in 4 years.
- MC** If the total interest earned on a \$6000 investment is \$600 after 4 years, then the annual interest is:

A. 10% **B.** 7.5% **C.** 5% **D.** 4% **E.** 2.5%
- WE8** Monique borrows \$5000 for 3 years at 8% per annum simple interest.
 - Calculate how much interest she is charged.
 - Calculate the total amount she must repay.
- Calculate the simple interest earned on an investment of \$15 000 at 5.2% p.a. over 30 months.
- For each loan in the table, calculate:
 - the simple interest
 - the amount repaid.

	Principal (\$)	Interest rate per annum	Time
a.	1000	5%	2 years
b.	4000	16%	3 years
c.	8000	4.5%	48 months
d.	2700	3.9%	2 years 6 months
e.	15 678	9.2%	42 months

10. Calculate the final value of each of the following investments.
- \$3000 for 2 years at 5% p.a.
 - \$5000 for 3 years at 4.3% p.a.
11. Hasim borrows \$14 950 to buy a used car. The bank charges a 9.8% p.a. flat rate of interest over 60 months.
- Calculate the total amount he must repay.
 - Calculate the monthly repayment.
12. Carla borrows \$5200 for an overseas trip at 8.9% p.a. simple interest over 30 months. If repayment is made in equal monthly instalments, calculate the amount of each instalment.



Understanding

13. Janan invested \$2000 at a simple interest rate of 4% p.a. Calculate how long he needs to invest it in order to earn \$200 in interest.
14. If Jodie can invest her money at 8% p.a., calculate how much she needs to invest to earn \$2000 in 2 years.
15. If the simple interest charged on a loan of \$9800 over 3 years is \$2352, determine the percentage rate of interest that was charged.
16. Calculate the missing quantity in each row of the table shown.

	Principal	Rate of interest p.a.	Time	Interest earned
a.	\$2000	6%		\$240.00
b.	\$3760	5.8%		\$545.20
c.		7%	3 years	\$126.00
d.		4.9%	1 year 9 months	\$385.88
e.	\$10 000		$1\frac{1}{2}$ years	\$1200.00
f.	\$8500		42 months	\$1041.25

17. **WE9** Mika is buying a used car for \$19 998. He has a deposit of \$3000 and will pay the balance in equal monthly payments over 4 years. The simple interest rate will be 12.9% p.a.

- Calculate how much money he needs to borrow.
- Calculate how much interest he will pay.
- Determine the total cost of the car.
- Determine how many payments he will make.
- Calculate the cost of each payment.



Reasoning

18. If a bank offers interest on its savings account of 4.2% p.a. and the investment is invested for 9 months, explain why 4.2 is not substituted into the simple interest formula as the interest rate.
19. Theresa invests \$4500 at 5.72% per annum in an investment that attracts simple interest for 6 months. Show that at the end of 6 months she should expect to have \$4628.70.
20. A \$269 000 business is purchased on an \$89 000 deposit with the balance payable over 5 years at an 8.95% p.a. flat rate.
- Calculate how much money is borrowed to purchase this business.
 - Calculate how much interest is charged.
 - Calculate the total amount that must be repaid.
 - Determine the size of each of the equal monthly repayments.
 - Explain two ways in which these payments could be reduced.
21. Nam has \$6273 in his bank account at a simple interest rate of 4.86% per annum. After 39 days he calculates that he will have \$6305.57 in his account. Did Nam calculate his interest correctly? Justify your answer by showing your calculations.

Problem solving

22. Giang is paid \$79.50 in interest for an original investment of \$500 for 3 years. Calculate the annual interest rate.
23. A loan is an investment in reverse: you borrow money from a bank and are charged interest. The value of a loan becomes its total cost.
Jitto wishes to borrow \$10 000 from a bank, which charges 11.5% interest per year. If the loan is over 2 years:
- calculate the total interest paid
 - calculate the total cost of the loan.
24. A new sound system costs \$3500, but it can be purchased for no deposit, followed by 48 equal monthly payments, at a simple interest rate of 16.2% p.a.
- Determine the total cost of the sound system.
 - Under a 'no deposit, no payment for 2 years' scheme, 48 payments are still required, but the first payment isn't made for 2 years. This will stretch the loan over 6 years. Calculate how much the system will cost using this scheme.
 - Determine the monthly payments under each of these two schemes.



25. For the following questions, assume that the interest charged on a home loan is simple interest.

a. Tex and Molly purchase their first home and arrange for a home loan of \$375 000. Their home loan interest rate rises by 0.25% per annum within the first 6 months of the loan.

Determine the monthly increase of their repayments. Give your answer to the nearest cent.

b. Brad and Angel's interest on their home loan is also increased by 0.25% per annum. Their monthly repayments increase by \$60.

Determine the amount of their loan. Give your answer in whole dollars.

26. Juanita sells her car for \$10 984. She invests $x\%$ of the money in a bank account at a simple interest rate of 6.68% per annum for 1.5 years. She spends the remainder of the money. At the end of the investment she has exactly enough money to buy a new car for \$11 002.

Determine the value of x , correct to 2 decimal places.



LESSON

8.5 Compound interest

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the final amount received on an investment with compound interest
- calculate the interest earned on an investment.



8.5.1 Compound interest

eles-4842

- Consider \$1000 invested for 3 years at 10% p.a. simple interest.
- Each year the value of the investment increases by \$100, reaching a total value of \$1300.
- The simple interest process can be summarised in the following table.

	Principal	Interest	Total value
Year 1	\$1000	\$100	\$1100
Year 2	\$1000	\$100	\$1200
Year 3	\$1000	\$100	\$1300

Total interest = \$300

- For **compound interest**, the value of the principal is increased by adding the amount of interest earned at the end of each year. In other words the principal and the interest are compounded annually.
- The compound interest process can be summarised in the following table.

	Principal	Interest	Total value
Year 1	\$1000	\$100	\$1100
Year 2	\$1100	\$110	\$1210
Year 3	\$1210	\$121	\$1331

Total interest = \$331



- The principal increases each year and therefore the amount of interest it earns does so as well.
- The difference between simple interest and compound interest can become enormous over many years.

WORKED EXAMPLE 10 Calculating compound interest

Complete the table to determine the interest received when \$5000 is invested at 11% p.a. compounded annually for 3 years.

Year	Principal	Interest	Total value
Year 1	\$5000		
Year 2			
Year 3			

Total interest =

THINK

- Interest for year 1 = 11% of \$5000
Calculate the principal for year 2 by adding the interest to the year 1 principal.
- Interest for year 2 = 11% of \$5550
Calculate the total value at the end of year 2.
This is the principal for year 3.
- Interest for year 3 = 11% of \$6160.50.
- Calculate the interest earned over 3 years by subtracting the year 1 principal from the final amount.
- Complete the table.

WRITE

$$11\% = \frac{11}{100} = 0.11$$

$$I = 0.11 \times 5000 = 550$$

$$5000 + 550 = 5550$$

$$0.11 \times 5550 = 610.50$$

$$5550 + 610.50 = 6160.50$$

$$0.11 \times 6160.50 = 677.66$$

$$6160.50 + 677.66 = 6838.16$$

$$6838.16 - 5000 = 1838.16$$

	Principal	Interest	Total value
Year 1	\$5000	\$550	\$5550
Year 2	\$5550	\$610.50	\$6160.50
Year 3	\$6160.50	\$677.66	\$6838.16

Total interest = \$1838.16

- There is a quicker way of calculating the total value of the investment. Look again at Worked example 10. The investment grows by 11% each year, so its value at the end of the year is 111% $\left(\frac{111}{100} = 1.11\right)$ of its value at the start of the year.

$$\begin{aligned} & 1.11\% \text{ of } 5000 \\ & = 1.11 \times 5000 \\ & = 5550 \end{aligned}$$

- This process is repeated each year for 3 years.

$$5000 \xrightarrow{\times 1.11} 5550 \xrightarrow{\times 1.11} 6160.50 \xrightarrow{\times 1.11} 6838.16$$

$$\text{or } 5000 \times 1.11 \times 1.11 \times 1.11 = 6838.16$$

$$\text{or } 5000 \times (1.11)^3 = 6838.16$$

- After 3 years the value of the investment is \$6838.16.

WORKED EXAMPLE 11 Calculating the value of an investment

An investment of \$2000 receives compounded interest at a rate of 8% p.a. Complete the table to calculate its value after 4 years.

Year	Start of year	End of year
Year 1	\$2000	
Year 2		
Year 3		
Year 4		

THINK

- Interest is compounded at 8%, so at the end of the first year the value is 108% of the initial value. The multiplying factor is 1.08.
- For the value at the end of year 2, calculate 108% of the amount accumulated in year 1, so calculate 108% of 2160.
- For the value at the end of year 3, calculate 108% of the amount accumulated in year 2, so calculate 108% of 2332.80.
- For the value at the end of year 4, calculate 108% of the amount accumulated in year 3, so calculate 108% of 2519.424.
- Complete the table.

WRITE

$$108\% = \frac{108}{100} = 1.08$$

$$1.08 \times 2000 = 2160$$

$$1.08 \times 2160 = 2332.80$$

$$1.08 \times 2332.80 = 2519.424$$

$$1.08 \times 2519.424 = 2720.98$$

Year	Start of year	End of year
Year 1	\$2000	\$2160
Year 2	\$2160	\$2332.80
Year 3	\$2332.80	\$2519.42
Year 4	\$2519.42	\$2720.98

TI | THINK

- In a new problem on a Calculator page type '2000'. Press ENTER.


WRITE



CASIO | THINK

- On the Main screen type '2000'. Press EXE.

WRITE



TI | THINK

- b. Press $\times 1.08$ and ENTER. This will multiply the previous entry of 2000 by 1.08 and display the amount of the investment after 1 year.

WRITE

2000	2000
2000 * 1.08	2160.

- c. Press ENTER to display the amount of the investment after 2 years.

2000	2000
2000 * 1.08	2160.
2160 * 1.08	2332.8

- d. Press ENTER 2 more times (making 4 times in total) to display the amount of the investment after 4 years.

2000	2000
2000 * 1.08	2160.
2160 * 1.08	2332.8
2332.8 * 1.08	2519.424
2519.424 * 1.08	2720.97792

CASIO | THINK

- b. Press $\times 1.08$ and EXE. This will multiply the previous entry of 2000 by 1.08 and display the amount of the investment after 1 year.

WRITE

2000	2000
ans*1.08	2160
□	

- c. Press EXE to display the amount of the investment after 2 years.

2000	2000
ans*1.08	2160
ans*1.08	2332.8
□	

- d. Press EXE 2 more times (making 4 times in total) to display the amount of the investment after 4 years.

2000	2000
ans*1.08	2160
ans*1.08	2332.8
ans*1.08	2519.424
ans*1.08	2720.97792
□	

- The repeated multiplication shown in Worked example 11 can be developed into a formula for compound interest.
- In Worked example 11 the principal (\$2000) was multiplied by 108% four times (because there were 4 years). The final amount, A , can be given by the formula:

$$\begin{aligned}
 A &= 2000 \times 108\% \times 108\% \times 108\% \times 108\% \\
 &= 2000(100\% + 8\%)^4 \\
 &= 2000\left(1 + \frac{8}{100}\right)^4 \\
 &= 2000(1.08)^4
 \end{aligned}$$

- With compound interest the final amount of an investment is calculated using the following formula.

Formula for compound interest

$$A = P(1 + i)^n$$

where:

- A = the final value of the investment
- P = the principal
- i = the interest rate as a percentage per annum (yearly), written as a decimal (e.g. 7.5% p.a. is equal to 0.075 p.a.)
- n = the duration of the investment in years.

- In future studies of financial mathematics, the duration of investment under consideration will include monthly, quarterly or weekly investments. At this stage, n will only be considered in terms of the number of years.
- To calculate the amount of compound interest earned or paid, subtract the principal from the final value of the investment.

Calculating compound interest earned

$$I = A - P$$

where:

- I = the amount of interest earned or paid
- A = the final value of the investment
- P = the principal.

WORKED EXAMPLE 12 Investing an amount earning compound interest

Peter invests \$40 000 for 8 years at 7.5% p.a., compounding annually.

- Calculate how much money Peter has in total at the end of the 8 years.
- Calculate how much interest Peter earned on the investment.

THINK

1. Write the compound interest formula.
2. State the values of the variables P , i and n .
3. Substitute the values into the formula and calculate the value of the final amount.
4. Write the answer in a sentence.

WRITE

$$\begin{aligned} \text{a. } A &= P(1 + i)^n \\ P &= 40\,000 \\ i &= 0.075 \\ n &= 8 \\ A &= 40\,000(1 + 0.075)^8 \\ &= 40\,000(1.075)^8 \\ &= 71\,339.113\,02 \end{aligned}$$

The final amount of Peter's investment was \$71 339.11 (to the nearest cent).

- b. 1. Write the relationship between the interest earned, the final amount and the principal.
2. State the values for A and P .
3. Substitute the values in the formula and Calculate the value of the interest.
4. Write the answer in a sentence.

$$b. I = A - P$$

$$A = 71\,339.11$$

$$P = 40\,000$$


$$I = 71\,339.11 - 40\,000 \\ = 31\,339.11$$

Peter earned \$31 339.11 in interest.


TI | THINK

- a. In a new problem on a Calculator page, type $40\,000\left(1 + \frac{7.5}{100}\right)^8$. Press ENTER.

WRITE




- b. To find the interest earned, type $-40\,000$. Press ENTER.




CASIO | THINK

- a. In a new problem on a Calculator page, type $40\,000\left(1 + \frac{7.5}{100}\right)^8$. Press EXE.

WRITE





- b. To find the interest earned, type $-40\,000$. Press EXE.



COLLABORATIVE TASK: Compound interest conga

1. Arrange yourself into teams of 3 or more players. The number of players in each team becomes the number of years of a compound interest problem.
2. One person from your class writes a principal value and an interest rate on the board.
3. Using the interest rate and the principal of the investment written on the board, the first member of each group calculates the value of the investment after 1 year.
4. The second member of each group takes the first member's numbers and calculates the value of the investment after 2 years.
5. The rest of the members of the group take it in turns to calculate the interest for the following years.
6. The first team in each round to get the correct answer gets 3 points. The second team to get the correct answer gets 1 point. The team with the most points after the agreed-upon number of rounds is the winner.
7. Repeat the game using different compounding periods (the compounding period is the period of time after which the interest earned is added to the principal).
8. As a class, discuss the effect of decreasing compounding periods on the value of the investment.

-  **eWorkbook** Topic 8 Workbook (worksheets, code puzzle and project) (ewbk-2008)
-  **Interactivities** Individual pathway interactivity: Compound interest (int-4527)
Compound interest (int-6075)

Exercise 8.5 Compound interest

8.5 Quick quiz **on**

8.5 Exercise

Individual pathways

PRACTISE

1, 3, 5, 7, 11, 12, 15

CONSOLIDATE

2, 6, 9, 10, 13, 16

MASTER

4, 8, 14, 17, 18

Fluency

1. **WE10** Complete the tables shown to determine the interest earned when:

- a. \$1000 is invested at 12% p.a. compounded annually for 3 years

	Principal	Interest	Total value
Year 1	\$1000		
Year 2			
Year 3			

Total interest =

- b. \$100 000 is invested at 9% p.a. compounded annually for 4 years.

	Principal	Interest	Total value
Year 1	\$100 000		
Year 2			
Year 3			
Year 4			

Total interest =

2. **WE11** Complete the tables shown to calculate the final value of each investment.

- a. \$5000 invested at 12% p.a. compounded annually for 3 years

	Start of year	End of year
Year 1	\$5000	
Year 2		
Year 3		

- b. \$200 000 invested at 7% p.a. compounded annually for 3 years

	Start of year	End of year
Year 1	\$200 000	
Year 2		
Year 3		

3. Calculate the final amount of an investment of \$5000 when compound interest is earned at the rate of:
- a. 4% p.a. for 3 years
 - b. 3% p.a. for 4 years
 - c. 2.75% p.a. for 6 years
 - d. 3.5% p.a. for 2 years.
4. Calculate the final value of the following investments when compound interest is earned:
- a. \$4650 at 4.5% p.a. for 3 years
 - b. \$12 500 at 6.2% p.a. for 5 years
 - c. \$3560 at 2.4% p.a. for 4 years
 - d. \$25 000 at 3.2% p.a. for 10 years.

Understanding

5. **WE12** For each of the following investments, use the compound interest formula to calculate:
- i. the total value
 - ii. the amount of interest paid.
 - a. \$8000 is invested for 8 years at 15% p.a. interest compounding annually.
 - b. \$50 000 is invested for 4 years at 6% p.a. interest compounding annually.
 - c. \$72 000 is invested for 3 years at 7.8% p.a. interest compounding annually.
6. For each of the following investments, use the compound interest formula to calculate:
- i. the total value
 - ii. the amount of interest paid.
 - a. \$150 000 is invested for 7 years at 6.3% p.a. interest compounding annually.
 - b. \$3500 is invested for 20 years at 15% p.a. interest compounding annually.
 - c. \$21 000 is invested for 10 years at 9.2% p.a. interest compounding annually.
7. Peter invests \$5000 for 3 years at 6% p.a. simple interest. Maria invests the same amount for 3 years at 5.8% p.a. compounding annually.
- a. Calculate the value of Peter's investment on maturity.
 - b. Calculate the value of Maria's investment on maturity.
 - c. Explain why Maria's investment is worth more, even though she received a lower interest rate.
8. Gianni invests \$8000 at 15% p.a. compounded annually. Dylan invests \$8000 at a 15% p.a. flat rate. Calculate how much more Gianni's investment is worth than Dylan's after:
- a. 1 year
 - b. 2 years
 - c. 5 years
 - d. 10 years.
9. When Kim's granddaughter was born Kim invested \$100 at the rate of 7% p.a. compounding annually. Kim plans to give it to her granddaughter on her eighteenth birthday. Calculate how much her granddaughter will receive when she turns 18.
10. Mai's investment account has compounded at a steady 9% for the last 10 years. If it is now worth \$68 000, determine how much it was worth:
- a. last year
 - b. 10 years ago.



Reasoning

- 11.** Two investment options are available to invest \$3000.
- Invest for 5 years at 5% p.a. compounding monthly.
 - Invest for 5 years at 5% p.a. compounding weekly.
- Explain which option you would choose and why.
- 12.** There are 3 factors that affect the value of a compound interest investment: the principal, the interest rate and the length of the investment.
- a.** If the compound interest rate is 10% p.a. and the length of the investment is 2 years, calculate the value of an investment of:
- i.** \$1000
 - ii.** \$2000
 - iii.** \$4000.
- b.** Comment on the effect that increasing the principal has on the value of the investment.
- 13. a.** If the principal is \$1000 and the compound interest rate is 10% p.a., calculate the value of an investment that lasts for:
- i.** 2 years
 - ii.** 4 years
 - iii.** 8 years.
- b.** Comment on the effect that increasing the length of the investment has on the value of the investment.
- 14. a.** If the principal is \$1000 and the length of investment is 5 years, calculate the value of an investment when the compound interest rate is:
- i.** 6% p.a.
 - ii.** 8% p.a.
 - iii.** 10% p.a.
- b.** Comment on the effect of increasing the interest rate on the value of the investment.

Problem solving


- 15.** A bank offers a term deposit for 3 years at an interest rate of 8% p.a. with a compounding period of 6 months. Calculate the end value of a \$5000 investment under these conditions.
- 16.** A building society offers term deposits at 9% p.a., compounded annually. A credit union offers term deposits at 10% but with simple interest only.
- a.** Determine which offer will result in the greatest value after 2 years.
 - b.** Determine which offer will result in the greatest value after 3 years.
 - c.** Determine how many years it will take for the compound interest offer to have the greater value.
- 17.** Chris and Jenny each invested \$10 000. Chris invested at 6.5% p.a. compounding annually. Jenny took a flat rate of interest.
- After 5 years, their investments had equal value.
- a.** Calculate the value of Chris's investment after 5 years.
 - b.** Calculate Jenny's interest rate.
 - c.** Calculate the values of Chris's and Jenny's investments after 6 years.
- 18.** One aspect of compound interest is of great importance to investors: how long does it take for an investment to double in value?
- Consider a principal of \$100 and an annual interest rate of 10% (compounding annually).
- a.** Determine how long it would take for this investment to be worth \$200.
 - b.** Evaluate how long it would take for the investment to be worth \$400 (to double, and then double again).



LESSON

8.6 Review




8.6.1 Topic summary



8.6.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
8.2	I understand the difference between a salary and a wage.			
	I can calculate weekly, fortnightly or monthly pays from a given annual salary.			
	I can calculate the amount earned from the hours worked and the hourly rate of pay.			
8.3	I can calculate total wages when overtime or penalty rates are included.			
	I can calculate hours paid when time-and-a-half or double time is involved.			
8.4	I can calculate the simple interest on a loan or an investment.			
	I can apply the simple interest formula to determine the time, the rate or the principal.			
8.5	I can calculate the final amount received on an investment with compound interest.			
	I can calculate the interest earned on an investment.			

8.6.3 Project

Australian currency

Since decimal currency was introduced in Australia in 1966, our notes and coins have undergone many changes. Only our 5c, 10c and 20c coins are still minted as they were back then. The 1c and 2c coins are no longer in circulation, the 50c coin is a different shape, the \$1 and \$2 notes have been replaced by coins, and our notes have changed from paper to a special type of plastic.

Coins have two sides: an obverse side and a reverse side. The obverse side of all Australian coins depicts the year in which the coin was minted and our reigning monarch at the time of minting. Most coins depict Queen Elizabeth II, however from 2023 onwards the coins will depict King Charles III. The reverse side depicts a typical Australian feature and sometimes a special commemorative event.



- Describe what is depicted on the reverse side of each Australian coin.

The following table includes information on Australia's coins currently in circulation. Use the table to answer questions 2 to 4.

Coin	Diameter (mm)	Mass (g)	Composition
5c	19.41	2.83	75% copper, 25% nickel
10c	23.60	5.65	75% copper, 25% nickel
20c	28.52	11.30	75% copper, 25% nickel
50c	31.51	15.55	75% copper, 25% nickel
\$1	25.00	9.00	92% copper, 6% aluminium, 2% nickel
\$2	20.50	6.60	92% copper 6% aluminium, 2% nickel

- Identify the heaviest and lightest type of coin. List every type of coin in order from lightest to heaviest.
- Identify which coin has the smaller diameter: the 5c coin or the \$2 coin. Calculate the difference in size.

The following table displays information on the currency notes currently circulating in Australia. The column on the far right compares the average life of the previously used paper notes with that of the current plastic notes. Use the table to answer questions 4 to 8.

Note	Date of issue	Size (mm)	Average life of notes (months)	
			Plastic	Paper
\$5	07/07/1992 24/04/1995 01/01/2001 01/09/2016	130 × 65	40	6
\$10	01/11/1993 20/09/2017	137 × 65	40	8
\$20	31/10/1994 09/10/2019	144 × 65	50	10
\$50	04/10/1995 18/10/2018	151 × 65	About 100	24
\$100	15/05/1996 29/10/2020	158 × 65	About 450	104

- List the denominations of the notes that are available in Australian currency.
- State the date on which Australia's first plastic note was issued and the denomination of the note.
- The \$5 note has been issued 4 times, while all other notes have only been issued twice. Suggest a reason for this.
- Suggest a reason why each note is a different size.
- The table clearly shows that the plastic notes last about 5 times as long as the paper notes we once used. Suggest a reason why the \$50 and \$100 notes last longer than the \$5 and \$10 notes.





eWorkbook Topic 8 Workbook (worksheets, code puzzle and project) (ewbk-2008)



Interactivities Crossword (int-2700)
Sudoku puzzle (int-3210)

Exercise 8.6 Review questions

Fluency

- Jane earns an annual salary of \$45 650. Calculate her fortnightly pay.
- Express \$638.96 per week as an annual salary.
- Minh works as a casual shop assistant and is paid \$8.20 an hour from 3:30 pm to 5:30 pm, Monday to Friday, and \$9.50 an hour from 7 am to 12 noon on Saturday. Calculate his total pay for the week.
- MC** Below are the pay details for 5 people. Choose who receives the most money.
 - Billy receives \$18.50 per hour for a 40-hour working week.
 - Jasmine is on an annual salary of \$38 400.
 - Omar receives \$1476.90 per fortnight.
 - Thuy receives \$3205 per month.
 - Mikke receives \$735 per week.
- Daniel earns \$10 an hour for a regular 38-hour week. If he works overtime, he is paid double time. Calculate how much he would earn if he worked 42 hours in one week.
- Bjorn earns \$468.75 per week award wage for a 38-hour week.
 - Calculate his standard hourly rate.
 - If he is paid time-and-a-half for normal overtime, calculate his pay for a week in which he works 41 hours.
- Phillipa is paid an annual salary of \$48 800. Calculate how much she would be paid each time if she was paid:
 - weekly
 - monthly.
- Deng works as a waiter and is paid \$12.50 per hour. If he works 40 hours per week, calculate Deng's gross pay.

Problem solving

- Calculate the simple interest on an investment of \$4000 for 9 months at 4.9% per annum.
 - Determine the final amount of the investment by the end of its term.
- Calculate the monthly instalment needed to repay a loan of \$12 500 over 40 months at a 9.75% p.a. flat interest rate.
- A simple interest loan of \$5000 over 4 years incurred interest of \$2300. Calculate the interest rate charged.

12. Complete the table shown to calculate the final value of \$100 000 invested at 8.5% p.a. compounded annually for 5 years.

	Start of year	End of year
Year 1	\$100 000	
Year 2		
Year 3		
Year 4		
Year 5		

13. Complete the table shown to calculate the final value of \$12 000 invested at 15% p.a. compounded annually for 4 years.

	Start of year	End of year
Year 1	\$12 000	
Year 2		
Year 3		
Year 4		

14. Daniela is to invest \$16 000 for 2 years at 9% p.a. with interest compounded annually. Calculate:
- the final value of the investment
 - the amount of interest earned.
15. Calculate the amount of interest paid on \$40 000 for 5 years at 6% p.a. with interest compounded every six months.
16. Calculate 3.5% of 900.
17. Milos works in a supermarket and earns \$11.70 per hour. If Milos works on Saturday he is paid at time-and-a-half. Determine the hourly rate for which Milos works on Saturday.
18. Frank invests \$1000 at 5% p.a. for 3 years with interest compounded annually. Calculate the final value of Frank's investment.
19. Calculate the final value of each of the following investments if the principal is \$1000.
- Interest rate = 8% p.a., compounding period = 1 year, time = 2 years.
 - Interest rate = 8% p.a., compounding period = 6 months, time = 2 years.
 - Interest rate = 8% p.a., compounding period = 3 months, time = 2 years.
20. Jenny is given \$5000 by her grandparents on her 18th birthday on the condition she invests the money until she is at least 21 years old, which she agrees to do. She finds the following investment options:
- Option 1: A local bank paying 2.5% p.a. interest compounding annually.
- Option 2: A local accountant paying a 2.7% p.a. flat rate of interest.
- Option 3: A building society that is offering a special deal of 2.4% p.a. interest, compounding annually with a 1% bonus added to the final value.
- Identify the option that Jenny should choose, giving a reason for your answer.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 8 Financial mathematics (extending)

8.1 Pre-test

1. a. \$1643.28 b. \$3286.57 c. \$7120.89
2. \$184
3. \$32.50
4. D
5. \$32.50 per hour
6. 35 hours
7. \$412.50
8. a. \$10 b. \$810
9. D
10. a. \$8372 b. \$31 072 c. \$553.58
11. C
12. 9% p.a
13. C
14. C
15. \$7728

8.2 Salaries and wages

1. a. \$1105.42 b. \$2210.85 c. \$4790.17
2. a. \$1198.08 b. \$2396.15 c. \$5191.67
3. a. \$19 136 b. \$46 410 c. \$68 684.20
4. a. \$25 870 b. \$42 183.96 c. \$100 498.06
5. a. \$3890 per month b. \$3200.58 per fortnight
6. a. \$19.75/h b. \$12.17/h
c. \$25.73/h d. \$39.06/h
7. a. \$459.04 b. Jenny earns \$211.66 more.
8. \$210.72
9. \$261.49
10. Rob earns more.
11. \$30.77
12. 38 hours
13. a. \$1605.77 b. \$2788.81
c. \$1100 d. \$8770
14. B, as the annual salary for option B is the highest.
15. Job A has a higher hourly rate.
16. Minh. Minh's annual salary is \$63 745.76, which is higher than Michelle's annual salary.
17. \$40 per hour. This is the option with the greatest pay.
18. \$1735.95, \$1833.75
19. a. \$29.81 b. \$25.64 c. \$72 080.58
20. 12 weeks

8.3 Penalty rates

1. a. \$23.94 b. \$47.80 c. \$43.50
2. a. \$906.10 b. \$794.33 c. \$833.56
3. a. \$25.16 b. \$113.22 c. \$1069.30

4. a. \$170.40 b. \$340.80 c. \$426.00
5. \$1156.96
6. a. \$30.20 b. \$45.30
c. \$60.40 d. \$135.90
7. a. \$32.93 b. \$197.55
8. a. \$11.70 b. \$15.60 c. \$19.50
9. a. Mon: 8, Tue: 8, Wed: 8, Thu: 8, Fri: 6

b.

Pay slip for Susan Jones	Week ending 17 August
Normal hours	38
Normal pay rate	\$25.60
Overtime hours	0
Overtime pay rate	\$38.40
Total pay	\$972.80

10. a. Mon: 8, Tue: 8, Thu: 8, Sat: 4

b.

Pay slip for Manu Taumata	Week ending 21 December
Normal hours	24
Normal pay rate	\$10.90
Overtime hours	4
Overtime pay rate	\$16.35
Total pay	\$327.00

11. a. Mon: 8, Tue: 8, Fri: 8, Sat: 8, Sun: 8

b.

Pay slip for Eleanor Rigby	Week ending 17 August
Normal hours	24
Normal pay rate	\$16.80
Time-and-a half hours	8
Time-and-a half pay rate	\$25.20
Double time hours	8
Double time pay rate	\$33.60
Total pay	\$873.60

12. a. \$754.40 b. \$1042.80
13. a. \$232.02 b. \$1211.66

14. $46\frac{1}{2}$

15. \$1155

16. No. Lin should have been paid \$1034.

17.

	Hours worked	Regular pay	Overtime pay	Total pay
a.	32	\$464	\$0	\$464
b.	38.5	\$522	\$54.38	\$576.38
c.	40.5	\$522	\$101.50	\$623.50
d.	47.2	\$522	\$295.80	\$817.80

8.4 Simple interest

- \$32
- \$945
 - \$495
 - \$675
 - \$900
- \$315
 - \$327.60
 - \$22.25
 - \$1500.80
- 3 years
 - 5 years
 - 1.5 years or 1 year and 6 months
 - 2.75 years or 2 year and 9 months
- 3% p.a.
 - 4.5% p.a.
 - 4.8% p.a.
 - 6.8% p.a.
- E
- \$1200
 - \$6200
- \$1950
- \$100
 - \$1920
 - \$1440
 - \$263.25
 - \$5048.32
 - \$1100
 - \$5920
 - \$9440
 - \$2963.25
 - \$20 726.32
- \$3300
 - \$5645
- \$22 275.50
 - \$371.26
- \$211.90
- 2.5 years
- \$12 500
- 8%
- 2 years
 - 2.5 years
 - \$600
 - \$4500.06
 - 8%
 - 3.5%
- \$16 998
 - \$8770.97
 - \$28 768.97
 - 48
 - \$536.85
- The interest rate in the simple interest formula needs to be converted from a percentage into a decimal: $4.2\% = 0.042$.
- Sample responses can be found in the worked solutions in the online resources.
- \$180 000
 - \$80 550
 - \$260 550
 - \$4342.50
 - Increase the size of the deposit or increase the length of time over which the loan can be repaid
- Yes
- 5.3%
- \$2300
 - \$12 300
- \$5768
 - \$6902
 - \$120.17, \$143.79
- \$78.13
 - \$288 000
- 91.04

8.5 Compound interest

- | | Principal | Interest | Total value |
|--------|-----------|----------|-------------|
| Year 1 | \$1000 | \$120 | \$1120 |
| Year 2 | \$1120 | \$134.40 | \$1254.40 |
| Year 3 | \$1254.40 | \$150.53 | \$1404.93 |

Total interest = \$404.93
- | | Principal | Interest | Total value |
|--------|--------------|-------------|--------------|
| Year 1 | \$100 000 | \$9000 | \$109 000 |
| Year 2 | \$109 000 | \$9810 | \$118 810 |
| Year 3 | \$118 810 | \$10 692.90 | \$129 502.90 |
| Year 4 | \$129 502.90 | \$11 655.26 | \$141 158.16 |

Total interest = \$41 158.16
- | | Start of year | End of year |
|--------|---------------|-------------|
| Year 1 | \$5000 | \$5600 |
| Year 2 | \$5600 | \$6272 |
| Year 3 | \$6272 | \$7024.64 |
- | | Start of year | End of year |
|--------|---------------|--------------|
| Year 1 | \$200 000 | \$214 000 |
| Year 2 | \$214 000 | \$228 980 |
| Year 3 | \$228 980 | \$245 008.60 |
- \$5624.32
 - \$5883.84
- \$5627.54
 - \$5356.13
- \$5306.42
 - \$3914.26
- \$16 886.23
 - \$34 256.03
- \$24 472.18
 - \$63 123.85
 - \$90 196.31
- \$16 472.18
 - \$13 123.85
 - \$18 196.31
- \$230 050.99
 - \$57 282.88
 - \$50 634.40
- \$80 050.99
 - \$53 782.88
 - \$29 634.40
- \$5900
 - \$5921.44
 - Maria's principal increases each year.
- 0
 - \$180
 - \$2090.86
 - \$12 364.46
- \$337.99
- \$62 385.32
 - \$28 723.93

11. The second option would be the best choice, as the shorter the time between the compounding periods, the greater the interest paid.
12. a. i. \$1210 ii. \$2420 iii. \$4840
 b. Increasing the principal will increase the value of the investment because it will have a higher value of interest.
13. a. i. \$1210 ii. \$1464.10 iii. \$2143.59
 b. Increasing the length of the investment will increase the value of the investment because it will have a higher value of interest.
14. a. i. \$1338.23 ii. \$1469.33 iii. \$1610.51
 b. Increasing the interest rate will increase the value of the investment because it will have a higher value of interest.
15. \$6326.60
16. a. Simple interest
 b. Simple interest
 c. 4 years
17. a. \$13 700.87
 b. 7.4%
 c. Chris, \$14 591.42; Jenny, \$14 440
18. a. 7.27 years b. 14.55 years

Project

1. 5c coin: echidna
 10c coin: lyrebird
 20c coin: platypus
 50c coin: coat of arms
 \$1 coin: five kangaroos
 \$2 coin: Walpiri-Anmatyerre Aboriginal Elder Gwoya Jungarai
2. 5c, 10c, \$2, \$1, 20c, 50c
3. The 5c coin has a smaller diameter. It is smaller by 1.09 mm.
4. \$5, \$10, \$20, \$50, \$100
5. 7 July 1992, \$5
6. Sample responses can be found in the worked solutions in the online resources.
7. The different sizes allow people with vision impairment to tell the difference between each note.
8. The \$50 and \$100 notes are used less frequently.

8.6 Review questions

1. \$1755.77
2. \$33 225.92
3. \$129.50
4. A
5. \$460
6. a. \$12.34 b. \$524.28
7. a. \$938.46 b. \$4066.67
8. \$500
9. a. \$147 b. \$4147
10. \$414.06
11. 11.5%

12.

	Start of year	End of year
Year 1	\$100 000	\$108 500
Year 2	\$108 500	\$117 722.50
Year 3	\$117 722.50	\$127 728.91
Year 4	\$127 728.91	\$138 585.87
Year 5	\$138 585.87	\$150 365.67

13.

	Start of year	End of year
Year 1	\$12 000	\$13 800
Year 2	\$13 800	\$15 870
Year 3	\$15 870	\$18 250.50
Year 4	\$18 250.50	\$20 988.08

14. a. \$19 009.60 b. \$3009.60
15. \$13 756.66
16. 31.5
17. \$17.55
18. \$1157.625
19. a. \$1166.40 b. \$1169.86 c. \$1171.66
20. The best option for Jenny would be Option 3 because it gives a larger final amount: \$5422.40.

9 Measurement

LESSON SEQUENCE

9.1 Overview	488
9.2 Measurement	492
9.3 Area	501
9.4 Area and perimeter of a sector	512
9.5 Surface area of rectangular and triangular prisms	519
9.6 Surface area of a cylinder	525
9.7 Volume of prisms and cylinders	530
9.8 Review	543



LESSON

9.1 Overview

9.1.1 Why learn this?

We live in a world surrounded by shapes and objects. Often, we ask questions such as ‘how long?’, ‘how far?’ or ‘how big?’. These questions are all answered using measurement. Measurements may be in one, two or three dimensions. Examples of measurements in one dimension would be length and perimeter, temperature and time. Measurements in two dimensions include areas of shapes including the areas of curved surfaces of objects such as drink cans. Measurements in three dimensions include volumes of objects and capacity, such as the number of millilitres in a drink can.

Since objects are all around us, measurement skills are used in many real-world situations. Carpenters, builders, concreters, landscape gardeners and construction workers are just a few of the many tradespeople who need to understand various aspects of measurement when ordering or working onsite and following plans. Designers, interior decorators and architects use measurement in their drawings and calculations. Chefs measure ingredients in their cooking. Nurses and health professionals follow instructions regarding the amount of a drug to administer to a patient. To maximise profits, manufacturers need to minimise the amount of raw materials used in production. This means knowing the measurements of various parts. Understanding the basic concepts involved in measurement is beneficial in many real-world situations.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept




Extra learning resources


Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck


1. Calculate the perimeter of this figure, giving your answer in centimetres.



2. Calculate the area of the triangle, correct to 2 decimal places.




3. Calculate the perimeter of the triangle shown.



4. **MC** 0.005 m^2 is equivalent to:


- A. $0.000\ 05 \text{ cm}^2$
- B. 0.5 cm^2
- C. 50 cm^2
- D. 0.005 cm^2
- E. 5 cm^2

5. Calculate the volume of the cuboid shown in mm^3 .




6. **MC** Select the correct formula to calculate the perimeter of the shape shown.

- A. $P = \pi \times 4^2$
- B. $P = 2 \times \pi \times 4^2$
- C. $P = 8 \times \pi$
- D. $P = \pi \times 4^2 + 8$
- E. $P = 6 \times \pi + 8$




7. **MC** Select the correct formula to calculate the area of the sector shown.

- A. $A = \frac{60}{360} \times \pi \times 5$
- B. $A = \frac{60}{360} \times \pi \times 25$
- C. $A = \frac{60}{300} \times \pi \times 25$
- D. $A = \frac{60}{360} \times 2 \times \pi \times 5$
- E. $A = \frac{60}{300} \times 2 \times \pi \times 5$




8. **MC** Select the perimeter of the running track shown, correct to 2 decimal places.

- A. 94.25 m
- B. 304.24 m
- C. 467.12 m
- D. 514.25 m
- E. 916.85 m




9. **MC** Select the correct formula to calculate the area of the annulus.

- A. $A = \pi(7^2 - 4^2)$
- B. $A = \pi(7^2 - 3^2)$
- C. $A = \pi \times 4^2$
- D. $A = 2 \times \pi \times (7 - 3)$
- E. $A = 2 \times \pi \times (7 - 4)$




10. Calculate the surface area (SA) of the rectangular prism, giving your answer correct to 2 decimal places.




11. **MC** Select the correct formula to calculate the surface area of the triangular prism.


- A. $SA = \frac{1}{2} \times 2 \times 2 \times 7$
 B. $SA = \frac{1}{2} \times 2 \times 2 \times 7 \times 3$
 C. $SA = \frac{1}{2} \times 2 \times 2 \times 2 + 3 \times 7 \times 2$
 D. $SA = \frac{1}{2} \times 2 \times 2 + 2 \times 7 \times 2$
 E. $SA = 2 \times \sqrt{3} + 3 \times 7 \times 2$




12. Convert 3.1 L into cm^3 .
 13. Calculate the capacity (in mL) of two-thirds of the soup can shown, correct to 1 decimal place.



14. A garden bed is designed in the shape of a trapezium with the dimensions shown. If the area of the garden bed is 30 m^2 , calculate its perimeter.



15. A label is made for a tin as shown in the diagram. The label is symmetrically positioned so that it is 3 cm from the top and bottom of the tin. Determine the area of paper required to make the label, correct to 1 decimal place.



LESSON

9.2 Measurement

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- convert between different units of length
- calculate the perimeter of a given shape
- calculate the circumference of a circle.

9.2.1 Length


eles-4851

- In the **metric system**, units of length are based on the metre. The following units are commonly used.

millimetre	(mm)	one-thousandth of a metre
centimetre	(cm)	one-hundredth of a metre
metre	(m)	one metre
kilometre	(km)	one thousand metres

Converting units of length

The following chart is useful when converting from one unit of length to another.



For example, $36\,000\text{ mm} = 36\,000 \div 10 \div 100\text{ m}$
 $= 36\text{ m}$

WORKED EXAMPLE 1 Converting units of length

Convert the following lengths into cm.

- 37 mm
- 2.54 km

THINK

- There will be fewer cm, so divide by 10.
km \rightarrow m: $\times 1000$
m \rightarrow cm: $\times 100$
- There will be more cm, so multiply.


WRITE

- $37 \div 10 = 3.7\text{ cm}$
- $2.54 \times 1000 = 2540\text{ m}$
 $2540 \times 100 = 254\,000\text{ cm}$

9.2.2 Perimeter

eles-4852

- The **perimeter** of a plane (flat) figure is the distance around the outside of the figure.
- If the figure has straight edges, then the perimeter can be found by simply adding all the side lengths.
- Ensure that all lengths are in the same unit.



$$\text{Perimeter} = a + b + c + d + e$$

Circumference


- **Circumference** is a special name given to the perimeter of a circle.
- Circumference is calculated using the formula:

Circumference of a circle

$C = \pi d$, where d is the **diameter** of the circle

or


$C = 2\pi r$, where r is the **radius** of the circle.



- Use the exact value of π unless otherwise directed.

WORKED EXAMPLE 2 Calculating the perimeter of a shape

Calculate the perimeter of this figure. Give your answer in centimetres.



THINK

1. State the lengths of the sides in the same unit of length.
Convert cm to mm: $4 \text{ cm} = 4 \times 10 = 40 \text{ mm}$
2. Add the side lengths together.
3. As the answer is required in cm, convert mm to cm by dividing 108 by 10.
4. Write the answer in the correct unit.

WRITE

23 mm, 45 mm, 40 mm

$$P = 45 + 40 + 23 \\ = 108 \text{ mm}$$


$$P = \frac{108}{10} \\ = 10.8 \text{ cm}$$

The perimeter is 10.8 cm.

WORKED EXAMPLE 3 Calculating the circumference of a circle

Calculate the circumference of the circle. Give your answer correct to 2 decimal places.

- Give your answer as an exact value.
- Give your answer correct to 2 decimal places.



THINK

- The radius is known, so apply the formula $C = 2\pi r$.
- Substitute $r = 2.5$ into $C = 2\pi r$.
- Write your answer as an exact value.
- Calculate the circumference to 4 decimal places and then round correct to 2 decimal places.

WRITE

$$C = 2\pi r$$

$$C = 2 \times \pi \times 2.5$$

$$C = 5\pi$$



$$C \approx 15.7079$$

$$C \approx 15.71 \text{ cm}$$

TI | THINK

- It is possible to measure the circumference of a circle using a CAS calculator. In a new problem, on a Graphs page, press ESC to hide the function entry line. To accurately draw a circle of radius 2.5 units with centre at the origin, we need to place a point on the x -axis at $(2.5, 0)$. Press:
 - MENU
 - 8: Geometry
 - 1: Points & Lines
 - 2: Points On.
 Then click on the x -axis and press ENTER, then press ESC.
- To see the coordinates of this point, hover over the point and press:
 - CTRL
 - MENU
 - 7: Coordinates and Equations.
 The coordinates of the point are displayed.



WRITE




CASIO | THINK

- It is possible to measure the circumference of this circle using a CAS calculator. On the Geometry screen, press the Grid icon. Then press:
 - Draw
 - Basic object
 - Circle
 Place the centre at $(0, 0)$ and press EXE. Drag the circle so that a second point sits at $x = 2.5$. Then press EXE.
- To measure the circumference of the circle, select the circle by placing the pointer on its circumference and then pressing EXE. Now place the pointer elsewhere on the circumference and press EXE.

WRITE




3. With the x -coordinate selected, click. In the text box that appears, delete the current value and type in 2.5, then press ENTER. The point will jump so that it is positioned at the correct point, (2.5, 0).



4. To see the equation of the circle, select the circle and press:

- CTRL
- MENU
- 7: Coordinates and Equations.


The equation of the circle is shown as $x^2 + y^2 = 2.5^2$.



To measure the circumference of the circle, press:

- MENU
- 8: Geometry
- 3: Measurement
- 1: Length.


Place the pointer over the circle and press ENTER. Then press ESC.



5. To check the ratio of the circumference of the circle to its radius, we will store the value of the circumference to a variable, c . To do this, select the text representing the value of the circumference and press:


- CTRL
- MENU
- 5: Store.

Delete the var and type in c , then press ENTER.



3. To measure the circumference of this circle, press:

- Draw
- Measurement
- Circumference



6. Open a Calculator page and store the variable r as 2.5. To do this, type: $r := 2.5$ and press ENTER.


Now determine the ratio $\frac{c}{2r}$. Notice that its value is the same, correct to 4 decimal places, as that of π .

Input	Output
$r = 2.5$	2.5000
$\frac{c}{2r}$	3.1416
π	π
π	3.1416

- Some shapes will contain sectors of circles. To calculate the perimeter of these shapes, determine what proportion of a circle is involved, and multiply this by the perimeter of the circle.

WORKED EXAMPLE 4 Determining the perimeter of a given shape

Determine the perimeter of the shape shown. Give your answer in cm correct to 2 decimal places.



THINK

- There are two straight sections and two semicircles. The two semicircles make up a full circle, the diameter of which is known, so apply the formula $C = \pi d$.
- Substitute $d = 2.8$ into $C = \pi d$ and give the answer to three decimal places.
- Convert 23 mm to cm ($23 \text{ mm} = 2.3 \text{ cm}$). The perimeter is the sum of all the outside lengths.
- Round to 2 decimal places.

WRITE

$$\begin{aligned}
 C &= \pi d \\
 &= \pi \times 2.8 \\
 &\approx 8.796 \text{ cm} \\
 P &= 8.796 + 2 \times 2.3 \\
 &= 13.396 \\
 P &\approx 13.40 \text{ cm}
 \end{aligned}$$

Resources



eWorkbook

Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)



Interactivities

Individual pathway interactivity: Measurement (int-4528)

Converting units of length (int-4011)

Perimeter (int-4013)

Circumference (int-3782)

Individual pathways

PRACTISE

1, 3, 6, 7, 9, 12, 14, 20, 24

CONSOLIDATE

2, 4, 10, 13, 15, 17, 21, 25

MASTER

5, 8, 11, 16, 18, 19, 22, 23, 26

Fluency

1. **WE1** Convert the following lengths to the units shown.

a. $5 \text{ cm} = \underline{\hspace{2cm}} \text{ mm}$

b. $1.52 \text{ m} = \underline{\hspace{2cm}} \text{ cm}$

c. $12.5 \text{ mm} = \underline{\hspace{2cm}} \text{ m}$

d. $0.0322 \text{ m} = \underline{\hspace{2cm}} \text{ mm}$

2. Convert the following lengths to the units shown.





a. $6.57 \text{ m} = \underline{\hspace{2cm}} \text{ km}$

b. $64 \text{ cm} = \underline{\hspace{2cm}} \text{ km}$




c. $0.000\ 014\ 35 \text{ km} = \underline{\hspace{2cm}} \text{ mm}$

d. $18.35 \text{ cm} = \underline{\hspace{2cm}} \text{ km}$




3. **WE2** Calculate the perimeter of the following figures, in millimetres.






4. Calculate the perimeter of each of the following figures. Give your answers in centimetres.






5. Calculate the perimeter of the following figures, in millimetres.






6. Calculate the perimeter of each of the squares.






7. **WE3** Calculate the circumference of the circles. Give your answer correct to 2 decimal places.






8. Calculate the circumference of the circles. Give your answer correct to 2 decimal places.



9. Calculate the perimeter of the rectangles shown.



10. Calculate the perimeter of the rectangles shown.



11. **MC** A circle has a radius of 34 cm. Its circumference, to the nearest centimetre, is:

- A. 102 cm B. 214 cm C. 107 cm D. 3630 cm E. 50 cm

Understanding

12. Timber is sold in standard lengths, which increase in 300-mm intervals from the smallest available length of 900 mm. (The next two standard lengths available are therefore 1200 mm and 1500 mm.)

- Write the next four standard lengths (after 1500 mm) in mm, cm and m.
- Calculate the number of pieces 600 mm in length that could be cut from a length of timber that is 2.4 m long.
- If I need to cut eight pieces of timber, each 41 cm long, determine the smallest standard length I should buy.

Note: Ignore any timber lost due to the cuts.






13. The world's longest bridge is the Akashi–Kaikyo Bridge, which links the islands of Honshu and Shikoku in Japan. Its central span covers 1.990 km.




- a. State the length of the central span, in metres.
- b. Calculate how much longer the span of the Akashi–Kaikyo Bridge is than the span of the Sydney Harbour Bridge, which is 1149 m.






14. **WE4** Determine the perimeter of the shapes shown. Give your answer correct to 2 decimal places.




15. Determine the perimeter of the shapes shown. Give your answer correct to 2 decimal places.




16. Determine the perimeter of the shapes shown. Give your answer correct to 2 decimal places.




17. Determine the perimeter of the racetrack shown in the plan.



18. Yacht races are often run over a triangular course, as shown. Determine the distance covered by the yachts if they completed 3 laps of the course.



19. Use Pythagoras' theorem to calculate the length of the missing side and, hence, the perimeter of the triangular frame shown.




Reasoning

20. The Hubble Space Telescope is over 13 m in length. It orbits the Earth at a height of 559 km, where it can take extremely sharp images outside the distortion of the Earth's atmosphere.
- If the radius of the Earth is 6371 km, show that the distance travelled by the Hubble Space Telescope in one orbit, to the nearest km, is 43 542 km.
 - If the telescope completes one orbit in 96 minutes, show that its speed is approximately 7559 m/s.




21. A bullet can travel in air at 500 m/s.
- Show how the bullet travels 50 000 cm in 1 second.
 - Calculate how long it takes for the bullet to travel 1 centimetre.
 - If a super-slow-motion camera can take 100 000 pictures each second, determine how many shots would be taken by this camera to show the bullet travelling 1 cm.
22. Edward is repainting all the lines of a netball court at the local sports stadium. The dimensions of the netball court are shown.




- Calculate the total length of lines that need to be repainted.
Edward starts painting at 8 pm when the centre is closing, and it takes him $1\frac{1}{2}$ minutes on average to paint each metre of line.
 - Show that it will take him 233 minutes to complete the job.
23. The radius of the Earth is accepted to be roughly 6400 km.
- Calculate how far, to the nearest km, you travel in one complete rotation of the Earth.
 - As the Earth spins on its axis once every 24 hours, calculate the speed you are moving at.
 - If the Earth is 150 000 000 km from the Sun, and it takes 365.25 days to circle around the Sun, show that the speed of the Earth's orbit around the Sun is 107 515 km/h.
Give your answer to the nearest whole number.

Problem solving

24. One-fifth of an 80 cm length of jewellery wire is cut off. A further 22 cm length is then removed. Evaluate whether there is enough wire remaining to make a 40 cm necklace.
25. A church needs to repair one of its hexagonal stained glass windows. Use the information given in the diagram to calculate the width of the window.



26. A spider is sitting in one top corner of a room that has dimensions 6 m by 4 m by 4 m. It needs to get to the corner of the floor that is diagonally opposite. The spider must crawl along the ceiling, then down a wall, until it reaches its destination.
- If the spider crawls first to the diagonally opposite corner of the ceiling, then down the wall to its destination, calculate the distance it would crawl.
 - Determine the shortest distance from the top back corner to the lower left corner.



LESSON

9.3 Area

LEARNING INTENTIONS



At the end of this lesson you should be able to:

- convert between units of area
- calculate areas of triangles, quadrilaterals and circles using formulas
- determine areas of composite shapes.

9.3.1 Area

eles-4853


- The diagram shows a square of side length 1 cm. By definition it has an area of 1 cm^2 (1 square centimetre).
Note: This is a 'square centimetre', not a 'centimetre squared'.
- Area tells us how many squares it takes to cover a figure, so the area of the rectangle shown is 12 cm^2 .



- Area is commonly measured in square millimetres (mm^2), square centimetres (cm^2), square metres (m^2), or square kilometres (km^2).

Converting units of area

The following chart is useful when converting between units of area.



$$\begin{aligned} \text{For example, } 54 \text{ km}^2 &= 54 \times 1000^2 \times 100^2 \\ &= 540\,000\,000\,000 \text{ cm}^2 \end{aligned}$$

- Another common unit is the **hectare** (ha), a $100 \text{ m} \times 100 \text{ m}$ square equal to $10\,000 \text{ m}^2$, which is used to measure small areas of land.

The hectare

$$\begin{aligned} 1 \text{ hectare (ha)} &= 100 \text{ m} \times 100 \text{ m} \\ &= 10\,000 \text{ m}^2 \end{aligned}$$

WORKED EXAMPLE 5 Converting units of area

Convert 1.3 km^2 into:

a. square metres

b. hectares.

THINK

- a. There will be more m^2 , so multiply by 1000^2 .
 b. Divide the result of part a by $10\,000$, as $1 \text{ ha} = 10\,000 \text{ m}^2$.

WRITE


- a. $1.3 \times 1000^2 = 1\,300\,000 \text{ m}^2$
 b. $1\,300\,000 \div 10\,000 = 130 \text{ ha}$

9.3.2 Using formulas to calculate area


eles-4854







- There are many useful formulas that can be used to calculate the areas of simple shapes. Some common formulas are summarised here.

Square




Rectangle




Parallelogram**Triangle****Trapezium****Kite****Circle****Rhombus****WORKED EXAMPLE 6** Calculating the areas of figures

Calculate the area of each of the following figures in cm^2 , correct to 1 decimal place.

a.



b.



THINK

a. 1. The figure is a circle. State the radius.

2. Apply the formula for area of a circle: $A = \pi r^2$.

3. Round the answer to 1 decimal place.

b. 1. The figure is a kite. State the lengths of the diagonals.

2. Apply the formula for area of a kite: $A = \frac{1}{2}xy$.

3. Round the answer to 1 decimal place.

WRITE

a. $r = 2.7$ cm

$$\begin{aligned} A &= \pi r^2 \\ &= \pi \times 2.7^2 \\ &\approx 22.90 \end{aligned}$$

$$A \approx 22.9 \text{ cm}^2$$

b. $x = 4.9$ cm, $y = 7.2$ cm


$$\begin{aligned} A &= \frac{1}{2}xy \\ &= \frac{1}{2} \times 4.9 \times 7.2 \end{aligned}$$

$$\begin{aligned} &= 17.64 \\ A &\approx 17.6 \text{ cm}^2 \end{aligned}$$

9.3.3 Composite shapes

eles-4855


- A composite shape is made up of smaller, simpler shapes. Here are two examples.



- In the second example, the two semicircles are subtracted from the square to obtain the shaded area on the left.

WORKED EXAMPLE 7 Calculating the area of composite shapes

Calculate the area of the figure shown, giving your answer correct to 1 decimal place.



THINK

1. Draw the diagram, divided into basic shapes.

2. A_1 is a semicircle of radius $\frac{50}{2} = 25$ mm.

The area of a semicircle is half the area of a complete circle.

3. Substitute $r = 25$ into the formula and evaluate, correct to 4 decimal places.


4. A_2 is a square of side length 50 mm. Write the formula.

5. Substitute $l = 50$ into the formula and evaluate A_2 .

6. Sum to calculate the total area.

7. Round the final answer correct to 1 decimal place.

WRITE/DRAW



$$A_1 = \frac{1}{2}\pi r^2$$

$$A_1 = \frac{1}{2} \times \pi \times 25^2 \\ \approx 981.7477 \text{ mm}^2$$

$$A_2 = l^2 \\ = 50^2 \\ = 2500 \text{ mm}^2$$

$$\text{Total area} = A_1 + A_2 \\ = 981.7477 + 2500 \\ = 3481.7477 \text{ mm}^2 \\ \approx 3481.7 \text{ mm}^2$$

on Resources



eWorkbook

Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)



Video eLesson

Composite area (eles-1886)



Interactivities

Individual pathway interactivity: Area (int-4529)

Conversion chart for area (int-3783)

Area of rectangles (int-3784)

Area of parallelograms (int-3786)

Area of rhombuses (int-3787)

Area of circles (int-3788)

Area of trapeziums (int-3790)

9.3 Quick quiz **on**

9.3 Exercise

Individual pathways

PRACTISE

1, 2, 4, 7, 9, 13, 17, 19, 22, 27

CONSOLIDATE




3, 5, 8, 10, 14, 15, 18, 20, 23, 24, 28

MASTER




6, 11, 12, 16, 21, 25, 26, 29, 30

Fluency




- MC** To convert an area measurement from square kilometres to square metres:
 - divide by 1000
 - multiply by 1000
 - divide by 1 000 000
 - multiply by 1 000 000
 - divide by 100
- WE5** Convert the following to the units shown.
 - $13\,400\text{ m}^2 = \underline{\hspace{2cm}}\text{ km}^2$
 - $0.04\text{ cm}^2 = \underline{\hspace{2cm}}\text{ mm}^2$
 - $3\,500\,000\text{ cm}^2 = \underline{\hspace{2cm}}\text{ m}^2$
 - $0.005\text{ m}^2 = \underline{\hspace{2cm}}\text{ cm}^2$
- Convert the following to the units shown.
 - $0.043\text{ km}^2 = \underline{\hspace{2cm}}\text{ m}^2$
 - $200\text{ mm}^2 = \underline{\hspace{2cm}}\text{ cm}^2$
 - $1.41\text{ km}^2 = \underline{\hspace{2cm}}\text{ ha}$
 - $3800\text{ m}^2 = \underline{\hspace{2cm}}\text{ ha}$
- WE6** Calculate the area of each of the following shapes. Where appropriate, give your answer correct to 2 decimal places.






- Calculate the area of each of the following shapes. Where appropriate, give your answer correct to 2 decimal places.






- Calculate the area of each of the following shapes. Where appropriate, give your answer correct to 2 decimal places.






7. Calculate the area of each of the following shapes. Where appropriate, give your answer correct to 2 decimal places.






8. Calculate the area of each of the following shapes. Where appropriate, give your answer correct to 2 decimal places.





9. **WE7** Calculate the areas of the composite shapes shown. Where appropriate, express your answers correct to 1 decimal place.





10. Calculate the areas of the composite shapes shown. Where appropriate, express your answers correct to 1 decimal place.



11. Calculate the areas of the composite shapes shown. Where appropriate, express your answers correct to 1 decimal place.




12. Calculate the areas of the composite shapes shown. Where appropriate, express your answers correct to 1 decimal place.




Understanding

13. Calculate the area of the regular hexagon shown by dividing it into two trapeziums.




14. Calculate the area of the regular octagon by dividing it into two trapeziums and a rectangle, as shown in the figure. Express your answer correct to 1 decimal place.




15. An annulus is a shape formed by two concentric circles (two circles with a common centre). Calculate the area of each of the annuli shown by subtracting the area of the smaller circle from the area of the larger circle. Give answers correct to 2 decimal places.

a.




b.




16. Calculate the area of each of the annuli shown (the shaded area). Give answers correct to 2 decimal places.

a.



b.




17. **MC** A pizza has a diameter of 30 cm. If your sister eats one-quarter, determine what area of pizza remains.

A. 168.8 cm^2 B. 530.1 cm^2 C. 706.9 cm^2 D. 176.7 cm^2 E. 531.1 cm^2

18. A circle has an area of 4500 cm^2 . Calculate its diameter to the nearest mm.

19. Determine the cost of covering the sportsground shown in the figure with turf, if the turf costs \$7.50 per square metre.




20. The Murray–Darling River Basin is Australia’s largest catchment. Irrigation of farms in the Murray–Darling Basin has caused soil degradation due to rising salt levels. Studies indicate that about 500 000 hectares of the basin could be affected in the next 50 years.

- a. Convert the possible affected area to square kilometres. ($1 \text{ km}^2 = 100$ hectares)
- b. The total area of the Murray–Darling Basin is about 1 million square kilometres, about one-seventh of the continent. Calculate what percentage of this total area may be affected by salinity.




21. The plan shows two rooms, which are to be refloored.




Calculate the cost if the flooring costs \$45 per square metre. Allow 10% more for wastage and round to the nearest \$10.

Reasoning

22. A sheet of paper measures 29.5 cm by 21.0 cm.
- Calculate the area of the sheet of paper.
 - Determine the radius of the largest circle that could be drawn on this sheet.
 - Calculate the area of this circle.
 - If the interior of the circle is shaded red, show that 56% of the paper is red.
23. A chessboard is made up of 8 rows and 8 columns of squares. Each square is 42 cm^2 in area. Show that the shortest distance from the upper right corner to the lower left corner of the chessboard is 73.32 cm.
24. Two rectangles of sides 15 cm by 10 cm and 8 cm by 5 cm overlap as shown. Show that the difference in area between the two non-overlapping sections of the rectangles is 110 cm^2 .



25. Answer the following questions using the information in the figure.
- Calculate the area of a square with side lengths 40 cm.
 - If the midpoints of each side of the previous square are joined by straight lines to make another square, calculate the area of the smaller square.
 - Now the midpoints of the previous square are also joined with straight lines to make another square. Calculate the area of this even smaller square.
 - This process is repeated again to make an even smaller square. Calculate the area of this smallest square.
 - Explain any patterns you observe.
 - Determine the percentage of the original square's area taken up by the smallest square.
 - Show that the area of the combined figure that is coloured pink is 1000 cm^2 .
26. Show that a square of perimeter $4x + 20$ has an area of $x^2 + 10x + 25$.




Problem solving


27. The area of a children's square playground is 50 m^2 .
- Calculate the exact length of the playground.
 - Pine logs 3 m long are to be laid around the playground. Determine how many logs need to be bought. (The logs can be cut into smaller pieces if required.)



28. A sandpit is designed in the shape of a trapezium, with the dimensions shown. If the area of the sandpit is 14 m^2 , determine its perimeter.



29. A rectangular classroom has a perimeter of 28 m and its length is 4 m shorter than its width. Determine the area of the classroom.
30. The playground equipment is half the length and half the width of a square kindergarten yard, as shown.



- Identify the fraction of the kindergarten yard that is occupied by the play equipment.
- During a working bee, the playground equipment area is extended 2 m in length and 1 m in width. If x represents the length of the kindergarten yard, write an expression for the area of the play equipment.
- Write an expression for the area of the kindergarten yard *not* taken up by the playground equipment.
- The kindergarten yard that is not taken up by the playground equipment is divided into 3 equal-sized sections:
 - A grassed section
 - A sandpit
 - A concrete section
- Write an expression for the area of the concrete section.
- The children usually spend their time on the play equipment or in the sandpit. Write a simplified expression for the area of the yard where the children usually play.

LESSON

9.4 Area and perimeter of a sector

LEARNING INTENTIONS



At the end of this lesson you should be able to:

- identify sectors, semicircles and quadrants
- calculate the area of a sector
- calculate the length of the arc and the perimeter of a sector.

9.4.1 Sectors

eles-4856

- A **sector** is the shape created when a circle is cut by two radii. The smaller area is called the minor sector and the larger area is called the major sector.
- A circle, like a pizza, can be cut into many sectors.
- Two important sectors that have special names are the **semicircle** (half-circle) and the **quadrant** (quarter-circle).




Semicircle



Quadrant

WORKED EXAMPLE 8 Calculating the area of a quadrant

Calculate the area enclosed by the figure, correct to 1 decimal place.



THINK

1. The figure is a quadrant or quarter-circle. Write the formula for its area.
2. Substitute $r = 11$ into the formula.
3. Evaluate and round the answer correct to 1 decimal place. Include the units.

WRITE

$$A = \frac{1}{4} \times \pi r^2$$

$$A = \frac{1}{4} \times \pi \times 11^2$$


$$\approx 95.03$$

$$\approx 95.0 \text{ cm}^2$$

9.4.2 Area of a sector

eles-4857

- Sectors are specified by the angle (θ) between the two radii.
For example, in a quadrant, $\theta = 90^\circ$, so a quadrant is $\frac{90}{360}$ or $\frac{1}{4}$ of a circle.



Area of a sector


For any value of θ , the area of the sector is given by:

$$A_{\text{sector}} = \frac{\theta}{360} \times \pi r^2$$

where r is the radius of the sector.

WORKED EXAMPLE 9 Calculating the area of a sector

Calculate the area of the sector shown, correct to 1 decimal place.



THINK

- This sector is $\frac{30}{360}$ of a circle. Write the formula for its area.
- Substitute $r = 5$ into the formula.
- Evaluate and round the answer correct to 1 decimal place. Include the units.

WRITE

$$A = \frac{30}{360} \pi r^2$$

$$A = \frac{30}{360} \times \pi \times 5^2$$


$$\approx 6.54$$

$$\approx 6.5 \text{ m}^2$$

9.4.3 Perimeter of a sector

eles-4858

- The perimeter, P , of a sector is the sum of the 2 radii and the curved section, which is called an arc of a circle.
- The length of the arc, l , will be $\frac{\theta}{360}$ of the circumference of the circle.



Arc length of a sector

For any value of θ , the arc length, l , of the sector is given by:

$$l = \frac{\theta}{360} \times 2\pi r$$

where r is the radius of the sector.

- The perimeter of a sector can therefore be calculated using the following formula.

Perimeter of a sector


The perimeter, P , of a sector is given by:

$$P = 2r + l$$

where r is the radius of the sector and l is the arc length.

WORKED EXAMPLE 10 Calculating the perimeter of a sector

Calculate the perimeter of the sector shown, correct to 1 decimal place.



THINK

- The sector is $\frac{80}{360}$ of a circle. Write the formula for the length of the curved side.
- Substitute $r = 3$ and evaluate l . Don't round off until the end.
- Add all the sides together to calculate the perimeter.
- Round the answer to 1 decimal place.

WRITE

$$l = \frac{80}{360} \times 2\pi r$$

$$l = \frac{80}{360} \times 2 \times \pi \times 3$$

$$\approx 4.189 \text{ cm}$$

$$P = 4.189 + 3 + 3$$

$$= 10.189 \text{ cm}$$

$$P \approx 10.2 \text{ cm}$$

on Resources



eWorkbook Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)



Interactivities Individual pathway interactivity: Area and perimeter of a sector (int-4530)

Area of a sector (int-6076)

Perimeter of a sector (int-6077)

Individual pathways

PRACTISE

1, 2, 7, 9, 13, 15, 20, 23

CONSOLIDATE





3, 4, 8, 10, 14, 16, 19, 21, 24

MASTER

5, 6, 11, 12, 17, 18, 22, 25

Fluency





1. Calculate the area of the semicircles shown, correct to 2 decimal places.




WE8 For each of the quadrants in questions 2–5, calculate to 1 decimal place:

a. the perimeter

b. the area enclosed.



6. **MC** Select the correct formula for calculating the area of the sector shown.



A. $A = \frac{1}{4} \pi r^2$

B. $A = \frac{3}{4} \pi r^2$

C. $A = \frac{1}{2} \pi r^2$




D. $A = \frac{1}{100} \pi r^2$




E. $A = \frac{1}{10} \pi r^2$

WE9&10 For each of the sectors in questions 7–12, calculate to 1 decimal place:

a. the perimeter


b. the area.






Understanding


13. A searchlight lights up the ground to a distance of 240 m. Calculate the area the searchlight illuminates if it can swing through an angle of 120° , as shown in the diagram. (Give your answer correct to 1 decimal place.)



14. Calculate the perimeter, correct to 1 decimal place, of the figure shown.




15. A goat is tethered by an 8.5 m rope to the outside of a corner post in a paddock, as shown in the diagram.




Calculate the area of grass (shaded) on which the goat is able to graze. (Give your answer correct to 1 decimal place.)

16. A sector has an angle of 80° and a radius of 8 cm; another sector has an angle of 160° and a radius of 4 cm. Determine the ratio of the area of the first sector to the area of the second sector.

17. A beam of light is projected onto a theatre stage as shown in the diagram.





- Calculate the illuminated area (correct to 1 decimal place) by first evaluating the area of the sector.
 - Calculate the percentage of the total stage area that is illuminated by the light beam.
18. The minute hand on a vintage clock is 20 centimetres long and the hour hand is 12 centimetres long.
- Calculate the distance travelled by the minute hand from midday until 2 pm.
 - During this same time, calculate how far the hour hand travelled. Give your answers correct to the nearest centimetre.
19. Four baseball fields are to be constructed inside a rectangular piece of land. Each field is in the shape of a sector of a circle, as shown in light green. The radius of each sector is 80 m.
- Calculate the area of one baseball field, correct to the nearest whole number.
 - Calculate the percentage, correct to 1 decimal place, of the total area occupied by the four fields.
 - The cost of the land is \$24 000 per hectare. Calculate the total purchase price of the land.



Reasoning

20. John and Jim are twins, and on their birthday they have identical birthday cakes, each with a diameter of 30 cm. Grandma Maureen cuts John's cake into 8 equal sectors. Grandma Mary cuts Jim's cake with a circle in the centre and then 6 equal portions from the rest.



- Show that each sector of John's cake makes an angle of 45° at the centre of the cake.
- Calculate the area of one slice of John's cake, correct to 1 decimal place.
- Calculate the area of the small central circular part of Jim's cake, correct to 1 decimal place.
- Calculate the area of one of the larger portions of Jim's cake, correct to 1 decimal place.
- If each boy eats one slice of the largest part of his own cake, does John eat more cake? Explain your answer.


21. A lighthouse has a light beam in the shape of a sector of a circle that rotates at 10 revolutions per minute and covers an angle of 40° . A person stands 200 m from the lighthouse and observes the beam. Show that the time between the end of one flash and the start of the next is approximately 5.33 seconds.

22. Answer the following questions. Where appropriate, give all answers to the nearest whole number.

- a. A donkey inside a square enclosure is tethered to a post at one of the corners. Show that a rope of length 120 m is required to prevent the donkey from eating more than half the grass in the enclosure.
- b. Suppose two donkeys are tethered at opposite corners of the square region shown. Calculate how long each rope should be so that the donkeys together can graze half of the area.
- c. This time four donkeys are tethered, one at each corner of the square region. Calculate how long each rope should be so that all the donkeys can graze only half of the area.
- d. Another donkey is tethered to a post inside an enclosure in the shape of an equilateral triangle. The post is at one of the vertices. Show that a rope of length 64 m is required so that the donkey eats only half of the grass in the enclosure.
- e. This time the donkey is tethered halfway along one side of the equilateral triangular region shown in the diagram. Calculate how long the rope should be so that the donkey can graze half of the area.



150 m




100 m

100 m


100 m

Problem solving

23. A new logo, divided into 6 equal sectors, has been designed based on two circles with the same centre, as shown in the diagram. The radius of the larger circle is 60 cm and the radius of the smaller circle is 20 cm. The shaded sections are to be coated with a special paint costing \$145 a square metre. The edge of the shaded sections is made from a special material costing \$36 a metre. Calculate the cost to make the shaded sections. Give your answer to the nearest dollar.



- 24. The metal washer shown has an inner radius of r cm and an outer radius of $(r + 1)$ cm.
 - a. State, in terms of r , the area of the circular piece of metal that was cut out of the washer.
 - b. State, in terms of r , the area of the larger circle.
 - c. Show that the area of the metal washer in terms of r is $\pi(2r + 1) \text{ cm}^2$.
 - d. If r is 2 m, what is the exact area of the washer?
 - e. If the area of the washer is $15\pi \text{ cm}^2$, show that the radius would be 7 cm.



25. The area of a sector of a circle is $\pi \text{ cm}^2$, and the length of its arc is 2 cm. Determine the radius of the circle (in terms of π).

LESSON

9.5 Surface area of rectangular and triangular prisms

LEARNING INTENTION


At the end of this lesson you should be able to:

- calculate the surface area of rectangular and triangular prisms.


9.5.1 Prisms and surface areas of prisms

eles-4859


- A **prism** is a solid object with a uniform (unchanging) **cross-section** and all sides flat.



Triangular prism




Rectangular prism (cuboid)




Hexagonal prism

- A prism can be sliced (cross-sectioned) in such a way that each 'slice' has an identical base.




'Slicing' a prism into pieces produces congruent cross-sections.


- The following objects are not prisms because they do not have uniform cross-sections.




Cone



Sphere




Square pyramid



Triangular pyramid

Surface area of a prism


- Consider the **triangular prism** shown at left.
- It has 5 faces: 2 bases, which are right-angled triangles, and 3 rectangular sides. The net of the prism is drawn at right.



- The area of the net is the same as the total surface area (SA) of the prism.

WORKED EXAMPLE 11 Calculating the surface area of a rectangular prism


Calculate the surface area of the rectangular prism (cuboid) shown.



THINK

- There are 6 faces: 2 rectangular bases and 4 rectangular sides. Draw diagrams for each pair of faces and label each region.

WRITE/DRAW



- Calculate the area of each rectangle by applying the formula $A = lw$, where l is the length and w is the width of the rectangle.
- The total surface area (SA) is the sum of the area of 2 of each shape. Write the answer.

$$B = 3 \times 5 \\ = 15$$


$$R_1 = 3 \times 8 \\ = 24$$

$$R_2 = 5 \times 8 \\ = 40$$

$$SA = 2 \times B + 2 \times R_1 + 2 \times R_2 \\ = 30 + 48 + 80 \\ = 158 \text{ cm}^2$$

WORKED EXAMPLE 12 Calculating the surface area of a triangular prism


Calculate the surface area of the right-angled triangular prism shown.



THINK

- There are 5 faces: 2 triangles and 3 rectangles.
Draw diagrams for each face and label each region.

WRITE/DRAW



- Calculate the area of the triangular base by applying the formula $A = \frac{1}{2}bh$.
- Calculate the area of each rectangular face by applying the formula $A = lw$.
- The total surface area is the sum of all the areas of the faces, including 2 bases.

$$B = \frac{1}{2} \times 3 \times 4$$

$$= 6 \text{ m}^2$$

$$R_1 = 3 \times 7 = 21$$

$$R_2 = 4 \times 7 = 28$$




$$R_3 = 5 \times 7 = 35$$

$$SA = 2 \times B + R_1 + R_2 + R_3$$

$$= 12 + 21 + 28 + 35$$

$$= 96 \text{ m}^2$$

on Resources

-  **eWorkbook** Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)
-  **Video eLesson** Total surface area of prisms (eles-1909)
-  **Interactivities** Individual pathway interactivity: Surface area of rectangular and triangular prisms (int-4531)
Cross-sections (int-6078)
Surface area of prisms (int-6079)

Exercise 9.5 Surface area of rectangular and triangular prisms **learnon**

9.5 Quick quiz **on**

9.5 Exercise

Individual pathways

PRACTISE

1, 3, 7, 9, 12, 14, 17

CONSOLIDATE




2, 4, 6, 8, 10, 15, 18

MASTER




5, 11, 13, 16, 19

Fluency




1. **WE11** Calculate the surface area of the following rectangular prisms (cuboids).






2. Calculate the surface area of the following rectangular prisms (cuboids).



3. **WE12** Calculate the surface area of each of the following triangular prisms.




4. Calculate the surface area of each of the triangular prisms shown.




5. A shipping company is planning to buy and paint the outside surface of one of these shipping containers. Determine how many cans of paint the company should buy if the base of the container is not painted and each can of paint covers about 40 m^2 .



6. The aim of the Rubik's cube puzzle is to make each face of the cube one colour. Calculate the surface area of the Rubik's cube if each small coloured square is 1.2 cm in length. Assume that there are no gaps between the squares.




7. Determine how many square metres of iron sheet are needed to construct the water tank shown.
8. Calculate the surface area of the tank in question 7 if no top is made.




Understanding

9. Calculate the area of cardboard that would be needed to construct a box to pack this prism, assuming that no overlap occurs.




10. An aquarium is a triangular prism with the dimensions shown. The top of the tank is open. Calculate the area of glass that was required to construct the tank. Give your answer correct to 2 decimal places.



11. A tent is constructed as shown. Calculate the area of canvas needed to make the tent, including the floor.




12. a. Determine how many square centimetres of cardboard are needed to construct the shoebox shown, ignoring the overlap on the top.
 b. Draw a sketch of a net that could be used to make the box.
 13. Calculate the surface area of a square-based prism of height 4 cm, given that the side length of its base is 6 cm.




Reasoning

14. a. Calculate the surface area of the toy block shown.




- b. If two of the blocks are placed together as shown, calculate the surface area of the prism that is formed.



- c. Calculate the surface area of the prism formed by three blocks.
 d. Use the pattern to determine the surface area of a prism formed by eight blocks arranged in a line. Explain your reasoning.
 15. A cube has a side length of 2 cm. Show that the least surface area of a solid formed by joining eight such cubes is 96 cm^2 .
 16. A prism has an equilateral triangular base with a perimeter of 12 cm. If the total surface area of the prism is 302 cm^2 , show that the length of the prism is approximately 24 cm.

Problem solving

17. A wedge in the shape of a triangular prism, as shown in the diagram, is to be painted.
- Draw a net of the wedge so that it is easier to calculate the area to be painted.
 - Calculate the area to be painted. (Do not include the base.)



18. Ken wants to paint his son's bedroom walls blue and the ceiling white. The room measures 3 m by 4 m, with a ceiling height of 2.6 m. There is one door that measures 1 m by 2 m and one window that measures 1.8 m by 0.9 m.

Each surface takes two coats of paint and 1 L of paint covers 16 m^2 on the walls and 12 m^2 on the ceiling. Cans of wall paint cost \$33.95 for 1 L, \$63.90 for 4 L, \$147 for 10 L and \$174 for 15 L. Ceiling paint costs \$24 for 1 L and \$60 for 4 L.

Determine the cheapest options for Ken to paint the room.

19. A swimming pool has a length of 50 m and a width of 28 m. The shallow end of the pool has a depth of 0.80 m, which increases steadily to 3.8 m at the deep end.
- Calculate the area of the floor of the pool to be painted.
 - If the pool is to be filled to the top, calculate how much water will be needed.



LESSON

9.6 Surface area of a cylinder

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- draw the net of a cylinder
- calculate the curved surface area of a cylinder
- calculate the total surface area of a cylinder.

9.6.1 Surface area of a cylinder

eles-4861

- A **cylinder** is a solid object with two identical flat circular ends and one curved side. A cylinder has a uniform cross-section.
- The net of a cylinder has two circular bases and one rectangular face. The rectangular face is the curved surface of the cylinder.
- Because the rectangle wraps around the circular base, the width of the rectangle is the same as the circumference of the circle. Therefore, the width is equal to $2\pi r$.
- The area of each base is πr^2 , and the area of the rectangle is $2\pi rh$.




Surface area of a cylinder

$$SA = 2\pi r^2 + 2\pi rh$$

$$= 2\pi r(r + h)$$

WORKED EXAMPLE 13 Calculating the surface area of a cylinder

- Use the formula $A = 2\pi rh$ to calculate the area of the curved surface of the cylinder, correct to 1 decimal place.
- Use the formula $SA = 2\pi rh + 2\pi r^2$ to calculate the surface area of the cylinder, correct to 1 decimal place.



THINK

- Write the formula for the curved surface area.
 - Identify the values of the pronumerals.
 - Substitute $r = 2$ and $h = 3$.
 - Evaluate, round to 1 decimal place and include units.
- Write the formula for the surface area of a cylinder.
 - Identify the values of the pronumerals.
 - Substitute $r = 2$ and $h = 3$.
 - Evaluate, round to 1 decimal place and include units.

WRITE

- $$A = 2\pi rh$$

$$r = 2, h = 3$$

$$A = 2 \times \pi \times 2 \times 3$$

$$\approx 37.69$$

$$\approx 37.7 \text{ m}^2$$

- $$SA = 2\pi rh + 2\pi r^2$$


$$r = 2, h = 3$$

$$SA = (2 \times \pi \times 2 \times 3) + (2 \times \pi \times 2^2)$$

$$\approx 62.83$$

$$\approx 62.8 \text{ m}^2$$

 **eWorkbook** Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)

 **Interactivities** Individual pathway interactivity: Surface area of a cylinder (int-4532)
Surface area of a cylinder (int-6080)

Exercise 9.6 Surface area of a cylinder

learn**on**

9.6 Quick quiz **on**

9.6 Exercise

Individual pathways

PRACTISE

1, 2, 7, 9, 13, 15, 18

CONSOLIDATE

3, 4, 8, 10, 14, 16, 19







MASTER

5, 6, 11, 12, 17, 20, 21

Fluency

WE13 For each of the cylinders in questions 1–6, answer the following questions.


- Use the formula $A = 2\pi rh$ to calculate the area of the curved surface of the cylinder.
- Use the formula $SA = 2\pi rh + 2\pi r^2$ to calculate the total surface area of the cylinder.




- A can of energy drink has a height of 130 mm and a radius of 24 mm.
 - Draw a net of the can.
 - Calculate the surface area of the can, to the nearest cm^2 .
- A cylinder has a radius of 15 cm and a height of 45 mm. Determine its surface area correct to 1 decimal place.

Understanding



9. A cylinder has a surface area of 2000 cm^2 and a radius of 8 cm . Determine the cylinder's height correct to 2 decimal places.
10. A can of asparagus is 137 mm tall and has a diameter of 66 mm ; a can of tomatoes is 102 mm tall and has a diameter of 71 mm ; and a can of beetroot is 47 mm tall with a diameter of 84 mm .
- Determine which can has the largest surface area and which has the smallest surface area.
 - Calculate the difference between the largest and smallest surface areas, correct to the nearest cm^2 .
11. A 13 m -high storage tank was constructed from stainless steel (including the lid and the base). The diameter is 3 metres , as shown.
- Calculate the surface area of the tank. Give your answer correct to 2 decimal places.
 - Determine the cost of the steel for the side of the tank if it comes in sheets 1 m wide that cost $\$60$ a metre.



12. The concrete pipe shown in the diagram has the following measurements: $t = 30 \text{ mm}$, $D = 18 \text{ cm}$, $l = 27 \text{ cm}$. Answer the following, giving your answers correct to 2 decimal places.
- Calculate the outer curved surface area.
 - Calculate the inner curved surface area.
 - Calculate the total surface area of both ends.
 - Hence calculate the surface area for the entire shape.



13. Wooden mouldings are made by cutting cylindrical dowels in half, as shown. Calculate the surface area of the moulding.
14. Kiara has a rectangular sheet of cardboard with dimensions 25 cm by 14 cm . She rolls the cardboard to form a cylinder so that the shorter side, 14 cm , is its height, and glues the edges together with a 1 cm overlap.



- Determine the radius of the circle Kiara needs to construct to put at the top of her cylinder. Give your answer correct to 2 decimal places.
- Determine the total surface area of her cylinder if she also makes the top and bottom of her cylinder out of cardboard. Give your answer correct to 1 decimal place.


Reasoning

15. If the radius of a cylinder is twice its height, write a formula for the surface area in terms of its height only.
16. Cylinder A has a 10% greater radius and a 10% greater height than Cylinder B. Show that the ratio of their surface areas is $121 : 100$.


17. Two identical cylinders (with height h cm and radius r cm) are modified slightly. The first cylinder's radius is increased by 10% and the second cylinder's height is increased by 20%. After these modifications have been made, the surface area of the second cylinder is greater than that of the first cylinder. Show that $h > 2.1r$.

Problem solving


18. An above-ground swimming pool has the following shape, with semicircular ends. Calculate how much plastic would be needed to line the base and sides of the pool.



19. A over-sized solid wooden die is constructed for a children's playground. The side dimensions of the die are 50 cm. The number on each side of the die will be represented by cylindrical holes that will be drilled out of each side. Each hole will have a diameter of 10 cm and depth of 2 cm. All surfaces on the die will be painted (including the die holes). Show that the total area required to be painted is 1.63 m^2 .
20. The following letterbox is to be spray-painted on the outside. Calculate the total area to be spray-painted. Assume that the end of the letterbox is a semicircle above a rectangle. The letter slot is open and does not require painting.



21. A timber fence is designed as shown. Evaluate how many square metres of paint are required to completely paint the fence front, back, sides and top with 2 coats of paint. Assume each paling is 2 cm in thickness and that the top of each paling is a semicircle.



LESSON

9.7 Volume of prisms and cylinders

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the volume of prisms and cylinders
- convert between units of volume
- convert between units of capacity and units of volume.




eles-4862

9.7.1 Volume and capacity

Volume


- The volume of a solid is the amount of space it fills or occupies.
- The diagram at far right shows a single cube of side length 1 cm. By definition, the cube has a **volume** of 1 cm^3 (1 cubic centimetre).
Note: This is a ‘cubic centimetre’, not a ‘centimetre cubed’.
- The volume of some solids can be found by dividing them into cubes with 1 cm sides.
- Volume is commonly measured in cubic millimetres (mm^3), cubic centimetres (cm^3), cubic metres (m^3) or cubic kilometres (km^3).



This solid (a cuboid) fills the same amount of space as 12 cubic centimetres. The volume of this cuboid is 12 cm^3 .

Converting units of volume

The following chart is useful when converting between units of volume.



$$\begin{aligned} \text{For example, } 3 \text{ m}^3 &= 3 \times 100^3 \times 10^3 \text{ mm}^3 \\ &= 3\,000\,000\,000 \text{ mm}^3 \end{aligned}$$

Capacity

- **Capacity** is another term for volume, which is usually applied to the measurement of liquids and containers.
- The capacity of a container is the volume of liquid that it can hold.
- The standard measurement for capacity is the litre (L).
- Other common units are the millilitre (mL), kilolitre (kL), and megalitre (ML), where:


$$1 \text{ L} = 1000 \text{ mL}$$

$$1 \text{ kL} = 1000 \text{ L}$$

$$1 \text{ ML} = 1\,000\,000 \text{ L}$$

Converting units of capacity

The following chart is useful when converting between units of capacity.



- The units of capacity and volume are related as follows:

Converting between units of volume and capacity

$$1 \text{ cm}^3 = 1 \text{ mL}$$

$$1000 \text{ cm}^3 = 1 \text{ L}$$

$$1 \text{ m}^3 = 1000 \text{ L} = 1 \text{ kL}$$

WORKED EXAMPLE 14 Converting units of volume and capacity

Convert:

a. 13.2 L into cm^3

c. 0.13 cm^3 into mm^3

b. 3.1 m^3 into litres

d. 3.8 kL into m^3 .

THINK

a. 1. $1 \text{ L} = 1000 \text{ mL}$, so multiply by 1000.

2. $1 \text{ mL} = 1 \text{ cm}^3$. Convert to cm^3 .

b. $1 \text{ m}^3 = 1000 \text{ L}$, so multiply by 1000.

c. There will be more mm^3 , so multiply by 10^3 .

d. $1 \text{ kL} = 1 \text{ m}^3$. Convert to m^3 .

WRITE

a. $13.2 \times 1000 = 13\,200 \text{ mL}$

$$= 13\,200 \text{ cm}^3$$

b. $3.1 \times 1000 = 3100 \text{ L}$

c. $0.13 \times 1000 = 130 \text{ mm}^3$

d. $3.8 \text{ kL} = 3.8 \text{ m}^3$

9.7.2 Volume of a prism

eles-4863

- The volume of a prism can be found by multiplying its cross-sectional area (A) by its height (h).

Volume of a prism


$$V_{\text{prism}} = A \times h$$

where:

A is the area of the base or cross-section of the prism

h is the perpendicular height of the prism.


- The cross-section (A) of a prism is often referred to as the base, even if it is not at the bottom of the prism.
- The height (h) is always measured perpendicular to the base, as shown in the following diagrams.



Volume of a cube or a cuboid


- A specific formula can be developed for the volumes of cubes and cuboids.

Cube



$$\begin{aligned}\text{Volume} &= \text{base area} \times \text{height} \\ &= l^2 \times l \\ &= l^3\end{aligned}$$


Cuboid (rectangular prism)



$$\begin{aligned}\text{Volume} &= \text{base area} \times \text{height} \\ &= lw \times h \\ &= lwh\end{aligned}$$

WORKED EXAMPLE 15 Calculating the volume of a prism

Calculate the volume of the hexagonal prism.



THINK


1. Write the formula for the volume of a prism.
2. Identify the values of the pronumerals.
3. Substitute $A = 40$ and $h = 8$ into the formula and evaluate. Include the units.

WRITE

$$\begin{aligned}V &= A \times h \\A &= 40, h = 8 \\V &= 40 \times 8 \\&= 320 \text{ cm}^3\end{aligned}$$

WORKED EXAMPLE 16 Calculating the volume of a prism

Calculate the volume of the prism.



THINK

1. The base of the prism is a triangle. Write the formula for the area of the triangle.
2. Substitute $b = 4$, $h = 3$ into the formula and evaluate.
3. Write the formula for the volume of a prism.
4. State the values of A and h .
5. Substitute $A = 6$ and $h = 8$ into the formula and evaluate. Include the units.

WRITE


$$\begin{aligned}A &= \frac{1}{2}bh \\&= \frac{1}{2} \times 4 \times 3 \\&= 6 \text{ cm}^2 \\V &= A \times h \\A &= 6, h = 8 \\V &= 6 \times 8 \\&= 48 \text{ cm}^3\end{aligned}$$

9.7.3 Volume of a cylinder

eles-4860

- A cylinder has a circular base and uniform cross-section.
- A formula for the volume of a cylinder is shown.

Volume of a cylinder



$$\begin{aligned}\text{Volume} &= \text{base area} \times \text{height} \\ &= \text{area of circle} \times \text{height} \\ &= \pi r^2 \times h \\ &= \pi r^2 h\end{aligned}$$


WORKED EXAMPLE 17 Calculating the capacity of a cylinder

Calculate the capacity, in litres, of a cylindrical water tank that has a diameter of 5.4 m and a height of 3 m. Give your answer correct to the nearest litre.

THINK

1. Draw a labelled diagram of the tank.
2. The base is a circle, so $A = \pi r^2$. Write the formula for the volume of a cylinder.
3. Recognise the values of r and h . Recall that the radius, r , is half the diameter. $r = \frac{5.4}{2} = 2.7$ m
4. Substitute $r = 2.7$, $h = 3$ into the formula and calculate the volume.
5. Convert this volume to litres (multiply by 1000).

WRITE/DRAW



$$V = \pi r^2 h$$


$$r = 2.7, h = 3$$

$$\begin{aligned}V &= \pi \times (2.7)^2 \times 3 \\ &\approx 68.706\,631 \text{ m}^3\end{aligned}$$

$$\begin{aligned}V &= 68.706\,631 \times 1000 \\ &= 68\,706.631 \\ &\approx 68\,707 \text{ L}\end{aligned}$$

WORKED EXAMPLE 18 Calculating the volume of a solid



Calculate the total volume of this solid.



THINK

- The solid is made from two objects.
Draw and label each object.
Let V_1 = volume of the square prism.
Let V_2 = volume of the cylinder.

WRITE/DRAW



- Calculate V_1 (square prism).

$$\begin{aligned} V_1 &= Ah \\ &= 16^2 \times 6 \\ &= 1536 \text{ cm}^3 \end{aligned}$$

- Calculate V_2 (cylinder).



$$\begin{aligned} V_2 &= \pi r^2 h \\ &= \pi \times 5^2 \times 12 \\ &\approx 942.478 \text{ cm}^3 \end{aligned}$$

- To calculate the total volume, add the volumes found above.

$$\begin{aligned} V &= V_1 + V_2 \\ &= 1536 + 942.478 \\ &= 2478.478 \\ &\approx 2478 \text{ cm}^3 \end{aligned}$$

WORKED EXAMPLE 19 Comparing volume and surface area

Calculate the volume and surface area of the following rectangular prisms, compare and comment on your results.



THINK

- Use the volume formula to calculate the volume of the first shape.

WRITE

$$\begin{aligned} V &= A \times h \\ &= 3 \times 2 \times 4 \\ &= 24 \text{ m}^3 \end{aligned}$$

2. Use the volume formula to calculate the volume of the second shape.

$$\begin{aligned}V &= A \times h \\ &= 6 \times 2 \times 2 \\ &= 24 \text{ m}^3\end{aligned}$$

3. Use the surface area formula to calculate the surface area of the first shape.

$$\begin{aligned}SA &= 2(wh + lh + lw) \\ &= 2(3 \times 2 + 4 \times 2 + 4 \times 3) \\ &= 52 \text{ m}^2\end{aligned}$$

4. Use the surface area formula to calculate the surface area of the second shape.

$$\begin{aligned}SA &= 2(wh + lh + lw) \\ &= 2(2 \times 6 + 2 \times 6 + 2 \times 2) \\ &= 56 \text{ m}^2\end{aligned}$$

5. Compare the volume of each shape.

Both shapes are rectangular prisms but with different dimensions. Even though they have different dimensions they have the same volume of 24 m^3 .

6. Compare the surface area of each shape.

Even though these two shapes have the same volume, they do not have the same surface area. For these shapes, their surface areas are 52 m^2 and 56 m^2 .

on Resources



eWorkbook Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)



Interactivities Individual pathway interactivity: Volume of prisms and cylinders (int-4533)
Volume (int-3791)
Prisms? (int-3792)

Exercise 9.7 Volume of prisms and cylinders

learn on

9.7 Quick quiz

9.7 Exercise

Individual pathways

PRACTISE

1, 4, 5, 8, 9, 12, 22, 25

CONSOLIDATE

2, 6, 10, 13, 14, 16, 17, 20, 23, 26

MASTER

3, 7, 11, 15, 18, 19, 21, 24, 27, 28

Fluency

1. **WE14** Convert the following units into mL.

a. 325 cm^3

b. 2.6 m^3

c. 5.1 L

d. 0.63 kL

2. Convert the following units into cm^3 .

a. 5.8 mL

b. 6.1 L

c. 3.2 m^3

d. 59.3 mm^3

3. Convert the following units into kL.

a. 358 L


b. 55.8 m^3

c. 8752 L


d. 5.3 ML

4. Calculate the volumes of the cuboids shown. Assume that each small cube has sides of 1 cm.


a.



b.




c.




5. WE15 Calculate the volumes of these objects.


a.



b.




c.




6. Calculate the volumes of these objects, correct to 1 decimal place where necessary.


a.



b.




c.




7. Calculate the volumes of these objects, correct to 1 decimal place.


a.



b.




c.




8. Calculate the volumes of these rectangular prisms.


a.



b.




c.




9. **WE16** Calculate the volumes of the prisms shown. Give your answers correct to 2 decimal places where necessary.


a.




b.



c.




d.




10. Calculate the volume of the following cylinders. Give your answers correct to 1 decimal place.


a.



b.




c.



11. Calculate the volume of the following cylinders. Give your answers correct to 1 decimal place.


a.




$$r = 2.4 \text{ m}$$

$$h = 1.7 \text{ m}$$

b.






c.






12. **WE17** Calculate the volume of water, in litres (L), that can fill a cylindrical water tank that has a diameter of 3.2 m and a height of 1.8 m.
13. Calculate the capacity of the Esky shown, in litres.



14. **WE18** Calculate the volume of each solid to the nearest cm^3 .



15. Calculate the volume of each solid to the nearest cm^3 .



Understanding


16. Calculate the capacity, in litres, of the cylindrical storage tank shown. Give your answer correct to 1 decimal place.



17. Calculate the capacity (in mL) of this cylindrical coffee plunger when it is filled to the level shown. Give your answer correct to 1 decimal place.
18. Sudhira is installing a rectangular pond in a garden. The pond is 1.5 m wide, 2.2 m long and has a uniform depth of 1.5 m.
- Calculate the volume of soil (in m^3) that Sudhira must remove to make the hole before installing the pond.
 - Calculate the capacity of the pond in litres. (Ignore the thickness of the walls in this calculation.)



19. Until its closure in 2001, Fresh Kills on Staten Island outside New York City was one of the world's biggest landfill garbage dumps. (New Yorkers throw out about 100 000 tonnes of refuse weekly.) Calculate the approximate volume (in m^3) of the Fresh Kills landfill if it covers an area of 1215 hectares and is about 240 m high. (Note: 1 hectare = 10 000 m^2 .)
20. Calculate the internal volume (in cm^3) of the wooden chest shown, correct to 2 decimal places. (Ignore the thickness of the walls.)




21. Answer the following questions using the given information.

- Calculate the volume, in m^3 , of the refrigerator shown.
- Determine the capacity of the refrigerator if the walls are all 5 cm thick.




Reasoning

- Calculate the volume of plastic needed to make the door wedge shown. Give your answer correct to 2 decimal places.



- The wedges can be packed snugly into cartons with dimensions $45 \text{ cm} \times 70 \text{ cm} \times 35 \text{ cm}$. Determine how many wedges fit into each carton.
23. A cylindrical glass is designed to hold 1.25 L. Show that its height is 132 mm if it has a diameter of 110 mm.
24. Mark is responsible for the maintenance of the Olympic (50 m) pool at an aquatic centre. The figure shows the dimensions of an Olympic pool.



- Identify the shape of the pool.
- Draw the cross-section of the prism and calculate its area.
- Show that the capacity of the pool is 1 650 000 L.
- Mark needs to replace the water in the pool every 6 months. If the pool is drained at 45 000 L per hour and refilled at 35 000 L per hour, calculate how long it will take to:
 - drain
 - refill (in hours and minutes).

Problem solving

- 25.** A cylindrical container of water has a diameter of 16 cm and is 40 cm tall.
Determine how many full cylindrical glasses can be filled from the container if the glasses have a diameter of 6 cm and are 12 cm high.
- 26.** An internal combustion engine consists of 4 cylinders. In each cylinder a piston moves up and down. The diameter of each cylinder is called the bore and the height that the piston moves up and down within the cylinder is called the stroke (stroke = height).
- a.** If the bore of a cylinder is 84 mm and the stroke is 72 mm, calculate the volume (in litres) of 4 such cylinders, correct to 2 decimal places.
- b.** When an engine gets old, the cylinders have to be 're-bored'; that is, the bore is increased by a small amount (and new pistons put in them).
If the re-boring increases the diameter by 1.1 mm, determine the increase in volume of the four cylinders (in litres).
Give your answer correct to 3 decimal places.
- 27.** Answer the following questions.
- a.** A square sheet of metal with dimensions 15 cm by 15 cm has a 1 cm square cut out of each corner. The remainder of the square is folded to form an open box.
Calculate the volume of the box.
- b.** Write a general formula for calculating the volume of a box created from a metal sheet of any size with any size square cut out of the corners.
- 28.** **WE19** Identify two prisms that have the same volume but different surface areas.
Show your working.





LESSON

9.8 Review

9.8.1 Topic summary


MEASUREMENT

Units of measurement

- **Length:**

- **Area:**

- **Volume:**

- **Capacity:**

- When converting between units of capacity and volume:
 - $1 \text{ cm}^3 = 1 \text{ mL}$
 - $1000 \text{ cm}^3 = 1 \text{ L}$
 - $1 \text{ m}^3 = 1000 \text{ L} = 1 \text{ kL}$


Area and perimeter of a sector

- The curved part of a sector is often labeled l . Its length can be calculated by the formula $l = \frac{\theta}{360} \times 2\pi r$, where r is the radius and θ is the angle between the two straight sides.
- The perimeter of a sector is: $P = 2r + l$
- The area of a sector is: $A = \frac{\theta}{360} \times \pi r^2$



Perimeter and circumference

- The **perimeter** is the distance around a shape.
- The perimeter of a circle is called the **circumference**.
- The circumference of a circle can be calculated using either of the following formulas, where r is the radius of the circle and d is the diameter of the circle:
 - $C = 2\pi r$
 - $C = \pi d$



Area

- The area of a shape is the amount of space it takes up.
- The areas of common shapes can be calculated using the following formulas:
 - Square: $A = l^2$
 - Rectangle: $A = lw$
 - Parallelogram: $A = bh$
 - Triangle: $A = \frac{1}{2}bh$
 - Trapezium: $A = \frac{1}{2}(a + b)h$
 - Kite: $A = \frac{1}{2}xy$
 - Circle: $A = \pi r^2$
 - Rhombus: $A = \frac{1}{2}xy$

Surface area

- The surface area of an object can be calculated by adding the areas of all its faces.
- The formula for the surface area of a cylinder is: $SA = 2\pi r^2 + 2\pi rh = 2\pi r(r + h)$




Volume

- The volume of a prism is equal to its cross-sectional area multiplied by its height: $V_{\text{prism}} = A \times h$
- The volume of a cube is: $V_{\text{cube}} = l^3$
- The volume of a cuboid is: $V_{\text{cuboid}} = lwh$
- The volume of a cylinder is: $V_{\text{cylinder}} = \pi r^2 h$
- Capacity is another term for volume, and is usually applied to the measurement of liquids and containers.

9.8.2 Success criteria

Tick a column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)


Lesson	Success criteria			
9.2	I can convert between different units of length.			
	I can calculate the perimeter of a given shape.			
	I can calculate the circumference of a circle.			
9.3	I can convert between units of area.			
	I can calculate areas of triangles, quadrilaterals and circles using formulas.			
	I can determine areas of composite shapes.			
9.4	I can identify sectors, semicircles and quadrants.			
	I can calculate the area of a sector.			
	I can calculate the length of the arc and the perimeter of a sector.			
9.5	I can calculate the surface area of rectangular and triangular prisms.			
9.6	I can draw the net of a cylinder.			
	I can calculate the curved surface area of a cylinder.			
	I can calculate the total surface area of a cylinder.			
9.7	I can calculate the volume of prisms and cylinders.			
	I can convert between units of volume.			
	I can convert between units of capacity and units of volume.			

9.8.3 Project

Areas of polygons

The area of plane figures can be found using a formula. For example, if the figure is a rectangle, its area is found by multiplying its length by its width. If the figure is a triangle, the area is found by multiplying half the base length by its perpendicular height. If the figures are complex, break them down into simple shapes and determine the area of each of these shapes. The area of a complex shape is the sum of the area of its simple shapes. Drawing polygons onto grid paper is one method that can be used to determine their area.

Consider the following seven polygons drawn on 1-cm grid paper.



1. By counting the squares or half-squares, determine the area of each polygon in cm^2 . In some cases, it may be necessary to divide some sections into half-rectangles in order to determine the exact area.



Formulas define a relationship between dimensions of figures. In order to search for a formula to calculate the area of polygons drawn on the grid paper, consider the next question.

2. For each of the polygons provided above, complete the following table. Count the number of dots on the perimeter of each polygon and count the number of dots that are within the perimeter of each polygon.

Polygon	Dots on perimeter (b)	Dots within perimeter (i)	Area of polygon (A)
A			
B			
C			
D			
E			
F			
G			

- Choose a pronumeral to represent the headings in the table. Investigate and determine a relationship between the area of each polygon and the dots on and within the perimeter. Test that the relationship determined works for each polygon. Write the relationship as a formula.
- Draw some polygons on grid paper. Use your formula to determine the area of each shape. Confirm that your formula works by counting the squares in each polygon. The results of both methods (formula and method) should be the same.



on Resources

-  **eWorkbook** Topic 9 Workbook (worksheets, code puzzle and project) (ewbk-2009)
-  **Interactivities** Crossword (int-2709)
Sudoku puzzle (int-3211)





Exercise 9.8 Review questions

learn on





Fluency

- MC** Identify which of the following is true.
 - 5 cm is 100 times as big as 5 mm.
 - 5 metres is 100 times as big as 5 cm.
 - 5 km is 1000 times as small as 5 metres.
 - 5 mm is 100 times as small as 5 metres.
 - 5 metres is 1000 times as big as 5 cm.
- MC** The circumference of a circle with a diameter of 12.25 cm is:
 - 471.44 cm
 - 384.85 mm
 - 76.97 cm
 - 117.86 cm
 - 384.85 cm
- MC** The area of the given shape is:
 - 216 m^2
 - 140 m^2
 - 150 m^2
 - 90 m^2
 - 141 m^2
- MC** The area of a circle with diameter 7.5 cm is:
 - 176.7 cm^2
 - 47.1 cm^2
 - 23.6 cm^2
 - 44.2 cm^2
 - 44.5 cm^2
- MC** The perimeter of the shape shown is:
 - 1075.2 cm
 - 55.5 cm
 - 153.2 cm
 - 66.1 cm
 - 661 cm
- MC** The surface area and volume of a cube with side length 7 m are respectively:
 - $294 \text{ m}^2, 343 \text{ m}^3$
 - $49 \text{ m}^2, 343 \text{ m}^3$
 - $147 \text{ m}^2, 49 \text{ m}^3$
 - $28 \text{ m}^2, 84 \text{ m}^3$
 - $295 \text{ m}^2, 343 \text{ m}^3$





7. **MC** The surface area of a rectangular box with dimensions 7 m, 3 m, 2 m is:
A. 41 m^2 **B.** 42 m^2 **C.** 72 m^2 **D.** 82 cm^2 **E.** 82 m^2
8. **MC** The surface area and volume of a cylinder with radius 35 cm and height 40 cm are:
A. $16\,493.36 \text{ cm}^2$ and $153\,938 \text{ cm}^3$ **B.** 8796.5 cm^2 and $11\,246.5 \text{ cm}^3$
C. $153\,938 \text{ cm}^2$ and $11\,246.5 \text{ cm}^3$ **D.** 8796.5 cm^2 and $153\,938 \text{ cm}^3$
E. $16\,493.36 \text{ m}^2$ and $153\,938 \text{ cm}^3$
9. Convert the following lengths into the required units.
a. $26 \text{ mm} = \underline{\hspace{2cm}} \text{ cm}$ **b.** $1385 \text{ mm} = \underline{\hspace{2cm}} \text{ cm}$
c. $1.63 \text{ cm} = \underline{\hspace{2cm}} \text{ mm}$ **d.** $1.5 \text{ km} = \underline{\hspace{2cm}} \text{ m}$
e. $0.077 \text{ km} = \underline{\hspace{2cm}} \text{ m}$ **f.** $2850 \text{ m} = \underline{\hspace{2cm}} \text{ km}$
10. Calculate the cost of 1.785 km of cable, if the cable costs \$4.20 per metre.
11. Calculate the circumference of circles with the following dimensions (correct to 1 decimal place).
a. Radius 4 cm **b.** Radius 5.6 m **c.** Diameter 12 cm
12. Calculate the perimeters of the following shapes (correct to 1 decimal place).







13. Calculate the perimeter of the following shapes (correct to 1 decimal place).







14. Calculate the area of the following shapes.





15. Calculate the area of the following shapes, correct to 1 decimal place where necessary.



16. Calculate the area of the following 2-dimensional shapes by dividing them into simpler shapes. (Where necessary, express your answer correct to 1 decimal place.)



17. Calculate the area of the sectors shown (correct to 2 decimal places).




18. Calculate the perimeter of the sectors in question 17, correct to 2 decimal places.






19. Calculate the inner surface area of the grape-collecting vat using the dimensions for length, width and depth shown.



20. Calculate the surface area of the triangular prism shown.




21. Calculate the volume of each of the following, giving your answers to 2 decimal places where necessary.



Problem solving


22. Calculate the cost of painting an outer surface (including the lid) of a cylindrical water tank that has a radius of 2.2 m and a height of 1.6 m, if the paint costs \$1.90 per square metre.
23. A harvester travels at 11 km/h. The comb (harvesting section) is 8.7 m wide. Determine how many hectares per hour can be harvested with this comb.

24. Nina decides to invite some friends for a sleepover. She plans to have her guests sleep on inflatable mattresses with dimensions 60 cm by 170 cm. A plan of Nina's bedroom is shown.





- Calculate the area of floor space available.
 - The area of an inflatable mattress is 1.02 m^2 . Nina's best friend suggests that, by dividing the available floor space by the area of one inflatable mattress, the number of mattresses that can fit into the bedroom can be calculated. Explain whether the friend is correct.
 - Determine how many self-inflatable mattresses can fit in the room as it is.
 - Explain how your answer to **c** would change if the bed could be moved within the bedroom.
25. A block of chocolate is in the shape of a triangular prism. The face of the prism is an equilateral triangle of 5 cm sides, and the length is 22 cm.
- Determine the area of one of the triangular faces.
 - Calculate the total surface area of the block.
 - Calculate the volume of the block.
 - If it cost \$0.025 per cm^3 to produce the block, calculate the total cost of production.

26. An A4 sheet of paper has dimensions $210 \text{ mm} \times 297 \text{ mm}$ and can be rolled two different ways (by rotating the paper) to make baseless cylinders, as illustrated. Compare the volumes to decide which shape has the greatest volume.



27. These congruent squares have shaded parts of circles inside them.



Compare the shaded area in figure **a** with that in figure **b**.

28. The widespread manufacture of soft drinks in cans only really started in the 1960s. Today cans of soft drink are very common. They can be purchased in a variety of sizes. You can also purchase them as a single can or bulk-buy them in larger quantities.

Does the arrangement of the cans in a multi-pack affect the amount of packaging used to wrap them? For this investigation we will look at the packaging of 375 mL cans. This size of can has a radius of approximately 3.2 cm and a height of approximately 12 cm.

We will consider packaging these cans in cardboard in the shape of a prism, ignoring any overlap for glueing.

- a. A 6-can pack could be packed in a single row of cans.
 - i. Draw a diagram to show what this pack would look like.
 - ii. Calculate the amount of cardboard required for the package.
- b. The cans could also be packaged in a 3×2 arrangement.
 - i. Show the shape of this pack.
 - ii. Calculate the amount of cardboard required for this pack.
- c. Evaluate which of the two 6-can pack arrangements is the more economical in terms of packaging.
- d. These cans can be packaged as a single layer of 4×3 . Calculate how much cardboard this would require.
- e. They could also have a two-level arrangement of 3×2 each layer. Determine what packaging this would require.
- f. It is also possible to have a three-level arrangement of 2×2 each layer. Determine how much packaging would be needed in this case.
- g. Comment on the best way to package a 12-can pack so that it uses the minimum amount of wrapping.
Provide diagrams and mathematical evidence to support your conclusion.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 9 Measurement

9.1 Pre-test

- 15.4 cm
- 12.48 cm^2
- 30 cm
- C
- 1440 mm^3
- E
- B
- D
- B
- 3.48 m^2
- E
- 3100 cm^3
- 678.6 mL
- 25 m
- 527.8 cm^2

9.2 Measurement

- a. 50 mm b. 152 cm c. 0.0125 m d. 32.2 mm
- a. 0.006 57 km b. 0.000 64 km
c. 14.35 mm d. 0.000 183 5 km
- a. 101 mm b. 54 mm c. 60 mm d. 103 mm
- a. 1060 cm b. 85.4 cm c. 206 cm
- a. 78.4 cm b. 113 cm c. 13 cm
- a. 9.6 cm b. 46 mm c. 31 km
- a. 25.13 cm b. 25.13 m c. 69.12 mm
- a. 44.61 cm b. 19 741.77 km c. 3314.38 mm
- a. 192 m b. 1220 mm c. 260 cm
- a. 74 mm b. 9.6 km c. 8 m
- B
- a. 1800 mm, 2100 mm, 2400 mm, 2700 mm, 180 cm, 210 cm, 240 cm, 270 cm, 1.8 m, 2.1 m, 2.4 m, 2.7 m
b. 4
c. 3300 mm
- a. 1990 m b. 841 m
- a. 127.12 cm b. 104.83 cm c. 61.70 cm
- a. 8 m b. 480 mm c. 405.35 cm
- a. 125.66 cm b. 245.66 m c. 70.41 cm
- 222.5 m
- 37.5 km
- 12 m
- a. $r = 6371 + 559$
 $= 6930 \text{ km}$
 $C = 2\pi r$
 $= 2 \times \pi \times 6930$
 $= 13\,860\pi$
 $= 43\,542.474$
 $\approx 43\,542 \text{ km}$

b. 96 minutes = 96×60
 $= 5760 \text{ seconds}$
 $43\,542 \text{ km} = 43\,542\,000 \text{ m}$
Speed = $43\,542\,000 \div 5760$
 $= 7559.375 \dots$
 $\approx 7559 \text{ m/s}$

- a. $500 \text{ m/s} = (500 \times 100) \text{ cm/s}$
 $= 50\,000 \text{ cm/s}$
b. 0.000 02 seconds
c. 2
- a. 155.62 m
b. $155.62 \text{ m} \times 1.5 \text{ minutes/metre} = 233.43 \text{ minutes}$
- a. 40 212 km
b. 1676 km/h
c. Sample responses can be found in the worked solutions in the online resources.
- Yes, 42 cm of wire remains.
- 69.3 cm
- a. $(2\sqrt{13} + 4) \text{ m}$ b. $2\sqrt{17} \text{ m}$

9.3 Area

- D
- a. $13\,400 \text{ m}^2 = 0.0134 \text{ km}^2$
b. $0.04 \text{ cm}^2 = 4 \text{ mm}^2$
c. $3\,500\,000 \text{ cm}^2 = 350 \text{ m}^2$
d. $0.005 \text{ m}^2 = 50 \text{ cm}^2$
- a. $0.043 \text{ km}^2 = 43\,000 \text{ m}^2$
b. $200 \text{ mm}^2 = 2 \text{ cm}^2$
c. $1.41 \text{ km}^2 = 141 \text{ ha}$
d. $3800 \text{ m}^2 = 0.38 \text{ ha}$
- a. 24 cm^2 b. 16 mm^2 c. 537.5 cm^2
- a. 149.5 cm^2 b. 16.32 m^2 c. 11.25 cm^2
- a. 292.5 cm^2 b. 2.5 cm^2 c. 1250 m^2
- a. 50.27 m^2 b. 3.14 mm^2 c. 36.32 m^2
- a. 15.58 cm^2 b. 4.98 cm^2 c. 65 m^2
- a. 6824.2 mm^2 b. 7.6 m^2 c. 734.2 cm^2
- a. 578.5 cm^2 b. 7086.7 m^2 c. 5.4 m^2
- a. 1143.4 m^2 b. 100.5 cm^2
- a. 821 cm^2 b. 661.3 mm^2
- 378 cm^2
- 19.3 cm^2
- a. 100.53 cm^2 b. 1244.07 m^2
- a. 301.59 cm^2 b. 103.67 mm^2
- B
- 75.7 cm or 757 mm
- \$29 596.51
- a. 5000 km^2 b. 0.5%
- \$10 150
- a. 619.5 cm^2
b. 10.5 cm
c. 346.36 cm^2
d. Sample responses can be found in the worked solutions in the online resources.

23. Sample responses can be found in the worked solutions in the online resources.
24. Sample responses can be found in the worked solutions in the online resources.
25. a. 1600 cm^2
 b. 800 cm^2
 c. 400 cm^2
 d. 200 cm^2
 e. The area halves each time.
 f. 12.5%

g. Sample responses can be found in the worked solutions in the online resources.

26. Sample responses can be found in the worked solutions in the online resources.

27. a. $\sqrt{50} = 5\sqrt{2} \text{ m}$ b. 10 logs

28. 17 m

29. 45 m^2

30. a. $\frac{1}{4}$ b. $\frac{1}{4}(x^2 + 6x + 8)$

c. $\frac{1}{4}(3x^2 - 6x - 8)$

d. i. $\frac{1}{12}(3x^2 - 6x - 8)$

ii. $\frac{1}{6}(3x^2 + 6x + 8)$

9.4 Area and perimeter of a sector

1. a. 14.14 cm^2 b. 157.08 cm^2
 c. 27.71 cm^2 d. 226.19 mm^2
2. a. 14.3 cm b. 12.6 cm^2
3. a. 43.6 cm b. 116.9 cm^2
4. a. 40.7 m b. 102.1 m^2
5. a. 5.4 m b. 1.8 m^2
6. E
7. a. 184.6 cm b. 1869.2 cm^2
8. a. 66.8 m b. 226.2 m^2
9. a. 27.4 cm b. 42.4 cm^2
10. a. 342.1 m b. 7243.6 m^2
11. a. 354.6 cm b. 7645.9 cm^2
12. a. 104.4 cm b. 201.1 cm^2
13. $60\,318.6 \text{ m}^2$
14. 303.4 cm
15. 170.2 m^2
16. 2 : 1
17. a. 14.8 m^2 b. 4.4%
18. a. 251 cm b. 13 cm
19. a. 5027 m^2 b. 78.5% c. \$61 440

20. a. Sample responses can be found in the worked solutions in the online resources.
 b. 88.4 cm^2
 c. 50.3 cm^2
 d. 109.4 cm^2
 e. No, Jim eats more cake

21. Sample responses can be found in the worked solutions in the online resources.

22. a. Sample responses can be found in the worked solutions in the online resources.

b. 85 m each

c. 60 m each

d. Sample responses can be found in the worked solutions in the online resources.

e. 37 m

23. \$250

24. a. $\pi r^2 \text{ cm}^2$

b. $\pi(r + 1)^2 \text{ cm}^2$

c. Sample responses can be found in the worked solutions in the online resources.

d. $5\pi \text{ cm}^2$


e. Sample responses can be found in the worked solutions in the online resources.

25. Radius = $\pi \text{ cm}$

9.5 Surface area of rectangular and triangular prisms

1. a. 66 cm^2 b. 62 m^2 c. 6.7 m^2
2. a. 4.44 m^2 b. $11\,572.92 \text{ cm}^2$ c. 1.9 m^2
3. a. 86 cm^2 b. 210.7 m^2 c. 8.37 cm^2
4. a. 840 cm^2 b. 191.08 cm^2 c. 2370 mm^2
5. 2 cans of paint
6. 77.76 cm^2
7. 26.44 m^2
8. 20.36 m^2
9. 261.5 cm^2
10. 2.21 m^2
11. 15.2 m^2
12. a. 2080 cm^2
 b. Sample responses can be found in the worked solutions in the online resources.
13. 168 cm^2
14. a. 150 cm^2 b. 250 cm^2 c. 350 cm^2 d. 850 cm^2
15. The solid formed is a cube with side length 4 cm.
16. Sample responses can be found in the worked solutions in the online resources.

17. a.




b. 315 cm^2

18. \$145.85 (2 L of ceiling paint, 1 L + 4 L for walls)

19. a. 1402.52 m^2 b. 3220 m^3

9.6 Surface area of a cylinder

- | | |
|-----------------------------|--------------------------|
| 1. a. 75.4 m^2 | b. 131.9 m^2 |
| 2. a. 28.3 cm^2 | b. 84.8 cm^2 |
| 3. a. 2010.6 m^2 | b. 3619.1 m^2 |
| 4. a. 1121.5 cm^2 | b. 1575.5 cm^2 |
| 5. a. 6.6 m^2 | b. 9.7 m^2 |
| 6. a. 25.6 m^2 | b. 61.8 m^2 |
7. a.



- b. 232 cm^2
8. 1837.8 cm^2
9. 31.79 cm
10. a. Asparagus is largest; beetroot is smallest.
b. 118 cm^2
11. a. 136.66 m^2 b. \$7351.33
12. a. 2035.75 cm^2 b. 1526.81 cm^2
c. 395.84 cm^2 d. 3958.41 cm^2
13. 154.73 cm^2
14. a. 3.82 cm b. 427.7 cm^2
15. $12\pi h^2$
16. Sample responses can be found in the worked solutions in the online resources.
17. Sample responses can be found in the worked solutions in the online resources.

18. 83.6 m^2

19. The area to be painted is 1.63 m^2 .

20. $11\,231.12 \text{ cm}^2$

21. 4.34 m^2

9.7 Volume of prisms and cylinders

- | | | | |
|--|---------------------------------------|--------------------------|------------------------|
| 1. a. 325 mL | b. 2 600 000 mL | | |
| c. 5100 mL | d. 630 000 mL | | |
| 2. a. 5.8 cm^3 | b. 6100 cm^3 | | |
| c. 3 200 000 cm^3 | d. 0.0593 cm^3 | | |
| 3. a. 0.358 kL | b. 55.8 kL | c. 8.752 kL | d. 5300 kL |
| 4. a. 36 cm^3 | b. 15 cm^3 | c. 72 cm^3 | |
| 5. a. 12 cm^3 | b. 68 cm^3 | c. 9.6 m^3 | |
| 6. a. 630 mm^3 | b. 420 cm^3 | c. 3152.7 mm^3 | |
| 7. a. 1319.5 mm^3 | b. 523.6 m^3 | c. 10.5 m^3 | |
| 8. a. 36 cm^3 | b. 30 m^3 | c. 1.144 m^3 | |
| 9. a. 7020 cm^3 | b. 6.91 m^3 | c. 300 m^3 | d. 8806 cm^3 |
| 10. a. $16\,085.0 \text{ m}^3$ | b. 4766.6 cm^3 | c. 2.3 m^3 | |
| 11. a. 30.8 m^3 | b. 30.6 m^3 | c. 56.5 mm^3 | |
| 12. $14\,476.5 \text{ L}$ | | | |
| 13. 176.4 L | | | |
| 14. a. $158\,169 \text{ cm}^3$ | b. 4092 cm^3 | c. 2639 cm^3 | |
| 15. a. 641 cm^3 | b. 1784 cm^3 | c. 1057 cm^3 | |
| 16. $75\,322.8 \text{ L}$ | | | |
| 17. 1145.1 mL | | | |
| 18. a. 4.95 m^3 | b. 4950 L | | |
| 19. $2.916 \times 10^9 \text{ m}^3$ | | | |
| 20. $234\,256.54 \text{ cm}^3$ | | | |
| 21. a. 1.1475 m^3 | b. 840 L | | |
| 22. a. 45.94 cm^3 | | | |
| | b. 2400 (to the nearest whole number) | | |
| 23. Sample responses can be found in the worked solutions in the online resources. | | | |
| 24. a. Prism | | | |
| b. 75 m^2 | | | |
| c. Sample responses can be found in the worked solutions in the online resources. | | | |
| d. i. 36 h 40 min | ii. 47 h 9 min | | |
| 25. 23 | | | |
| 26. a. 1.60 L | b. 0.042 L | | |
| 27. a. 169 cm^3 | | | |
| b. $V = x(l - 2x)(w - 2x)$ | | | |
| 28. There are many possible answers. An example of one possible solution. | | | |
| Shape 1: Length = 6 m, Width = 4 m, Height = 2 m | | | |
| Shape 2: Length = 8 m, Width = 4.5 m, Height = 2 m | | | |

Project



- | | | | |
|-------------------------|------------------------|------------------------|-----------------------|
| 1. A. 15 cm^2 | B. 20 cm^2 | C. 6 cm^2 | D. 8.5 cm^2 |
| E. 10.5 cm^2 | F. 11.5 cm^2 | G. 12.5 cm^2 | |

Polygon	Dots on perimeter (b)	Dots within perimeter (i)	Area of polygon (A)
A	16	8	15
B	24	9	20
C	4	6	6
D	7	6	8.5
E	11	6	10.5
F	13	6	11.5
G	15	6	12.5

3. Sample responses can be found in the worked solutions in the online resources.
4. Sample responses can be found in the worked solutions in the online resources.

9.8 Review questions

- B
- B
- B
- D
- D
- A
- E
- A
- $26 \text{ mm} = 2.6 \text{ cm}$
 - $1385 \text{ mm} = 138.5 \text{ cm}$
 - $1.63 \text{ cm} = 16.3 \text{ mm}$
 - $1.5 \text{ km} = 1500 \text{ m}$
 - $0.077 \text{ km} = 77 \text{ m}$
 - $2850 \text{ m} = 2.85 \text{ km}$
- \$7497
- 25.1 cm
 - 35.2 m
 - 37.7 cm
- 70.4 cm
 - 30.4 mm
 - 34 m
 - 240 mm or 24 cm
- 13.1 m
 - 308.5 m
 - 97.1 cm
 - 192.0 mm or 19.2 cm
- 25 m^2
 - 950 cm^2
 - 387.5 cm^2
 - 6120 m^2
- 135 mm^2
 - 2170 cm^2
 - 279.4 cm^2
 - 6939.8 mm^2

- 1486.9 cm^2
 - 362.5 m^2
 - 520.4 cm^2
 - 473.2 cm^2
- 628.32 m^2
 - 59.45 cm^2
 - 197.92 cm^2
 - $10\,734.47 \text{ m}^2$
- 102.83 m
 - 31.07 cm
 - 57.99 cm
 - 470.27 m
- $48\,920 \text{ cm}^2$ or 4.892 m^2
- 3550 cm^2
- 343 cm^3
 - 672 cm^3
 - $153\,938.04 \text{ cm}^3$
 - 1.45 m^3
 - 1800 cm^3
- \$99.80
- 9.57 hectares
- 10.75 m^2
 - No
 - 8
 - 9 mattresses could fit in the room.
- 10.83 cm^2
 - 351.65 cm^2
 - 238.26 cm^3
 - \$5.95
- The cylinder of height 210 mm has the greater volume.
- The shaded parts have the same area.
- 
 - 1566.72 cm^2
 - 
 - 1259.52 cm^2
 - 3×2 arrangement
 - 2058.24 cm^2
 - 2027.52 cm^2
 - Yes, 2170.88 cm^2
 - The smallest amount of packaging required is 3×2 , with two levels of packaging.

10 Probability

LESSON SEQUENCE

10.1 Overview	558
10.2 Theoretical probability	564
10.3 Experimental probability	571
10.4 Venn diagrams and two-way tables	580
10.5 Compound events	597
10.6 Review	610



LESSON

10.1 Overview

10.1.1 Why learn this?

New technologies which predict human behaviour are all based on probability. On 11 May 1997, IBM's supercomputer Deep Blue made history by defeating chess grandmaster Gary Kasparov in a six-game match under standard time controls. This was the first time that a computer had defeated the highest ranked chess player in the world. Deep Blue won by evaluating millions of possible positions each second and determining the probability of victory from each possible choice. This application of probability allowed artificial intelligence to defeat a human who had spent his life mastering the game.

Combining advanced computing with probability is no longer used only to play games. From the daily convenience of predictive text, through to assisting doctors with cancer diagnosis, companies spend vast amounts of money on developing probability-based software. Self-driving cars use probability functions to predict the behaviour of other cars on the road and pedestrians. The more accurate the predictive functions, the safer self-driving cars can become. Politicians use campaign data analysis to develop models that produce predictions about individual citizens' likelihood of supporting specific candidates and issues, and the likelihood of these citizens changing their support if they're targeted with various campaign interventions. Even ChatGPT uses probability to create the responses it gives.

As technology improves, so will its predictive power in determining the likelihood of certain outcomes occurring. It is important that we study and understand probability so we know how technology is being used and the impact it will have on our day-to-day life.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 10.1 Pre-test

1. **MC** A six-sided die is rolled, and the number uppermost is noted. The event of rolling an even number is:

A. {1, 2, 3, 4, 5, 6} B. {0, 1, 2, 3, 4, 5, 6} C. {2, 4, 6}
 D. {1, 3, 5} E. {2, 4, 5, 6}

2. The coloured spinner shown is spun once and the colour noted. Written in its simplest form, state the probability of spinning:
 a. an orange and a blue b. an orange or a pink.



3. A coin is tossed in an experiment and the outcomes recorded.


Outcome	Heads	Tails
Frequency	72	28

- a. Identify how many trials there were.
 b. Calculate the experimental probability for tossing a Tail, giving your answer in simplest form.
 c. Calculate the theoretical probability for tossing a Tail with a fair coin, giving your answer in simplest form.
 d. In simplest form, calculate the difference in this experiment between the theoretical and experimental probabilities for tossing a Tail.
4. A random number is selected from:
 $\xi = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30\}$
 Calculate the exact probability of selecting a number that is a multiple of 3, giving your answer in simplest form.


5. Identify whether the following statement is True or False.
 If a large number of trials is conducted in an experiment, the relative (experimental) frequency of each outcome will be very close to its theoretical probability.

6. **MC** Consider the Venn diagram shown.
 If $\xi = \{\text{numbers between 1 and 20 inclusive}\}$, identify which of the following set B is equal to.

A. $B = \{\text{multiples of 3 numbers between 1 and 20}\}$
 B. $B = \{\text{number 1}\}$
 C. $B = \{\text{even numbers between 1 and 20}\}$
 D. $B = \{\text{odd numbers between 1 and 20}\}$
 E. $B = \{\text{prime numbers between 1 and 20}\}$



7. **MC** The information presented in the Venn diagram can be shown on a two-way frequency table as:



A.

	A	A'	Total
B	5	3	8
B'	0	6	6
Total	5	9	14

B.

	A	A'	Total
B	3	6	9
B'	5	0	5
Total	8	6	14

C.

	A	A'	Total
B	3	5	8
B'	6	0	6
Total	9	5	14


D.

	A	A'	Total
B	3	2	5
B'	3	3	6
Total	6	5	11

E.

	A	A'	Total
B	3	3	6
B'	2	3	5
Total	8	6	11

8. **MC** If $\xi = \{\text{numbers between 1 and 20 inclusive}\}$, identify the complement of the event $A = \{\text{multiples of 5 and prime numbers}\}$.
- A.** $\{1, 4, 6, 8, 9, 12, 14, 16, 18\}$ **B.** $\{5, 10, 15, 20\}$
C. $\{2, 3, 5, 7, 11, 13, 17, 19\}$ **D.** $\{2, 3, 5, 7, 10, 11, 13, 15, 17, 19, 20\}$
E. $\{4, 6, 8, 9, 12, 14, 16, 18\}$
9. The jacks and aces from a deck of cards are shuffled, then two cards are drawn. Calculate the exact probability, in simplest form, that two aces are chosen:
- a.** if the first card is replaced **b.** if the first card is not replaced.
10. The Venn diagram shows the results of a survey where students were asked to indicate whether they prefer drama (D) or comedy (C) movies. In simplest form, determine the probability that a student selected at random prefers drama movies but does not like comedy.








11. From a bag of mixed lollies, students can choose from three different types of lollies: red frogs, milk bottles or jelly babies. The bag contains five of each type of lolly. If Mahsa chooses two lollies without looking into the bag, calculate the probability that she will choose two different types of lollies.

12. The students in a class were asked about their sport preferences — whether they played basketball or tennis or neither.

The information was recorded in a two-way frequency table.





	Basketball (B)	No basketball (B')
Tennis (T)	25	20
No tennis (T')	10	5

a. **MC** The student sport preferences recorded in the two-way frequency table is represented on a Venn diagram as:



b. If one student is selected at random, calculate the probability that the student plays tennis only, correct to 2 decimal places.

13. Match the following Venn diagrams to the correct set notation for the shaded regions.

Venn diagram	Set notation
<p>a.</p> 	<p>A. $A \cap B$</p>
<p>b.</p> 	<p>B. A'</p>
<p>c.</p> 	<p>C. $A \cap B'$</p>
<p>d.</p> 	<p>D. $A \cup B$</p>

14. A survey of Year 8, 9 and 10 students asked them to choose dinner options for their school camp.

Year level	Lasagna	Stir-fry
8	82	75
9	67	90
10	89	45
Total	240	210


Calculate the following probabilities.

- The probability that a randomly selected student chose lasagne.
- The probability that a randomly selected student was in Year 10.
- The probability that a randomly selected student was in Year 8 and chose stir-fry.
- The probability that a randomly selected student who chose lasagna was in Year 9.


15. **MC** A bag contains three blue balls and two red balls. A ball is taken at random from the bag and its colour noted. Then a second ball is drawn, without replacing the first one.

Identify the tree diagram that best represents this sample space.


A.




B.




C.



D.



E.



LESSON

10.2 Theoretical probability

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- estimate whether events are impossible, unlikely, likely or certain
- use key probability terminologies such as trials, outcomes, sample space and events
- calculate the theoretical probability of an event.

10.2.1 The language of probability


eles-4877

- The **probability** of an event is a measure of the likelihood that the event will take place.
- If an event is **certain** to occur, then it has a probability of **1**.
- If an event is **impossible**, then it has a probability of **0**.
- The probability of any other event taking place is given by a number between 0 and 1.
- An event is likely to occur if it has a probability between 0.5 and 1.
- An event is unlikely to occur if it has a probability between 0 and 0.5.



WORKED EXAMPLE 1 Placing events on a probability scale

On the probability scale given, insert each of the following events at appropriate points.




- If today is Saturday, then tomorrow is Sunday.
- You will come to school the next Monday during a school term.
- It will snow in Victoria this year, during summer and winter.

THINK

- Carefully read the given statement and label its position on the probability scale.
 - Write the answer and provide reasoning.
- Carefully read the given statement and label its position on the probability scale.


WRITE/DRAW

a.



It is certain that if the current day is Saturday, then the next day will be Sunday.


b.



2. Write the answer and provide reasoning.

It is very likely but not certain that I will come to school on a Monday during term. Circumstances such as illness or public holidays may prevent me from coming to school on a specific Monday during a school term.

c. 1. Carefully read the given statement and label its position on the probability scale.



2. Write the answer and provide reasoning.

It is highly likely but not certain that it will snow in Victoria during winter. The chance of snow falling in Victoria in summer is highly unlikely but not impossible.

10.2.2 Key terms of probability

eles-4878

- The study of probability uses many special terms.
- **Chance experiment:** A chance experiment is a process that can be repeated many times. For example, rolling a die.
- **Trial:** A trial is one performance of an experiment to get a result. For example, each roll of the die is called a trial.
- **Outcome:** The outcome is the result obtained when the experiment is conducted. For example, when a normal six-sided die is rolled the outcome can be 1, 2, 3, 4, 5 or 6.
- **Sample space:** The set of all possible outcomes is called the sample space and is given by the symbol ξ . For the example of rolling a die, $\xi = \{1, 2, 3, 4, 5, 6\}$.
- **Event:** An event is the favourable outcome of a trial and is often represented by a capital letter. For example, when a die is rolled, A could be the event of getting an even number; $A = \{2, 4, 6\}$.
- **Favourable outcome:** A favourable outcome for an event is any outcome that belongs to the event. For event A above (rolling an even number), the favourable outcomes are 2, 4 and 6.

WORKED EXAMPLE 2 Identifying sample space, events and outcomes

For the chance experiment of rolling a die:

- list the sample space
- list the events for:
 - rolling a 4
 - rolling an even number
 - rolling at least 5
 - rolling at most 2
- list the favourable outcomes for:
 - $\{4, 5, 6\}$
 - not rolling 5
 - rolling 3 or 4
 - rolling 3 and 4.



THINK

- a. The outcomes are the numbers 1 to 6.
- b. i. This describes only one outcome.
 ii. The possible even numbers are 2, 4 and 6.
 iii. 'At least 5' means 5 is the smallest.
 iv. 'At most 2' means 2 is the largest.
- c. i. The outcomes are shown inside the brackets.
 ii. 'Not 5' means everything except 5.
 iii. The event is {3, 4}.
 iv. There is no number that is both 3 and 4.

WRITE

- a. $\xi = \{1, 2, 3, 4, 5, 6\}$
- b. i. {4}
 ii. {2, 4, 6}
 iii. {5, 6}
 iv. {1, 2}
- c. i. 4, 5, 6
 ii. 1, 2, 3, 4, 6
 iii. 3, 4
 iv. There are no favourable outcomes.

10.2.3 Theoretical probability

eles-4879

- When a coin is tossed, there are two possible outcomes, Heads or Tails. That is, $\xi = \{H, T\}$.
- We assume the coin is unbiased, meaning each outcome is **equally likely** to occur.
- Since each outcome is equally likely, they must have the same probability.
- The total of all probabilities is 1, so the probability of an outcome is given by:

$$\bullet \text{Pr}(\text{outcome}) = \frac{1}{\text{number of outcomes}}$$

$$\bullet \text{Pr}(\text{Heads}) = \frac{1}{2} \text{ and } \text{Pr}(\text{Tails}) = \frac{1}{2}$$

- The probability of an event A is found by adding up all the probabilities of the favourable outcomes in event A and dividing by the total number of outcomes.



Theoretical probability

When determining probabilities of equally likely outcomes, use the following:

- $\text{Pr}(\text{outcome}) = \frac{1}{\text{total number of outcomes}}$
- $\text{Pr}(\text{event } A) = \frac{\text{number of favourable outcomes}}{\text{total number of outcomes}}$

WORKED EXAMPLE 3 Calculating theoretical probability

A die is rolled and the number uppermost is noted. Determine the probability of each of the following events.

a. $A = \{1\}$

b. $B = \{\text{odd numbers}\}$

c. $C = \{4 \text{ or } 6\}$

THINK

There are six possible outcomes.

WRITE

a. A has one favourable outcome.

$$\text{a. } \Pr(A) = \frac{1}{6}$$

b. B has three favourable outcomes: 1, 3 and 5.

$$\begin{aligned} \text{b. } \Pr(B) &= \frac{3}{6} \\ &= \frac{1}{2} \end{aligned}$$

c. C has two favourable outcomes.

$$\begin{aligned} \text{c. } \Pr(C) &= \frac{2}{6} \\ &= \frac{1}{3} \end{aligned}$$

on Resources



eWorkbook Topic 10 Workbook (worksheets, code puzzle and project) (ewbk-2010)



Interactivities Individual pathway interactivity: Theoretical probability (int-4534)

Probability scale (int-3824)

Theoretical probability (int-6081)

Exercise 10.2 Theoretical probability

learn on

10.2 Quick quiz on

10.2 Exercise

Individual pathways

PRACTISE

1, 3, 5, 8, 10, 14, 15, 19, 22

CONSOLIDATE

2, 4, 6, 9, 11, 16, 20, 23

MASTER

7, 12, 13, 17, 18, 21, 24, 25

Fluency

1. **WE1** On the given probability scale, insert each of the following events at appropriate points.



Indicate the chance of each event using one of the following terms: certain, likely, unlikely, impossible.

- The school will have a lunch break on Friday.
 - Australia will host two consecutive Olympic Games.
2. On the given probability scale, insert each of the following events at appropriate points.



Indicate the chance of each event using one of the following terms: certain, likely, unlikely, impossible.

- At least one student in a particular class will obtain an A for Mathematics.
- Australia will have a swimming team in the Commonwealth Games.

3. On the given probability scale, insert each of the following events at appropriate points.
Indicate the chance of each event using one of the following terms: certain, likely, unlikely, impossible.



- a. Mathematics will be taught in secondary schools.
- b. In the future most cars will run without LPG or petrol.

4. On the given probability scale, insert each of the following events at appropriate points.
Indicate the chance of each event using one of the following terms: certain, likely, unlikely, impossible.



- a. Winter will be cold.
- b. Bean seeds, when sown, will germinate.

5. **WE2a** For each chance experiment below, list the sample space.

- a. Rolling a die.
- b. Tossing a coin.
- c. Testing a light bulb to see whether it is defective or not.
- d. Choosing a card from a normal deck and noting its colour.
- e. Choosing a card from a normal deck and noting its suit.



6. **WE2b** A normal six-sided die is rolled. List each of the following events.

- a. Rolling a number less than or equal to 3.
- b. Rolling an odd number.
- c. Rolling an even number or 1.
- d. Not rolling a 1 or 2.
- e. Rolling at most a 4.
- f. Rolling at least a 5.

7. **WE2c** A normal six-sided die is rolled. List the favourable outcomes for each of the following events.

- | | |
|---|---|
| a. $A = \{3, 5\}$ | b. $B = \{1, 2\}$ |
| c. $C = \text{'rolling a number greater than 5'}$ | d. $D = \text{'not rolling a 3 or a 4'}$ |
| e. $E = \text{'rolling an odd number or a 2'}$ | f. $F = \text{'rolling an odd number and a 2'}$ |
| g. $G = \text{'rolling an odd number and a 3'}$ | |

8. A card is selected from a normal deck of 52 cards and its suit is noted.

- a. List the sample space.
- b. List each of the following events.
 - i. Drawing a black card.
 - ii. Drawing a red card.
 - iii. Not drawing a heart.
 - iv. Drawing a black or a red card.

9. Determine the number of outcomes there are for:


- a. rolling a die
- b. tossing a coin
- c. drawing a card from a standard deck
- d. drawing a card and noting its suit
- e. noting the remainder when a number is divided by 5.

10. A card is drawn at random from a standard deck of 52 cards.

Note: 'At random' means that every card has the same chance of being selected.


Calculate the probability of selecting:

- an ace
- a king
- the 2 of spades
- a diamond.



11. **WES** A card is drawn at random from a deck of 52. Determine the probability of each event below.

- $A = \{5 \text{ of clubs}\}$
- $B = \{\text{black card}\}$
- $C = \{5 \text{ of clubs or queen of diamonds}\}$
- $D = \{\text{hearts}\}$
- $E = \{\text{hearts or clubs}\}$



12. A card is drawn at random from a deck of 52. Determine the probability of each event below.


- $F = \{\text{hearts and } 5\}$
- $G = \{\text{hearts or } 5\}$
- $H = \{\text{aces or kings}\}$
- $I = \{\text{aces and kings}\}$
- $J = \{\text{not a } 7\}$

13. A letter is chosen at random from the letters in the word PROBABILITY. Determine the probability that the letter is:

- B
- not B
- a vowel
- not a vowel.

14. The following coloured spinner is spun and the colour is noted. Determine the probability of each of the events given below.


- $A = \{\text{blue}\}$
- $B = \{\text{orange}\}$
- $C = \{\text{orange or pink}\}$
- $D = \{\text{orange and pink}\}$
- $E = \{\text{not blue}\}$



Understanding

15. A bag contains four purple balls and two green balls.

- If a ball is drawn at random, then calculate the probability that it will be:
 - purple
 - green.
- Design an experiment like the one in part **a** but where the probability of drawing a purple ball is 3 times that of drawing a green ball.



16. Design spinners (see question 14) using red, white and blue sections so that:

- each colour has the same probability of being spun
- red is twice as likely to be spun as either of the other two colours
- red is twice as likely to be spun as white and three times as likely to be spun as blue.

17. A bag contains red, green and blue marbles. Calculate how many marbles there must be in the bag for the following to be true when a single marble is selected at random from the bag.
- Each colour is equally likely to be selected and there are at least six red marbles in the bag.
 - Blue is twice as likely to be selected as the other colours and there are at least five green marbles in the bag.
 - There is a $\frac{1}{2}$ chance of selecting red, a $\frac{1}{3}$ chance of selecting green and a $\frac{1}{6}$ chance of selecting blue from the bag when there is between 30 and 40 marbles in the bag.
18. a. A bag contains seven gold and three silver coins. If a coin is drawn at random from the bag, calculate the probability that it will be:
- gold
 - silver.
- b. After a gold coin is taken out of the bag, a second coin is then selected at random. Assuming the first coin was not returned to the bag, calculate the probability that the second coin will be:
- gold
 - silver.

Reasoning


19. Do you think that the probability of tossing Heads is the same as the probability of tossing Tails if your friend tosses the coin? Suggest some reasons that it might not be.
20. If the following four probabilities were given to you, explain which two you would say were not correct.

0.725, -0.5, 0.005, 1.05


21. A coin is going to be tossed five times in a row. During the first four flips the coin comes up Heads each time. What is the probability that the coin will come up Heads again on the fifth flip? Justify your answer.

Problem solving

22. Consider the spinner shown. Discuss whether the spinner has an equal chance of falling on each of the colours.
23. A box contains two coins. One is a double-headed coin, and the other is a normal coin with Heads on one side and Tails on the other. You draw one of the coins from a box and look at one of the sides. It is Heads. Determine the probability that the other side also shows Heads.
24. 'Unders and Overs' is a game played with two normal six-sided dice. The two dice are rolled, and the numbers uppermost added to give a total. Players bet on the outcome being 'under 7', 'equal to 7' or 'over 7'.
If you had to choose one of these outcomes, which would you choose? Explain why.



25. Justine and Mary have designed a new darts game for their Year 9 Fete Day. Instead of a circular dart board, their dart board is in the shape of two equilateral triangles. The inner triangle (bullseye) has a side length of 3 cm, while the outer triangle has side length 10 cm. Given that a player's dart falls in one of the triangles, determine the probability that it lands in the bullseye. Write your answer correct to 2 decimal places.



LESSON

10.3 Experimental probability

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- determine the relative frequency of an outcome
- understand the difference between theoretical probability and experimental probability.

10.3.1 Relative frequency

eles-4880

- A die is rolled 12 times and the outcomes are recorded in the table shown.

Outcome	1	2	3	4	5	6
Frequency	3	1	1	2	2	3

In this chance experiment there were 12 trials.

The table shows that the number 1 was rolled 3 times out of 12.

- So the **relative frequency** of 1 is 3 out of 12, or $\frac{3}{12} = \frac{1}{4}$.

As a decimal, the relative frequency of 1 is equal to 0.25.

Relative frequency

The relative frequency of an outcome is given by:

$$\text{Relative frequency} = \frac{\text{frequency of an outcome}}{\text{total number of trials}}$$

- As the number of trials becomes larger, the relative frequency of each outcome will become very close to the theoretical probability.

WORKED EXAMPLE 4 Calculating relative frequency

For the chance experiment of rolling a die, the following outcomes were noted.

Outcome	1	2	3	4	5	6
Frequency	3	1	4	6	3	3



- Calculate the number of trials.
- Identify how many threes were rolled.
- Calculate the relative frequency for each number written as a decimal.

THINK

- Adding the frequencies will give the number of trials.
- The frequency of 3 is 4.

WRITE

- $1 + 3 + 4 + 6 + 3 + 3 = 20$ trials
- 4 threes were rolled.

- c. Add a relative frequency row to the table and complete it. The relative frequency is calculated by dividing the frequency of the outcome by the total number of trials.

Outcome	1	2	3	4	5	6
Frequency	3	1	4	6	3	3
Relative frequency	$\frac{3}{20} = 0.15$	$\frac{1}{20} = 0.05$	$\frac{4}{20} = 0.2$	$\frac{6}{20} = 0.3$	$\frac{3}{20} = 0.15$	$\frac{3}{20} = 0.15$

10.3.2 Experimental probability

eles-4881


- When it is not possible to calculate the theoretical probability of an outcome, carrying out simulations involving repeated trials can be used to determine the **experimental probability**.
- The relative frequency of an outcome is the experimental probability.

Experimental probability

- The experimental probability of an outcome is given by:

$$\text{Experimental probability} = \frac{\text{frequency of an outcome}}{\text{total number of trials}}$$

- The spinner shown is not symmetrical, and the probability of each outcome cannot be determined theoretically.
- The experimental probability of each outcome can be found by using the spinner many times and recording the outcomes. As more trials are conducted, the experimental probability will become more accurate and closer to the true probability of each section.



WORKED EXAMPLE 5 Calculating experimental probabilities

The spinner shown above was spun 100 times and the following results were achieved.

Outcome	1	2	3	4
Frequency	7	26	9	58

- Calculate the number of trials.
- Calculate the experimental probability of each outcome.
- Calculate the sum of the four probabilities.

THINK

- Adding the frequencies will determine the number of trials.

WRITE

a. $7 + 26 + 9 + 58 = 100$ trials

b. The experimental probability equals the relative frequency. This is calculated by dividing the frequency of the outcome by the total number of trials.

$$\text{b. Pr}(1) = \frac{7}{100}$$

$$= 0.07$$

$$\text{Pr}(2) = \frac{26}{100}$$

$$= 0.26$$

$$\text{Pr}(3) = \frac{9}{100}$$

$$= 0.09$$

$$\text{Pr}(4) = \frac{58}{100}$$

$$= 0.58$$

c. Add the probabilities (they should equal 1).

$$\text{c. } 0.07 + 0.26 + 0.09 + 0.58 = 1$$



eles-4882

10.3.3 Expected number of results

- If we tossed a coin 100 times, we would expect there to be 50 Heads, since $\text{Pr}(\text{Heads}) = \frac{1}{2}$.

Expected number of results

The expected number of favourable outcomes from a series of trials is found from:

$$\text{Expected number} = \text{probability of outcome} \times \text{number of trials}$$

- The probability of an outcome can be the theoretical probability or an experimental probability.

WORKED EXAMPLE 6 Expected number of outcomes

Calculate the expected number of results in the following situations.

- The number of Tails after flipping a coin 250 times.
- The number of times a one comes up after rolling a dice 120 times.
- The number of times a royal card is picked from a deck that is reshuffled with the card replaced 650 times.

THINK

a. The probability of getting a Tail is $\text{Pr}(\text{Tails}) = \frac{1}{2}$.

b. The probability of getting a one is $\text{Pr}(\text{one}) = \frac{1}{6}$.

WRITE

$$\begin{aligned} \text{a. Number of Tails} &= \text{Pr}(\text{Tails}) \times 250 \\ &= \frac{1}{2} \times 250 \\ &= 125 \text{ times} \end{aligned}$$

$$\begin{aligned} \text{b. Number of ones} &= \text{Pr}(\text{one}) \times 120 \\ &= \frac{1}{6} \times 120 \\ &= 20 \text{ times} \end{aligned}$$

c. A deck of cards has 52 cards and 12 of them are royal cards. This means the probability of getting a royal is $\text{Pr}(\text{royal}) = \frac{12}{52} = \frac{3}{13}$.

c. Number of royals = $\text{Pr}(\text{royal}) \times 650$
 $= \frac{3}{13} \times 650$
 $= 150$ times

Digital technology

There are digital tools available to conduct probability simulations, including calculator technology.





TI | THINK

In a new document on a Calculator page, select:

- Menu
- 9 Functions & Programs
- 1 Program Editor
- 1 New

Call the program 'Dice' and select LibPub (Show in Catalog) for Library Access. Press OK.

WRITE/DISPLAY





Localize a die, namely the green die. This is to make the variables used only in this program. More dice can be added by naming them with different names.

Initialise them, to make sure no other values are already in a variable. The arrow is found by sto-> (ctl+var).


Store a randomly generated integer from 1 to 6 into the die. Use $\text{int}(6 \cdot \text{rand} + 1)$ or $\text{randInt}(1,6)$.

CASIO | THINK

In a Main screen select the drop-down box in the  menu and choose the probability simulator by selecting 


The following menu will appear as shown.

WRITE/DISPLAY



To simulate rolling a six sided die once, select OK. This simulation resulted in a 4.

The number of trials and number of faces can be adjusted to suit and there is also an option to change the experiment to simulate picking items out of a container.



TI | THINK

To display results, enter the following:

Disp green, "green"

To check syntax and store, go to Menu and select:

- 2 Check Syntax & Store
- 1 Check Syntax & Store

WRITE/DISPLAY

```
* dice 4/4
Define LibPub dice()=
Prgm
Local green
0 → green
randInt(1,6) → green
Disp green, "green"
EndPrgm
```

```
1 Actions
2 Chec1 Check Syntax & Store (Ctrl+B)
3 Defin2 Check Syntax
4 Cont3 Run (Ctrl+R)
5 Transfers
6 I/O
7 Mode
8 Hub
9 Draw
EndPrgm
```

```
"dice" stored successfu
Define LibPub dice()=
Prgm
Local green
0 → green
randInt(1,6) → green
Disp green, "green"
EndPrgm
```

Enter the following on the left side of the document page:

dice ()

Press Enter.

This simulation will generate values by pressing Enter and the program will keep on generating the values.

The number of trials and number of faces can be adjusted to suit.

```
"dice" stored successfu
Define LibPub dice()=
Prgm
Local green
0 → green
randInt(1,6) → green
Disp green, "green"
EndPrgm
6 green
Done
```

```
"dice" stored successfu
Define LibPub dice()=
Prgm
Local green
0 → green
randInt(1,6) → green
Disp green, "green"
EndPrgm
1 green
Done
```

CASIO | THINK

WRITE/DISPLAY

COLLABORATIVE TASK

Construct an irregular spinner using cardboard and a toothpick. By carrying out a number of trials, estimate the probability of each outcome.

on Resources

- eWorkbook** Topic 10 Workbook (worksheets, code puzzle and project) (ewbk-2010)
- Interactivities** Individual pathway interactivity: Experimental probability (int-4535)
Experimental probability (int-3825)

10.3 Quick quiz **on**

10.3 Exercise

Individual pathways

PRACTISE

1, 3, 8, 10, 11, 15, 17, 19, 22

CONSOLIDATE

2, 4, 6, 9, 12, 18, 20, 23, 24

MASTER

5, 7, 13, 14, 16, 21, 25, 26, 27

Fluency

1. **WE4** Each of the two tables shown contains the results of a chance experiment (rolling a die).

For each table, calculate:

- the number of trials held
- the number of fives rolled
- the relative frequency for each outcome, correct to 2 decimal places
- the sum of the relative frequencies.

a.

Number	1	2	3	4	5	6
Frequency	3	1	5	2	4	1

b.

Number	1	2	3	4	5	6
Frequency	52	38	45	49	40	46

2. A coin is tossed in two chance experiments. The outcomes are recorded in the tables shown. For each experiment, calculate:

- the relative frequency of both outcomes
- the sum of the relative frequencies.

a.

Outcome	H	T
Frequency	22	28

b.

Outcome	H	T
Frequency	31	19

3. **WE5** An unbalanced die was rolled 200 times and the following outcomes were recorded.

Number	1	2	3	4	5	6
Frequency	18	32	25	29	23	73

Using these results, calculate:

- $\text{Pr}(6)$
- $\text{Pr}(\text{odd number})$
- $\text{Pr}(\text{at most } 2)$
- $\text{Pr}(\text{not } 3)$.

4. A box of matches claims on its cover to contain 100 matches.
A survey of 200 boxes established the following results.

Number of matches	95	96	97	98	99	100	101	102	103	104
Frequency	1	13	14	17	27	55	30	16	13	14

If you were to purchase a box of these matches, calculate the probability that:

- the box would contain 100 matches
 - the box would contain at least 100 matches
 - the box would contain more than 100 matches
 - the box would contain no more than 100 matches.
5. A packet of chips is labelled as weighing 170 grams. This is not always the case and there will be some variation in the weight of each packet.

A packet of chips is considered underweight if its weight is below 168 grams. Chips are made in batches of 1000 at a time. A sample of the weights from a particular batch are shown below.

Weight (grams)	166	167	168	169	170	171	172	173	174
Frequency	2	3	13	17	27	18	9	5	1

Calculate the probability that:

- a packet of chips is its advertised weight
- a packet of chips is above its advertised weight
- a packet of chips is underweight.

A batch of chips is rejected if more than 50 packets in a batch are classed as underweight.
d. Explain if the batch that this sample of chips is taken from should be rejected.




Understanding

6. Here is a series of statements based on experimental probability. If a statement is not reasonable, give a reason why.
- I tossed a coin five times and there were four Heads, so $\text{Pr}(\text{Heads}) = 0.8$.
 - Sydney Roosters have won 1064 matches out of the 2045 that they have played, so $\text{Pr}(\text{Sydney will win their next game}) = 0.52$.
 - $\text{Pr}(\text{the sun will rise tomorrow}) = 1$
 - At a factory, a test of 10 000 light globes showed that 7 were faulty. Therefore, $\text{Pr}(\text{faulty light globe}) = 0.0007$.
 - In Sydney it rains an average of 143.7 days each year, so $\text{Pr}(\text{it will rain in Sydney on the 17th of next month}) = 0.39$.
7. At a birthday party, some cans of soft drink were put in a container of ice. There were 16 cans of Coke, 20 cans of Sprite, 13 cans of Fanta, 8 cans of Sunkist and 15 cans of Pepsi.
If a can was picked at random, calculate the probability that it was:
- a can of Pepsi
 - not a can of Fanta.
8. **WE6** Calculate the expected number of Tails if a fair coin is tossed 400 times.
9. Calculate the expected number of threes if a fair die is rolled 120 times.

10. **MC** In Tattslotto, six numbers are drawn from the numbers 1, 2, 3, ... 45. The number of different combinations of six numbers is 8 145 060. If you buy one ticket, select from the following the probability that you will win the draw.

- A. $\frac{1}{8\,145\,060}$ B. $\frac{1}{45}$
 C. $\frac{45}{8\,145\,060}$ D. $\frac{1}{6}$
 E. $\frac{6}{8\,145\,060}$



11. **MC** A survey of high school students asked 'Should Saturday be a normal school day?' 350 students voted yes, and 450 voted no. From the following, select the probability that a student chosen at random said no.

- A. $\frac{7}{16}$ B. $\frac{9}{16}$ C. $\frac{7}{9}$ D. $\frac{9}{14}$ E. $\frac{1}{350}$

12. In a poll of 200 people, 110 supported party M, 60 supported party N and 30 were undecided. If a person is chosen at random from this group of people, calculate the probability that he or she:

- a. supports party M b. supports party N
 c. supports a party d. is not sure what party to support.

13. A random number is picked from $N = \{1, 2, 3, \dots, 100\}$. Calculate the probability of picking a number that is:
- a. a multiple of 3 b. a multiple of 4 or 5 c. a multiple of 5 and 6.

14. The numbers 3, 5 and 6 are combined to form a three-digit number such that no digit may be repeated.

- a. i. Determine how many numbers can be formed.
 ii. List them.
 b. Determine $\text{Pr}(\text{the number is odd})$.
 c. Determine $\text{Pr}(\text{the number is even})$.
 d. Determine $\text{Pr}(\text{the number is a multiple of 5})$.

15. Gorri was a disc-bowling game played in Central Australia which involved the throwing of a spear at a moving target. If a person had success hitting the target 2 out of every 10 times, calculate the number of times they are expected to hit the target in 25 games.



16. Svetlana, Sarah, Leonie and Trang are volleyball players. The probabilities that they will score a point on serve are 0.6, 0.4, 0.3 and 0.2 respectively. Calculate how many points on serve are expected from each player if they serve 10 times each.

17. **MC** A survey of the favourite leisure activity of 200 Year 9 students produced the following results.

Activity	Playing sport	Fishing	Watching TV	Video games	Surfing
Number of students	58	26	28	38	50

The probability (given as a percentage) that a student selected at random from this group will have surfing as their favourite leisure activity is:

- A. 50% B. 100% C. 25% D. 0% E. 29%

18. The numbers 1, 2 and 5 are combined to form a three-digit number, allowing for any digit to be repeated up to three times.
- Determine how many different numbers can be formed.
 - List the numbers.
 - Determine $\Pr(\text{the number is even})$.
 - Determine $\Pr(\text{the number is odd})$.
 - Determine $\Pr(\text{the number is a multiple of 3})$.

Reasoning

19. John has a 12-sided die numbered 1 to 12 and Lisa has a 20-sided die numbered 1 to 20. They are playing a game where the first person to get the number 10 wins. They are rolling their dice individually.
- Calculate $\Pr(\text{John gets a 10})$.
 - Calculate $\Pr(\text{Lisa gets a 10})$.
 - Explain whether the game is fair.
20. At a supermarket checkout, the scanners have temporarily broken down and the cashiers must enter in the bar codes manually. One particular cashier overcharged 7 of the last 10 customers she served by entering the incorrect bar code.
- Based on the cashier's record, determine the probability of making a mistake with the next customer.
 - Explain if another customer should have any objections with being served by this cashier.
21. If you flip a coin six times, determine how many of the possible outcomes could include a Tail on the second toss.

Problem solving

22. In a jar, there are 600 red balls, 400 green balls, and an unknown number of yellow balls. If the probability of selecting a green ball is $\frac{1}{5}$, determine how many yellow balls are in the jar.
23. In a jar there are an unknown number of balls, N , with 20 of them green. The other colours contained in the jar are red, yellow and blue, with $\Pr(\text{red or yellow}) = \frac{1}{2}$, $\Pr(\text{red or green}) = \frac{1}{4}$ and $\Pr(\text{blue}) = \frac{1}{3}$. Determine the number of red, yellow and blue balls in the jar.
24. The biological sex of babies in a set of triplets is simulated by flipping three coins. If a coin lands Tails up, the baby is male. If a coin lands Heads up, the baby is female. In the simulation, the trial is repeated 40 times. The following results show the number of Heads obtained in each trial:
0, 3, 2, 1, 1, 0, 1, 2, 1, 0, 1, 0, 2, 0, 1, 0, 1, 2, 3, 2, 1, 3, 0, 2, 1, 2, 0, 3, 1, 3, 0, 1, 0, 1, 3, 2, 2, 1, 2, 1
- Calculate the probability that exactly one of the babies in a set of triplets is female.
 - Calculate the probability that more than one of the babies in the set of triplets is female.
25. Use your calculator to generate three random sets of numbers between 1 and 6 to simulate rolling a dice. The number of trials in each set will be:
- 10
 - 25
 - 50
- Calculate the relative frequency of each outcome (1 to 6) and comment on what you notice about experimental and theoretical probability.

26. A survey of the favourite foods of Year 9 students is recorded, with the following results.

Meal	Tally
Hamburger	45
Fish and chips	31
Macaroni and cheese	30
Lamb souvlaki	25
BBQ pork ribs	21
Cornflakes	17
T-bone steak	14
Banana split	12
Corn-on-the-cob	9
Hot dogs	8
Garden salad	8
Veggie burger	7
Smoked salmon	6
Muesli	5
Fruit salad	3

- Estimate the probability that macaroni and cheese is the favourite food of a randomly selected Year 9 student.
- Estimate the probability that a vegetarian dish is a randomly selected student's favourite food.
- Estimate the probability that a beef dish is a randomly selected student's favourite food.

27. A spinner has six sections of different sizes. Steven conducts an experiment and finds the following results:

Section	1	2	3	4	5	6
Frequency	6	24	15	30	48	12

Determine the angle size of each section of the spinner using the results above.

LESSON

10.4 Venn diagrams and two-way tables

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify the complement of an event
- create and interpret Venn diagrams and two-way tables
- use a Venn diagram or two-way table to determine $A \cap B$
- use a Venn diagram or two-way table to determine $A \cup B$.




10.4.1 The complement of an event


eles-4883

- Suppose that a die is rolled. The sample space is given by: $\xi = \{1, 2, 3, 4, 5, 6\}$.
- If A is the event 'rolling an odd number', then $A = \{1, 3, 5\}$.

- There is another event called ‘the **complement** of A' , or not A . This event contains all the outcomes that do not belong to A . It is given by the symbol A' .
- In this case $A' = \{2, 4, 6\}$.
- A and A' can be shown on a **Venn diagram**.



A is shaded.



A' (not A) is shaded.

Complementary events

- For the event A , the complement is denoted A' , and the two are related by the following:

$$\Pr(A) + \Pr(A') = 1$$

WORKED EXAMPLE 7 Determining the complement

For the sample space $\xi = \{1, 2, 3, 4, 5\}$, list the complement of each of the following events.

- $A = \{\text{multiples of } 3\}$
- $B = \{\text{square numbers}\}$
- $C = \{1, 2, 3, 5\}$

THINK

- The only multiple of 3 in the set is 3. Therefore $A = \{3\}$. A' is every other element of the set.
- The only square numbers are 1 and 4. Therefore $B = \{1, 4\}$. B' is every other element of the set.
- $C = \{1, 2, 3, 5\}$. C' is every other element of the set.

WRITE


- $A' = \{1, 2, 4, 5\}$
- $B' = \{2, 3, 5\}$
- $C' = \{4\}$




10.4.2 Venn diagrams: the intersection of events

eles-4884


- A Venn diagram consists of a rectangle and one or more circles.
- A Venn diagram contains all possible outcomes in the sample space and will have the ξ symbol in the top left corner.
- A Venn diagram is used to illustrate the relationship between sets of objects or numbers.
- All outcomes for a given event will be contained within a specific circle.
- Outcomes that belong to multiple events will be found in the overlapping region of two or more circles.
- The overlapping region of two circles is called the intersection of the two events and is represented using the \cap symbol.



The circle on the left contains all outcomes in event A .




The circle on the right contains all outcomes in event B .



The overlap or intersection of the two circles contains the outcomes that are in event A 'and' in event B . This is denoted by $A \cap B$.

- A Venn diagram for two events A and B has four distinct regions.
- $A \cap B'$ contains the outcomes in event A and not in event B .
- $A \cap B$ contains the outcomes in event A and in event B .
- $A' \cap B$ contains the outcomes not in event A and in event B .
- $A' \cap B'$ contains the outcomes not in event A and not in event B .



WORKED EXAMPLE 8 Setting up a Venn diagram

In a class of 20 students, 5 study Art, 9 study Biology and 2 students study both.

Let $A = \{\text{students who study Art}\}$ and $B = \{\text{students who study Biology}\}$.

- Create a Venn diagram to represent this information.
- Identify the number of students represented by the following and state what these regions represent:
 - $A \cap B$
 - $A \cap B'$
 - $A' \cap B$
 - $A' \cap B'$

THINK

- Draw a sample space with events A and B .




Place a 2 in the intersection of both circles since we know 2 students take both subjects.

Since 5 study Art and there are 2 already in the middle, place a 3 in the remaining section of circle A .


Since 9 study Biology and there are 2 already in the middle, place a seven in the remaining section of circle B .

The total number inside the three circles is $3 + 2 + 7 = 12$. This means there must be $20 - 12 = 8$ outside of the two circles. Place an 8 outside the circles, within the rectangle.


WRITE/DRAW




- b. i. From the Venn diagram, $A \cap B = 2$. These are the students in both A and B .




- ii. From the Venn diagram, $A \cap B' = 3$. These are the students in A and not in B .



- iii. From the Venn diagram, $A' \cap B = 7$. These are the students not in A and in B .



- iv. From the Venn diagram, $A' \cap B' = 8$. These are the students not in A and not in B .



- b. i. There are 2 students who study Art and Biology.

- ii. There are 3 students who study Art and not Biology, i.e. 3 students study Art only.

- iii. There are 7 students who do not study Art and do study Biology, i.e. 7 students study Biology only.


- iv. There are 8 students who do not study Art and do not study Biology.

10.4.3 Two-way tables

eles-4885

- The information in a Venn diagram can also be represented using a **two-way table**. The relationship between the two is shown below.

	Event B	Event B'	Total
Event A	$A \cap B$	$A \cap B'$	A
Event A'	$A' \cap B$	$A' \cap B'$	A'
Total	B	B'	



WORKED EXAMPLE 9 Creating a two-way table

In a class of 20 students, 5 study Art, 9 study Biology and 2 students study both.
Create a two-way table to represent this information.

THINK

1. Create an empty two-way table.

WRITE/DRAW

	Biology	Not Biology	Total
Art			
Not Art			
Total			

2. Fill the table in with the information provided in the question.

- 2 students study both subjects
- 5 in total take Art
- 9 in total take Biology
- 20 students in the class

	Biology	Not Biology	Total
Art	2		5
Not Art			
Total	9		20

3. Use the totals of the rows and columns to fill in the gaps in the table.

- $9 - 2 = 7$ Biology and not Art
- $5 - 2 = 3$ Art and not Biology
- $20 - 9 = 11$ not Biology total
- $20 - 5 = 15$ not Art total

	Biology	Not Biology	Total
Art	2	3	5
Not Art	7	8	15
Total	9	11	20

4. The last value, not Biology and not Art, can be found from either of the following:

- $11 - 3 = 8$
- $15 - 7 = 8$

10.4.4 Number of outcomes


eles-4886

- If event A contains seven outcomes or members, this is written as $n(A) = 7$.
- So $n(A \cap B') = 3$ means that there are three outcomes that are in event A and not in event B .

WORKED EXAMPLE 10 Determining the number of outcomes in an event


For the Venn diagram shown, write down the number of outcomes in each of the following.

- a. M b. M' c. $M \cap N$ d. $M \cap N'$ e. $M' \cap N'$




THINK

a. Identify the regions showing M and add the outcomes.

WRITE/DRAW


$$n(M) = 6 + 11 = 17$$

b. Identify the regions showing M' and add the outcomes.




$$n(M') = 4 + 15 = 19$$

c. $M \cap N$ means 'M and N'. Identify the region.




$$n(M \cap N) = 11$$

d. $M \cap N'$ means 'M and not N'. Identify the region.



$$n(M \cap N') = 6$$


e. $M' \cap N'$ means 'not M and not N'. Identify the region.



$$n(M' \cap N') = 4$$

WORKED EXAMPLE 11 Using a Venn diagram to create a two-way table

Show the information from the Venn diagram on a two-way table.



THINK

1. Draw a 2×2 table and add the labels A , A' , B and B' .
2. There are 7 elements in A and B .
There are 3 elements in A and 'not B '.
There are 2 elements in 'not A ' and B .
There are 5 elements in 'not A ' and 'not B '.
3. Add in a column and a row to show the totals.

WRITE/DRAW

	A	A'
B		
B'		

	A	A'
B	7	2
B'	3	5

	A	A'	Total
B	7	2	9
B'	3	5	8
Total	10	7	17

WORKED EXAMPLE 12 Using a two-way table to create a Venn diagram

Show the information from the two-way table on a Venn diagram.

	Left-handed	Right-handed
Blue eyes	7	20
Not blue eyes	17	48



THINK

Draw a Venn diagram that includes a sample space and events L for left-handedness and B for blue eyes. (Right-handedness = L' .)


$$n(L \cap B) = 7$$

$$n(L \cap B') = 17$$

$$n(L' \cap B) = 20$$

$$n(L' \cap B') = 48$$

WRITE/DRAW



WORKED EXAMPLE 13 Probability from a two-way table

In a class of 30 students, 15 swim for exercise and 20 run for exercise and 5 participate in neither activity.

- Create a two-way table to represent this information.
- Calculate the probability that a randomly chosen student from this class does running and swimming for exercise.

THINK

1. Create an empty two-way table.
 2. Fill the table in with the information provided in the question.
 - 5 students do neither activity.
 - 15 in total swim.
 - 20 in total run.
 - 30 students are in the class.
 3. Use the totals of the rows and columns to fill in the gaps in the table.

- The probability a student runs and swims is given by:

$$\Pr(\text{run} \cap \text{swim}) = \frac{\text{number in run} \cap \text{swim}}{\text{total in the class}}$$

WRITE/DRAW

- | | Swim | Not swim | Total |
|---------|------|----------|-------|
| Run | | | |
| Not run | | | |
| Total | | | |

- | | Swim | Not swim | Total |
|---------|------|----------|-------|
| Run | | | 20 |
| Not run | | 5 | |
| Total | 15 | | 30 |

- | | Swim | Not swim | Total |
|---------|------|----------|-------|
| Run | 10 | 10 | 20 |
| Not run | 5 | 5 | 10 |
| Total | 15 | 15 | 30 |


- $$\begin{aligned} \Pr(\text{run} \cap \text{swim}) &= \frac{\text{number in run} \cap \text{swim}}{\text{total in the class}} \\ &= \frac{10}{30} \\ &= \frac{1}{3} \end{aligned}$$




eles-4887

10.4.5 Venn diagrams: the union of events

- The **intersection** of two events ($A \cap B$) is all outcomes in event A 'and' in event B .
- The **union** of two events ($A \cup B$) is all outcomes in events A 'or' in event B .



The intersection of the two circles contains the outcomes that are in event A 'and' in event B . This is denoted by $A \cap B$.




Everything contained within the two circles is an outcome that is in event A 'or' in event B . This is denoted by $A \cup B$.

WORKED EXAMPLE 14 Calculating the union of two events

Use the Venn diagram shown to calculate the value of the following:

- $n(A)$
- $\Pr(B)$
- $n(A \cup B)$
- $\Pr(A \cap B)$



THINK

- Identify the number of outcomes in the A circle.
- Identify the number of outcomes in the B circle.

$$\Pr(B) = \frac{\text{number of favourable outcomes in } B}{\text{total number of outcomes}}$$

- Identify the number of outcomes in either of the two circles.
- Identify the number of outcomes in the intersection of the two circles.

$$\Pr(A \cap B) = \frac{\text{favourable outcomes in } A \cap B}{\text{total number of outcomes}}$$

WRITE

$$\text{a. } n(A) = 29 + 16 = 45$$

$$\begin{aligned} \text{b. } \Pr(B) &= \frac{\text{favourable outcomes in } B}{\text{total number of outcomes}} \\ &= \frac{n(B)}{\text{total number of outcomes}} \\ &= \frac{16 + 31}{29 + 16 + 31 + 44} \\ &= \frac{47}{120} \end{aligned}$$

$$\text{c. } n(A \cup B) = 29 + 16 + 31 = 76$$

$$\begin{aligned} \text{d. } \Pr(A \cap B) &= \frac{\text{number of favourable outcomes in } A \cap B}{\text{total number of outcomes}} \\ &= \frac{n(A \cap B)}{\text{total number of outcomes}} \\ &= \frac{16}{29 + 16 + 31 + 44} \\ &= \frac{16}{120} \\ &= \frac{2}{15} \end{aligned}$$

WORKED EXAMPLE 15 Completing a Venn diagram

- Place the elements of the following sets of numbers in their correct position in a single Venn diagram.

$$\xi = \{\text{Number 1 to 20 inclusive}\}$$

$$A = \{\text{Multiples of 3 from 1 to 20 inclusive}\}$$

$$B = \{\text{Multiples of 2 from 1 to 20 inclusive}\}$$

- Use this Venn diagram to determine the following:

- $A \cap B$

- $A \cup B$

- $A \cap B'$

- $A' \cup B'$

THINK

- a. Write out the numbers in each event A and B :

$$A = \{3, 6, 9, 12, 15, 18\}$$


$$B = \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20\}$$

Identify the numbers that appear in both sets. In this case it is 6, 12 and 18. These numbers will be placed in the overlap of the two circles for A and B .

All numbers not in A or B are placed outside the two circles.

After placing the numbers in the Venn diagram, check that all numbers from 1 to 20 are written down.

- b. i. $A \cap B$ are the numbers in A 'and' in B .
 ii. $A \cup B$ are the numbers in A 'or' in B .
 iii. $A \cap B'$ are the number in A and not in B . Refer to the four sections of the Venn diagram to locate this region.
 iv. $A' \cup B'$ is any number that is not in A 'or' not in B . This ends up being any number not in $A \cap B$.

WRITE/DRAW

- b. i. $A \cap B = \{6, 12, 18\}$
 ii. $A \cup B = \{2, 3, 4, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20\}$
 iii. $A \cap B' = \{3, 9, 15\}$
 iv. $A' \cup B' = \{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 19, 20\}$


WORKED EXAMPLE 16 Calculating the probability of the union of two events

In a class of 24 students, 11 students play basketball, 7 play tennis, and 4 play both sports.


- a. Show the information on a Venn diagram.
 b. If one student is selected at random, then calculate the probability that:
 i. the student plays basketball
 ii. the student plays tennis or basketball
 iii. the student plays tennis or basketball but not both.

THINK


- a. 1. Draw a sample space with events B and T .

WRITE/DRAW

2. $n(B \cap T) = 4$
 $n(B \cap T') = 11 - 4 = 7$
 $n(T \cap B') = 7 - 4 = 3$
 So far, 14 students out of 24 have been placed.
 $n(B' \cap T') = 24 - 14 = 10$




- b. i. Identify the number of students who play basketball.




$$\Pr(B) = \frac{\text{number of favourable outcomes}}{\text{total number of outcomes}}$$

- ii. Identify the number of students who play tennis or basketball.



- iii. Identify the number of students who play tennis or basketball but not both.



- b. i. $\Pr(B) = \frac{\text{number of students who play basketball}}{\text{total number of students}}$

$$= \frac{n(B)}{24}$$

$$= \frac{11}{24}$$

- ii. $\Pr(T \cup B) = \frac{n(T \cup B)}{24}$

$$= \frac{14}{24}$$

$$= \frac{7}{12}$$

- iii. $n(B \cap T') + n(B' \cap T) = 3 + 7$
 $= 10$




Pr(tennis or basketball but not both)

$$= \frac{10}{24}$$

$$= \frac{5}{12}$$

DISCUSSION

How will you remember the difference between when one event **and** another occurs and when one event **or** another occurs?

-  **eWorkbook** Topic 10 Workbook (worksheets, code puzzle and project) (ewbk-2010)
-  **Video eLesson** Venn diagrams (eles-1934)
-  **Interactivities** Individual pathway interactivity: Venn diagrams and two-way tables (int-4536)
Venn diagrams (int-3828)
Two-way tables (int-6082)

Exercise 10.4 Venn diagrams and two-way tables

10.4 Quick quiz **on**

10.4 Exercise

Individual pathways

PRACTISE

1, 4, 5, 10, 13, 16, 17, 20, 24

CONSOLIDATE


2, 6, 7, 9, 11, 14, 18, 21, 25

MASTER

3, 8, 12, 15, 19, 22, 23, 26, 27

Fluency

1. **WE7** For the sample space $\xi = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$, list the complement of each of the following events.
 - a. $A = \{\text{evens}\}$
 - b. $B = \{\text{multiples of 5}\}$
 - c. $C = \{\text{squares}\}$
 - d. $D = \{\text{numbers less than 8}\}$
2. If $\xi = \{11, 12, 13, 14, 15, 16, 17, 18, 19, 20\}$, list the complement of each of the following events.
 - a. $A = \{\text{multiples of 3}\}$
 - b. $B = \{\text{numbers less than 20}\}$
 - c. $C = \{\text{prime numbers}\}$
 - d. $D = \{\text{odd numbers or numbers greater than 16}\}$
3. If $\xi = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20\}$, list the complement of each of the following events.
 - a. $A = \{\text{multiples of 4}\}$
 - b. $B = \{\text{primes}\}$
 - c. $C = \{\text{even and less than 13}\}$
 - d. $D = \{\text{even or greater than 13}\}$
4. **WE10** For the Venn diagram shown, write down the number of outcomes in:



- a. ξ
 - b. S
 - c. T
 - d. $T \cap S$
 - e. $T \cap S'$
 - f. $S' \cap T'$
5. **WE11** Show the information from question 4 on a two-way table.




6. **WE12** Show the information from this two-way table on a Venn diagram.

	S	S'
V	21	7
V'	2	10

7. For each of the following Venn diagrams, use set notation to write the name of the region coloured in:

i. blue

ii. pink.



8. **WE9&13** The membership of a tennis club consists of 55 men and 45 women. There are 27 left-handed people, including 15 men.

- Show the information on a two-way table.
- Show the information on a Venn diagram.
- If one member is chosen at random, calculate the probability that the person is:
 - right-handed
 - a right-handed man
 - a left-handed woman.




9. **WE14** Using the information given in the Venn diagram, if one outcome is chosen at random, determine:

a. $\Pr(L)$

b. $\Pr(L')$

c. $\Pr(L \cap M)$

d. $\Pr(L \cap M')$



10. **WE15** Place the elements of the following sets of numbers in their correct position in a single Venn diagram.

$\xi = \{\text{numbers between 1 to 10 inclusive}\}$
 $A = \{\text{odd numbers from 1 to 10}\}$
 $B = \{\text{sqared numbers between 1 to 10 inclusive}\}$

11. Place the elements of the following sets of numbers in their correct position in a single Venn diagram.

$\xi = \{\text{numbers between 1 to 25 inclusive}\}$
 $A = \{\text{multiples of 3 from 1 to 25}\}$
 $B = \{\text{numbers that are odd or over 17 from 1 to 25 inclusive}\}$

12. Place the elements of the following sets of numbers in their correct position in a single Venn diagram.

$A = \{\text{prime numbers from 1 to 20}\}$

$B = \{\text{even numbers from 1 to 20}\}$

$C = \{\text{multiples of 5 from 1 to 20}\}$

$\xi = \{\text{numbers between 1 and 20 inclusive}\}$

13. Using the information given in the table, if one family is chosen at random, calculate the probability that they own:

Pets owned by families		
	Cat	No cat
Dog	4	11
No dog	16	9

- a. a cat
 b. a cat and a dog
 c. a cat or a dog or both
 d. a cat or a dog but not both
 e. neither a cat nor a dog.
14. Using the information given in the table, if a person is chosen at random, calculate the probability that for exercise, this person:

Type of exercise		
	Cycling	No cycling
Running	12	19
No running	13	6


- a. cycles
 b. cycles and runs
 c. cycles or runs
 d. cycles or runs but not both.
15. A barista decides to record what the first 45 customers order on a particular morning. The information is partially filled out in the table shown.

	Croissant	No croissant	Total
Coffee	27		
No coffee		7	
Total	36		

- a. Fill in the missing information in the table.
 b. Using the table, calculate the probability that the next customer:
 i. orders a coffee and a croissant
 ii. order a coffee or a croissant
 iii. orders a coffee or a croissant but not both.
 c. The barista serves an average of 315 customers a day.
 If the café is open six days a week, calculate how many coffees the barista will make each week.

Understanding

16. A group of athletes was surveyed and the results were shown on a Venn diagram. $S = \{\text{sprinters}\}$ and $L = \{\text{long jumpers}\}$.



- Write down how many athletes were included in the survey.
- If one of the athletes is chosen at random, calculate the probability that the athlete competes in:
 - long jump
 - long jump and sprints
 - long jump or sprints
 - long jump or sprints but not both.

17. **WE8a,16** In a class of 40 students, 26 take a train to school, 19 take a bus and 8 take neither of these. Let $T = \{\text{takes the train}\}$ and $B = \{\text{takes the bus}\}$.

- Show the information on a Venn diagram.
- If one student is selected at random, calculate the probability that:
 - the student takes the bus
 - the student takes the train or the bus
 - the student takes the train or the bus but not both.

18. **WE8b** If $\xi = \{\text{children}\}$, $S = \{\text{swimmers}\}$ and $R = \{\text{runners}\}$, describe in words each of the following.

- S'
- $S \cap R$
- $R' \cap S'$
- $R \cup S$


19. A group of 12 students was asked whether they liked hip hop (H) and whether they liked classical music (C). The results are shown in the table.

	C	H
Ali	✓	✓
Anu		
Chris		✓
George		✓
Imogen		✓
Jen	✓	✓
Luke	✓	✓
Pam	✓	
Petra		
Roger	✓	
Seedevi		✓
Tomas		

- Show the results on:
 - a Venn diagram
 - a two-way table.
- If one student is selected at random, calculate:
 - $\Pr(H)$
 - $\Pr(H \cup C)$
 - $\Pr(H \cap C)$
 - $\Pr(\text{student likes classical or hip hop but not both})$

Reasoning

20. One hundred Year 9 Maths students were asked to indicate their favourite topic in mathematics. Sixty chose Probability, 50 chose Measurement and 43 chose Algebra. Some students chose two topics: 15 chose Probability and Algebra, 18 chose Measurement and Algebra, and 25 chose Probability and Measurement. Five students chose all three topics.




- Copy and complete the Venn diagram shown.
- Calculate how many students chose Probability only.
- Calculate how many students chose Algebra only.
- Calculate how many students chose Measurement only.
- Calculate how many students chose any two of the three topics.

A student is selected at random from this group. Calculate the probability that this student has chosen:

- Probability
 - Algebra
 - Algebra and Measurement
 - Algebra and Measurement but not Probability
 - all of the topics.
21. Create a Venn diagram using two circles to accurately describe the relationships between the following quadrilaterals: rectangle, square and rhombus.

22. Use the Venn diagram shown to write the numbers of the correct regions for each of the following problems.

- $A' \cup (B' \cap C)$
- $A \cap (B \cap C')$
- $A' \cap (B' \cap C')$
- $A \cup (B \cap C)'$



23. A recent survey taken at a cinema asked 90 teenagers what they thought about three different movies. In total, 47 liked 'Hairy Potter', 25 liked 'Stuporman' and 52 liked 'There's Something About Fred'.

16 liked 'Hairy Potter' only.

4 liked 'Stuporman' only.

27 liked 'There's Something About Fred' only.

There were 11 who liked all three films and 10 who liked none of them.

- Construct a Venn diagram showing the results of the survey.
- Calculate the probability that a teenager chosen at random liked 'Hairy Potter' and 'Stuporman' but not 'There's Something About Fred'.

Problem solving

24. 120 children attended a school holiday program during September. They were asked to select their favourite board game from Cluedo, Monopoly and Scrabble. They all selected at least one game, and four children chose all three games.

In total, 70 chose Monopoly and 55 chose Scrabble.

Some children selected exactly two games — 12 chose Cluedo and Scrabble, 15 chose Monopoly and Scrabble, and 20 chose Cluedo and Monopoly.

- Draw a Venn diagram to represent the children's selections.
- Calculate the probability that a child selected at random did not choose Cluedo as a favourite game.

25. Valleyview High School offers three sports at Year 9: baseball, volleyball and soccer. There are 65 students in Year 9.
 2 have been given permission not to play sport due to injuries and medical conditions.
 30 students play soccer.
 9 students play both soccer and volleyball but not baseball.
 9 students play both baseball and soccer (including those who do and don't play volleyball).
 4 students play all three sports.
 12 students play both baseball and volleyball (including those who do and don't play soccer).
 The total number of players who play baseball is one more than the total of students who play volleyball.
- Determine the number of students who play volleyball.
 - If a student was selected at random, calculate the probability that this student plays soccer and baseball only.
26. A Venn diagram consists of overlapping ovals which are used to show the relationships between sets. Consider the numbers 156 and 520. Show how a Venn diagram could be used to determine their:
- HCF
 - LCM.
27. A group of 200 shoppers was asked which type of fruit they had bought in the last week. The results are shown in the table.

Fruit	Number of shoppers
Apples (A) only	45
Bananas (B) only	34
Cherries (C) only	12
A and B	32
A and C	15
B and C	26
A and B and C	11

- Display this information in a Venn diagram.
- Calculate $n(A \cap B' \cap C)$.
- Determine how many shoppers purchased apples and bananas but not cherries.
- Calculate the relative frequency of shoppers who purchased:
 - apples
 - bananas or cherries.
- Estimate the probability that a shopper purchased cherries only.



LESSON

10.5 Compound events

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- create the sample space for compound events using arrays and tree diagrams
- solve problems using tree diagrams for experiments with and without replacement.

10.5.1 The sample space of compound events

eles-4888

- A two-step experiment involves two actions. It may be the same action repeated (tossing a coin twice) or two separate actions (tossing a coin and rolling a dice).
- Imagine two bags that contain coloured counters. The first bag has a mixture of black and white counters, and the second bag holds blue, magenta (pink) and purple counters. In a probability experiment, one counter is to be selected at random from each bag and its colour noted.




- The sample space for this experiment can be found using a table called an **array** that systematically displays all the outcomes.

		Second action			
		Bag 2			
		B	M	P	
First action	Bag 1	B	B B	B M	B P
		W	W B	W M	W P

The sample space, $\xi = \{BB, BM, BP, WB, WM, WP\}$.

- The sample space can also be found using a **tree diagram**.



WORKED EXAMPLE 17 Using an array to create a sample space

Two dice are rolled and the numbers uppermost are noted.
List the sample space in an array.

- Determine how many outcomes there are.
- Calculate how many outcomes contain at least one 5.
- Calculate $\text{Pr}(\text{at least one } 5)$.



THINK

Draw an array (table) showing all the possible outcomes.

WRITE/DRAW

		Second die					
		1	2	3	4	5	6
First die	1	1, 1	1, 2	1, 3	1, 4	1, 5	1, 6
	2	2, 1	2, 2	2, 3	2, 4	2, 5	2, 6
	3	3, 1	3, 2	3, 3	3, 4	3, 5	3, 6
	4	4, 1	4, 2	4, 3	4, 4	4, 5	4, 6
	5	5, 1	5, 2	5, 3	5, 4	5, 5	5, 6
	6	6, 1	6, 2	6, 3	6, 4	6, 5	6, 6

- The table shows 36 outcomes.
 - Count the outcomes that contain 5.
The cells are shaded in the table.
 - There are 11 favourable outcomes and 36 in total.
- There are 36 outcomes.
 - Eleven outcomes include 5.
 - $\text{Pr}(\text{at least one } 5) = \frac{11}{36}$

WORKED EXAMPLE 18 Using a tree diagram for two step experiments

Two coins are tossed and the outcomes are noted.
Show the sample space on a tree diagram.


- Determine how many outcomes there are.
- Calculate the probability of tossing at least one Head.




THINK

1. Draw a tree representing the outcomes for the toss of the first coin.


WRITE/DRAW



2. For the second coin the tree looks like this:



Add this tree to both ends of the first tree.



3. List the outcomes in the sample space column.

4. Count the outcomes in the sample space.

There are four outcomes (HH, HT, TH, TT).

b. Three outcomes have at least one Head.

b. $\Pr(\text{at least one Head}) = \frac{3}{4}$

WORKED EXAMPLE 19 Using a tree diagram for three step experiments


Three coins are tossed and the outcomes are noted.

- Show the outcomes on a tree diagram.
- Calculate the probability of getting three tails.
- Calculate the probability of getting at least one head.


THINK

Draw a tree representing the outcomes for the toss of the first coin. Write the probability on the branch.


WRITE/DRAW



1. For the second trial and third trials the tree is the same. Add this tree to both ends of the first tree, and all ends of the second tree.



2. List all the outcomes.



b. Count the outcomes. There is one outcome that has three tails out of the total 8 outcomes.

b. $\Pr(TTT) = \frac{1}{8}$

c. Count the outcomes. There are seven outcomes that have at least one head.

c. $\Pr(\text{at least one Head}) = \frac{7}{8}$

10.5.2 Two step probabilities

eles-4889

- When a coin is tossed, $\Pr(H) = \frac{1}{2}$, and when a die is rolled, $\Pr(3) = \frac{1}{6}$.



- If a coin is tossed and a die is rolled, what is the probability of getting a Head *and* a 3?
- Consider the sample space.

	1	2	3	4	5	6
H	H, 1	H, 2	H, 3	H, 4	H, 5	H, 6
T	T, 1	T, 2	T, 3	T, 4	T, 5	T, 6

There are 12 outcomes, and $\Pr(\text{Head and } 3) = \frac{1}{12}$.

- In this case, $\Pr(\text{Head and } 3) = \Pr(H) \times \Pr(3)$; that is, $\frac{1}{12} = \frac{1}{2} \times \frac{1}{6}$.

Two-step probabilities

If A is the outcome of the first action and B is the outcome of the second action in a two-step experiment, then:

- $A \cap B$ is the outcome of A followed by B
- $\Pr(A \cap B) = \Pr(A) \times \Pr(B)$.

Note: This formula is only applied when the two events are independent (if the occurrence of one event does not affect the chances of the occurrence of the other event).

WORKED EXAMPLE 20 Calculating two-step probabilities

In one cupboard Joe has two black T-shirts and one yellow one. In his drawer there are three pairs of white socks and one black pair. If he selects his clothes at random, calculate the probability that his socks and T-shirt will be the same colour.

THINK

If they are the same colour then they must be black.

$$\Pr(\text{black T-shirt}) = \Pr(B_t) = \frac{2}{3}$$

$$\Pr(\text{black socks}) = \Pr(B_s) = \frac{1}{4}$$

WRITE

$$\begin{aligned}\Pr(B_t \cap B_s) &= \Pr(B_t) \times \Pr(B_s) \\ &= \frac{2}{3} \times \frac{1}{4} \\ &= \frac{1}{6}\end{aligned}$$



eles-4890


10.5.3 Experiments with replacement

- If an experiment requires an object to be selected, say from a bag, the person doing the selecting has two options after the first selection.
 - Place the object bag back in the bag, meaning the experiment is being carried out **with replacement**. In this case, the number of objects in the bag remains constant.
 - Permanently remove the object from the bag, meaning the experiment is being carried out **without replacement**. In this situation, the number of the objects in the bag is reduced by 1 after every selection.

WORKED EXAMPLE 21 Calculating two-step experiments with replacement

A bag contains three pink and two blue counters. A counter is taken at random from the bag, its colour is noted, then it is returned to the bag and a second counter is chosen.

- Show the outcomes on a tree diagram.
- Calculate the probability of each outcome.
- Calculate the sum of the probabilities.




THINK


1. Draw a tree for the first trial. Write the probability on the branch.

Note: The probabilities should sum to 1.

WRITE/DRAW



2. For the second trial the tree is the same. Add this tree to both ends of the first tree, then list the outcomes.



- b. For both draws $\Pr(P) = \frac{3}{5}$ and $\Pr(B) = \frac{2}{5}$.

Use the rule $\Pr(A \cap B) = \Pr(A) \times \Pr(B)$ to determine the probabilities.

b. $\Pr(P \cap P) = \Pr(P) \times \Pr(P)$

$$= \frac{3}{5} \times \frac{3}{5}$$

$$= \frac{9}{25}$$

$$\Pr(P \cap B) = \Pr(P) \times \Pr(B)$$

$$= \frac{3}{5} \times \frac{2}{5}$$

$$= \frac{6}{25}$$

$$\Pr(B \cap P) = \Pr(B) \times \Pr(P)$$

$$= \frac{2}{5} \times \frac{3}{5}$$

$$= \frac{6}{25}$$

$$\Pr(B \cap B) = \Pr(B) \times \Pr(B)$$

$$= \frac{2}{5} \times \frac{2}{5}$$

$$= \frac{4}{25}$$

- c. Add the probabilities.


c. $\frac{9}{25} + \frac{6}{25} + \frac{6}{25} + \frac{4}{25} = 1$

10.5.4 Experiments without replacement

eles-4891


- Let us consider again the situation described in Worked example 21, and consider what happens if the first counter is not replaced.
- Initially the bag contains three pink and two blue counters, and either a pink counter or a blue counter will be chosen.
- In Worked example 21, $\Pr(P) = \frac{3}{5}$ and $\Pr(B) = \frac{2}{5}$ for both trials.

This would not be true if a counter is selected *but not replaced*.




- If the counter is not replaced, then the sample space is affected as follows:

If the first counter randomly selected is **pink**, then the sample space for the second draw looks like this:



$$\text{So } \Pr(P) = \frac{2}{4} \text{ and } \Pr(B) = \frac{2}{4}.$$


If the first counter randomly selected is **blue**, then the sample space for the second draw looks like this:



$$\text{So } \Pr(P) = \frac{3}{4} \text{ and } \Pr(B) = \frac{1}{4}.$$

WORKED EXAMPLE 22 Calculating two-step experiments without replacement

A bag contains three pink and two blue counters. A counter is taken at random from the bag and its colour is noted, then a second counter is drawn, without replacing the first one.




- Show the outcomes on a tree diagram.
- Calculate the probability of each outcome.
- Calculate the sum of the probabilities.

THINK

- Draw a tree diagram, listing the probabilities.

WRITE/DRAW



- Use the rule $\Pr(A \cap B) = \Pr(A) \times \Pr(B)$ to determine the probabilities.

b. $\Pr(P \cap P) = \Pr(P) \times \Pr(P)$

$$= \frac{3}{5} \times \frac{2}{4}$$

$$= \frac{6}{20}$$

$$= \frac{3}{10}$$

$$\Pr(P \cap B) = \Pr(P) \times \Pr(B)$$

$$= \frac{3}{5} \times \frac{2}{4}$$

$$= \frac{6}{20}$$

$$= \frac{3}{10}$$

$$\Pr(B \cap P) = \Pr(B) \times \Pr(P)$$

$$= \frac{2}{5} \times \frac{3}{4}$$

$$= \frac{6}{20}$$

$$= \frac{3}{10}$$

$$\Pr(B \cap B) = \Pr(B) \times \Pr(B)$$

$$= \frac{2}{5} \times \frac{1}{4}$$

$$= \frac{2}{20}$$

$$= \frac{1}{10}$$

c. Add the probabilities.

$$\text{c. } \frac{3}{10} + \frac{3}{10} + \frac{3}{10} + \frac{1}{10} = 1$$

WORKED EXAMPLE 23 Language of two-step experiments

Consider the situation presented in Worked example 22. Use the tree diagram to calculate the following probabilities:

- Pr(a pink counter and a blue counter)
- Pr(a pink counter then a blue counter)
- Pr(a matching pair)
- Pr(different colours)

THINK

- Think of the outcomes that have a pink and a blue counter. There are two: **PB** and **BP**. Add their probabilities together to find the answer.
- In this case, we have to take the order the counters are selected into account. The only outcome that has a pink then a blue is **PB**.
- A matching pair implies both counters are the same colour. These outcomes are **BB** and **PP**.

WRITE

- Pr(a pink counter and a blue counter)
= $\Pr(\text{PB}) + \Pr(\text{BP})$

$$= \frac{3}{10} + \frac{3}{10}$$

$$= \frac{3}{5}$$

- Pr(a pink counter then a blue counter)
= $\Pr(\text{PB})$

$$= \frac{3}{10}$$

- Pr(a matching pair)
= $\Pr(\text{BB}) + \Pr(\text{PP})$

$$= \frac{1}{10} + \frac{3}{10}$$

$$= \frac{2}{5}$$




d. One way to approach this is to think that if the colours are different, this is all outcomes not counted in part c. Thus, we can find the answer by subtracting the answer for part c from 1.

$$\begin{aligned} \text{d. Pr(different colours)} &= 1 - \text{Pr(a matching pair)} \\ &= 1 - \frac{2}{5} \\ &= \frac{3}{5} \end{aligned}$$

COLLABORATIVE TASK

Calculate the probability of drawing an ace, then a king, then a queen of the same suit both with replacement and without replacement.

Resources

-  **eWorkbook** Topic 10 Workbook (worksheets, code puzzle and project) (ewbk-2010)
-  **Video eLesson** Tree diagrams (eles-1894)
-  **Interactivities** Individual pathway interactivity: Two-step experiments (int-4537)
Two-step experiments (int-6083)

Exercise 10.5 Compound events

learn **on**

10.5 Quick quiz **on**

10.5 Exercise

Individual pathways

■ PRACTISE

1, 2, 4, 5, 9, 10, 13, 16, 17, 20, 23, 24

■ CONSOLIDATE

3, 7, 8, 11, 14, 18, 21, 25, 26

■ MASTER

6, 12, 15, 19, 22, 27, 28, 29

Fluency

1. **WE17** If two dice are rolled and their *sum* is noted, complete the array below to show the sample space.

		Die 1					
		1	2	3	4	5	6
Die 2	1	2					
	2					7	
	3						
	4						
	5						
	6			9			

- a. Calculate $\text{Pr}(\text{rolling a total of } 5)$.
- b. Calculate $\text{Pr}(\text{rolling a total of } 1)$.
- c. Determine the most probable outcome.

2. In her cupboard Rosa has three scarves (red, blue and pink) and two beanies (brown and purple). If she randomly chooses one scarf and one beanie, show the sample space in an array.



3. A ten-sided die is rolled and then a coin is flipped.

- a. Use an array to determine all the outcomes.
- b. Calculate:
 - i. $\text{Pr}(\text{even and a Head})$
 - ii. $\text{Pr}(\text{even or a Head})$
 - iii. $\text{Pr}(\text{a Tail and a number greater than 7})$
 - iv. $\text{Pr}(\text{a Tail or a number greater than 7})$


4. One box contains red and blue pencils, and a second box contains red, blue and green pencils. If one pencil is chosen at random from each box and the colours are noted, draw a tree diagram to show the sample space.



5. **WE18** A bag contains three discs labelled 1, 3 and 5, and another bag contains two discs, labelled 2 and 4, as shown below. A disc is taken from each bag and the larger number is recorded.



a. Complete the tree diagram below to list the sample space.



b. Calculate:

- i. $\text{Pr}(2)$
- ii. $\text{Pr}(1)$
- iii. $\text{Pr}(\text{odd number})$

6. Two dice are rolled and the difference between the two numbers is found.

- a. Use an array to find all the outcomes.
- b. Calculate:
 - i. $\text{Pr}(\text{odd number})$
 - ii. $\text{Pr}(0)$
 - iii. $\text{Pr}(\text{a number more than 2})$
 - iv. $\text{Pr}(\text{a number no more than 2})$

7. **WE19** Two coins and one die are tossed.
- List all possible outcomes.
 - Calculate the probability of getting a head, then a tail, then rolling a 2.
 - Calculate the probability of getting no heads.
 - Calculate the probability of getting no heads and an even number.
8. **WE20** A die is rolled twice. Calculate the probability of rolling:
- a 6 on the first roll
 - a double 6
 - an even number on both dice
 - a total of 12.
9. A coin is tossed twice.
- Show the outcomes on a tree diagram.
 - Calculate:
 - $\Pr(2 \text{ Tails})$
 - $\Pr(\text{at least } 1 \text{ Tail})$
10. **WE21** A bag contains three red counters and one blue counter. A counter is chosen at random. A second counter is drawn with replacement.
- Show the outcomes and probabilities on a tree diagram.
 - Calculate the probability of choosing:
 - a red counter then a blue counter
 - two blue counters.
11. A bag contains five red, six green and four blue counters. A counter is chosen at random. A second counter is drawn with replacement. Use a tree diagram to calculate the probability of choosing:
- a red and a green counter
 - a red counter then a green counter
 - a red or green counter
 - a matching pair of the same colour.
12. A bag contains eight black, nine white and three red counters. A counter is chosen at random. A second counter is drawn with replacement. Use a tree diagram to calculate the probability of choosing:
- at least one black counter
 - a black counter and a white counter
 - a black counter or a white counter
 - a matching pair of the same colour
 - different colours.
13. **WE22** A bag contains three black balls and two red balls. If two balls are selected, randomly, without replacement:
- show the outcomes and their probabilities on a tree diagram
 - calculate $\Pr(2 \text{ red balls})$.
14. **WE23** A bag contains five red, three blue and two green balls. If two balls are randomly selected without replacement, calculate:
- $\Pr(\text{two red balls})$
 - $\Pr(\text{a red ball then a green ball})$
 - $\Pr(\text{a red ball and a green ball})$
 - $\Pr(\text{the same colour selected twice})$
15. A bag contains five black, one white and six red balls. If two balls are randomly selected without replacement, calculate:
- $\Pr(\text{two red balls})$
 - $\Pr(\text{a red ball then a white ball})$
 - $\Pr(\text{a red ball and a black ball})$
 - $\Pr(\text{the same colour selected twice})$

Understanding

16. The kings and queens from a deck of cards are shuffled, then two cards are chosen. Calculate the probability that two kings are chosen:
- if the first card is replaced
 - if the first card is not replaced.
17. A coin is flipped and two dice are tossed and their values summed. Calculate the probability that:
- there is a Head and two 1s
 - the dice sum to 12
 - there is a Tail.
18. The 12 royal cards from a deck of cards are shuffled together, then two cards are chosen. Determine the probability that:
- two queens are chosen if the first card is replaced
 - a matching pair (same value, not suit) is chosen if the first card is not replaced
 - two cards from different suits are chosen if the first card is not replaced.
19. Each week John and Paul play two sets of tennis against each other. They each have an equal chance of winning the first set.



When John wins the first set, his probability of winning the second set rises to 0.6, but if he loses the first set, he has only a 0.3 chance of winning the second set.

- Show the possible outcomes on a tree diagram.
- Calculate:
 - $\Pr(\text{John wins both sets})$
 - $\Pr(\text{Paul wins both sets})$
 - $\Pr(\text{they win one set each})$

Reasoning

20. A bag contains four red and six yellow balls. If the first ball drawn is yellow, explain the difference in the probability of drawing the second ball if the first ball was replaced compared to not being replaced.
21. James has six different ties, five different shirts and three different suits that he can choose from when getting ready for work.
- Determine how many days he can go without repeating an item of clothing.
 - Determine how many possible combinations of clothing there are.
 - James receives a new shirt and tie for his birthday. Determine how many more combinations of clothing are now possible.
22. Three dice are tossed and the total is recorded.
- State the smallest and largest possible totals.
 - Calculate the probabilities for all possible totals.

Problem solving

23. You draw two cards, one after the other without replacement, from a deck of 52 cards. Calculate the probability of:
- drawing two aces
 - drawing two face cards (J, Q, K)
 - getting a 'pair' (22, 33, 44 ... QQ, KK, AA).
24. A chance experiment involves flipping a coin and rolling two dice. Determine the probability of obtaining Tails and two numbers whose sum is greater than 4.
25. In a jar there are 10 red balls and 6 green balls. Jacob takes out two balls, one at a time, without replacing them. Calculate the probability that both balls are the same colour.
26. A coin is being tossed repeatedly. Determine how many possible outcomes there will be if it is tossed:
- 3 times
 - 5 times
 - 4 times
 - x times.
27. A box of chocolates contains milk chocolates and dark chocolates.



- The probability of selecting a dark chocolate from a full box is $\frac{1}{6}$. Once a dark chocolate has been taken from the box, the chance of selecting a second dark chocolate drops to $\frac{1}{7}$.
- Calculate how many chocolates are in the box altogether.
 - If two chocolates were randomly selected from the box, calculate the probability of getting two of the same type of chocolate.
28. Claire's maths teacher decided to surprise the class with a four-question multiple-choice quiz. If each question has four possible options, calculate the probability the Claire passes the test given she guesses every question.
29. In the game of 'Texas Hold'Em' poker, five cards are progressively placed face up in the centre of the table for all players to use.
At one point in the game there are three face-up cards (two hearts and one diamond). You have two diamonds in your hand for a total of three diamonds. Five diamonds make a flush.
Given that there are 47 cards left, determine the probability that the next two face-up cards are both diamonds.

LESSON

10.6 Review

10.6.1 Topic summary

PROBABILITY

Probability of an outcome

- The probability of any outcome always falls between 0 and 1.
- An outcome that is **certain** has a probability of 1.
- An outcome that is **impossible** has a probability of 0.
- An outcome that is **likely** has a probability between 0.5 and 1.
- An outcome that is **unlikely** has a probability between 0 and 0.5.

Key terms

- Trial:** A single performance of an experiment to produce a result, such as rolling a die.
- Sample space:** The set of all possible outcomes, represented by the symbol ξ . When rolling a 6-sided die, $\xi = \{1, 2, 3, 4, 5, 6\}$.
- Event:** A set of favourable outcomes. For example, A can represent the event of rolling an even number on a 6-sided die: $A = \{2, 4, 6\}$.
- Complement:** All outcomes that are not part of an event. The complement of event A above is denoted by A' . $A' = \{1, 3, 5\}$.

Experimental probability

- Experimental probability** is used when it is difficult or impossible to determine the theoretical probability.
- Experimental probability** = Relative frequency
- Relative frequency** = $\frac{\text{frequency of outcome}}{\text{total number of trials}}$
- The more trials conducted, the more accurate the experimental probability will be.
- Expected number** = probability of outcome \times number of trials

Venn diagrams and two-way tables

- These are two different methods that can be used to represent a sample space and visualise the interaction of different events.
- For example, in a class of 20 students, 5 study Art (event A), 9 study Biology (event B) and 2 study both subjects.
 - Venn diagram:
 - Two-way table:

	B	B'	Total
A	2	3	5
A'	7	8	15
Total	9	11	20
- Using these we can see:
 - $n(A \cap B) = 2$
 - $\Pr(A \cap B') = \frac{3}{20}$

Two-step experiments

- Two-step experiments involve two separate actions performed one after the other.
- The sample space of a two-step experiment can be represented with an array or a tree diagram.
- An experiment can be conducted with replacement or without replacement.
- If an experiment is performed without replacement, the probabilities will change for the second action.

		Bag 2		
		B	M	P
Bag 1	B	BB	BM	BP
	W	WB	WM	WP

First selection	Second selection	Sample space
<div style="display: flex; align-items: center;"> <div style="margin-right: 5px;">B</div> <div style="display: flex; flex-direction: column; gap: 5px;"> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">B</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">M</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">P</div> </div> </div> </div>	<div style="display: flex; flex-direction: column; gap: 5px;"> <div style="font-size: 0.8em;">BB</div> <div style="font-size: 0.8em;">BM</div> <div style="font-size: 0.8em;">BP</div> </div>	
<div style="display: flex; align-items: center;"> <div style="margin-right: 5px;">W</div> <div style="display: flex; flex-direction: column; gap: 5px;"> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">B</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">M</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></div> <div style="font-size: 0.8em;">P</div> </div> </div> </div>	<div style="display: flex; flex-direction: column; gap: 5px;"> <div style="font-size: 0.8em;">WB</div> <div style="font-size: 0.8em;">WM</div> <div style="font-size: 0.8em;">WP</div> </div>	

Sections of a Venn diagram

- A Venn diagram can be split into four distinct sections, as shown.



Intersection and union

- The intersection of two events A and B is written $A \cap B$. These are the outcomes that are both in A and in B .
- The union of two events A and B is written $A \cup B$. These are the outcomes that are either in A or in B .

10.6.2 Success criteria

Tick a column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
10.2	I can determine whether an event is likely or unlikely.			
	I am familiar with key terms such as <i>trials</i> , <i>outcomes</i> , <i>events</i> and <i>sample space</i> .			
	I can calculate theoretical probability of an event.			
10.3	I can calculate the relative frequency and experimental probability of an outcome.			
	I can explain the difference between theoretical and experimental probability.			
10.4	I can identify the complement of an event A .			
	I can create and interpret a Venn diagram.			
	I can create and interpret a two-way table.			
	I can use a Venn diagram or two-way table to find $A \cap B$.			
	I can use a Venn diagram or two-way table to find $A \cup B$.			
10.5	I can display the sample space for a two-step experiment using an array and a tree diagram.			
	I can solve two-step probabilities for experiments with replacement.			
	I can solve two-step probabilities for experiments without replacement.			

10.6.3 Project


Sand-rings

A class of students and their teacher spent a day at the beach as part of a school excursion.




Part of the day was devoted to activities involving puzzles in the sand. One of the popular — and most challenging — puzzles was ‘sand-rings’. Sand-rings involves drawing three rings in the sand, as shown in the diagram.

The first sand-rings puzzle requires eight shells to be arranged inside the circles, so that four shells appear inside circle *A*, five shells appear inside circle *B* and six shells appear inside circle *C*. The overlapping of the circles shows that the shells can be counted in two or three circles. One possible arrangement is shown below. Use this diagram to answer questions 1 to 4.



1. How many shells appear inside circle *A*, but not circle *B*?
2. How many shells appear in circles *B* and *C*, but not circle *A*?



A shell is selected at random from the sand.



3. Calculate the probability it came from circle *A*.
4. Calculate the probability it was not in circle *C*.
5. The class was challenged to find the rest of the arrangements of the eight shells. (*Remember*: Four shells need to appear in circle *A*, five in circle *B* and six in circle *C*.)

After completing the first puzzle, the students are given new rules. The number of shells to be arranged in the circles is reduced from eight to six. However, the number of shells to be in each circle remains the same; that is, four shells in circle *A*, five shells in circle *B* and six shells in circle *C*.

6. Using six shells, calculate how many ways the shells can be arranged so that there are four, five and six shells in the three circles.
7. Explain the system or method you used to determine your answer to question 6 above. Draw diagrams to show the different arrangements.
8. Using seven shells, calculate how many ways the shells can be arranged so that there are four, five and six shells in the three circles.
9. Again, explain the system or method you used to determine your answer to question 8. Draw diagrams to show the different arrangements.
10. Determine the minimum number of shells required to play sand-rings, so that there are four, five and six shells in the three circles.
11. Modify the rules of this game so that different totals are required for the three circles. Challenge your classmates to find all possible solutions to your modified game.



on Resources

-  **eWorkbook** Topic 10 Workbook (worksheets, code puzzle and project) (ewbk-2010)
-  **Interactivities** Crossword (int-2712)
Sudoku puzzle (int-3212)

Exercise 10.6 Review questions

learn on

Fluency

1. **MC** In a trial, it was found that a drug cures $\frac{2}{5}$ of those treated by it. If 700 sufferers are treated with the drug, calculate how many of them are not expected to be cured.
A. 280 **B.** 420 **C.** 140 **D.** 350 **E.** 100
2. **MC** If a die is rolled and a coin tossed, determine the probability of a 6–Heads result.
A. $\frac{1}{6}$ **B.** $\frac{1}{2}$ **C.** $\frac{1}{8}$ **D.** $\frac{1}{12}$ **E.** $\frac{1}{3}$

3. **MC** Twelve nuts are taken from a jar containing macadamias and cashews. If three macadamias are obtained, the experimental probability of obtaining a cashew is:
- A. $\frac{1}{12}$ B. $\frac{1}{4}$ C. $\frac{1}{3}$ D. $\frac{3}{4}$ E. $\frac{11}{12}$
4. **MC** From a normal pack of 52 playing cards, one card is randomly drawn and replaced. If this is done 208 times, the number of red or picture cards expected to turn up is:
- A. 150 B. 130 C. 128 D. 120 E. 144
5. **MC** A cubic die with faces numbered 2, 3, 4, 5, 6 and 6 is rolled. The probability of rolling an even number is:
- A. $\frac{1}{3}$ B. $\frac{2}{3}$ C. $\frac{1}{6}$ D. $\frac{1}{2}$ E. 1
6. **MC** The probability of rolling an odd number or a multiple of 2 using the die in question 5 is:
- A. 1 B. $\frac{1}{3}$ C. $\frac{1}{4}$ D. $\frac{3}{4}$ E. $\frac{4}{3}$

The following information should be used to answer questions 7 and 8.

Students in a Year 9 class chose the following activities for a recreation day.

Activity	Tennis	Fishing	Golf	Bushwalking
Number of students	8	15	5	7

7. **MC** If a student is selected at random from the class, the probability that the student chose fishing is:
- A. $\frac{1}{7}$ B. $\frac{2}{7}$ C. $\frac{3}{7}$ D. $\frac{4}{7}$ E. $\frac{5}{7}$
8. **MC** If a student is selected at random, the probability that the student did not choose bushwalking is:
- A. $\frac{1}{35}$ B. $\frac{2}{5}$ C. $\frac{3}{5}$ D. $\frac{4}{5}$ E. $\frac{4}{35}$
9. The mass of 40 students in a Year 9 Maths class was recorded in a table.

Mass (kg)	Less than 50	50– < 55	55– < 60	60– < 65	65 and over
Number of students	4	6	10	15	5

Calculate the experimental probability of selecting a student who has:


- a. a mass of 55 kg or more, but less than 60 kg
 b. a mass less than 50 kg
 c. a mass of 65 kg or greater.
10. Calculate the following expected values.
- a. The number of Heads in 80 tosses of a coin.
 b. The number of sixes in 200 rolls of a die.
 c. The number of hearts if a card is picked from a reshuffled pack and replaced 100 times.
11. A normal six-sided die is rolled. Calculate the probability of getting an odd number or a multiple of 4.
12. A card is drawn from a pack of 52 cards. Calculate the probability that the card is a heart or a club.

13. Insert each of the letters **a** to **d** to represent the following events at appropriate places on the probability scale shown.




- You will go to school on Christmas Day.
 - All Year 9 students can go to university without doing Year 10.
 - Year 9 students will study Maths.
 - An Australian TV channel will telecast the news at 6:00 pm.
14. Indicate the set that each of the shaded regions represents.


a. Subject preference



b. Leisure activity



c. Favourite drinks




15. An equal number of red (R), black (B) and yellow (Y) counters makes up a total of 30 in a bag.
- List the sample space for picking a counter from the bag.
 - Event A is 'draw a yellow counter, then randomly draw another counter from the bag'. List the sample space of event A .
 - Explain whether 'choosing a green counter' is an outcome.

Problem solving

16. Teachers at a school opted for the choice of morning recess refreshments shown in the Venn diagram.

- State how many teachers are in the set 'cake \cap coffee'.
- State the total number of teachers surveyed.
- If a teacher is selected at random, calculate the probability that the teacher:
 - chose tea
 - chose coffee only
 - chose milk
 - did not choose tea, coffee, cake or milk
 - did not choose coffee.



17. Thirty-two students ordered fried rice, chicken wings and dim sims for lunch. Four students ordered all three, two ordered fried rice and chicken wings only, three ordered fried rice and dim sims only, and three ordered chicken wings and dim sims only. When the waiter organised the orders, he found that 16 students ordered fried rice and 18 students ordered dim sims.
- Show this information on a Venn diagram.
 - Calculate how many students ordered chicken wings only.
 - If a student is picked at random, calculate the probability that the student has:
 - ordered chicken wings and dim sims
 - ordered fried rice
 - not ordered dim sims.

18. The following are options for dorm rooms at a university. You are required to choose one option from each of the four categories.

- Upstairs or downstairs rooms
- Single or double rooms
- Male or female roommates
- A choice of 10 different locations on campus

Calculate how many different combinations of rooms there are to choose from.

19. A future king is the oldest male child. The future king of Mainland has two siblings. Determine the probability that he has an older sister.

20. A witness described a getaway car as having a NSW registration plate starting with TLK. The witness could not remember the three digits that followed, but recalled that all three digits were different. Calculate how many cars in NSW could have a registration plate with these letters and numbers.


21. There are 12 people trying out for a badminton team. Five of them are girls. Calculate the probability that a team chosen at random to play is a mixed doubles team.

22. An ace is chosen from a deck of standard cards and not replaced. A king is then chosen from the deck. Calculate the probability of choosing an ace and a king in this order.

23. If you randomly select one number from 1 to 500 (inclusive), calculate the probability that the selected number will have at least one 4 in the digits.

24. If you flip a coin eight times, calculate how many of the possible outcomes would you expect to have a Head on the second toss.

25. Use the Venn diagram to calculate each of the following.



a. $A' \cup (B' \cap C')$

c. $A \cap (B' \cup C')$

b. $A' \cap (B \cap C')$

d. $(A \cup B \cup C)' \cap (A \cup B \cup C)$



To test your understanding and knowledge of this topic go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 10 Probability

10.1 Pre-test

- C
- a. Zero b. $\frac{3}{8}$ or 0.375
- a. 100 b. $\frac{7}{25}$ c. $\frac{1}{2}$ d. $\frac{11}{50}$
- $\frac{1}{3}$
- True
- D
- C
- A
- a. $\frac{1}{4}$ b. $\frac{3}{14}$
- $\Pr(\text{drama not comedy}) = \frac{2}{5}$
- $\frac{5}{7}$
- a. B b. 0.33
- a. C b. D c. A d. B
- a. $\frac{8}{15}$ b. $\frac{67}{225}$ c. $\frac{1}{6}$ d. $\frac{67}{240}$
- D

10.2 Theoretical probability

- a. Certain b. Unlikely
- a. Likely b. Certain
- a. Certain b. Likely
- a. Likely b. Likely
- $\{1, 2, 3, 4, 5, 6\}$
 - $\{H, T\}$
 - $\{\text{defective, not defective}\}$
 - $\{\text{red, black}\}$
 - $\{\text{hearts, clubs, diamonds, spades}\}$
- $\{1, 2, 3\}$
 - $\{1, 3, 5\}$
 - $\{1, 2, 4, 6\}$
 - $\{3, 4, 5, 6\}$
 - $\{1, 2, 3, 4\}$
 - $\{5, 6\}$

- 3, 5
 - 1, 2
 - 6
 - 1, 2, 5, 6
 - 1, 2, 3, 5
 - No favourable outcomes
 - 3
- $\{\text{hearts, clubs, diamonds, spades}\}$
 - $\{\text{clubs, spades}\}$
 - $\{\text{hearts, diamonds}\}$
 - $\{\text{clubs, diamonds, spades}\}$
 - $\{\text{hearts, clubs, diamonds, spades}\}$
- 6
 - 2
 - 52
 - 4
 - 5
- $\frac{1}{13}$
 - $\frac{1}{13}$
 - $\frac{1}{52}$
 - $\frac{1}{4}$
- $\frac{1}{52}$
 - $\frac{1}{2}$
 - $\frac{1}{26}$
 - $\frac{1}{4}$
 - $\frac{1}{2}$
- $\frac{1}{52}$
 - $\frac{4}{13}$
 - $\frac{2}{13}$
 - 0
 - $\frac{12}{13}$
- $\frac{2}{11}$
 - $\frac{9}{11}$
 - $\frac{4}{11}$
 - $\frac{7}{11}$
- $\frac{1}{4}$
 - $\frac{1}{8}$
 - $\frac{5}{16}$
 - 0
 - $\frac{3}{4}$
- $\frac{2}{3}$
 - $\frac{1}{3}$
 - Sample responses can be found in the worked solutions in the online resources.
- Sample responses can be found in the worked solutions in the online resources.
- 18
 - 20
 - 36
- $\frac{7}{10}$
 - $\frac{3}{10}$
 - $\frac{2}{3}$
 - $\frac{1}{3}$
- Sample responses can be found in the worked solutions in the online resources.
- Probabilities must be between 0 and 1, so -0.5 and 1.05 can't be probabilities.
- $\frac{1}{2}$
- The coloured portions outside the arc of the spinner shown are of no consequence. The four colours within the arc of the spinner are of equal area (each $\frac{1}{4}$ circle), so there is equal chance of falling on each of the colours.
- $\frac{2}{3}$
- There are 36 outcomes, 15 under 7, 6 equal to 7 and 15 over 7. So, you would have a greater chance of winning if you chose 'under 7' or 'over 7' rather than 'equal to 7'.
- 0.09

10.3 Experimental probability

1. a. i. 16
ii. 4

iii.

Outcome	1	2	3	4	5	6
Relative frequency	0.19	0.06	0.31	0.13	0.25	0.06

iv. 1

- b. i. 270
ii. 40

iii.

Outcome	1	2	3	4	5	6
Relative frequency	0.19	0.14	0.17	0.18	0.15	0.17

iv. 1

2. a. i. r.f. (H) = 0.44, r.f. (T) = 0.56
ii. 1
b. i. r.f. (H) = 0.62, r.f. (T) = 0.38
ii. 1

3. a. 0.365 b. 0.33 c. 0.25 d. 0.875

4. a. 0.275 b. 0.64 c. 0.365 d. 0.635

5. a. $\frac{27}{95}$ b. $\frac{33}{95}$

c. $\frac{1}{19}$ d. Yes, reject this batch.

6. a. Not reasonable; not enough trials were held.
b. Not reasonable; the conditions are different under each trial.
c. Reasonable; the Sun rises every morning, regardless of the weather or season.
d. Reasonable; enough trials were performed under the same conditions.
e. Not reasonable; monthly rainfall in Sydney is not consistent throughout the year.

7. a. $\frac{5}{24}$ b. $\frac{59}{72}$

8. 200

9. 20

10. A

11. B

12. a. $\frac{11}{20}$ b. $\frac{3}{10}$ c. $\frac{17}{20}$ d. $\frac{3}{20}$

13. a. $\frac{33}{100}$ b. $\frac{40}{100} = \frac{2}{5}$ c. $\frac{3}{100}$

14. a. i. 6

ii. {356, 365, 536, 563, 635, 653}

b. $\frac{2}{3}$

c. $\frac{1}{3}$

d. $\frac{1}{3}$

15. 5

16. Svetlana 6, Sarah 4, Leonie 3, Trang 2

17. C

18. a. 27

b. $\left\{ \begin{array}{l} 111, 112, 115, 121, 122, 125, 151, 152, 155, 211, 212, \\ 215, 221, 222, 225, 251, 252, 255, 511, 512, 515, 521, \\ 522, 525, 551, 552, 555 \end{array} \right\}$

c. $\frac{1}{3}$

d. $\frac{2}{3}$

e. $\frac{1}{3}$

19. a. $\frac{1}{12}$

b. $\frac{1}{20}$

c. No, because John has a higher probability of winning.

20. a. $\frac{7}{10}$

b. Yes, far too many mistakes

21. 32

22. 1000 balls

23. Red = 10, yellow = 50, blue = 40

24. a. $\frac{7}{20}$

b. $\frac{2}{5}$

25. Sample responses can be found in the worked solutions in the online resources.

26. a. $\frac{30}{241}$

b. $\frac{91}{241}$

c. $\frac{59}{241}$

27. 16, 64, 40, 80, 128 and 32 degrees.

10.4 Venn diagrams and two-way tables

1. a. $A' = \{1, 3, 5, 7, 9\}$

b. $B' = \{1, 2, 3, 4, 6, 7, 8, 9\}$

c. $C' = \{2, 3, 5, 6, 7, 8, 10\}$

d. $D' = \{8, 9, 10\}$

2. a. $A' = \{11, 13, 14, 16, 17, 19, 20\}$

b. $B' = \{20\}$

c. $C' = \{12, 14, 15, 16, 18, 20\}$

d. $D' = \{12, 14, 16\}$

3. a. $A' = \{1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19\}$

b. $B' = \{1, 4, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20\}$

c. $C' = \{1, 3, 5, 7, 9, 11, 13, 14, 15, 16, 17, 18, 19, 20\}$

d. $D' = \{1, 3, 5, 7, 9, 11, 13\}$

4. a. 27

b. 11

c. 13


d. 6

e. 7

f. 9

5.


	T	T'
S	6	5
S'	7	9



7. a. i. W' ii. None
 b. i. $A \cap B'$ ii. $A \cap B$
 c. i. $A' \cap B'$ ii. $B \cap A'$




8. a.

	Left-handed	Right-handed
Male	15	40
Female	12	33



- c. i. $\frac{73}{100}$ ii. $\frac{2}{5}$ iii. $\frac{3}{25}$

9. a. $\frac{8}{25}$ b. $\frac{17}{25}$ c. $\frac{1}{5}$ d. $\frac{3}{25}$




13. a. $\frac{1}{2}$ b. $\frac{1}{10}$ c. $\frac{31}{40}$
 d. $\frac{27}{40}$ e. $\frac{9}{40}$

14. a. $\frac{1}{2}$ b. $\frac{6}{25}$
 c. $\frac{22}{25}$ d. $\frac{16}{25}$

15. a.


	Croissant	No croissant	Total
Coffee	27	2	29
No coffee	9	7	16
Total	36	9	45

- b. i. $\frac{3}{5}$ ii. $\frac{38}{45}$ iii. $\frac{11}{45}$
 c. 1218
 16. a. 20
 b. i. $\frac{1}{4}$ ii. $\frac{3}{20}$ iii. $\frac{1}{2}$ iv. $\frac{7}{20}$



- b. i. $\frac{29}{40}$ ii. $\frac{4}{5}$ iii. $\frac{19}{40}$


18. a. Children who are not swimmers
 b. Children who are swimmers and runners
 c. Children who neither swim nor run
 d. Children who swim or run or both



ii.

	H	H'
C	3	2
C'	4	3


- b. i. $\frac{7}{12}$ ii. $\frac{3}{4}$ iii. $\frac{1}{4}$ iv. $\frac{1}{2}$




- b. 25 c. 15 d. 12

- e. 43 f. $\frac{3}{5}$ g. $\frac{43}{100}$
 h. $\frac{9}{50}$ i. $\frac{13}{100}$ j. $\frac{1}{20}$


21. Quadrilaterals



22. a. 2, 5, 6, 7, 8 b. 4
 c. 8 d. 1, 2, 3, 4, 5, 7, 8




b. $\frac{8}{90} = \frac{4}{45}$




b. $\frac{7}{12}$

25. a. 31 students b. $\frac{1}{13}$



- a. $HCF = 2 \times 2 \times 13 = 52$
 b. $LCM = 3 \times 2 \times 2 \times 13 \times 2 \times 5 = 1560$



- b. 4
 c. 21
 d. i. $\frac{81}{200}$ ii. $\frac{97}{200}$
 e. $\frac{3}{50}$

10.5 Compound events

1. See table at the foot of the page.*
 a. $\frac{1}{9}$ b. 0 c. A total of 7

2.

		Scarves		
		R	Bl	Pi
Beanies	Br	Br, R	Br, Bl	Br, Pi
	Pu	Pu, R	Pu, Bl	Pu, Pi



3. a. See table at the foot of the page.*
 b. i. $\frac{1}{4}$ ii. $\frac{3}{4}$ iii. $\frac{3}{20}$ iv. $\frac{13}{20}$

*1.

		Die 1					
		1	2	3	4	5	6
Die 2	1	2	3	4	5	6	7
	2	3	4	5	6	7	8
	3	4	5	6	7	8	9
	4	5	6	7	8	9	10
	5	6	7	8	9	10	11
	6	7	8	9	10	11	12

*3. a.


		Dice roll									
		1	2	3	4	5	6	7	8	9	10
Coin toss	H	1, H	2, H	3, H	4, H	5, H	6, H	7, H	8, H	9, H	10, H
	T	1, T	2, T	3, T	4, T	5, T	6, T	7, T	8, T	9, T	10, T



- b. i. $\frac{1}{6}$ ii. 0 iii. $\frac{1}{2}$

6. a. See table at the foot of the page.*


- b. i. $\frac{1}{2}$ ii. $\frac{1}{6}$ iii. $\frac{1}{3}$ iv. $\frac{2}{3}$



b. $\Pr(\text{HT2}) = \frac{1}{24}$ c. $\Pr(\text{no heads}) = \frac{1}{4}$

d. $\Pr(\text{no heads}) = \frac{1}{8}$


8. a. $\frac{1}{6}$ b. $\frac{1}{36}$ c. $\frac{1}{4}$ d. $\frac{1}{36}$



*6. a.

		Die 1					
		1	2	3	4	5	6
Die 2	1	0	1	2	3	4	5
	2	1	0	1	2	3	4
	3	2	1	0	1	2	3
	4	3	2	1	0	1	2
	5	4	3	2	1	0	1
	6	5	4	3	2	1	0

10. a.



b. i. $\frac{3}{16}$

ii. $\frac{1}{16}$

11. a. $\frac{4}{15}$

b. $\frac{2}{15}$

c. $\frac{209}{225}$

d. $\frac{77}{225}$

12. a. $\frac{16}{25}$


b. $\frac{9}{25}$

c. $\frac{391}{400}$

d. $\frac{77}{200}$

e. $\frac{123}{200}$

13. a.



b. $\frac{1}{10}$

14. a. $\frac{2}{9}$

b. $\frac{1}{9}$

c. $\frac{2}{9}$

d. $\frac{14}{45}$

15. a. $\frac{5}{22}$

b. $\frac{1}{22}$

c. $\frac{5}{11}$

d. $\frac{25}{66}$

16. a. $\frac{1}{4}$

b. $\frac{3}{14}$

17. a. $\Pr(H11) = \frac{1}{72}$

b. $\Pr(\text{dice sum is 12}) = \frac{1}{36}$


c. $\Pr(\text{Tail}) = \frac{1}{2}$

18. a. $\frac{1}{9}$

b. $\frac{3}{11}$

c. $\frac{8}{11}$

19. a.



b. i. 0.3

ii. 0.35

iii. 0.35

20. If the first ball is replaced, the probability of drawing a yellow ball stays the same on the second draw, i.e. $\left(\frac{3}{5}\right)$.
If the first ball isn't replaced, the probability of drawing a yellow ball on the second draw decreases, i.e. $\left(\frac{5}{9}\right)$.

21. a. 3 days

b. 90 combinations

c. 36 new combinations

22. a. Smallest total: 3, largest total: 18

b.

Total	Probability
3	$\frac{1}{216}$
4	$\frac{3}{216}$
5	$\frac{6}{216}$
6	$\frac{10}{216}$
7	$\frac{15}{216}$
8	$\frac{21}{216}$
9	$\frac{25}{216}$
10	$\frac{27}{216}$
11	$\frac{27}{216}$
12	$\frac{25}{216}$
13	$\frac{21}{216}$
14	$\frac{15}{216}$
15	$\frac{10}{216}$
16	$\frac{6}{216}$
17	$\frac{3}{216}$
18	$\frac{1}{216}$

23. a. $\frac{1}{221}$

b. $\frac{11}{221}$

c. $\frac{1}{17}$

24. $\frac{5}{12}$

25. $\frac{1}{2}$

26. a. 8

b. 32

c. 16

d. 2^x

27. a. 36

b. $\frac{5}{7}$


28. $\frac{67}{256}$

29. $\frac{45}{1081}$


Project

1. 3
2. 4
3. $\frac{1}{2}$
4. $\frac{1}{4}$
5. 18
6. 2
7. Sample responses can be found in the worked solutions in the online resources.
8. 8
9. Sample responses can be found in the worked solutions in the online resources.
10. 6
11. Sample responses can be found in the worked solutions in the online resources.

10.6 Review questions

1. B
2. D
3. D
4. C
5. B
6. A
7. C
8. D
9. a. $\frac{1}{4}$ b. $\frac{1}{10}$ c. $\frac{1}{8}$
10. a. 40 b. 33 c. 25
11. $\frac{2}{3}$
12. $\frac{1}{2}$
13. b


14. a. Students who do not like music.
 b. People who like surfing and golf as leisure activities, but not reading.
 c. People who like Coke or Fanta or both, but not Sprite.
15. a. {R, B, Y}
 b. {YR, YY, YB}
 c. No, there is no green counter.
16. a. 12
 b. 60
 c. i. $\frac{3}{10}$ ii. $\frac{17}{60}$ iii. $\frac{1}{10}$
 iv. $\frac{1}{12}$ v. $\frac{31}{60}$



- b. 5
- c. i. $\frac{7}{32}$ ii. $\frac{1}{2}$ iii. $\frac{7}{16}$
18. 80
19. $\frac{3}{7}$
20. 720
21. $\frac{35}{66}$
22. $\frac{4}{663}$
23. $\frac{176}{500}$
24. 128
25. a. {1, 5, 6, 7, 8} b. {5}
 c. {1, 2, 4} d. ϕ

11 Statistics

Lesson sequence

11.1 Overview	626
11.2 Collecting data	630
11.3 Organising data	639
11.4 Displaying data	649
11.5 Measures of centre	664
11.6 Measures of spread	680
11.7 Review	689



LESSON

11.1 Overview

11.1.1 Why learn this?

The science of statistics is concerned with understanding the world around us. It presents a way of processing data about people, places and things in order to make that data useful and easier to understand.

One example of statistics is the data that social media companies collect about how you use their platforms. This data is statistically analysed and used to adapt these platforms to the way you use them (and to help the company maximise profits). Another example of statistics is the Australian Bureau of Meteorology — they collect data about the weather, analyse the statistics from that data and then use that knowledge to help predict what the weather will be like tomorrow, next week and even next year.

A good understanding of statistics can help you develop critical analysis skills so that you are not tricked by bad or misleading statistics. It is also important because statistics are a valuable tool used in a broad range of careers including product design, science, sports coaching and finance. Next time you are reading an article or watching a news report, keep your eye out for statistics and you'll start to notice that they are everywhere!



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

1. Match and classify the following different types of data.

Example	Data type
a. Attendance at a concert	i. nominal
b. Mass of bag of lollies	ii. ordinal
c. Eye colour	iii. discrete
d. The quality of food at a restaurant	iv. continuous

2. Thirty Year 9 students from a particular school were surveyed on the number of hours they spend each week using social media on their phones.
State whether this type of data is a population or a sample.
3. Fifty members of a tennis club were selected to participate in a survey. The club has 80 male adult, 70 female adult, 55 male adolescent and 45 female adolescent members.
State how many members are required from each of these four groups to perform a stratified sample survey.
4. The stem and leaf plot shown represents the speeds of cars recorded by a roadside camera outside a school where the speed limit is: 40 km/h.

Key: 1 | 9 = 19 km/h

Stem	Leaf
1	9
2	5 7 7 9 9
3	0 0 4 5 6 7 8 8
4	0 0 0 0 2 4 5 5
5	2 3
6	0

- a. State how many cars had their speed recorded.
b. Determine how many cars were travelling at a speed over 40 km/h.
5. The dot plot shown displays the results of a maths test for a Year 9 class.
The results show the students' marks out of 20.




- a. State how many students are in the class.
b. Determine how many students scored more than 15 out of 20 for the test.


6. Match the different types of data shown in the table with the most appropriate type of graph.

Data type	Type of graph
a. Categorical data	i. Box plot
b. Numerical continuous data	ii. Stem plot
c. Numerical discrete data	iii. Bar chart

7. **MC** Based on the pie chart shown, the percentage of students who catch public transport to school is approximately:



- A. 40% B. 43% C. 52% D. 60% E. 65%
8. The histogram shows the weights of passengers' luggage for a flight to Perth.



- a. State the type of data shown the histogram.
 b. State how many passengers exceeded the maximum limit of 20 kg.
 c. Explain whether the data distribution is skewed or symmetrical.

9. **MC** Identify the correct meaning of the spreadsheet function =RANDBETWEEN(1, 5).

- A. Randomly select a real number between 1 and 5.
 B. Randomly select an integer between 1 and 5.
 C. Randomly select a number 5 times.
 D. Randomly select 5 numbers.
 E. Randomly select a number between 2 and 4.
10. **MC** Megan wanted to select a random number in the range 10 to 20. Select the command she used to generate the number in her spreadsheet.
- A. = range (10, 20) B. = ran (10, 20)
 C. = randbetween (10, 20) D. = rand (10, 20)
 E. = random (10, 20)

11. Calculate the median for the frequency table shown.

Score	Frequency
0	3
1	8
2	10
3	14
4	5
5	2

12. Examine the grouped frequency table shown and answer the following questions.

Score	Frequency
0– < 20	3
20– < 40	4
40– < 60	8
60– < 80	15
80– < 100	7

- a. Calculate the mean for the grouped frequency table, rounded to 1 decimal place.
 b. **MC** Select the modal class for the grouped frequency table.
 A. 0– < 20 B. 20– < 40 C. 40– < 60
 D. 60– < 80 E. 80– < 100
13. Determine the missing value from the set of numbers {8, 8, 10, 12, 15} if the mean is 10, the mode is 8 and the median is 9.

14. **MC** Select the range and interquartile range for the stem plot shown, which presents the weight of school bags.
 A. Range = 36 and interquartile range = 10
 B. Range = 36 and interquartile range = 17
 C. Range = 34 and interquartile range = 9
 D. Range = 34 and interquartile range = 10
 E. Range = 34 and interquartile range = 11

Key: 1 | 3 = 13 kg

Stem	Leaf
0	2 3 4 5 8
1	0 0 1 1 2 2 2 3
1	5 5 5 6 7 7 8 8 9 9 9
2	0 0 1 1 2 2 4
2	5 5 8
3	1 3
3	6

15. **MC** Select the correct statement to describe symmetrical distributions.
 A. The data on the right have more low values.
 B. Distribution has two peaks.
 C. Mean has a smaller value than median.
 D. Mean has a larger value than median.
 E. Mean, median and mode have similar values.

LESSON

11.2 Collecting data

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify different types of data
- determine if a data set is a sample or a population
- generate random numbers.


11.2.1 Types of data

eles-4909

- Data can be **categorical** or **numerical**.
- There are two types of categorical data:
 - **nominal data**
 - **ordinal data**.
- There are two types of numerical data:
 - **discrete data**
 - **continuous data**.

Determining types of data

- The information in the flowchart shown can be used to determine the type of data being considered.



WORKED EXAMPLE 1 Classifying data as categorical or numerical

Classify each of the following type of data as categorical data or numerical data.

- Favourite sport watched on TV.
- Quantity of books carried in a school bag by 20 Year 9 students.
- Cars passing your home each hour in a day.
- Eye colour of 100 preschool children.



THINK

- Decide how the data would initially be recorded, as either numbers or categories. Examples of sports watched on TV could include football, netball, soccer or tennis.
- Decide how the data would be recorded, as either numbers or categories. The quantity of books carried in a school bag by 20 Year 9 students will be collected by counting the number of books.
- Decide how the data would be recorded, as numbers or categories. Data for cars passing your home each hour of a day would be collected by counting the number of cars.
- Decide how the data would be recorded, as numbers or categories. Eye colour of 100 preschool children could include blue, brown, green or hazel.

WRITE

- The sports watched on TV are collected into categories so this is categorical data.
- Because the recorded data would be collected by counting the number of books in a school bag, this is numerical data.
- Because the recorded data would be collected by counting the number of cars, this is numerical data.
- Because the eye colour of 100 preschool children would be collected into categories, this is categorical data.

WORKED EXAMPLE 2 Classifying types of data

Classify each of the following types of data into the correct groups, stating whether it is categorical or numerical. If it is categorical, state whether it is nominal or ordinal.

If it is numerical, state whether it is discrete or continuous.

- Type of transport used to go to work
- Daily attendance at a zoo
- Hair colour of 100 Year 9 students
- Mass of an individual Easter egg
- A student's behavioural report

THINK

- Types of transport used to go to work could include car, bicycle, tram, bus or train.

WRITE

- The possible transport modes are categories; therefore, they are categorical data.

2. For categorical data, decide whether the categories can be ranked in a logical order (categorical ordinal) or not (categorical nominal).
- b. 1. Attendance at a zoo each day would be counted and recorded as a whole number.
2. For numerical data, decide whether it is collected by counting (numerical discrete data) or by measuring (numerical continuous data).
- c. 1. Hair colour of 100 Year 9 students could include black, brown, blonde, brunette or redhead.
2. For categorical data, decide if the categories can be ranked in a logical order (categorical ordinal) or not (categorical nominal).
- d. 1. Mass of an individual Easter egg would be measured on a scale.
2. For numerical data, decide whether it is collected by counting (numerical discrete data) or by measuring (numerical continuous data).
- e. 1. A student's behavioural report could be described as excellent, good, satisfactory, improvement needed or unsatisfactory.
2. For categorical data, decide whether the categories can be ranked in a logical order (categorical ordinal) or not (categorical nominal).

Because types of transport cannot be logically ranked, the data can be classified as categorical nominal data.

- b. The recorded data would be the number of visitors at a zoo; therefore, this is numerical data.

The visitors would be counted; therefore, it is classified as numerical discrete data.

- c. The hair colours are categories; therefore, this is categorical data.

Because hair colours cannot be logically ranked, the data can be classified as categorical nominal data.

- d. Because the recorded data would be the mass of an individual egg, this is numerical data.

The mass of each egg would be measured. Therefore, it is numerical continuous data.


- e. A student's behavioural report uses categories; therefore, it is categorical data.

Because the student's behavioural report can be logically ranked, it can be classified as categorical ordinal data.

11.2.2 Collecting data


eles-4910

- Data is collected to obtain information and draw conclusions.
- Data can be collected using:
 - a census of the entire **population**, or
 - a survey of a **sample** of the population




Samples and populations

- Data is classified as a *population* if data is collected from an entire group. Examples of population data include surveying all students at a school or measuring the size of all frogs in a pond.



In this example, 11 out of 11 people in the group were surveyed.

- Data is classified as a *sample* if data is collected from parts of a group. Examples of sample data include surveying only some students at a school or measuring only some of the frogs in a pond.



In this example, 5 out of 11 people in the group were surveyed.

WORKED EXAMPLE 3 Classifying the data collection methods

For the following data collection activities, define whether the data collected is a sample or a population.

- Sixty Year 7 students collected data on the amount of money they donated to a charity, then used this data to calculate the average donation from the 1523 students at their school.
- A data set of the force (measured in Newtons) needed to break wooden ice-block sticks.
- The average age of the members of your own family.

THINK

- Only 60 Year 7 students were surveyed out of a total of 1523.
- An unknown number of wooden ice-block sticks are required. It is impossible to collect data on the entire population (which would mean every wooden ice-block stick in the world).
- Every family member is surveyed to find the average age of the group.

WRITE

- This is a sample, because only 60 of the 1523 students were surveyed.
- This is a sample, because there is an unknown number of wooden ice-block sticks.
- This is a population, because every member of the group has been surveyed.

11.2.3 Sampling

eles-4911

- Sampling is the process of selecting a sample of a population to provide an estimate of the entire population.
- When a sample of data is collected, it is important to ensure that the sample is indicative of the population and is selected at random. If the sample is not large enough and is not selected at random, it may provide data that is biased towards one particular group of people.
- **Simple random sampling** is the basic method of selecting a sample. This type of sampling ensures that each individual in a population has an equal chance of being selected for the sample.

Digital technology

Digital technology often features random number generators that can be used to create simple random samples.

- In a spreadsheet, click on a cell and type `=RANDBETWEEN(<lowest number>,<highest number>)`.
- Press **ENTER**.
- A random number will appear in the cell.
- If you want a list of random numbers, use **Fill Down** to get more numbers.

fx	=RANDBETWEEN(1,10)	
	A	B
2	9	
3		

For example, in the screenshot shown, a random number in a sample of 20 is generated by:

- clicking on a cell
- typing `=RANDBETWEEN(1,20)`
- pressing **ENTER**.

1	4				
2	1				
3	19				
4	18				
5	14				
6	14				
7	11				
8	14				
9	1				
10	3				
11	19				
12	19				
13	7				
14	10				
15	3				
16	14				
17	4				
18	5				
19	9				
20	15				

WORKED EXAMPLE 4 Selecting a sample using random number generator

Use a random number generator on a spreadsheet to select a sample for the following surveys.

- Randomly select 4 days from an entire year.
- Randomly select 6 months that fall between January 2022 and December 2023

THINK

1. Use the Julian date system, which associates each day of the year with the numbers 1 to 365.
2. Click on a cell.
Type `=RANDBETWEEN(1,365)`.
Click on the bottom right corner of the cell.
Fill down by dragging your cursor down four cells.
Note: If any numbers are repeated, fill down as required to generate another number to replace the repeated number.

WRITE

a.

fx	=RANDBETWEEN(1,365)	
	A	B
1	206	
2	188	
3	214	
4	114	
5		
6		

3. Use the Julian calendar to convert each number to a calendar date.

Day 206 is 25 July
 Day 188 is 7 July
 Day 214 is 2 August
 Day 114 is 24 April

b. 1. Convert the months to numbers by creating a numbered list.

b. Label the months numerically as:

1. January 2022
2. February 2022
- ...
23. November 2023
24. December 2023.

2. Click on a cell.
 Type =RANDBETWEEN (1, 24)
 Press ENTER.
 Click on the bottom right corner of the cell.
 Fill down by dragging your cursor down six cells.
Note: If any numbers are repeated, fill down an extra cell to replace the repeated numbers.

	A	B
3	15	
4	14	
5	6	
6	21	
7	22	
8	12	
9		

3. Convert the numbers back to months to write the answer.

15 is March 2023
 14 is February 2023
 6 is June 2022
 21 is September 2023
 22 is October 2023
 12 is December 2022


TI | THINK

a. On a Calculator page, press:

- Menu
- 5. Probability
- 4. Random
- 2. Integers

Complete the entry lines as:
 randInt(1,365)
 Press Enter.
 Random numbers are generated every time the entry line is completed.

DISPLAY/WRITE




CASIO | THINK

a. A random number in a sample of 365 is generated by:

- selecting **rand**(from the Catalog tab in the Keyboard
- typing 1, 365 to indicate the lowest number and highest number
- pressing **EXE**.

DISPLAY/WRITE



COLLABORATIVE TASK: Designing a survey

1. Working in pairs, think of a topic on which you would like to gather your classmates' opinions. Some suggestions for things to ask them about include pets, favourite movies, favourite social platform or school subjects.
2. Think of five questions you would like to ask your classmates about this topic, ensuring that you collect both categorical and numerical data.
3. When coming up with your questions, you should consider what sorts of answers you want. Do you want yes and no answers, or would you like questions that require people to make a judgement on a scale? Your questions should be easy to understand and must be relevant to your topic.
4. Write down how you intend to collect the data, the type of data you expect to collect, and how you think it is best to represent this data. Think about the possible responses you may get. It is important to consider the methods used to collect data, because they could possibly bias the results of the analysis, which would lead to incorrect conclusions.
5. As a class, listen to and offer constructive advice on the questions that each pair has developed. Your feedback might include comments about the type of responses you might expect, how you could collect the data, whether the questions make sense (or if they could give a biased result), and whether they are relevant to the topic.
6. Once you have received your feedback, rewrite some of your questions and rethink the way you will collect the data.
7. Collect your data and keep it ready to use again during this topic.

on Resources



eWorkbook Topic 11 Workbook (worksheets, code puzzle and project) (ewbk-2011)



Interactivities Individual pathway interactivity: Sampling (int-4540)

Types of data (int-6086)

Selecting samples (int-3811)

Random number generator (int-0089)

Exercise 11.2 Collecting data

learn on

11.2 Quick quiz on

11.2 Exercise

Individual pathways

PRACTISE

1, 4, 7, 8, 10, 13, 16

CONSOLIDATE

2, 5, 11, 14, 17, 18

MASTER

3, 6, 9, 12, 15, 19, 20

Fluency

1. **WE1** Classify each of the following data sets according to whether they are numerical or categorical data.
 - a. Favourite TV programs of each student in your year level.
 - b. Shoe size of the top ten international models.
2. Classify each of the following data sets according to whether they are numerical or categorical data.
 - a. Hours of games played at home each week by each student in your class.
 - b. Birth weight of each baby in a mothers' group.

3. **WE2** Classify each of the following numerical data sets according to whether they are discrete or continuous.
 - a. Length of a stride of each golfer in a local golf club.
 - b. Distance travelled on one full tank from the top ten most popular vehicles.
4. Classify each of the following numerical data sets according to whether they are discrete or continuous.
 - a. Rainfall recorded for each day during spring.
 - b. Cars owned by each family of the students in your class.
5. Classify each of the following categorical data sets as nominal or ordinal.
 - a. Satisfaction levels of customers (Satisfied, Indifferent, Dissatisfied).
 - b. Favourite magazine of each student in your year level.
6. Classify each of the following categorical data sets as nominal or ordinal.
 - a. Sales ranking of the top ten magazines in Australia.
 - b. Age group of the readers of the top ten magazines in Australia.
7. Classify each of the following categorical data sets as nominal or ordinal.
 - a. Types of fish caught while fishing off a pier.
 - b. Genres of movies on a streaming service.
8. **WE3** Classify each of the following data collections according to whether they are samples or populations.
 - a. Favourite TV programs of secondary school students, determined by surveying students in Year 8.
 - b. Size of shoes to be stocked in a department store, determined by the shoe sizes of the top ten international models.
 - c. Number of hours your classmates spend playing games at home, determined by surveying each student in your class.
9. Classify each of the following data collections according to whether they are samples or populations.
 - a. Survey of first 40 customers at the opening of a new store to gauge customer satisfaction.
 - b. The most popular duco colours for cars sold in the last month in Victoria, taken from the registration board database of all 2310 cars registered last month.



Understanding

10. Identify the missing word from each of the following sentences.
 - a. When writing questions for a survey, it is important to consider which type of _____ will be useful for you to collect.
 - b. If you _____ an entire population, this data can be used for the purpose of a _____.
 - c. If the data you collect is grouped numerical data, it is best to use a _____ to visually represent the data.
11. **WE4** Complete the following.
 - a. Use a random number generator to select six days from the month of March.
 - b. Use a random number generator to select five students from your mathematics class.
 - c. State whether your selections from **a** and **b** are samples or populations.
12.
 - a. Randomly select five days of a non-leap year.
 - b. State whether this data is a sample or population.

Reasoning

13. The following are 2020 statistics for motorcycle and car fatalities in Australia.
- Motorcycle fatalities: 190
 - Car fatalities: 732
 - Registered motorcycles: 880 881
 - Registered cars: 14 679 249
 - Kilometres travelled by motorcycles (millions): 1683
 - Kilometres travelled by cars (millions): 162 983
 - Population of Australia: 25 693 059
- a. State whether motorcycles or cars are the safer mode of transport, according to the data. Explain your answer.
 - b. Calculate the ratios or percentages that could be extracted from this data to reinforce your conclusion.
 - c. State the information that you would include if you wanted to argue for the safety of riding a motorcycle. State the information that you would leave out in order to make this argument.
 - d. Write a sentence to convince your readers that a motorcycle is a safer mode of transport.
 - e. Write a sentence to convince your readers that a car is a safer mode of transport.
14. Explain how data is classified into different types, giving examples for each type.
15. Identify and justify two sets of information a person might want to collect from each of the following sources.
- a. A maternity hospital.
 - b. An airport.
 - c. Shoppers at a supermarket.

Problem solving

16. Questions are being written for a survey about students' views on sports. Select the questions from the following list that you would use. Explain your choices.
- a. What do you think about sport?
 - b. List these sports in order from your most favourite to least favourite: football; tennis; netball; golf.
 - c. How often do you watch sports on television?
Choose from these options: once a week; once a day; once a month; once a year; never.
 - d. Why do you like sports?
 - e. How many live sporting events do you attend each year?
 - f. Can you play a sport?
 - g. Which sports do you watch on the television?
Choose from these options: football; tennis; netball; golf; all of these; none of these.
17. Businesses collect and analyse all sorts of data.
- a. Suggest some types of data that businesses might want to collect.
 - b. Suggest some ways in which businesses might collect data.
 - c. Provide some examples of how businesses might use the data that they collect.
18. The local government wishes to survey people about expanding a large windfarm in the Western Downs region of Queensland.
- a. Explain if a census or survey would be most appropriate.
 - b. Explain how survey respondents should be selected to participate.

19. A company involved in constructing a desalination plant in the Wonthaggi region of Victoria wishes to gather some data on public opinion about the presence of the desalination plant in the area and its possible effects on the environment.
The company has decided to conduct a survey and has asked you to plan and implement the survey and then analyse the data and interpret the results.
- Suggest three questions you could include in the survey.
 - Explain what you are trying to find out by asking these questions.
 - State the types of data you would collect.
 - Suggest what the company could use this data for.
 - The company conducts the survey on residents of Wonthaggi.
Discuss the appropriateness of this sample.
20. A large supermarket surveyed shoppers in their store between 7 pm and 9 pm each day to ask their thoughts on the ban on single-use plastics and found that most shoppers did not support the ban.
Discuss the reasonableness of this inference.

LESSON

11.3 Organising data

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- organise data in frequency tables
- display data in dot plots
- display data in stem-and-leaf plots.



11.3.1 Frequency tables

eles-4912

- Data needs to be recorded and organised so that graphs and tables can be drawn and analysis conducted.
- A **frequency table** can be used to record and organise data.
- Data in frequency tables can be numerical or categorical data, and can even include two sets of data.
- A frequency table looks like this:

The tally is recorded from the data.

This is the value/category.

Person's age	Tally	Number of people (f)
0–29		5
30–59		10
60–89		3
		$\Sigma f = 18$

This is the frequency. It is the total number given by the tally. It is labelled f .

This is the sum of values in this column.
Note: Σf means 'sum of f '.

WORKED EXAMPLE 5 Drawing frequency tables with values

The following is a result of a survey of 30 students showing the number of people in their family living at home.

Draw a frequency table to summarise the data.

2, 4, 5, 6, 4, 3, 4, 3, 2, 5, 6, 6, 5, 4, 5,
3, 3, 4, 5, 8, 7, 4, 3, 4, 2, 5, 4, 6, 4, 5

THINK

1. Identify the data as numerical discrete data.
2. There are only seven different scores so the data can be left ungrouped.
3. Set up a frequency table. The first column is for the possible scores and can be labelled 'Number of members in the family'.

The last column is for the frequency (f) and can be labelled 'Number of families'. An optional column between these two can be used to keep a tally for each category.

4. Add up the frequency column ($\sum f$) to confirm that 30 results have been recorded.

WRITE/DRAW

Number of members in the family	Tally	Number of families (f)
2		3
3		5
4		9
5		7
6		4
7		1
8		1
		$\sum f = 30$

WORKED EXAMPLE 6 Drawing a frequency table with categories

The final scores for 30 golfers were recorded as follows. Summarise the data as a frequency table, grouping the scores in intervals of 2, starting at 62.

73, 69, 75, 79, 68, 85, 78, 76, 72, 73,
71, 70, 81, 73, 74, 78, 68, 75, 76, 72,
63, 72, 74, 71, 70, 75, 66, 82, 87, 78



THINK

1. Identify the data as numerical discrete data.
2. The scores range from 63 to 87. This is a range of 24 shots.
3. For this example the data, starting at 62, will be grouped in intervals of 2.

WRITE/DRAW


Golf scores (x)	Tally	Number of players (f)
62–63		1
64–65		0
66–67		1
68–69		3
70–71		4
72–73	 	6
74–75	 	5
76–77		2
78–79		4
80–81		1
82–83		1
84–85		1
86–87		1
		$\Sigma f = 30$

4. Add up the frequency column to check that all 30 scores have been recorded.

11.3.2 Dot plots

eles-4913

- A **dot plot** is another way of displaying data.
- Data in a dot plot can be categorical or numerical.
- A dot plot looks like this:



Person's age (x)	Tally	Number of people (f)
0–29	 	5
30–59	 	10
60–89		3
		$\Sigma f = 18$

The x column categories are listed along the base of the plot.

The f column values indicate the number of dots.

WORKED EXAMPLE 7 Drawing a dot plot

A survey was conducted, asking students about the type of vehicle driven by their parents. Their responses were collected and recorded as shown.

Display the data as a frequency table and as a dot plot.


Sedan	4WD	Sedan	Sedan
4WD	SUV	SUV	Station wagon
SUV	Convertible	Station wagon	4WD
Sedan	Sports car	Convertible	Station wagon
Sedan	Station wagon	SUV	Station wagon
SUV	SUV	Sedan	Sedan

THINK

- Identify this data as categorical ordinal data.
- There are six different scores (or categorical labels).
- Use a frequency table of two (or three) columns.
The first column is for the possible scores (x) and can be labelled as 'Type of vehicle'.
The last column is for the frequency (f) and can be labelled 'Number of vehicles'.
An optional middle column can be used to keep a tally for each category.
- Add up the frequency column ($\sum f$) to confirm that 24 scores have been recorded.
- Place a dot for each vehicle type in the appropriate column.

WRITE/DRAW

Type of vehicle (x)	Tally	Number of vehicles (f)
4WD		3
SUV		6
Sedan		7
Station wagon		5
Sports car		1
Convertible		2
		$\sum f = 24$



11.3.3 Stem-and-leaf plots

eles-4914

- A **stem-and-leaf plot**, or stem plot, is a graph that displays numerical data.
- Stem plots have two columns:
 - the stem shows the leading part of the value
 - the leaf shows the trailing part of the value.

- The following stem plot shows the sizes of 20 students' book collections.

Book collection size for 20 students

Key: 2 | 5 = 25 books

<i>Stem</i>	<i>Leaf</i>
0	4
1	4 6
2	2 3 5 8
3	0 4 4 6 7 7 9
4	1 3 5 9
5	4 6

- From the stem plot shown, the sizes collected were 4, 14, 16, 22, 23, 25, 28, 30, 34, 34, 36, 37, 37, 39, 41, 43, 45, 49, 54, 56.

WORKED EXAMPLE 8 Summarising data in a stem plot

The following scores represent the distance (in km) travelled by a group of people over a particular weekend.

Summarise the data in a stem plot.

48, 67, 87, 2, 34, 105, 34, 45, 63, 98, 12, 23, 35,
54, 65, 41, 34, 23, 12, 38, 18, 58, 53, 44, 39, 29

THINK

- Write a title and key and set up a stem plot.
- The scores range from 2 to 105. Grouping them in ranges of 10 will give 11 categories. The stem will hold the tens place value and the leaf will hold the unit/ones place value (the last digit) for each score.

WRITE/DRAW

Distance travelled over one weekend
Key: 4 | 7 = 47 km

<i>Stem</i>	<i>Leaf</i>
0	2
1	2 2 8
2	3 3 9
3	0 4 4 5 8 9
4	1 4 5 8
5	3 4 8
6	3 5 7
7	
8	7
9	8
10	5

- Transfer each score to the stem plot by recording its last digit in the leaf column alongside the row of the stem column that corresponds to its tens place digit.
- Rearrange each row in the leaf column so that the digits are in ascending order from left to right.

- Sometimes the leaves of a stem plot can become too long. This can be fixed by breaking the stems into smaller intervals. The second part of the interval is marked with an asterisk (*).

WORKED EXAMPLE 9 Representing data in a stem plot with smaller stem intervals

The heights of 30 students are measured (in cm) and recorded as follows.

125, 143, 119, 136, 127, 131, 139, 122, 140, 118,
120, 123, 132, 134, 127, 129, 124, 131, 138, 133,
122, 128, 130, 135, 141, 139, 121, 138, 131, 126

Represent the data in a stem plot.

THINK

1. Write a title and key, then draw up the stem plot with the numbers in each row of the leaf column arranged in ascending order from the stem.

WRITE/DRAW

Heights of 30 students (in cm)

Key: 11 | 8 = 118 cm

Stem	Leaf
11	8 9
12	0 1 2 2 3 4 5 6 7 7 8 9
13	0 1 1 1 2 3 4 5 6 8 8 9 9
14	0 1 3

2. The leaves of the two middle stem values are long. They would be easier to interpret with the stem broken into smaller intervals, for example intervals of 5. The stem of 12 would then include the numbers from 120 to 124 while the stem of 12* would include the numbers from 125 to 129. Reconstruct the stem plot according to this new set of intervals.

Heights of 30 students (in cm)

Key: 12* | 5 = 125 cm

Stem	Leaf
11	8 9
12	0 1 2 2 3 4
12*	5 6 7 7 8 9
13	0 1 1 1 2 3 4
13*	5 6 8 8 9 9
14	0 1 3

on Resources



eWorkbook Topic 11 Workbook (worksheets, code puzzle and a project) (ewbk-2011)



Interactivities Individual pathway interactivity: Collecting data (int-4541)

Collecting data (int-3807)

Stem plots (int-6242)

Individual pathways

PRACTISE

1, 4, 8, 9, 12, 15

CONSOLIDATE

2, 5, 6, 10, 13, 16

MASTER

3, 7, 11, 14, 17, 18

Fluency

- WE5&7** The following set of data represents the scores achieved in a multiple choice Maths test.

4, 6, 8, 3, 6, 9, 1, 3, 5, 6, 4, 7, 5, 9, 3, 2, 7, 8, 9, 6, 5, 4, 6, 5, 3, 5, 7, 6, 7, 7

 - Present the data in a frequency table.
 - Present the data as a dot plot.
- The following set of data shows the types of pets kept by 18 people.

cat, dog, fish, cat, cat, cat, dog, dog, bird, turtle, dog, cat, turtle, cat, dog, dog, turtle, bird

 - Present the data in a frequency table.
 - Present the data as a dot plot.
 - Explain whether this data can be presented in a stem plot.
- WE6** Present the data shown in the table as a frequency table by grouping the data in intervals of 10 (0–9, 10–19, ...).

 - WE8** Present the data as a stem plot.

Death by motor vehicle accidents: Males aged 15–24 years (rates per 100 000 people)				
State	2008	2009	2010	2011
NSW	14	13	18	9
Vic.	19	25	14	12
Qld	25	17	26	15
SA	23	34	22	26
WA	22	24	36	27
Tas.	42	18	36	42
NT	43	60	71	45
ACT	0	11	14	21

- A sample of students surveyed by the Australian Bureau of Statistics were asked the colour of their eyes. The results were as follows:

Brown Hazel Blue Brown Brown Green Green Brown Brown Hazel
 Brown Brown Brown Brown Blue Brown Green Blue Blue Blue
 Brown Brown Brown Brown Hazel Blue Hazel Blue Brown Blue
 Brown Blue Green Hazel Hazel Hazel Green Blue Blue Brown

 - Set up a frequency table for the data.
 - Construct a dot plot for the data.
 - Comment on the proportion of people who have brown eyes.

5. Draw a stem-and-leaf plot for each of the following sets of data. Comment on each distribution.
- 18, 22, 20, 19, 20, 21, 19, 20, 21
 - 24, 19, 31, 43, 20, 36, 26, 19, 27, 24, 31, 42, 29, 25, 38
 - 346, 353, 349, 368, 371, 336, 346, 350, 359, 362

6. **WE9** Redraw the stem plots in question 5 so that the stems are arranged in intervals of 5 rather than intervals of 10.

7. The following times (to the nearest minute) were achieved by 40 students during a running race.

23	45	25	48	21	56	33	34	63	43	42	41	26	44
45	41	40	39	37	53	26	55	48	39	29	52	57	33
31	32	71	60	49	52	32	28	47	42	37	33		

- Construct a stem plot with intervals of 5 minutes.
 - Construct a frequency table. Use intervals of 5 minutes.
 - If the top 10 runners were chosen for the representative team, identify the qualifying time as given by:
 - the stem plot
 - the frequency table.
 - Identify the number of times that the finish time of 33 minutes was recorded. Explain which summary (stem plot or frequency table) was able to give this information.
8. The ages of participants in a weights class at a gym are listed below:

17, 21, 36, 38, 23, 45, 32, 53, 18, 25, 14, 29, 42, 26, 18, 27, 37, 19, 34, 20, 35

Display the data as a stem plot using intervals of 5.

9. **MC** Consider the stem plot shown.

Key $7|6 = 7.6$

Stem	Leaf
7	8
8	0 8 9
9	1 6 7 8
10	3 5 8
11	2

Identify which of the following data sets matches the stem plot.

- 78, 80, 88, 89, 91, 96, 97, 98, 103, 105, 108, 112
- 8, 0, 8, 9, 1, 6, 7, 8, 3, 5, 8, 2
- 7.8, 8.0, 8.8, 8.9, 9.1, 9.6, 9.7, 9.8, 1.03, 1.05, 1.08, 1.12
- 7.8, 8.0, 8.8, 8.9, 9.1, 9.6, 9.7, 9.8, 10.3, 10.5, 10.8, 11.2
- 78, 8089, 91 678, 10 358, 112

Understanding

10. The following data represent the life expectancy (in years) of Australians from 40 different age groups.

83.7	84.5	84.7	85.2	84.6	84.5	85.9	84.9	88.3	86.5
84.1	84.5	84.8	84.4	84.7	84.5	86.0	85.0	88.3	86.6
84.1	84.5	84.8	84.4	84.7	85.7	87.3	85.0	88.8	86.6
84.1	84.5	84.9	84.5	84.7	85.8	87.6	85.1	88.8	86.9

- Present the data as a frequency table.
- Present the data as a stem plot.


11. Choose the most appropriate technique for summarising the data in the table. Justify your choice.

Median age of Australian females and males from 2002 to 2009		
Year	Females	Males
2002	36.6	35.2
2003	36.9	35.3
2004	37.1	35.5
2005	37.3	35.7
2006	37.4	35.9
2007	37.6	36.1
2008	37.6	36.1
2009	37.7	36.1

Reasoning

12. The dot plot shown is a Year 9 student's attempt to display the values:

8, 10, 10, 11, 11, 12, 12, 15, 15, 15, 15, 17, 18, 19



- a. List the mistakes that the student made in drawing the dot plot.
 b. Redraw the dot plot so that it correctly displays all of the data given.
13. a. Sometimes you are asked to represent two sets of data on separate stem-and-leaf plots. Suggest how you could represent two sets of data on a single stem plot while keeping them separate. Demonstrate your method using the data sets below that show the heights (in cm) of 12 boys and 12 girls in a Year 8 class:



Boys	154	156	156	158	159	160	162	164	168	171	172	180
Girls	145	148	152	152	155	156	160	161	161	163	165	168

- b. Consider the stem-and-leaf plot below. Explain how you could make this plot easier to read and use.

Stem	Leaf
1	0 2 2 2 5 7 8 9
2	0 0 0 0 0 1 1 2 2 2 2 2 4 4 7 7 7 7 8 8 8 8 9 9
3	0 2 2 3 3 3 5 5 7 8 9

14. The dot plots below display Maths test results for two Year 9 classes. The results show the marks out of 20.

- a. Determine the number of students in each class.
 b. For each class, determine the number of students who scored 15 out of 20 on the test.
 c. For each class, determine the number of students who scored more than 10 on the test.
 d. Use the dot plots to describe the performance of each class on the test.



Problem solving

15. The following data shows the speeds of 30 cars recorded by a roadside camera outside a school, where the speed limit is 40 km/h.

20, 27, 30, 36, 45, 39, 15, 22, 29, 30, 30, 38, 40, 40, 40,
40, 42, 44, 20, 45, 29, 30, 34, 37, 40, 45, 60, 38, 35, 32

- Present the data as an ordered stem-and-leaf plot.
 - Write a paragraph for the school newsletter about the speed of cars outside the school.
16. A Year 9 Science class recorded the masses of mice (measured in grams) as shown below.

25.1, 24.8, 25.1, 27.3, 25.3, 29.5, 24.5, 26.7, 24.0, 26.3, 25.4,
26.3, 23.9, 25.8, 25.4, 24.6, 25.1, 23.9, 33.2, 24.5, 28.1, 27.3

- Draw an ordered stem-and-leaf plot for this data.
- Calculate the range of this data.
- Draw an ordered stem-and-leaf plot for the masses of the mice rounded to the nearest gram. Discuss whether a stem-and-leaf plot is suitable for the rounded data. Compare the stem-and-leaf plot with the original data. Explain your conclusions.

17. A doughnut graph can be used to compare two or more sets of data using the basic principles of a pie graph. A doughnut graph comparing students' methods of travel to school on Monday, Tuesday and Wednesday is shown.

- List the similarities and differences between a doughnut graph and a pie graph.
- Write a series of instructions for drawing a doughnut graph.
- Test your instructions on the following data by creating a doughnut graph to represent it.


Method of travel	Monday	Tuesday	Wednesday
Walking	5	7	25
Cycling	6	8	23
Car	8	2	3
Bus	3	3	2
Train	2	2	1

- Compare your doughnut chart with the one shown.
- Suggest some advantages and disadvantages of using a doughnut graph instead of a multiple column graph to compare two or more sets of data.

18. Analysis of written English prose reveals the proportion of vowels and consonants to be roughly the same in every large sample — approximately 7:11. The letter *e* accounts for one-third of the occurrence of vowels. In a passage consisting of 10 000 letters, determine approximately how many times you would expect the letter *e* to occur.



Method of travel to school
Monday (inner ring)
Tuesday (middle ring)
Wednesday (outer ring)



LESSON

11.4 Displaying data

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- display data in bar charts, pie charts and histograms
- present data as back-to-back stem-and-leaf plots.

11.4.1 Bar charts and pie charts


eles-4915

- **Bar charts** and **pie charts** are used to display categorical data.


Displaying data in bar charts and pie charts

- A bar chart for the following data is shown as an example.

Items purchased from canteen	
Item purchased	Number (f)
Sandwich	35
Juice box	10
Chocolate bar	12
Fruit salad	5



- A pie chart for the same data is shown as an example.



WORKED EXAMPLE 10 Displaying a frequency table as a bar chart and a pie chart

The Chalemet family is having a major reunion. The age groups of the family members who are attending have been recorded in the table below.

Age group	Number
Baby	20
Toddler	24
Child	34
Teenager	36
Adult	50
Pensioner	36

Display the frequency table as a:

a. bar chart

b. pie chart.

THINK

a. 1. Give the bar chart a suitable title.

2. Identify the category names and use them to label the horizontal axis.

It is the convention to leave a space between each bar on a bar chart.


3. Label the vertical axis with a suitable scale and title.

4. Draw each bar so that its height matches the frequency of the category it represents.

WRITE/DRAW

a.

Chalemet family age groups



b. 1. Add one extra column to the frequency table to calculate the size of the sector.

The size of each sector is determined by its angle. To calculate this angle, use the rule


$$\text{angle} = \frac{f}{\sum f} \times 360^\circ.$$

b.

Age group	Number (f)	Size of sector
Baby	20	$\frac{20}{200} \times 360^\circ = 36^\circ$
Toddler	24	$\frac{24}{200} \times 360^\circ = 43.2^\circ \approx 43^\circ$
Child	34	$\frac{34}{200} \times 360^\circ = 61.2^\circ \approx 61^\circ$
Teenager	36	$\frac{36}{200} \times 360^\circ = 64.8^\circ \approx 65^\circ$
Adult	50	$\frac{50}{200} \times 360^\circ = 90^\circ$
Pensioner	36	$\frac{36}{200} \times 360^\circ = 64.8^\circ \approx 65^\circ$
Σ	$\Sigma f = 200$	360°

2. Give the pie chart a suitable title.
3. Create a legend with a suitable colour for each category.
4. Draw a circle.
5. Using a protractor, measure and draw the correct angle of each sector.
6. Colour the sectors using the legend.

Chalemet family age groups



- Adult
- Teenager
- Pensioner
- Child
- Toddler
- Baby

TI | THINK


- a. 1. Open a Lists & Spreadsheet page. Label column A as agegroup and column B as number. Enter the data as shown in the table.

WRITE

a.

agegroup	number
baby	20
toddler	24
child	34
teenager	36
adult	50

2. To create a summary plot, press:
 - MENU
 - 3: Data
 - 8: Summary Plot.
 Set the X List (category) as agegroup and the Summary List as number. You have the option of displaying the summary plot in a split screen or in a new page. Choose New Page for Display On. Use TAB to move to OK. Press ENTER. Hover the pointer over each bar to see the percentage frequency.



CASIO | THINK


- a. 1. Open a new Spreadsheet screen. Enter the data into columns A and B.

WRITE

a.

	A	B	C
1	Baby	20	
2	Toddler	24	
3	Child	34	
4	Teenager	36	
5	Adult	50	
6	Pensioner	36	


2. Highlight columns A and B. Press:
 - Graph
 - Bar
 - Stacked.
 Note: This is a frequency bar graph. To draw a percentage frequency bar graph, change the entered data to percentages.



TI | THINK

- b. To draw the pie chart, press:
- MENU
 - 1: Plot Type
 - 9: Pie Chart.
- Hover the pointer over each bar to see the percentage frequency.


WRITE



CASIO | THINK

- b. To draw the pie chart, highlight the numerical data only. Then press:
- Graph
 - Pie.
- Hover the pointer over each sector to see the percentage frequency.

WRITE



11.4.2 Histograms


eles-4916

- **Histograms** are used to display numerical data.
- They are used most commonly to display frequency data.
- The horizontal axis of a histogram can be labelled with:
 - values
 - intervals
 - mid points of intervals.

Displaying data in histograms


Histograms with values along the horizontal axis

- The *x*-axis of this type of histogram shows the values or results of the data collected.
- The *y*-axis of this type of histogram shows the frequency of each result.



Histograms with intervals along the horizontal axis


- The information along the horizontal axis are intervals (or groups) of values.



Values show the start of each interval.

Displaying data with midpoints along the horizontal axis

- The midpoint of a class interval is calculated by taking the average of the two extreme values of that class interval.



Values show the midpoint of each interval.

- For example, the midpoint of the 60–70 class interval is $\frac{60 + 70}{2} = 65$.

WORKED EXAMPLE 11 Constructing a histogram to represent data

A sample of 40 people was surveyed regarding the number of hours per week they spent on their phones. The results are listed below.

12.9, 18.3, 9.9, 17.1, 20, 7.8, 24.2, 16.7, 9.1, 27,
 7.2, 16, 26.5, 15, 7.4, 28, 11.4, 20, 9, 11,
 23.7, 19.8, 29, 12.6, 19, 12.5, 16, 21, 8.3, 5,
 16.4, 20.1, 17.5, 10, 24, 21, 5.9, 13.8, 29, 25

- Calculate the range for the data.
- Organise the data into five class intervals and create a frequency distribution table that displays class intervals, midpoints and frequencies.
- Construct a histogram to represent the data, using midpoints.

THINK

- Determine the range of the number of hours spent on phones.
- Determine the size of the class intervals. Class intervals have been recorded as 5– <10, 10– <15 and so on, to accommodate the continuous data.
 - Rule a table with three columns, headed ‘Number of hours on phone each week’ (class interval), ‘Midpoint’ (class centre) and ‘Frequency’.
 - Tally the scores and enter the information into the frequency column.
 - Calculate the total of the frequency column.


WRITE/DRAW

- Range = largest value – smallest value
 $= 29 - 5$
 $= 24$
- Class intervals of 5 hours will create 5 groups, and cover a range of 25 numbers.

Number of hours on phone each week	Midpoint	Frequency (f)
5– < 10	7.5	9
10– < 15	12.5	7
15– < 20	17.5	10
20– < 25	22.5	8
25– < 30	27.5	6
	Total	40

- Rule a set of axes on graph paper. Label the horizontal axis ‘Number of hours on phone’ and the vertical axis ‘Frequency’.
 - Draw the first column so that it starts and finishes halfway between class intervals and reaches a vertical height of 9 people. Make sure you leave space between the y-axis and the first column.
 - Draw the columns for each of the other scores.

c.



WORKED EXAMPLE 12 Creating a histogram from a stem plot

A stem plot was used to display the weight of airline luggage for flights to the Gold Coast.

Convert the stem plot shown into a histogram.

Weights of airline luggage


Key: 21 | 3 = 21.3 kg

Stem	Leaf
18	5 8 9
19	2 3 6 7
20	5 7 8 8 9 9
21	0 2 6 9
22	0 1 9
23	2

THINK

1. Identify the data as numerical continuous data with six groups or class intervals. Label the x -axis as 'Weight of airline luggage'. Leave a half-interval gap at the beginning and use an appropriate scale for the x -axis.
2. Label the y -axis as 'Frequency'.
3. Draw the first column to a height of its frequency.
4. Repeat for all class intervals.



WRITE/DRAW



11.4.3 Misleading graphs

- Some graphs misrepresent data, which can lead to incorrect conclusions.
- Data can be misrepresented by:
 - adjusting scales
 - omitting values
 - shortening or lengthening axes
 - distorting images.

Consider the following two graphs depicting the rate of growth of pollutants.





Both of these graphs show the same numerical information, but graph (a) has a linear scale on the vertical axis and graph (b) does not. Graph (a) emphasises the ever-increasing rate of growth of pollutants, whereas graph (b) suggests a slower, linear growth.

WORKED EXAMPLE 13 Determining whether a graph is misleading

The following data give wages and profits for a certain company. All figures are in millions of dollars.

Year	2015	2016	2017	2018
Wages	6	9	13	20
% increase in wages	25	50	44	54
Profits	1	1.5	2.5	5
% increase in profits	20	50	66	100



- State whether the graphs accurately reflect the data.
- Suggest which graph would you rather have published if you were:
 - an employer dealing with employees requesting pay increases
 - an employee negotiating with an employer for a pay increase.

THINK

- Look at the scales on both axes. Both scales are linear.
 - Look at the units on both axes. Graph (a) has its y-axis in \$, whereas graph (b) has its y-axis in %.
- Compare wage increases with profit increases.
 - The employer wants a graph that shows wages are already increasing exponentially to argue against pay increases.

WRITE

- The graphs do represent data accurately. However, quite a different picture of wage and profit increases is painted by graphing with different units on the y-axis.
- The employer would prefer graph (a) because he/she could argue that employees' wages were increasing at a greater rate than profits.

- ii. 1. Consider again the increases in wages and profits.
- 2. The employee doesn't like to see profits increasing at a much greater rate than wages.
- ii. The employee would choose graph (b), arguing that profits were increasing at a great rate while wage increases clearly lagged behind.

 **11.4.4 Back-to-back stem-and-leaf plots**

eles-4917

- **Back-to-back stem-and-leaf plots** are used to display two related numerical data sets.
- The leaf values start in the centre and count outwards.

WORKED EXAMPLE 14 Displaying data as a back-to-back stem-and-leaf plot

The ages of experienced and inexperienced bowlers using a ten-pin bowling centre are listed.

Experienced: 65, 15, 50, 15, 54, 16, 57, 16, 16, 21, 17, 28, 17, 27, 17, 35, 18, 19, 22, 30, 34, 22, 22, 31, 43, 23, 48, 23, 46, 25, 30, 21.

Inexperienced: 16, 60, 16, 52, 17, 38, 38, 43, 20, 17, 45, 18, 45, 36, 21, 34, 19, 32, 29, 21, 23, 32, 23, 22, 23, 31, 25, 28.

Display the data as a back-to-back stem-and-leaf plot and comment on the distribution.



THINK

1. Rule three columns, headed Leaf (Inexperienced), Stem and Leaf (Experienced).
2. Make a note of the smallest and largest values of both sets of data (15 and 65). List the stems in ascending order in the middle column.
3. Beginning with the experienced bowlers, work through the given data and enter the leaf (unit component) of each value in a row beside the appropriate stem.
4. Repeat step 3 for the set of data for inexperienced bowlers.
5. Include a key to the plot that informs the reader of the meaning of each entry.

WRITE/DRAW

<i>Leaf (Inexperienced)</i>	<i>Stem</i>	<i>Leaf (Experienced)</i>
987766	1	5566677789
8532331910	2	1872223351
1224688	3	50410
553	4	386
2	5	047
0	6	5

6. Redraw the stem-and-leaf plot so that the numbers in each row of the leaf columns are in ascending order.
Note: The smallest values should be closest to the stem column and increase as they move away from the stem.

Key: 1 | 5 = 15

<i>Leaf (Inexperienced)</i>	<i>Stem</i>	<i>Leaf (Inexperienced)</i>
987766	1	5566677789
9853332110	2	1122233578
8864221	3	00145
553	4	368
2	5	047
0	6	5

7. Comment on any interesting features.

The youngest experienced bowler attending the ten-pin bowling centre is 15 and the oldest is 65. The youngest and oldest inexperienced bowlers attending the ten-pin bowling centre are 16 and 60 respectively. There were an equal amount of experienced and inexperienced bowlers in their 20s. The next largest number of experienced bowlers were in their teens, while the next largest number of inexperienced bowlers were in their 30s.

on Resources



eWorkbook Topic 11 Workbook (worksheets, code puzzle and project) (ewbk-2011)



Interactivity Individual pathway interactivity: Displaying data (int-4542)

Exercise 11.4 Displaying data

learn on

11.4 Quick quiz **on**

11.4 Exercise

Individual pathways

PRACTISE

1, 2, 5, 8, 12, 15, 18

CONSOLIDATE

3, 6, 7, 9, 11, 13, 16, 19

MASTER

4, 10, 14, 17, 20

Fluency

1. **WE10** The number of participants in various classes at the local gym is given as:

Aerobics — 120 members

Pilates — 60 members

Aqua aerobics — 30 members

Boxing — 15 members

Spin — 45 members

Step — 90 members.

Represent this information as a:

- a. bar chart
- b. pie chart.

2. **WE11** In a survey, 40 people were asked about the number of hours a week they spent on their phone. The results are listed below.

10.3, 13, 7.1, 12.4, 16, 11, 6, 14, 6, 11.1, 5.8, 14, 12.9, 8.5, 27.5, 17.3, 13.5, 8, 14, 10.2, 13, 7.9, 15, 10.7, 16, 8.4, 18.4, 14, 21, 28, 9, 12, 11.5, 13, 9, 13, 29, 5, 24, 11

- State whether this data is continuous or discrete.
 - Determine the range of the data.
 - Organise the data into class intervals and create a frequency distribution table that displays the class intervals, midpoints and frequencies.
 - Construct a histogram.
3. The number of phone calls made per week in a sample of 56 people is listed below.
- 21, 50, 8, 64, 33, 58, 35, 61, 3, 51, 5, 62, 16, 44, 56, 17, 59, 23, 34, 57, 49, 2, 24, 50, 27, 33, 55, 7, 26, 52, 17, 54, 78, 69, 53, 2, 42, 52, 25, 48, 63, 12, 72, 36, 66, 15, 28, 67, 13, 23, 10, 72, 72, 89, 80, 62
- Organise the data into a grouped frequency distribution table using a suitable class interval.
 - Display the data as a combined histogram.
4. For the following data, construct:
- a frequency distribution table
 - a histogram.



Class sizes at Jacaranda Secondary College									
38	24	20	23	27	27	22	17	30	26
25	16	29	26	15	26	19	22	13	25
21	19	23	18	30	20	23	16	24	18
12	26	22	25	14	21	25	21	31	25

5. **WE12** The number of bananas sold over a 5-week period at a high-school canteen is shown in the following stem-and-leaf plot. Represent this information as a histogram.

Key: 0* | 7 = 7


Stem	Leaf
0	1 1 1 2 2 2 2 2 3 3 3 4 4 4 4 4
0*	5 5 5 6 6 7 7 8

6. The distances 300 students have to travel to attend their school are summarised in the table below.

Distance km	Number of students
0– < 2	112
2– < 4	65
4– < 6	56
6– < 8	44
8– < 10	15
10– < 12	8

Represent this information a histogram.

7. **MC** Examine the bar chart shown and state the number of students who catch public transport to school.



- A. 80 B. 100 C. 120 D. 200 E. 310

Understanding

8. An investigation into transport needs for an outer suburb community recorded the number of passengers boarding a bus during each of its journeys.

12, 43, 76, 24, 21, 46, 24, 46, 24, 21, 46, 54, 109, 87, 23,
78, 37, 22, 139, 65, 78, 89, 52, 23, 30, 54, 56, 32, 66, 49

Construct a histogram using class intervals of 20.

9. A drama teacher is organising his DVD collection into categories based on the genre of the movie or TV show. He recorded the following for the first 20 DVDs he categorised.

musical, musical, drama, comedy, horror,
musical, comedy, action, drama, comedy,
drama, drama, musical, drama, comedy,
romance, drama, musical, romance, drama

- a. Show the data set as a bar chart.

He continued with the task, and the classification for his collection of 200 DVDs is summarised in the frequency table shown below.

Genre	Number of DVDs	Size of sector for a pie chart
Musical	52	
Drama	98	
Comedy	30	
Horror	10	
Action	5	
Romance	5	
Total	200	360°

- b. Copy and complete the frequency table.
c. Construct a pie chart for the whole 200 DVD collection.


10. The height, in centimetres, of 32 students in Year 9 was recorded as follows.

167, 162.9, 157, 166.7, 146.5, 163, 156.1, 168, 159.4, 170.1,
 152.1, 152.1, 174.5, 156, 163, 157.8, 161, 178, 151, 148, 166.7,
 154.6, 150.3, 166, 160, 155.9, 164, 157, 171.1, 168, 158, 162

- State whether the data is continuous or discrete.
- Organise the data into seven classes, starting at 145 cm, and create a frequency distribution table.
- Display the data as a histogram.
- Explain whether it is possible to obtain the range of heights from the frequency distribution table without being given any other information.
- MC** Determine how many students stood at least 160 cm tall.
 - 15
 - 16
 - 17
 - 7
 - 8
- MC** Calculate what percentage of students were under the minimum height of 165 cm required to qualify for the basketball team.
 - 21%
 - 30%
 - 87%
 - 70%
 - 13%
- Reorganise the data into class intervals of 4 cm (for example, 145–148 cm).
- Draw a new histogram and compare it to your histogram from part c. Discuss any advantages or disadvantages of having a smaller class interval.



11. The following histogram illustrates the number of days that members of a tennis team practised in a week.




- Describe what the frequency in this problem represents.
- Calculate the average number of days that the team members practised in a week.

12. **WE13** Examine this graph of employment growth in a company.



Explain why this graph is misleading.

13. This graph shows how the \$27 that a buyer pays for a new album by their favourite singer-songwriter is distributed among the departments of a major recording company involved in its production and marketing.



You are required to find out whether or not the graph is misleading, to explain fully your reasoning, and to support any statements that you make.

- Comment on the shape of the graph and how it could be obtained.
 - Does your visual impression of the graph support the figures? Explain your response.
14. The back-to-back stem-and-leaf plot displays the heights of a group of Year 9 students.

- a. **MC** The total number of Year 9 students is:

- 13
- 17
- 30
- 36
- 171

Key: 13 | 7 = 137 cm

Leaf (males)	Stem	Leaf (females)
98	13	78
98876	14	356
988	15	1237
7665	16	356
876	17	1

- b. **MC** The tallest male and shortest female heights respectively are:

- 186 cm and 137 cm
- 171 cm and 148 cm
- 137 cm and 188 cm
- 178 cm and 137 cm
- 178 cm and 171 cm

- c. Compare the heights of boys and girls.

Reasoning

15. **WE14** The number of goals scored in football matches by Mitch and Yani were recorded as follows:

Mitch	0	3	1	0	1	2	1	0	0	1
Yani	1	2	0	1	0	1	2	2	1	1

- Display the data as a back-to-back stem-and-leaf plot and comment on the distribution.
- Calculate how many times Mitch and Yani scored more than a single goal.
- Determine who scored the most goals in any single match.
- Determine who scored the greatest number of goals overall.
- State who is the most consistent performer.
Explain your answer.

16. Percentages in a Mathematics exam for two classes were as follows:

9A	32	65	60	54	85	73	67	65	49	96	57	68
9B	46	74	62	78	55	73	60	75	73	77	68	81

- Construct a back-to-back stem-and-leaf plot of the data.
 - Calculate the percentage of each group that scored above 50.
Give your answers to the nearest whole number.
 - Identify the group with more scores over 80.
 - Compare the clustering for each group.
 - Comment on extreme values.
 - Calculate the average percentage for each group.
17. Suggest two reasons stem plots rather than frequency tables are preferred by statisticians.

Problem solving

18. When reading the menu at the local Chinese restaurant, you notice that the dishes are divided into sections. The sections are labelled chicken, beef, duck, vegetarian and seafood.
- Identify the type of data that this menu presents.
 - Suggest the best way to represent this data.




19. You are working in the marketing department of a company that makes sunscreen and your boss wants to know if there are any unusual results from the market research you completed recently.
You completed a survey of people who use your sunscreen, asking them how many times a week they used it.
Their answers are given in the data set shown.

7, 2, 5, 4, 7, 5, 7, 2, 5, 4, 3, 5, 7, 7, 4, 5, 21, 5, 5, 2

Analyse the data set to work out what you will say to your boss, and try to explain any unusual variables.

20. The pie chart shown displays data collected about the types of roses in the local botanical gardens.
- Carefully measure the angle of each section and record the results in a table.
 - Determine the smallest possible total number of rose bushes in the original data.
 - Create a bar chart displaying the data, assuming the total number of bushes is your answer from b.



LESSON

11.5 Measures of centre

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- calculate the mean, median and mode of raw data
- calculate the mean, median and mode of data in tables and plots
- calculate the mean, median and mode of grouped data
- determine whether a distribution is symmetrical, skewed or bimodal.

11.5.1 Mean, median and mode

eles-4918

- There are three values commonly used to measure the centre of data. These are the **mean**, **median** and **mode**.
 - The mean is also known as the average.
 - The median is the middle of the data.
 - The mode is the most common value in the data.

Calculating the mean

- For raw data, calculate the mean using the formula:

$$\bar{x} = \frac{\sum x}{n} = \frac{\text{sum of all data values}}{\text{number of all data values}}$$

For example, for the data 2, 2, 2, 3, 3, 7, 12, 13, 19:

$$\text{sum of data values} = 2 + 2 + 2 + 3 + 3 + 7 + 12 + 13 + 19 = 63$$

$$\text{number of data values} = 9$$

$$\text{mean} = \frac{63}{9} = 7$$

- For data in frequency tables, calculate the mean using the formula:

$$\bar{x} = \frac{\sum xf}{\sum f} = \frac{\text{sum of (data values} \times \text{frequency)}}{\text{sum of frequencies}}$$

Calculating the median

- Start with data in order from smallest to largest.
- The median is the middle of the data.
- The middle value will be the $\frac{n+1}{2}$ th score, where n is the number of scores or pieces of data.

For example, for the data 2, 2, 2, 3, 3, 7, 12, 13, 19 there are 9 scores, so the middle value will be the 5th score (because $\frac{9+1}{2} = 5$).

2, 2, 2, 3, 3, 7, 12, 13, 19

median = 3

Note: If n is even, find the median by averaging the two middle scores.

Calculating the mode

To calculate the mode, determine the most common value in the data.

For example, for the data 2, 2, 2, 3, 3, 7, 12, 13, 19:

the most common score is 2

mode = 2

Note: It is possible to have more than one mode.

WORKED EXAMPLE 15 Calculating the median, mean and mode

Consider the data set shown.

3, 7, 4, 5, 6, 2, 8, 3, 3, 5

Calculate the:

a. median

b. mean

c. mode.

THINK

a. 1. The median is the middle of the data.
Organise the data from smallest to largest.

2. The median, or middle score, is found by first counting how many scores there are. If there is an even number of scores, then the median is the average of the two middle scores. Use the rule: median score = $\frac{n+1}{2}$ th score where n is the number of scores.

b. 1. The mean is the average of the data values
Add up all the data values.

2. Divide the result by the total number of scores.

c. The mode is the score that occurs the most frequently in the list.

WRITE

a. 2 3 3 3 4 5 5 6 7 8

$$\begin{aligned}\text{Median score} &= \frac{n+1}{2} \text{th score} \\ &= \frac{10+1}{2} \\ &= 5.5 \text{th score}\end{aligned}$$

The median score is between the 5th and 6th scores, which is between 4 and 5.

\therefore Median = 4.5

b. Mean = \bar{x}

$$\begin{aligned}&= \frac{\sum x}{n} = \frac{\text{sum of all data values}}{\text{number of data values}} \\ \bar{x} &= \frac{3+7+4+5+6+2+8+3+3+5}{10} \\ \bar{x} &= \frac{46}{10} \\ &= 4.6\end{aligned}$$

c. Mode = 3

WORKED EXAMPLE 16 Calculating median, mean and mode from a frequency table

The combined number of goals scored in each of the 48 first round games of this year's World Cup is summarised in the frequency table below.

Total combined goals scored	Number of matches
0	6
1	13
2	12
3	9
4	5
5	2
6	0
7	1
Σf	48

Calculate the:

- median number of goals scored
- mean number of goals scored
- modal number of goals scored.

THINK

- The median is the middle value of the data set.
The sum of the frequencies is the number of scores, n . If there is an even number of scores, then the median is the average of the 2 middle scores.
 - Determine where the 24th and 25th scores lie.
- The mean is the average value.
Set up a third column on the frequency table and multiply the score and its frequency.
 - Add up all of the products of xf .

WRITE

$$\begin{aligned}
 \text{a. Median score} &= \frac{n+1}{2} \text{th match} \\
 &= \frac{48+1}{2} \\
 &= 24.5 \text{th match}
 \end{aligned}$$

This score is between the 24th match and the 25th match.

The 24th score is 2 combined goals.

The 25th score is 2 combined goals.

Median = 2 combined goals

- | Combined goals (x) | Number of matches (f) | $x \times f$ |
|------------------------|---------------------------|-------------------|
| 0 | 6 | 0 |
| 1 | 13 | 13 |
| 2 | 12 | 24 |
| 3 | 9 | 27 |
| 4 | 5 | 20 |
| 5 | 2 | 10 |
| 6 | 0 | 0 |
| 7 | 1 | 7 |
| Σf | 48 | $\Sigma xf = 101$ |

3. Divide the result by the sum of the frequencies, Σf .

$$\begin{aligned}\bar{x} &= \frac{\Sigma xf}{\Sigma f} = \frac{\text{sum of (scores} \times \text{frequency)}}{\text{sum of frequencies}} \\ &= \frac{101}{48} \\ &= 2.1\end{aligned}$$

- c. The mode is the score with the highest frequency. c. Mode = 1 combined goal scored


- Mean, median and mode can be calculated from plots.

WORKED EXAMPLE 17 Calculating the median from a plot

Calculate the median of each data set.

- a. Key : 1|8 = 18

Stem	Leaf
1	8 9
2	2 2 5 7 7 8
3	0 1 4 6 7
4	0 5



THINK

- a. 1. Check that the scores are arranged in ascending order.
2. Locate the position of the median using the rule $\frac{n+1}{2}$, where $n = 15$.
Note: The position of the median is $\frac{15+1}{2} = 8$ (the eighth score).
3. Write the answer in a sentence.
- b. 1. Locate the position of the median using the rule $\frac{n+1}{2}$, where $n = 12$.
Note: The position of the median is $\frac{12+1}{2} = 6.5$. This means it is between the sixth and seventh scores.
3. Calculate the average of the two middle scores.
4. Write the answer in a sentence.

WRITE

- a. The scores are arranged in ascending order.

Key : 1|8 = 18

Stem	Leaf
1	8 9
2	2 2 5 7 7 8
3	0 1 4 6 7
4	0 5

The median of the set of data is 18.

- b. The 6th score is 19.
 The 7th score is 19.

$$\text{Median} = \frac{19+19}{2}$$

$$\begin{aligned}&= \frac{38}{2} \\ &= 19\end{aligned}$$

The median of the set of data is 19.

11.5.2 Grouped data

eles-4919

- **Grouped data** is given when a frequency table shows intervals instead of a value.
- To calculate the mean of a grouped data set, multiply the midpoint of each class interval by the corresponding frequency, and then divide the sum of these products by the sum of the frequencies.
- The median of a grouped data set is the middle value when the values are in order.
- The mode of a grouped data set is the most common class interval. This is known as the **modal class**.
For example, let us calculate the mean, median and mode for the following data.

Interval	Frequency (f)
10– < 15	3
15– < 20	8
20– < 25	6
Σf	17

- To calculate the mean, add a column to the data table showing midpoints, and another column showing the product of frequency and midpoint.

Interval	Frequency (f)	Midpoint (x_{mid})	$f \times x_{\text{mid}}$
10 – < 15	3	12.5	$3 \times 12.5 = 37.5$
15 – < 20	8	17.5	$8 \times 17.5 = 140$
20 – < 25	6	22.5	$6 \times 22.5 = 135$
Σf	17	$\Sigma f \times x_{\text{mid}}$	$37.5 + 140 + 135 = 312.5$

- Mean = $\frac{312.5}{17} = 18.38$ (rounded to 2 decimal places)
- To calculate the median: since there are 17 data values, the middle value will be the 9th value, because $\frac{17+1}{2} = 9$.

Interval	Frequency (f)
10 – < 15	3
15 – < 20	8
20 – < 25	6
Σf	17

← First 3 values — need 6 more to get to 9

← The median is the 6th value in this group of 8

- To calculate the median, look at the interval group that contains the median and use the rule:
 - median = $15 + \frac{6}{8} \times 5 = 18.75$
- To calculate the modal class, determine the interval with the highest frequency.
 - Modal class = 15– < 20.

WORKED EXAMPLE 18 Calculating the median, mean and mode from grouped data

The actual diagonal lengths of 45 televisions in a store are presented as a frequency table. Calculate the:

- a. median (to the nearest inch) b. mean (to the nearest inch) c. modal class.

Diagonal length (inches)	Number of TVs (f)
10– < 20	1
20– < 30	3
30– < 40	8
40– < 50	19
50– < 60	9
60– < 70	3
70– < 80	2
Σf	45

THINK

- a. 1. The median is the middle value. The sum of the frequencies is the number of values, n . If there is an odd number of values, then the median is the middle value.
2. Determine where the 23rd value lies in the table by looking at the cumulative frequencies.
- < 20 inches = 1 TV
 - < 30 inches = 4 TVs
 - < 40 inches = 12 TVs
 - < 50 inches = 31 TVs
- The 23rd value is in the 40– < 50 inches group. It is the 11th value in that group of 19 TVs.
- b. 1. The mean is the average value. Set up a third and fourth column in the frequency table.
2. Record the midpoint as the middle value of each interval, for example $\frac{10 + 20}{2} = 15$.
3. Multiply the midpoint of each interval, x_{mid} and the interval's frequency, f .
4. Add up all the products of $x_{\text{mid}} \times f$.

WRITE/DISPLAY

a. Median value = $\frac{n + 1}{2}$ th value

$$= \frac{45 + 1}{2}$$

$$= 23\text{rd value}$$

The 23rd value is in the 40– < 50 inches class. It is the 11th value in a group of 19 values.

$$\frac{11}{19} \times 10 \text{ inches} = 5.79 \text{ inches}$$

$$\text{Median} = 40 + 5.79 \text{ inches}$$

$$= 45.79 \text{ inches}$$

$$\approx 46 \text{ inches}$$

b.

Diagonal length (inches)	Number of TVs (f)	Class midpoint x_{mid} (inches)	Product of x_{mid} and f
10– < 20	1	15	$1 \times 15 = 15$
20– < 30	3	25	$3 \times 25 = 75$
30– < 40	8	35	$8 \times 35 = 280$
40– < 50	19	45	$19 \times 45 = 855$
50– < 60	9	55	$9 \times 55 = 495$
60– < 70	3	65	$3 \times 65 = 195$
70– < 80	2	75	$2 \times 75 = 150$
Σf	= 45	$\Sigma x_{\text{mid}} \times f$	= 2065

5. Divide $x_{\text{mid}} \times f$ by the sum of the frequencies, Σf .

$$\begin{aligned}\bar{x} &= \frac{\Sigma xf}{\Sigma f} = \frac{\text{data values (score} \times \text{frequency)}}{\text{sum of frequencies}} \\ &= \frac{2065}{45} \\ &= 45.89 \approx 46 \text{ inches}\end{aligned}$$

6. Write the answer in a sentence.

The mean TV size is 46 inches.


c. The modal class is the class interval with the highest frequency.

c. The modal class is 40 to less than 50 inches.

11.5.3 Types of distributions



eles-4921

- The type of distribution describes the shape of the data.
- Symmetrical distributions** have a peak in the centre and evenly balanced data above and below the centre.
 - In symmetrical distributions, the mean, median and mode have similar values.




Stem	Leaf
0	7
1	2 3
2	2 4 5 7 9
3	0 2 3 6 8 8
4	4 7 8 9 9
5	2 7 8
6	1 3

- Skewed distributions** are not symmetrical.
 - Positively skewed distributions** (or right-skewed distributions) have a tail towards the positive direction, or right-hand side of the histogram.
 - In positively skewed distributions, the mean has a larger value than the median.
 - Negatively skewed distributions** (or left-skewed distributions) have a tail towards the negative direction, or left-hand side of the histogram.
 - In negatively skewed distributions, the mean has a smaller value than the median.



- **Bimodal distributions** have two peaks (the prefix *bi-* means *two*).




Types of distribution and measures of centre

Type of distribution	Most appropriate measure of centre
Symmetrical	Mean or Median
Skewed	Median


WORKED EXAMPLE 19 Identifying distribution type and calculating measure of centre

For each of the distributions shown, describe the type of distribution, determining its measure of centre using the most appropriate measures.

a. Number of TVs in a household



b. Number of family members



THINK

- a. 1. Describe the shape of the distribution.
2. Select an appropriate measure of centre.
3. Determine the mean.
4. Write the answer in a sentence.

WRITE

- a. The graph is roughly symmetrical.
Measure of centre: mean

$$\begin{aligned} \text{Mean} = \bar{x} &= \frac{\sum x}{n} = \frac{\text{sum of all data values}}{\text{number of data values}} \\ &= \frac{120}{28} \\ &= 4.3 \end{aligned}$$

The mean number of TVs per household is 4.3.

- b. 1. Describe the shape of the distribution.
 2. Select the appropriate measures of centre.
 3. Determine the median by first determining the number of data values.
 $25 + 45 + 30 + 10 + 5 + 1 + 1 + 5 = 121$

4. Write the answer in a sentence.

- b. The graph is positively skewed.
 Measure of centre: median

$$\begin{aligned} \text{Median} &= \frac{n+1}{2} \text{th data value} \\ &= \frac{122+1}{2} \\ &= 61.5 \end{aligned}$$

The median is the average of the 61st and 62nd data values.

$$\begin{aligned} \text{Median} &= \frac{2+2}{2} \\ &= 2 \end{aligned}$$

The median number of members in a family is 2.

WORKED EXAMPLE 20 Determining the measures of centre from a stem plot

Determine the best measures of centre for the heights of 25 people shown in the stem plot. Give your answer to the nearest centimetre.

Key: 17 | 8 = 178 cm

Stem	Leaf
14	8
15	3 6
16	1 4 7
17	0 0 7 8 9
18	1 2 2 4 6 8 8
19	3 4 7 7
20	0 2
21	2

THINK

- Describe the distribution.
- State the most appropriate measure of centre.
- Calculate the mean.

4. Write the answer in a sentence.

WRITE

The distribution is symmetrical.

The most appropriate measure of centre is the mean.




$$\begin{aligned} \text{Mean} = \bar{x} &= \frac{\sum x}{n} = \frac{\text{sum of all data values}}{\text{number of data values}} \\ &= \frac{4509}{25} \\ &= 180.36 \end{aligned}$$

The mean height of the 25 people is 180 cm.

DISCUSSION

It is important to understand what the numbers on an exam report mean, as these are used to make decisions about student learning. A student in a Year 9 exam receives a mean, median and range of scores. In pairs, discuss and write down what each of these measures suggests about the score of an individual student in comparison to the class.

on Resources

-  **eWorkbook** Topic 11 Workbook (worksheets, code puzzle and a project) (ewbk-2011)
-  **Video eLesson** Mean and median (eles-1905)
-  **Interactivities** Individual pathway interactivity: Measures of central tendency (int-4543)
 - Mean (int-3818)
 - Median (int-3819)
 - Mode (int-3820)

Exercise 11.5 Measures of centre

learn **on**

11.5 Quick quiz **on**

11.5 Exercise

Individual pathways

■ PRACTISE

1, 2, 3, 6, 10, 11, 14, 16, 19, 20, 25

■ CONSOLIDATE

4, 7, 8, 12, 15, 17, 21, 22, 26

■ MASTER

5, 9, 13, 18, 23, 24, 27, 28

Fluency

1. **WE15** Calculate the median, mean and mode for the data sets given. Confirm your answers for the median and mean using a graphics calculator. Where necessary, give your answers to 2 decimal places.
 - a. 27, 32, 45, 48, 53, 55, 55, 57, 59, 61, 75, 81
 - b. 8, 8, 2, 8, 8, 8, 8, 8, 9, 9, 1, 9, 5, 9, 7, 10, 0
2. **WE16** Calculate the median, mean (to 2 decimal places) and mode for the ungrouped frequency table shown.

Score	Frequency
0	2
1	5
2	12
3	10
4	7
5	2

3. **WE17** Calculate the median of each of the following data sets.

a. Key: $1|0 = 10$

Stem	Leaf
1	0 2
2	1 3 3 5
3	
4	4

b.




4. Calculate the median of each of the following data sets.

a. Key: $10|0 = 100$

Stem	Leaf
10	0
11	0 2 2 2
12	0 4 6 6
13	3

b.




5. Calculate the median of each of the following data sets.

a. Key: $6.1|8 = 6.18$

Stem	Leaf
6.1	8 8 9
6.2	0 5 6 8
6.3	0 1 2 4 4 4

b.



6. For the stem plot given below, calculate:

- the median
- the mean (to 2 decimal places)
- the mode.

Key: $3^*|7 = 37$

Stem	Leaf
1	3 4
1*	5 6 7 9
2	0 2 3 3 4
2*	5 5 5 6 6 7 8
3	1 2 3 4
3*	5 7
4	1

7. **WE18** For the following grouped frequency tables, calculate the:

- median
- mean
- modal class.

Where necessary, give your answers to 1 decimal place.

a.

Score x	Frequency f
120–129	2
130–139	8
140–149	10
150–159	5
160–169	1

b.

Score	Frequency
120–129	2
130–139	8
140–149	10
150–159	16
160–169	5
170–179	1

Hint: The midpoint of the first class interval is $\frac{120 + 129}{2} = 124.5$.

8. **MC** Select the correct mean, median and mode for the following set of data.

1, 5, 2, 8, 7, 3, 5, 7, 2, 8, 7, 6, 4, 5, 5

- A. 5, 6, 5 B. 6, 3, 5 C. 5, 5, 5 D. 5, 5, 7 E. 6, 5, 6

9. **MC** Select the median number of members per family surveyed as presented in this frequency table.

Number of family members	Frequency
3	19
4	53
5	21
6	4
7	2
8	1







- A. 3 B. 4 C. 4.5 D. 4.6 E. 53

10. **MC** Select the best estimate of the mean score for the grouped frequency table shown.

Score	Frequency
0 – < 50	1
50 – < 100	4
100 – < 150	10
150 – < 200	3
200 – < 250	1
250 – < 300	1

- A. 100 B. 150 C. 125 D. 130 E. 145

11. **WE19** For each of the following histograms, state whether the distribution is symmetrical, positively skewed or negatively skewed.



12. State the type of distribution shown by each of the following stem plots.

a. Key: $14 | 6 = 14.6$

Stem	Leaf
12	0
13	89
14	0899
15	1566678
16	3578
17	02
18	35

b. Key: $14 | 6 = 146$

Stem	Leaf
13	1246789
14	0456899
15	156678
16	378
17	02
18	3


c. Key: $14 | 6 = 1.46$

Stem	Leaf
12	78
13	6789
14	09
15	156
16	37899
17	02
18	3

d. Key: $14 | 6 = 1.46$

Stem	Leaf
1	1
2	
3	69
4	049
5	13568
6	237899
7	046778

13. **WE19** This histogram represents the results of data collected from households in a suburban street. Describe the type of distribution, determining its centre using the most appropriate measure.




14. **WE20** The number of customers served at a cafe each hour for the past 24-hour period is summarised in the stem plot shown.
- State the type of data and type of distribution.
 - Choose the most appropriate measure of centre.
 - Calculate the centre chosen in **b**.

Key: 2 | 4 = 24 customers

Stem	Leaf
0	0014
0*	5588999
1	123444
1*	5557
2	02
2*	
3	4

Understanding

15. Consider the distribution shown.
- State whether the data is symmetrical.
 - State the modal class.
 - Explain whether the mean and median can be seen from the graph. If they can, state their values.



16. The grouped frequency table shown is a summary of 40 golf scores recorded at the local 9-hole golf course. For the data given, determine the:
- type of data
 - median
 - mean
 - modal class.

Golf score	Number of games
30–39	4
40–49	32
50–59	2
60–69	1
70–79	1

17. The frequency table shown is a summary of 100 long-jump attempts by a group of primary school students during a recent school sports carnival. For the data given, determine the:
- type of data
 - median
 - mean
 - mode.

Where necessary, round your answers to the nearest whole number.

Long jump distance (cm)	Number of jumps
120 – < 140	22
140 – < 160	44
160 – < 180	17
180 – < 200	14
200 – < 220	2
220 – < 240	0
240 – < 260	0
260 – < 280	1



18. A survey of the number of people living in each house on a residential street produced the following data:

2, 5, 1, 6, 2, 3, 2, 1, 4, 3, 4, 3, 1, 2, 2, 0, 2, 4

- Prepare a frequency distribution table with an $f \times x$ column and use it to calculate the average (mean) number of people per household. Give your answer to 1 decimal place.
- Draw a dot plot of the data and use it to find the median number of people per household.
- Calculate the modal number of people per household.
- Explain which of the measures would be most useful to:
 - a real-estate agent renting out houses
 - a government population survey
 - an ice-cream truck owner.

Reasoning

19. A Geography teacher with a class of 20 students commented that test results for their class out of a score of 50 were negatively skewed. Show and explain a data set for the possible results obtained by the 20 students.

20. If the mean = 11, mode = 4, and the median = 9 and the missing values are positive integers:

- determine the missing two values in the following set of numbers:
4, ____, 19, ____, 21, 4
- calculate two different values that meet the stated criteria
- explain whether you can substitute 9 for both missing values.

21. A class of 26 students had a median mark of 54 in Mathematics. In reality, though, no-one actually obtained this result.

- Explain how this is possible.
- Explain how many students must have scored below 54.

22. A small business pays the following wages (in thousands of dollars) to its employees:

18, 18, 18, 18, 26, 26, 26, 35, 80 (boss)

- MC** Select the wage earned by most workers at this business.
A. \$29 400 **B.** \$18 000 **C.** \$29 000 **D.** \$26 000 **E.** \$19 000
- Calculate the average wage at this business.
- MC** Select the median of this distribution of wages.
A. \$26 000 **B.** \$26 500 **C.** \$18 000 **D.** \$18 500 **E.** \$26 400
- Explain which measure might be used in wage negotiations by:
 - the union, representing the employees (other than the boss)
 - the boss.

23. a. Create a data set of 8 numbers that meet the following criteria.

- Data values are positive integers
- Mean = 9.5
- Median = 7
- Mode = 6
- Maximum value in data set = 21
- Minimum value in data set = 1

b. Explain whether there is only one data set that can meet this criteria

24. A tyre manufacturer selects 48 tyres at random from the production line for testing. The total distance travelled during the safe life of each tyre is shown in the table.

Distance in km ('000)	82	78	56	52	50	46
Number of tyres	2	4	10	16	12	4

- Calculate the mean, median and mode.
- Explain which measure best describes average tyre life.
- Recalculate the mean with the 6 longest-lasting tyres removed. By how much is it lowered?
Give your answer to the nearest km.
- Determine what tyre life has the highest probability of being selected if you selected a tyre at random.
- In a production run of 10 000 tyres, determine how many could be expected to last for a maximum of 50 000 km.
- As the manufacturer, state the distance for which you would be prepared to guarantee your tyres.
Explain your answer.



Problem solving

25. Answer the following questions.
- Create a set of five values with a mean of 4 and a median of 5.
Demonstrate that your data set meets the given criteria.
 - Change only one of the numbers in your set so that the mean is now 5 and the median remains 5.
Demonstrate that your modified set meets the new criteria.
26. Seven students completed a challenging Science test. The test was scored out of 15. Six of the students got the following marks: 4, 4, 5, 5, 5, 6. Determine the mark that the 7th student obtained if:
- the mean score is the same as the median score
 - the mean score is greater than the median score
 - the mean score is equal to the modal score
 - the mean score is greater than 6.
27. Calculate the mean for this data set.

Score	4	8	12	16	20	24
Frequency	$4a$	$3a$	$2a$	a	$\frac{a}{2}$	$\frac{a}{4}$

28. On a piece of paper is a list of six 2-digit prime numbers. They have a mean and median of 39. Their mode is 31 and the smallest number is 13. What are the six numbers?
Explain how you got your answer.

LESSON

11.6 Measures of spread

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- calculate range and interquartile range
- use statistical measures to analyse data.

11.6.1 Range and interquartile range

eles-4920

- The **range** is the difference between the maximum and minimum values in a data set.
- To calculate interquartile range, the data is first organised into four sections called **quartiles**, each containing 25% of the data.
- The word ‘quartiles’ comes from the word ‘quarter’.
- The **interquartile range (IQR)** is the difference between the first quartile and the third quartile.

Range



- To calculate range, use the following formula:

$$\text{Range} = \text{maximum value} - \text{minimum value}$$

Interquartile range (IQR)

- To calculate interquartile range (IQR), use the following formula:

$$\text{IQR} = Q_3 - Q_1$$

where:

Q_1 is the median of the first half of the data

Q_3 is the median of the second half of the data.

WORKED EXAMPLE 21 Calculating the range and interquartile range of a data set

Calculate the range and interquartile range of the following data set.

5, 6, 8, 4, 12, 5, 7, 8, 4, 10, 9, 6, 5

THINK

1. Arrange the data from minimum to maximum value.
2. Calculate the range.
3. Calculate the median by using the rule:
median = $\frac{n+1}{2}$ th data value.
4. Calculate the first quartile (Q_1) from the 6 values in the lower half of the data set.
5. Calculate the third quartile (Q_3) from the 6 values in the upper half of the data set.
6. Subtract the first quartile from the third quartile to find the interquartile range.

WRITE

4 4 5 5 5 6 6 7 8 8 9 10 12

$$\begin{aligned} \text{Range} &= \text{maximum value} - \text{minimum value} \\ &= 12 - 4 \\ &= 8 \end{aligned}$$

$$\begin{aligned} \text{Median} &= \frac{n+1}{2} \text{th data value} \\ &= \frac{13+1}{2} \\ &= 7 \text{th data value} \end{aligned}$$

Median is the 7th data value.

$$\therefore \text{Median} = 6$$

4 4 5 5 5 6 6 7 8 8 9 10 12

Q_1 is between 3rd and 4th data values

$$Q_1 = 5$$

Q_3 is between 3rd last and 4 last data values.

$$\begin{aligned} Q_3 &= \frac{8+9}{2} \\ &= 8.5 \end{aligned}$$

$$\begin{aligned} \text{IQR} &= Q_3 - Q_1 \\ &= 8.5 - 5 \\ &= 3.5 \end{aligned}$$

TI | THINK

1. In a new problem, open a Lists & Spreadsheet page. Label column A as x values. Enter the data.

WRITE/DISPLAY

A	xvalues	B	C	D
10	10			
11	9			
12	6			
13	5			
14				


CASIO | THINK

1. Open a new Statistics screen. Enter the data into List1.

WRITE/DISPLAY

list 1	list 2	list 3
3		
7		
4		
5		
6		
2		
8		
3		
3		
5		

2. Press:
- MENU
 - 4: Statistics
 - 1: Stat Calculations
 - 1: One-Variable Statistics.
- Set Number of Lists = 1.
Press OK.
Complete the table as shown.
Press ENTER.




The summary statistics will appear as shown. Scroll down to find the measures required.

A	xvalues	B	C	D
=			=OneVar(
8	8	MinX	4.	
9	4	Q ₁ X	5.	
10	10	MedianX...	6.	
11	9	Q ₃ X	8.5	
12	6	MaxX	12.	
13	Σx = 8.5			

$$\begin{aligned} \text{Range} &= \text{MaxX} - \text{MinX} \\ &= 12 - 4 \\ &= 8 \\ \text{IQR} &= \text{Q}_3\text{X} - \text{Q}_1\text{X} \\ &= 8.5 - 5 \\ &= 3.5 \end{aligned}$$

2. Press:
- Calc
 - One-Variable
- Fill in the Set Calculation table as shown:
- Xlist: list1
 - Freq: 1
- Press OK.



The summary statistics will appear as shown. Scroll down to find the measures required.

Stat Calculation	
One-Variable	
XList:	list1
Freq:	1
OK Cancel	
1/	18
Calc	
[11] =	
Deg	Auto Standard

Stat Calculation	
One-Variable	
σ _x	= 2.3483596
s _x	= 2.442502
n	= 13
minX	= 4
Q ₁	= 5
Med	= 6
Q ₃	= 8.5
maxX	= 12
Mode	= 5
ModeN	= 1
OK Cancel	
1/	18
Calc	
[14] =	
Deg	Auto Standard

$$\begin{aligned} \text{Range} &= \text{Max}_x - \text{Min}_x \\ &= 12 - 4 \\ &= 8 \\ \text{IQR} &= \text{Q}_3 - \text{Q}_1 \\ &= 8.5 - 5 \\ &= 3.5 \end{aligned}$$

WORKED EXAMPLE 22 Calculating the range and interquartile range from a plot

For the stem plot shown, determine:

- the range
- the interquartile range.

Key: 4|7 = 4.7 kg

Stem	Leaf
1	9
2	2 7
3	0 1 3 7
4	2 4 5 7 8
5	1 2 8
6	0

THINK

- Calculate the range by finding the difference between the minimum and maximum value.

WRITE

- Range = maximum value – minimum value
= 6.0 – 1.9
= 4.1 kg

b. 1. Count the number of data values, n .

b. $n = 16$

2. Calculate the median by using the rule

$$\text{median} = \frac{n+1}{2} \text{th data value.}$$

$$\text{Median} = \frac{n+1}{2} \text{th data value}$$

$$= \frac{16+1}{2}$$

$$= 8.5 \text{th data value}$$

The median is between the 8th and 9th data values, i.e. between 4.2 and 4.4.

$$\therefore \text{Median} = 4.3 \text{ kg}$$

3. Calculate the first quartile, Q_1 (1st quartile) from the 8 values in the lower half of the data set.

Stem	Leaf
1	9
2	2 7
3	0 1 3 7
4	2 4 5 7 8
5	1 2 8
6	0

Q_1 is between the 4th value (3.0) and the 5th value (3.1).

$$\begin{aligned} \therefore Q_1 &= \frac{3.0 + 3.1}{2} \\ &= 3.05 \text{ kg} \end{aligned}$$

4. Calculate the third quartile, Q_3 (3rd quartile) from the 8 values in the upper half of the data set.

Q_3 is between the 4th last value (5.1) and the 5th last value (4.8), that is 4.95 kg.

$$\begin{aligned} Q_3 &= \frac{4.8 + 5.1}{2} \\ &= 4.95 \text{ kg} \end{aligned}$$

5. Determine the interquartile range by subtracting the first quartile from the third quartile.

$$\begin{aligned} \text{IQR} &= Q_3 - Q_1 \\ &= 4.95 - 3.05 \\ &= 1.9 \text{ kg} \end{aligned}$$

11.6.2 Analysing data

- The summary statistics examined can be used to analyse data. These include:
 - mean
 - median
 - mode
 - range
 - IQR.

WORKED EXAMPLE 23 Determining the statistical measure

State the statistical measure that is referred to in each of the following statements.

- The ages of fans at a Pink concert varied from 12 to 55.
- The majority of students surveyed preferred holidays to school time.

THINK

- Look for any keywords in the statement. The information gives the *highest* and *lowest* ages.
- Look for any keywords in the statement. The information refers to the *majority* of students.

WRITE

- The statement refers to the *range* of ages at the concert.
- Majority implies most, so the statement refers to the *mode*.

WORKED EXAMPLE 24 Analysing statistical data

A weather report delivered the following statement:

We had a large temperature range today: from 12°C to 37°C . This was by far the hottest day we've had this month. In fact the maximum was 14°C above the monthly average. It also happens that 12°C is the most common low temperature for the month.

- Explain what these statistics mean.
- Determine the mean monthly temperature.
- Calculate the range in temperature for this day.



THINK

- Look for the statistical terms, 'range', 'average' and 'most common'.
Explain what these terms mean.
- The average monthly temperature was $37^{\circ}\text{C} - 14^{\circ}\text{C}$.
- The range is the difference between the highest and lowest temperatures: $37^{\circ}\text{C} - 12^{\circ}\text{C}$.

WRITE

- Range is the difference between the highest and lowest temperatures.
The average is the mean.
 12°C was the modal minimum temperature.
- Monthly mean (average) = $37^{\circ}\text{C} - 14^{\circ}\text{C}$
 $= 23^{\circ}\text{C}$
- Range = $37^{\circ}\text{C} - 12^{\circ}\text{C}$
 $= 25^{\circ}\text{C}$

Resources

-  **eWorkbook** Topic 11 Workbook (worksheets, code puzzle and project) (ewbk-2011)
-  **Interactivities** Individual pathway interactivity: Measures of spread (int-4622)
Range (int-3822)
The interquartile range (int-4813)
Skewness (int-3823)

Individual pathways

PRACTISE

1, 4, 5, 7, 10, 13

CONSOLIDATE

3, 6, 8, 11, 14

MASTER

2, 9, 12, 15

Fluency

- WE21** Calculate the range and interquartile range for the following raw data sets.
 - 18, 22, 25, 27, 29, 31, 35, 37, 40, 42, 44
 - 120, 124, 125, 135, 135, 137, 141, 145, 151
- WE22** This stem plot gives the spread of 25 cars caught by a roadside speed camera. Determine:
 - the range
 - the interquartile range.
- For the following raw data sets, calculate the range and interquartile range.
 - 20, 45, 73, 65, 85, 127, 101
 - 1.22, 1.28, 1.48, 1.52, 1.62, 1.72, 1.85, 1.95, 2.03
- For the ungrouped frequency table shown, calculate:
 - the spread — the range and interquartile range
 - the measures of centre — the mode, the median and mean (correct to 2 decimal places).

Key: 8 | 2 = 82 km/h
8* | 6 = 86 km/h

Stem	Leaf
8	2 2 4 4 4 4
8*	5 5 6 6 7 9 9 9
9	0 1 1 2 4
9*	5 6 9
10	0 2
10*	
11	4

Score	Frequency
0	24
1	17
2	12
3	7
4	2
5	1

- What are summary statistics and how can they be used to analyse data?
- WE23** Explain which statistical measure is referred to in each of the following statements.
 - The majority of people surveyed were in favour of a public holiday on Anzac Day.
 - The average Year 8 student in Victoria spends an hour and a half doing homework every night.
 - The ages of people at the AFL grand final varied from 5 to 85.
 - There was a 20 °C variation in the temperature on Thursday.
 - On average, students in Year 8 miss 5.7 days of school over the year.
 - Most often, a pair of Nike runners will cost over \$150.
 - The average income for Australian workers is \$675 per week.
 - A middle income family earns about \$45 000 per year.

Understanding

7. **WE24** Read the following paragraph and explain what statistics are represented and what they mean.
It's been an exciting day at the races today. There was a record fast time of 38 seconds, and also a record slow time of 4 minutes and 52 seconds. We had an unbelievable number of people who ran the race in exactly 1 minute. Despite this, the average time was well over 2 minutes, due to the injury of a few key players.
8. Daily temperatures (in °C) were collected in Mildura over 3 weeks. The results are shown on the photo of Mildura at right.
- Analyse these temperatures using summary statistics.
 - Write a paragraph describing the weather in Mildura over these 3 weeks.



9. The following photo shows the number of goals a soccer team kicked in each game over a year.



Prepare a set of summary statistics and use them to write a paragraph describing the soccer team's performance over the year.

Reasoning

10. A classmate made the following statements. Is she correct? Help her with anything she may be confused about.
- The mean and the median try to predict the average of a data set.
 - The range of a data set is the mean absolute difference minus the mean.
 - The mode is the most common number.
11. Show a data set for which the range and interquartile range are the same. Explain what factors cause this.
12. Explain the effect on the range of a data set when a set of scores has:
- each score increased by 5
 - each score doubled
 - each score increased by 25%.

Problem solving

13. Write a story to represent the following statistics: mean = 45.6, median = 34, mode = 24 and range = 58.
14. This back-to-back stem-and-leaf plot displays the speeds of 25 cars caught by a speed camera on a major highway on Easter Sunday and Christmas Day.

Key: 6 | 3 = 63 km/h

<i>Easter Sunday</i>		<i>Christmas Day</i>
8 6 3	6	2 3 6 9 9
8 7 4 1	7	1 2 2 4 5 5 7 8
6 6 5 4 2 1	8	0 0 3 4 6 7
9 8 7 7 5 2 1	9	1 2 3 3
7 7 5 2 1	10	2 4

The police commissioner was heard to comment that he was surprised to find drivers behaved better on Christmas Day than they did on Easter Sunday.

- The mean speed for Easter Sunday was 87.68 km/h and the mean speed for Christmas Day was 79.92 km/h. Explain what this tells us about the drivers on these different days.
 - The interquartile range on Easter Sunday was 21 and on Christmas Day it was 17.5. Explain what this tells us about drivers on these different days.
 - Do your results from part **a** and **b** support the police commissioner's statement? Explain your answer.
15. Waterway Secondary has one junior and one senior rowing team. Each team has eight rowers and one cox (the cox steers the boat and keeps the rhythm, but doesn't row). The heights of all the rowing team members in centimetres is shown in the following stem-and-leaf plot.

Key: 16 | 4 = 164

<i>Stem</i>	<i>Leaf</i>
13	5 9
14	
15	9
16	2 4 5 5 8
17	0 1
18	0 1 2 2 4 5
19	0 4



- a. This data set has two outliers. Identify these outliers and state their value.
- b. Excluding the two outliers, calculate the mean of the data set.
- c. Excluding the two outliers, identify the data set's distribution type.
- d. Assuming that the junior rowers are shorter than the senior rowers, separate the data into junior team and senior team heights. Present the data in two stem-and-leaf plots.
- e. Calculate the mean height of the:
 - i. junior team
 - ii. senior team.
- f. Using your answers to **b** and **e**, explain why it is better to separate bimodal data before finding the mean.

COLLABORATIVE TASK: Surveys in the media

At times, reports of surveys in the media can be used to lead people to draw conclusions by representing graphs and statistics in a way that supports a certain point of view. Find a statistical report in the media and analyse by linking claims made in the report to displays, statistics and sampling methods.



LESSON

11.7 Review

11.7.1 Topic summary

Types of data

- Categorical data is expressed using labelled categories.
 - Nominal data cannot be ranked in a logical order; e.g. favourite colours, pet types.
 - Ordinal data can be ranked in a logical order; e.g. AFL players' salaries.
- Numerical data is expressed using numbers.
 - Discrete data is collected by counting; e.g. number of pens, number of days.
 - Continuous data is collected by measuring; e.g. the mass of eggs, the distance between school and each student's home.

Size of data set

- Population:** Data collected from an entire group; e.g. when all members of a family are surveyed.
- Sample:** Data collected from only parts of a group; e.g. when only Year 9 students are surveyed at a school.

Random number generator

- To create random data on a spreadsheet, use the command =RANDBETWEEN()
- Then enter the lowest and highest desired value within the brackets, separated by a comma.

STATISTICS

Displaying data

- Methods of displaying data include:
 - bar charts

Measure of centre

- The **mean** is also known as the average.
 - For raw data: $\bar{x} = \frac{\sum x}{n}$ = $\frac{\text{sum of all data values}}{\text{number of data values}}$
 - For data in a frequency table: $\bar{x} = \frac{\sum xf}{\sum f}$ = $\frac{\text{sum of (data values} \times \text{frequency)}}{\text{sum of frequencies}}$
- The **median** is the middle value of the data. $\frac{n+1}{2}$ th data value, where n is the number of pieces of data or scores.
- The **mode** is the most common value in the data.

Collecting data

- Methods of collecting data include:
 - frequency tables

Person's age (x)	Tally	Number of people (f)
0–29		5
30–59		10
60–89		3
		$\Sigma f = 18$

- dot plots

- stem-and-leaf plots.

Heights of 30 students (in cm)
Key: 11|8 = 118 cm

Stem	Leaf
11	8 9
12	0 1 2 2 3 4 5 6 7 7 8 9
13	0 1 1 1 2 3 4 5 6 8 8 9 9
14	0 1 3

- pie charts

Transport to school

Spread of data

- Range = maximum value – minimum value
- Interquartile range (IQR) = $Q_3 - Q_1$

- histograms

Mass of dolphin

Types of distribution




Types of distribution include:

- symmetrical distribution, which has a peak in the centre and evenly balanced data above and below the centre.
- skewed data, which is not evenly balanced.
 - For positively skewed data, the mean has a larger value than the median.
 - For negatively skewed data, the mean has a smaller value than the median.
- bimodal data, which has a distribution pattern with two peaks.

11.7.2 Success criteria

Tick the column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
11.2	I can identify different types of data.			
	I can determine if a data set is a sample or a population.			
	I can generate random numbers.			
11.3	I can organise data in frequency tables.			
	I can display data in dot plots.			
	I can display data in stem-and-leaf plots.			
11.4	I can display data in bar charts, pie charts and histograms.			
	I can present data as a back-to-back stem-and-leaf plots.			
11.5	I can calculate the mean, median and mode of raw data.			
	I can calculate the mean, median and mode of data in tables and plots.			
	I can calculate the mean, median and mode of grouped data.			
11.6	I can calculate range and interquartile range.			
	I can determine whether a distribution is symmetrical, skewed or bimodal.			

11.7.3 Project

Graphical displays of data

Statistical data can be communicated using words, tables or graphs.


When data is presented in words, it can be difficult to identify and compare trends.

When data is presented in table form, additional calculations are often needed to gain a complete understanding.

When data is presented as a graph it has an immediate visual impact, but the impression that graphs give can often be misleading, particularly if the graph is not looked at carefully. It is easy to manipulate graphs to produce a desired effect.



Consider the two graphs shown.



These graphs both display the same data (the value of a company's shares over the same period of time), but they look quite different. If the objective of the graph is to create the impression that the value of the shares has dropped quite dramatically, the first graph would serve this purpose. On the other hand, if we are to be convinced that the shares have not changed very much in value, the second graph reflects this point of view.

1. What technique is used to create two quite opposing impressions in the two graphs?



The following table shows the times for the men's and women's 100-m sprints in the Summer Olympic Games. A sports journalist wants to convince readers that, since 1928, women have been more successful in reducing their time than men.

Year	100 m men	Men's time (seconds)	100 m women	Women's time (seconds)
1928	Percy Williams, CAN	10.80	Betty Robinson, USA	12.20
1932	Eddie Tolan, USA	10.30	Stella Walsh, POL	11.90
1936	Jesse Owens, USA	10.30	Helen Stephens, USA	11.50
1948	Harrison Dillard, USA	10.30	Fanny Blankers-Koen, HOL	11.90
1952	Lindy Remigino, USA	10.40	Marjorie Jackson, AUS	11.50
1956	Bobby Morrow, USA	10.50	Betty Cuthbert, AUS	11.50
1960	Armin Hary, GER	10.20	Wilma Rudolph, USA	11.00
1964	Bob Hayes, USA	10.00	Wyomia Tyus, USA	11.40
1968	Jim Hines, USA	9.95	Wyomia Tyus, USA	11.08
1972	Valery Borzov, URS	10.14	Renate Stecher, GDR	11.07
1976	Hasely Crawford, TRI	10.06	Annegret Richter, FRG	11.08
1980	Allan Wells, GBR	10.25	Lyudmila Kondratyeva, URS	11.06
1984	Carl Lewis, USA	9.99	Evelyn Ashford, USA	10.97
1988	Carl Lewis, USA	9.92	Florence Griffith Joyner, USA	10.54
1992	Linford Christie, GBR	9.96	Gail Devers, USA	10.82
1996	Donovan Bailey, CAN	9.84	Gail Devers, USA	10.94
2000	Maurice Greene, USA	9.87	Marion Jones, USA	10.75
2004	Justin Gatlin, USA	9.85	Yuliya Nesterenko, BLR	10.93
2008	Usain Bolt, JAM	9.69	Shelley-Ann Fraser, JAM	10.78
2012	Usain Bolt, JAM	9.63	Shelly-Ann Fraser-Pryce, JAM	10.75
2016	Usain Bolt, JAM	9.81	Elaine Thompson, JAM	10.71

Note: The Summer Olympic Games were not held in 1940 and 1944.

2. Using graph paper, draw two graphs to support the sports journalist's view.
3. Describe the features of your graphs that helped you to present the journalist's view effectively.
4. Imagine you are the sports journalist and you are about to publish the graphs in the local newspaper. Write a short article to accompany the graphs.

on Resources

-  **eWorkbook** Topic 11 Workbook (worksheets, code puzzle and a project) (ewbk-2011)
-  **Interactivities** Crossword (int-2715)
Sudoku puzzle (int-3213)

Exercise 11.7 Review questions

learn on

Fluency

1. **MC** Recognise which one of the following is an example of numerical discrete data.
 - A. Your favourite weeknight television show
 - B. The speed of a car recorded on a speed camera
 - C. The number of home runs by a baseball player
 - D. Placing in a field of eight swimmers of a 100 m freestyle race
 - E. The heights of different animals at a zoo

The following information relates to questions 2 and 3.

Number of bedrooms	1	2	3	4	5	6
Number of homes	4	9	32	36	7	2

2. **MC** For the data given, select the most likely type of distribution.
 - A. A negatively skewed distribution
 - B. A positively skewed distribution
 - C. Skewed to the left
 - D. A symmetrical distribution
 - E. A bimodal distribution
3. **MC** Select the mode for the data collected.
 - A. 36
 - B. 5
 - C. 4
 - D. 3
 - E. 90
4. **MC** The results of a Science test marked out of 60 are represented by the stem plot shown. The median, mean and mode, respectively, are:
 - A. 60, 42, 52
 - B. 40, 45, 60
 - C. 45, 60, 45
 - D. 45, 44, 60
 - E. 42, 44, 45

Key: 2 | 3 = 23

Stem	Leaf
0	8
1	8
2	3 5
3	0 3 6
4	2 4 5 5 5 8
5	4 7 8 8 9
6	0 0 0 0

5. **MC** Select the best estimate of the mean score for the grouped frequency table shown.

Score	Frequency
0 – < 50	12
50 – < 100	24
100 – < 150	23
150 – < 200	4

- A. 63
B. 90
C. 92
D. 100
E. 85
6. **MC** Identify the range and IQR for the following set of data: 46, 46, 49, 53, 61, 63, 67, 72, 81, 84, 93.
A. 46 and 34 B. 46 and 65 C. 47 and 63 D. 47 and 32 E. 93 and 63
7. For the following surveys or studies, decide what type of data would be collected and how it should be displayed.
- The blood type of all workers at a shoe factory.
 - A survey of 500 Australians' views on bringing back corporal punishment (strongly agree, agree, disagree, strongly disagree).
 - Number of phones in the households of the students in your class.
 - The top 20 fastest recorded times for the women's 100 metres butterfly.
8. Classify each of the following data collections according to whether they are samples or populations.
- A survey of 1000 travellers on the quality of train service provided.
 - A survey of all the students at a secondary college about the state government's rules on underage alcohol laws.
 - Performance in Mathematics exams by this year's Year 9 cohort.
9. The following set of data shows the types of phones owned by the first 30 shoppers surveyed.
iPhone, Samsung, Motorola, iPhone, iPhone, iPhone, Samsung, Samsung, Nokia, Motorola, Samsung, iPhone, Motorola, iPhone, Samsung, Samsung, Motorola, iPhone, Samsung, Nokia, Samsung, Motorola, iPhone, iPhone, iPhone, iPhone, Samsung, Samsung, Nokia, Motorola
- Suggest two appropriate ways to display this data.
 - Present this data in the ways suggested in part a.
10. The following data represent the number of songs on people's favourite Spotify playlists. An initial analysis of 25 playlists resulted in the following data being collected.
- 47, 103, 45, 111, 67, 54, 59, 44, 31, 89, 34, 2, 43, 96, 70, 90, 66, 48, 67, 82, 74, 56, 35, 9, 77
- Present the data as a stem-and-leaf plot.
 - Comment on the shape of distribution.
 - Present the data as a frequency table grouped in intervals of 20.
 - State the type of data and type of distribution, and give the most appropriate measure of centre.

11. The following table shows the salaries for AFL players in 2019.

AFL players' salaries	Number of players
0–\$100 000	146
\$100 001–\$200 000	188
\$200 001–\$300 000	107
\$300 001–\$400 000	57
\$400 001–\$500 000	24
\$500 001–\$600 000	12
\$600 001–\$700 000	4
\$700 001–\$800 000	0
\$800 001–\$900 000	4
Total number of players	542

- Draw a histogram to display the data.
- Describe the type of distribution the histogram displays.
- Calculate the median salary of these football players (to the nearest thousand).
- Calculate the modal salary for AFL footballers in 2019.

Problem solving

12. The workers in an office are trying to obtain a salary rise. In the previous year, all ten people who work in the office each received a 2% salary rise, while the company CEO received a 42% salary rise.
- Calculate the mean salary rise received last year. Give your answer to 1 decimal place.
 - Calculate the median percentage salary rise received last year.
 - Calculate the modal percentage salary rise received last year.
 - The company is trying to avoid paying the salary rise. Explain the statistics you think they would quote about last year's rise. Give your reason for using these statistics.
 - Explain the statistics that you think the trade union would quote about salary rises. Give your reason for using these statistics.
 - Identify which statistic you think is the most 'honest' reflection of last year's salary rises. Explain your reasoning.
13. A data set with the values 8, ____, 25, ____, 21, 8 has a mean of 15, mode of 8 and a median of 14. If the missing values are positive integers:
- calculate two values for the set
 - calculate two different values that meet the given criteria.
14. Create a data set of 9 numbers that meet the following criteria.
- Data values are positive integers
 - Mean = 8
 - Median = 7
 - Mode = 10
 - Maximum value in data set = 14
 - Minimum value in data set = 3



To test your understanding and knowledge of this topic go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 11 Statistics

11.1 Pre-test

- a.** iii **b.** iv **c.** i **d.** ii
- Sample
- 16 male adult, 14 female adult, 11 male adolescent, 9 female adolescent
- a.** 25 **b.** 7
- a.** 20 **b.** 10
- a.** ii **b.** iii **c.** i
- B
- a.** Discrete **b.** 17 **c.** Skewed
- B
- C
- 2.5
- a.** 60.3 **b.** D
- 7
- D
- E

11.2 Collecting data

- a.** Categorical **b.** Numerical
- a.** Numerical **b.** Numerical
- a.** Continuous **b.** Continuous
- a.** Continuous **b.** Categorical
- a.** Ordinal **b.** Nominal
- a.** Ordinal **b.** Ordinal
- a.** Nominal **b.** Nominal
- a.** Sample **b.** Sample **c.** Population
- a.** Sample **b.** Population
- a.** data **b.** survey, census **c.** graph
- a.** RANDBETWEEN(1, 31)
b. RANDBETWEEN(1, class size)
c. Samples
- a.** RANDBETWEEN(1, 365)
b. Sample
- a.** Car
b. Sample responses can be found in the worked solutions in the online resources.
c. Sample responses can be found in the worked solutions in the online resources. Answers could include the lower number of motor cycle fatalities as compared to the number of car fatalities.
d. According to the data, there were 190 motorcycle fatalities in 2020, compared to 732 car fatalities.
e. Sample responses can be found in the worked solutions in the online resources.
- Sample responses can be found in the worked solutions in the online resources. Answers will vary but should include the following groups.

Categorical data can be:

 - nominal — arranged in categories, for example by colour or size
 - ordinal — arranged in categories that have an order, for example, karate ability, which ranges from white belt (beginner) to black belt (accomplished).

Numerical data can be:

 - discrete — data can have only particular values, for example amounts of money or shoe sizes
 - continuous — data can have any value within a range, for example the time it takes a person to run around an oval.
- a.** Sample responses can be found in the worked solutions in the online resources. Answers could include:
Data that might be collected from a maternity hospital:
 - birth weights of babies
 - length of time in labour.**b.** Sample responses can be found in the worked solutions in the online resources. Answers could include:
Data that might be collected from an airport:
 - number of delays
 - time spent waiting for flights.**c.** Sample responses can be found in the worked solutions in the online resources. Answers could include:
Data that might be collected from shoppers in a supermarket:
 - shoppers' favourite brands of detergent
 - the amount of money spent on an average weekly shop.
- a.** Sample responses can be found in the worked solutions in the online resources. Answers could include:
'What do you think of sport?' — This question should not be used because it is too broad.
b. Sample responses can be found in the worked solutions in the online resources. Answers could include:
'List these sports in order from your most favourite to your least favourite: football, tennis, netball, golf.'
c. Sample responses can be found in the worked solutions in the online resources.
d. Sample responses can be found in the worked solutions in the online resources. Answers could include:
'Why do you like sports?' — This question should not be used because it is too broad.
e. Sample responses can be found in the worked solutions in the online resources.
f. Sample responses can be found in the worked solutions in the online resources. Answers could include
'Can you play a sport?' — This question should not be used because it is too broad. It would be better to compile a list of sports and ask about those specifically.
g. Sample responses can be found in the worked solutions in the online resources.
'Which of these sports do you watch on the television? Football, tennis, netball, golf, all of these, none of these.' — This is a good question, though it must be noted that results may vary depending on the selection of the sample population.
- a.** Sample responses can be found in the worked solutions in the online resources.


- b. Sample responses can be found in the worked solutions in the online resources. Answers could include: Data can be collected manually, or by using technology to capture data as specific events occur. For example, a computer system might be set up to automatically record the number of sales made per month.
- c. Sample responses can be found in the worked solutions in the online resources. Answers could include: Data is used for forward planning and budgeting purposes. For example, it can be used to identify the most or least popular items that the company sells; this data can then be used to see which items require updating or which items should no longer be sold.
18. a. A survey would be most appropriate as it would require fewer participants, take less time and cost less to survey.
- b. Survey respondents should be selected by simple random sampling, which could include digital technologies to assist in random number generation.
19. a. Sample responses can be found in the worked solutions in the online resources.
- b. Sample response: We are trying to find out whether the population approves or disapproves of the desalination plant and we are also trying to capture data about their reasons.
- c. Sample response: You are collecting categorical data in both instances.
- d. Sample responses can be found in the worked solutions in the online resources.
- e. Sample responses can be found in the worked solutions in the online resources.
20. This inference is not reasonable as the survey was not a simple random sample and only included shoppers who were at the supermarket in the evenings. This could have created biased results.

11.3 Organising data

1. a. Frequency of Maths test results

Score x	Frequency f
1	1
2	1
3	4
4	3
5	5
6	6
7	5
8	2
9	3


b. Multiple choice Maths test results



2. Frequency of type of pet

a.

Type of pet x	Frequency f
Cat	6
Dog	6
Fish	1
Bird	2
Turtle	3



- c. No, only numeric data can be presented in a stem plot.

3. a.

Deaths by motor vehicle accidents (rates per 100 000 people)	Frequency f
0 – < 10	2
10 – < 20	11
20 – < 30	10
30 – < 40	3
40 – < 50	4
50 – < 60	0
60 – < 70	1
70 – < 80	1
	$\Sigma f = 32$

- b. Deaths by motor vehicle accidents (rates per 100 000 people)


Key: 7 | 3 = 73 deaths per 100 000 people

Stem	Leaf
0	09
1	1 2 3 4 4 4 5 7 8 8 9
2	1 2 2 3 4 5 5 6 6 7
3	4 6 6
4	2 2 3 5
5	
6	0
7	1

4. a. Frequency of eye colour in a sample of 40 people.

Score x	Frequency f
Brown	17
Blue	11
Hazel	7
Green	5
	$f = 40$

b. Eye colour in a sample of 40 people



c. 17 out of 40 people (or 42.5% of the people) have brown eyes.

5. a. Key: 1 | 8 = 18

Stem	Leaf
1	8 9 9
2	0 0 0 1 1 2

b. Key: 1 | 9 = 19

Stem	Leaf
1	9 9
2	0 4 4 5 6 7 9
3	1 1 6 8
4	2 3

c. Key: 33 | 6 = 336

Stem	Leaf
33	6
34	6 6 9
35	0 3 9
36	2 8
37	1

6. a. Key: 1* | 8 = 18 2 | 0 = 20

Stem	Leaf
1*	8 9 9
2	0 0 0 1 1 2

b. Key: 1* | 9 = 19 2 | 0 = 20

Stem	Leaf
1*	9 9
2	0 4 4
2*	5 6 7 9
3	1 1
3*	6 8
4	2 3

c. Key: 33* | 6 = 336 35 | 0 = 350

Stem	Leaf
33*	6
34	
34*	6 6 9
35	0 3
35*	9
36	2
36*	8
37	1

7. a. Running times for 40 students

Key: 4 | 3 = 43 minutes, 4* | 8 = 48 minutes

Stem	Leaf
2	1 3
2*	5 6 6 8 9
3	1 2 2 3 3 3 4
3*	7 7 9 9
4	0 1 1 2 2 3 4
4*	5 5 7 8 8 9
5	2 2 3
5*	5 6 7
6	0 3
6*	
7	1

b.

Run times (minutes) x	Number of runners f
20 – < 25	2
25 – < 30	5
30 – < 35	7
35 – < 40	4
40 – < 45	7
45 – < 50	6
50 – < 55	3
55 – < 60	3
60 – < 65	2
65 – < 70	0
70 – < 75	1
$\Sigma f =$	40

c. i. 32 min

ii. 30 min – < 35 min

d. 3, stem plot

8. Key: $1 \mid 4 = 141 \mid 7 = 17$

Stem	Leaf
1	4
1*	7 8 8 9
2	0 1 3
2*	5 6 7 9
3	2 4
3*	5 6 7 8
4	2
4*	5
5	3

9. D

10. a.

Life expectancy (years) x	Number of age groups f
83– < 84	1
84– < 85	21
85– < 86	7
86– < 87	5
87– < 88	2
88– < 89	4
89– < 90	0
$\Sigma f =$	40

b. Life expectancy of Australians across 40 different age groups.

Key: $86 \mid 1 = 86.1$ years

Stem	Leaf
83	7
84	1 1 1 4 4 5 5 5 5 5 5 6 7 7 7 7 8 8 9 9
85	0 0 1 2 7 8 9
86	0 5 6 6 9
87	3 6
88	3 3 8 8
89	

11. Median age of Australian females and males from 2002 to 2009


Key: $36 \mid 1 = 36.1$ years

Stem	Leaf
35	2 3
35*	5 7 9
36	1 1 1
36*	6 9
37	1 3 4
37*	6 6 7

Note: The alternative method — showing categories for 35, 36 and 37 years — is not appropriate because it only allows for 3 groups, which is below the 5 minimum groups allowed.

Score x	Frequency f
35– < 36	5
36– < 37	5
37– < 38	6
	16

12. a. Dots are not aligned vertically. The horizontal scale is not equally spaced.



13. a. Key: $15 \mid 4 = 154$ cm

Leaf (boys)	Stem	Leaf (girls)
	14	5 8
9 8 6 6 4	15	2 2 5 6
8 4 2 0	16	0 1 1 3 5 8
2 1	17	
0	18	

b. The data could be grouped into smaller intervals, such as 5, instead of 10. For example, the stem value of 2 could be broken into two groups, encompassing values 20–24 and 25–29.

14. a. Class 1 : 25, class 2 : 27

b. Class 1 : 2, class 2 : 3

c. Class 1 : 24, class 2 : 20

d. A total of 11 students in class 2 scored better than class 1, but 6 students scored worse than class 1. Class 2's results are more widely spread than class 1's results.

15. a. Key: $2 \mid 1 = 21$ km/h

Stem	Leaf
1	5
2	0 0 2 7 9 9
3	0 0 0 0 2 4 5 6 7 8 8 9
4	0 0 0 0 0 2 4 5 5 5
5	
6	0

b. Sample responses can be found in the worked solutions in the online resources. A possible answer could include that there were a high number of speeds ending in 0 (possibly indicating a machine malfunction), and most of the speeds (24 out of 30) were within the legal speed limit.

16. a. Key: $25 \mid 1 = 25.1$ g

Stem	Leaf
23	9 9
24	0 5 5 6 8
25	1 1 1 3 4 4 8
26	3 3 7
27	3 3
28	1
29	5
30	
31	
32	
33	2


- b. The range is $33.2 - 23.9 = 9.3$.
- c. Sample responses can be found in the worked solutions in the online resources. Answers could mention that the plot for rounded values provides less detail.

Key: $2 \mid 4 = 24 \text{ g}$

$2^* \mid 6 = 26 \text{ g}$

Stem	Leaf
2	4 4 4
2*	5 5 5 5 5 5 5 5 5 6 6 6 7 7 7 8
3	0 3

17. a. Sample responses can be found in the worked solutions in the online resources. Answers could mention that a doughnut graph and a pie graph are both circular, and both use colour to emphasise different categories.
- b. Draw circles of different size with the same centre. Then use the same instructions as for creating a pie chart, but mark only the lines between the two circles.
- c. **Methods of travel to school**




- d. They are the same graph.
- e. Sample responses can be found in the worked solutions in the online resources. Answers could include that a disadvantage of using a doughnut graph could be that you are restricted in the number of circles you can use, so this type of graph would not be suitable for large data sets.

18. 1296

11.4 Displaying data

1. a. See the figure at the foot of the page.*

b. **Pie chart of gym class participation**




2. a. Continuous
b. 24

c.


Class interval	Midpoint (class centre)	Tally	Frequency
5– < 10	7.5		11
10– < 15	12.5		19
15– < 20	17.5		5
20– < 25	22.5		2
25– < 30	27.5		3
		Total	40

d. **Hours of phone use per week**



*1. a.

Gym class participation



3. a.

Class interval	Midpoint (class centre)	Tally	Frequency
0 – 9	4.5		6
10 – 19	14.5		7
20 – 29	24.5		8
30 – 39	34.5		5
40 – 49	44.5		4
50 – 59	54.5		12
60 – 69	64.5		8
70 – 79	74.5		4
80 – 89	84.5		2
		Total	56

b. See the figure at the foot of the page.*


4. a.

Class interval	Midpoint (class centre)	Tally	Frequency
12 – 15	13.5		4
16 – 19	17.5		7
20 – 23	21.5		11
24 – 27	25.5		13
28 – 31	29.5		4
32 – 35	33.5	—	0
36 – 39	37.5		1
		Total	40

b. See the figure at the foot of the page.*


*3. b.

Number of phone calls made per week




*4. b.


Class sizes at Jacaranda SC



5. Number of bananas sold at recess




6. Distances travelled




7. D

8. Number of passengers boarding a bus




9. a. Bar chart of DVD genres (20 DVDs)



b.

Genre	Number of DVDs	Size of sector for a pie chart
Musical	52	93.6°
Drama	98	176.4°
Comedy	30	54°
Horror	10	18°
Action	5	9°
Romance	5	9°
Total	200	360°

c. DVD genres (sample size 200)




10. a. Continuous

b.

Class interval	Midpoint (class centre)	Tally	Frequency
145 – < 150	147.5		2
150 – < 155	152.5		5
155 – < 160	157.5		8
160 – < 165	162.5		7
165 – < 170	167.5		6
170 – < 175	172.5		3
175 – < 180	177.5		1
		Total	32

c. Heights of Year 9 students



- d. No. Individual scores are lost when data are grouped into class intervals.
 e. C
 f. D

g.

Class interval	Midpoint (class centre)	Tally	Frequency
145– < 148	146.5		1
148– < 151	149.5		2
151– < 154	152.5		3
154– < 157	155.5		4
157– < 160	158.5	/	5
160– < 163	161.5		4
163– < 166	164.5		3
166– < 169	167.5	/	6
169– < 172	170.5		2
172– < 175	173.5		1
175– < 178	176.5	—	0
178– < 181	179.5		1
		Total	32

- h. See the figure at the foot of the page.*
 Advantages: resolution is better — more details gained.
 Disadvantages: requires more space to display.
11. a. Number of players b. 5
12. Horizontal axis uses same division for 5 and 7 year periods.
13. a. It is a circle viewed on an angle to produce an ellipse.
 b. No, because it causes some angles to be larger and others to be smaller.
14. a. C
 b. D
 c. In general the boys are slightly taller than the girls, with the tallest 3 students all being boys.

15. a. Key: 0|0 = 0

Leaf (Mitch)	Stem	Leaf (Yani)
3 2 1 1 1 1 0 0 0 0	0	0 0 1 1 1 1 1 2 2 2

Mitch scored between 0 and 3 goals inclusive.
 Yani scored between 0 and 2 goals inclusive.


- b. Mitch: 2, Yani: 3
 c. Mitch
 d. Yani: 11
 e. Yani, because he scored goals in more games.

16. a. Key: 3|2 = 32


Leaf (9A)	Stem	Leaf (9B)
2	3	
9	4	6
7 4	5	5
8 7 5 5 0	6	0 2 8
3	7	3 3 4 5 7 8
5	8	1
6	9	

- b. 9A: 83%, 9B: 92%
 c. 9A: 2
 d. Sample responses can be found in the worked solutions in the online resources.
 e. Sample responses can be found in the worked solutions in the online resources.
 f. 9A: 64.25%, 9B: 68.5%
17. Sample responses can be found in the worked solutions in the online resources.
18. a. Categorical
 b. A column graph or a pie chart
19. The average amount of times per week people used the sunscreen was 5.6. The minimum amount of times was 2 and the maximum amount was 21, although this value was much higher than the rest of the data. 75% of the people surveyed used the sunscreen between 4 and 7 times per week.
20. a. Floribunda: 36°, Mister Lincon: 72°, Pillar: 108°, Hybrid tea: 144°
 b. 10

*10. h.



c.



11.5 Measures of centre

- 55, 54, 55
 - 8, 6.88, 8
- 2.5, 2.55, 2
- 23
 - 5
- 116
 - 82
- 6.28
 - 2.55
- 25
 - 25.24
 - 25
- 143.5, 142.6, 140–149
 - 150.9, 148.5, 150–159
- C
- B
- D
- Symmetrical
 - Symmetrical
 - Symmetrical
 - Negatively skewed
 - Negatively skewed
 - Positively skewed
- Symmetrical
 - Positively skewed
 - Bimodal
 - Negatively skewed with an outlier
- The data is positively skewed with an outlier. This indicates that the mean will not be a suitable measure of centre. The median, with a value of 2, is an appropriate measure of centre because it is not affected by the presence of an outlier.
- Numerical discrete data with a positively skewed distribution
 - Median is the best measure of centre for a skewed distribution.
 - Median = 11.5 customers
- No
 - 31 – 40
 - The mean and median cannot be seen directly from the graph itself. You would need to calculate the mean and the median using a frequency distribution table, which can be constructed from the distribution shown.
- Numerical discrete
 - 45.2
 - 45.25
 - 40–49

17. a. Numerical continuous


- 153
- 157
- 140 – < 160

18. a.

Score (x)	Frequency (f)	Freq \times score ($f \times x$)
0	1	0
1	3	3
2	6	12
3	3	9
4	3	12
5	1	5
6	1	6
	$n = 18$	$\Sigma fx = 47$

$$\bar{x} = 2.6$$

b.



c. 2

d. i. Median ii. Mean iii. Mode

- Possible answers could include that for data to be negatively skewed, the data needs to be a set with few low scores spread over a large range to the left. The most common scores are at the right. Assuming that the horizontal axis has test results in increasing order along the horizontal axis, it could be assumed that many of his students achieved good results.
- Sample responses can be found in the worked solutions in the online resources. Possible answers include 7, 11 and 8, 10.
 - Sample responses can be found in the worked solutions in the online resources. Possible answers include 7, 11 and 8, 10.
 - No, 4 is the mode.
- The median was calculated by taking the average of the 2 middle scores.
 - 13
- B
 - \$29 444
 - A
 - Sample responses can be found in the worked solutions in the online resources.
- Sample responses can be found in the worked solutions in the online resources.
 - No
- 55 250 km, 52 000 km, 52 000 km
 - Discuss in class.
 - 51 810 km (it is reduced by 3440 km)
 - 52 000 km

- e. 3333
 f. 50 000 km (92% of tyres last that distance or more)
25. Sample responses can be found in the worked solutions in the online resources. Possible answers include:
- a. $1, 2, 5, 6, 6 \frac{1+2+5+6+6}{5} = \frac{20}{5} = 4$
 b. $1, 2, 5, 6, 11 \frac{1+2+5+6+11}{5} = \frac{25}{5} = 5$
26. a. 6 b. > 6 c. 6 d. > 13
27. 8.93
28. 13, 31, 31, 47, 53 and 59

11.6 Measures of spread

1. a. 26, 15 b. 31, 18.5
2. a. 32 km/h b. 10 km/h
3. a. 107, 56 b. 0.81, 0.52
4. a. 5, 2 b. 0, 1, 1.19
5. Summary statistics measure the spread and centre of a set of data. They can be used to compare two or more sets of data.
6. a. Mode b. Mean c. Range
 d. Range e. Mean f. Mode
 g. Mean h. Median
7. The range was 4 minutes 14 seconds (fastest time of 38 seconds, slowest time of 4 minutes 52 seconds). The most common time (mode) was 1 minute. The average (mean) time was more than 2 minutes.
8. a. Mean = 34.9; median = 35.4; mode = 36.2; range = 8.5; IQR = 4.5
 b. The weather in Mildura over the three weeks was fairly consistent. The highest temperature was 39.4 °C and the lowest temperature was 30.9 °C. The average temperature was 34.9 °C and the most common temperature was 36.2 °C.
9. Over the year the soccer team performed well with an average of 4 goals per game, and their most frequent amount of goals being 2 and 3 goals per game. The middle amount of goals scored was 3. The range between the highest and lowest scoring games was 11, with the IQR being 4.5.
10. a. Correct
 b. Incorrect. The range of a data set is the maximum value minus the minimum value.
 c. Correct

11. Answers will vary. Possible data sets will have the highest score = Q_3 and the lowest score = Q_1 .
12. a. No change
 b. Range will double
 c. Range will increase by 25%
13. Sample responses can be found in the worked solutions in the online resources.
14. a. Sample responses can be found in the worked solutions in the online resources.
 b. Sample responses can be found in the worked solutions in the online resources.
 c. The Police Commissioner's claim is supported, because all statistical measures were lower on Christmas Day.
15. a. 135 cm and 139 cm
 b. 175 cm
 c. Bimodal
 d. Heights of junior rowing team

Key: 15 | 9 = 159 cm

Stem	Leaf
15	9
16	2 4 5 5 8
17	0 1

Heights of senior rowing team

Key: 18 | 0 = 180 cm


Stem	Leaf
18	0 1 2 2 4 5
19	0 4

- e. Junior team: 165.5 cm, senior team: 184.75 cm,
 f. Sample responses can be found in the worked solutions in the online resources.

Project

1. Stretching the axes vertically gives the impression of a fall in the value of the shares. Stretching the axes horizontally gives the impression the value of the shares has not changed very much.

2. See figure at the foot of the page.*




- For the women's graph, stretching the vertical axis and reducing the distance between the values on the horizontal axis gives the impression women's times have fallen dramatically. To show that the men's times have not fallen greatly, you can reduce the distance between the values on the vertical axis and stretch the values on the horizontal axis.
- Sample responses can be found in the worked solutions in the online resources.

11.7 Review questions


- C
- D
- C
- D
- B
- D
- Categorical nominal; frequency table, dot plot, bar chart or pie chart
 - Categorical ordinal; frequency table, dot plot, bar chart or pie chart
 - Numerical discrete; stem plot
 - Numerical continuous: box plot
- Sample
 - Sample
 - Population
- This is categorical nominal data. It could be displayed as a frequency table, dot plot, bar chart or pie chart.
 - Examples of a dot plot and frequency table are shown. Pie or bar charts are also acceptable.

Mobiles owned by 30 shoppers



Brand of mobile	Number of mobiles
iPhone	11
Samsung	10
Motorola	6
Nokia	3

*2



10. a. Key: 1 | 2 = 12 songs

Stem	Leaf
0	29
1	
2	
3	145
4	34578
5	469
6	677
7	047
8	29
9	06
10	3
11	1

b. Positively skewed with outliers

c. Number of songs on favourite Spotify playlist

Number of songs	Number of playlists
0-19	2
20-39	3
40-59	8
60-79	6
80-99	4
100-119	2

d. Numerical discrete, symmetrical distribution, mean = 60 (rounded to the nearest whole number) and standard deviation = 27.3

11. a. See figure at the foot of the page.*

b. Positively skewed

c. The median is \$166 755, which is \$167 000 to the nearest thousand.

d. The modal class is \$100 001–\$200 000

12. a. 5.6%

b. 2%

c. 2%

d. In order to give the impression that the pay rise is generous, the company would quote the mean.

e. The trade union would quote the median or the mode, because they would be arguing that the wage rise is too low.

f. The mode is probably the most 'honest' reflection of last year's wage rise, because it represents the situation of most of the workers.

13. a. 8, 20

b. 9, 19

14. Sample response: 3, 4, 5, 6, 7, 10, 10, 13, 14

*11. a.

2019 AFL player salaries



12 Quadratic equations and graphs

LESSON SEQUENCE

12.1	Overview	708
12.2	Quadratic equations (extending)	710
12.3	The Null Factor Law (extending)	715
12.4	Solving quadratic equations with two terms (extending)	719
12.5	Factorising and solving monic quadratic equations (extending)	723
12.6	Graphs of quadratic functions	729
12.7	Sketching parabolas of the form $y = ax^2$	738
12.8	Sketching parabolas of the form $y = (x - h)^2 + k$	744
12.9	Sketching parabolas of the form $y = (x + a)(x + b)$ (extending)	756
12.10	Applications (extending)	760
12.11	Review	766



LESSON

12.1 Overview

12.1.1 Why learn this?

Galileo did a lot of important things in his lifetime. One of these was showing that objects launched into the air at an angle follow a parabolic path as they fall back to the ground. You can see this for yourself every time you throw a basketball, kick a football or hit a golf ball. As a ball moves through the air, gravity acts on it, causing it to follow a parabolic curved path. The parabolic curve that it follows can be described mathematically by quadratic equations. Every ball that is thrown, every arrow shot from a bow and every rocket launched into space has a quadratic equation that describes that object's position in space and also in time.



Quadratic equations also describe the parabolic shape of radio telescope dishes like the one at Parkes in New South Wales. The curved shape of these telescope dishes allows radio waves from space objects like quasars, galaxies and nebulae to be focused at a receiver to get a stronger signal. It was even used to relay transmissions between NASA and Apollo 11 during the first moon landing. On a much smaller scale, parabolic satellite dishes allow us to get clear internet and television. The bend in a banana, the loop of a rollercoaster, the curve of some suspension bridges or the path of a droplet of water from a fountain — these are all examples of the parabolas and quadratic equations that are all around us.

Hey students! Bring these pages to life online



Watch videos



Engage with interactivities




Answer questions and check solutions



Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets


Questions with immediate feedback, and fully worked solutions to help students get unstuck

Exercise 12.1 Pre-test


- MC** The expression $(a - 4)^2 - 25$ factorises to:

 - $(a - 9)(a + 1)$
 - $(a + 9)(a - 1)$
 - $(a - 7)(a + 3)$
 - $(a - 4)(a + 4) - 25$
 - $(a - 4)(a + 4) - 5$
- State whether the following is True or False.
A monic quadratic trinomial is an expression in the form $ax^2 + bx + c$, where a, b and c are constants.
- Factorise $m^2 + 2m - 8$.
- Factorise the equation $2w^2 - 10w + 12$ in its simplest form.
- MC** Select the solutions to the quadratic equation $-(x - 3)^2 + 4 = 0$.

 - $x = 3$ and $x = -4$
 - $x = -3$ and $x = 4$
 - $x = 3$ and $x = 4$
 - $x = -5$ and $x = -1$
 - $x = 5$ and $x = 1$
- Identify the coordinate of the turning point for the graph shown.



- MC** Identify the x -intercepts of the graph shown.



- | | | |
|-------------------|-------------------|-------------------|
| A. 0 and 1 | B. 0 and 2 | C. 0 and 4 |
| D. 1 and 2 | E. 1 and 4 | |

8. **MC** Select the equation that matches this table of values.

x	-3	-2	-1	0	1	2
y	-8	-3	0	1	0	-3

- A. $y = (1 - x)^2$ B. $y = x^2 + x$ C. $y = x^2 - 1$ D. $y = 1 - x^2$ E. $y = x^2 + 1$
9. **MC** The graph of $y = 2x^2$ is:
A. wider than $y = x^2$ B. the same as $y = x^2$
C. narrower than $y = x^2$ D. a reflection of $y = x^2$ in the x -axis
E. a reflection of $y = x^2$ in the y -axis.
10. **MC** Select the equation for a parabola that has a maximum turning point and is narrower than $y = x^2$.
A. $y = 2x^2$ B. $y = \frac{1}{2}x^2$ C. $y = (x - 2)^2 + 1$ D. $y = -2x^2$ E. $y = -\frac{1}{2}x^2$
11. **MC** Select the equation of a quadratic relation in the form $y = ax^2$ which passes through $(-2, 1)$.
A. $y = 4x^2$ B. $y = 2x^2$ C. $y = -x^2$ D. $y = -2x^2$ E. $y = \frac{1}{4}x^2$
12. For the graph of the equation $y = -9 - x^2$, state the coordinate of the turning point.
13. **MC** The graph of $y = x^2$ has been translated 3 units to the right and 2 units down. The new equation is:
A. $y = (x + 3)^2 + 2$ B. $y = (x - 3)^2 - 2$ C. $y = (x - 2)^2 + 3$
D. $y = (x + 2)^2 - 3$ E. $y = (x + 2)^2 + 3$
14. **MC** Identify the axis of symmetry for the parabola with equation $y = (x - 4)(x + 2)$.
A. $x = -2$ B. $x = -1$ C. $x = 1$ D. $x = 4$ E. $y = -8$
15. State whether the following is True or False.
The equation for the graph $y = 2(x - 1)^2 + 3$ is the same as $y = 2x^2 - 4x + 5$.

LESSON

12.2 Quadratic equations (extending)

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify equations that are quadratic
- rearrange quadratic equations into the general form $y = ax^2 + bx + c$
- solve quadratic equations of the form $ax^2 + c = 0$.

12.2.1 Quadratic equations

eles-4929

- **Quadratic equations** are equations that contain an x^2 term. They may also contain the variable x and constants, but cannot contain any higher powers of x such as x^3 .
- Examples of quadratic equations include $x^2 + 2x - 7 = 0$, $2x^2 = 18$ and $x^2 = 5x$.
- The general form of a quadratic equation is $y = ax^2 + bx + c$.
- Any quadratic equation can be expressed in general form by rearranging and combining like terms.

WORKED EXAMPLE 1 Rearranging equations into general quadratic form

Rearrange the following quadratic equations so that they are in general form and state the values of a , b and c .

a. $5x^2 - 2x + 3 = 2x^2 + 4x - 12$ b. $\frac{x^2}{2} - \frac{1}{6} = x\left(\frac{x}{3}\right) - 4$ c. $x(3 - 2x) = 4(x - 6)$

THINK

- a. 1. Write the equation.
2. Subtract $2x^2$ from both sides of the equation.
3. Subtract $4x$ from both sides of the equation.
4. Add 12 to both sides of the equation.
The equation is now in the general form $ax^2 + bx + c = 0$.
5. Write the values of a , b and c .

- b. 1. Write the equation.
2. Expand the bracket.
Lowest common denominator (LCD) = 6.
3. Multiply through by 6.
4. Simplify each fraction.
5. Collect like terms on the left-hand side.
The equation is now in the general form $ax^2 + bx + c = 0$.
6. Write the values of a , b and c .

- c. 1. Write the equation.
2. Expand the brackets.
3. To collect like terms on the left-hand side of the equation, subtract $4x$ from both sides.
4. Add 24 to both sides.
5. Multiply all terms by -1 to make the x^2 term positive. The equation is now in the general form $ax^2 + bx + c = 0$.
6. Write the values of a , b and c .

WRITE

a. $5x^2 - 2x + 3 = 2x^2 + 4x - 12$

$$3x^2 - 2x + 3 = 4x - 12$$

$$3x - 6x + 3 = -12$$

$$3x^2 - 6x + 15 = 0$$

$$a = 3, b = -6, c = 15$$

b. $\frac{x^2}{2} - \frac{1}{6} = x\left(\frac{x}{3}\right) - 4$

$$\frac{x^2}{2} - \frac{1}{6} = \frac{x^2}{3} - 4$$

$$\frac{x^2}{2} \times \frac{6}{1} - \frac{1}{6} \times \frac{6}{1} = \frac{x^2}{3} \times \frac{6}{1} - 4 \times 6$$

$$3x^2 - 1 = 2x^2 - 24$$

$$x^2 - 1 = -24$$

$$x^2 + 23 = 0$$

$$a = 1, b = 0, c = 23$$

c. $x(3 - 2x) = 4(x - 6)$

$$3x - 2x^2 = 4x - 24$$

$$-2x^2 - x = -24$$

$$-2x^2 - x + 24 = 0$$

$$2x^2 + x - 24 = 0$$

$$a = 2, b = 1, c = -24$$



eles-4930

12.2.2 Solving equations of the form $ax^2 + c = 0$

- Solving a quadratic equation means determining the values of x that satisfy the equation.
- Some quadratic equations have two solutions, some have only one solution, and some have no solutions.

- Quadratic equations of the form $ax^2 + c = 0$ can be solved using square roots. To solve such an equation, rearrange it so that the x^2 term is isolated on one side of the equation. Then take the square root of both sides.

WORKED EXAMPLE 2 Solving equations of the form $ax^2 + c = 0$

Solve the following equations.

a. $2x^2 - 18 = 0$

b. $x^2 + 9 = 0$

c. $3x^2 + 4 = 4$

THINK

- a. 1. Write the equation. The aim is to make x the subject.
 2. Add 18 to both sides.
 3. Divide both sides by 2.
 4. Take the square root of both sides.
 5. Write the solutions.
- b. 1. Write the equation. The aim is to make x the subject.
 2. Take 9 from both sides.
 3. Take the square root of both sides.
 4. Write the solutions.
- c. 1. Write the equation. The aim is to make x the subject.
 2. Take 4 from both sides.
 3. Divide both sides by 3.
 4. Take the square root of both sides. Zero has only one square root.
 5. Write the solutions.

WRITE

a. $2x^2 - 18 = 0$

$$2x^2 = 18$$

$$x^2 = 9$$

$$x = 3, -3$$

Two solutions: $x = 3, x = -3$

b. $x^2 + 9 = 0$

$$x^2 = -9$$

-9 has no square root.

The equation has no real solution.

c. $3x^2 + 4 = 4$

$$3x^2 = 0$$

$$x^2 = 0$$


$$x = 0$$

One solution: $x = 0$

TI | THINK

- a–c. On a Calculator page, press:
- Menu
 - 3. Algebra
 - 1. Solve
- Complete the entry lines as:
- solve ($2x^2 - 18 = 0, x$)
 solve ($x^2 + 9 = 0, x$)
 solve ($3x^2 + 4 = 4, x$)
- Press ENTER after each entry.

DISPLAY/WRITE




- a. $x = -3, x = 3$
 b. No solution.
 c. $x = 0$

CASIO | THINK

- a–c. On a Main screen, complete the entry lines as:
- solve ($2x^2 - 18 = 0$)
 solve ($x^2 + 9 = 0$)
 solve ($3x^2 + 4 = 4$)
- Press EXE after each entry.

DISPLAY/WRITE



- a. $x = -3, x = 3$
 b. No solution.
 c. $x = 0$

Confirming the solutions to quadratic equations

- Solutions to quadratic equations can be checked by substitution.
- Substitute the solution(s) into the equation, and if the RHS of the equation is equal to the LHS, the solutions are correct.

WORKED EXAMPLE 3 Confirming solutions to quadratic equations

Determine whether any of the following are solutions to the quadratic equation $x^2 = 4x - 3$.

a. $x = 1$

b. $x = 2$

c. $x = 3$

THINK

- a. 1. Substitute $x = 1$ into both sides of the equation.
2. Evaluate the left-hand side.
3. Evaluate the right-hand side.
4. Since the left-hand side equals the right-hand side, $x = 1$ is a solution. Write the answer.
- b. 1. Substitute $x = 2$ into both sides of the equation.
2. Evaluate the left-hand side.
3. Evaluate the right-hand side.
4. The left-hand side does not equal the right-hand side, so $x = 2$ is not a solution. Write the answer.
- c. 1. Substitute $x = 3$ into both sides of the equation.
2. Evaluate the left-hand side.
3. Evaluate the right-hand side.
4. The left-hand side equals the right-hand side, so $x = 3$ is a solution. Write the answer.

WRITE

a. $x^2 = 4x - 3$

$$\text{LHS} = (1)^2 \\ = 1$$

$$\text{RHS} = 4(1) - 3 \\ = 4 - 3 \\ = 1$$

$x = 1$ is a solution to the equation.

b. $x^2 = 4x - 3$

$$\text{LHS} = (2)^2 \\ = 4$$

$$\text{RHS} = 4(2) - 3 \\ = 8 - 3 \\ = 5$$

$x = 2$ is not a solution to the equation.

c. $x^2 = 4x - 3$

$$\text{LHS} = (3)^2 \\ = 9$$

$$\text{RHS} = 4(3) - 3 \\ = 12 - 3 \\ = 9$$

$x = 3$ is a solution to the equation.

Resources



eWorkbook Topic 12 workbook (worksheets, code puzzle and a project) (ewbk-2012)



Interactivity Individual pathway interactivity: Quadratic equations (int-8349)

12.2 Quick quiz **on**

12.2 Exercise

Individual pathways

PRACTISE

1, 5, 7, 11, 14

CONSOLIDATE

2, 3, 8, 10, 12, 15

MASTER

4, 6, 9, 13, 16

Fluency

- WE1** Rearrange each of the following quadratic equations so that they are in the form $ax^2 + bx + c = 0$.

a. $3x - x^2 + 1 = 5x$ b. $5(x - 2) = x(4 - x)$ c. $x(5 - 2x) = 6(5 - x^2)$
- Rearrange each of the following quadratic equations so that they are in the form $ax^2 + bx + c = 0$.

a. $4x^2 - 5x + 2 = 6x - 4x^2$ b. $5(x - 12) = x(4 - 2x)$ c. $x^2 - 4x - 16 = 1 - 4x$
- MC** Select which of the following is a quadratic equation.

A. $2x - 1 = 0$ B. $2^x - 1 = 0$ C. $x^2 - x = 1 + x^2$
 D. $x^2 - x = 1 - x^2$ E. $x^3 + 2x = 0$
- MC** Select which of the following is not a quadratic equation.

A. $x(x - 1) = 2x - 1$ B. $-3x^2 + 2x = 1$ C. $3(x + 2) + 5(x + 3) = 2(x + 1)$
 D. $2(x - 1) + 3x = x(2x - 3)$ E. $2x^3 + x = 2 - 2x^2$
- WE2** Solve the following quadratic equations.

a. $x^2 - 16 = 0$ b. $2x^2 + 18 = 0$ c. $3x^2 + 2x = x(x + 2)$
- Solve the following quadratic equations.

a. $2(x^2 + 7) = 16$ b. $x^2 + 17 = 13$ c. $-3x^2 + 17 = 5$

Understanding

- WE3** Determine whether $x = 4$ is a solution to the following equations.

a. $x^2 = x + 12$ b. $x^2 = 3x + 1$ c. $x^2 = 4x$
- Determine whether $x = -3$ is a solution to the following equations.


a. $x^2 = x + 12$ b. $x^2 = 3x + 1$ c. $x^2 = 4x$
- Determine whether $x = 0$ is a solution to the following equations.

a. $x^2 = x + 12$ b. $x^2 = 3x + 1$ c. $x^2 = 4x$
- Determine whether the following are solutions to the equation $5(x - 1)^2 + 7 = 27$.

a. $x = -1$ b. $x = 1$ c. $x = 0$


Reasoning

- Is $(x + 4)^2$ equal to $x^2 + 16$? Explain using a numerical example.
- a. Show that the shaded area can be represented by $A = x^2(4 - \pi)$.
 b. If the shaded area = 10 cm^2 , determine the value of x correct to 3 significant figures.
 c. Why can't x be equal to -3.41 ?
- Explain why $x^2 + 7x + 4 = 7x$ has no solutions.



Problem solving

14. Five squares of increasing size and five area formulas are given below.
- Use factorisation to determine the side length that correlates to each area formula.
 - Using the area given and side lengths found, match the squares below with the appropriate algebraic expression for their area.



$$A_a = x^2 + 6x + 9$$

$$A_b = x^2 + 10x + 25$$

$$A_c = x^2 + 16x + 64$$

$$A_d = x^2 - 6x + 9$$

$$A_e = x^2 + 12x + 36$$

- If $x = 5$ cm, use the formula given to calculate the area of each square.
15. The sum of 5 and the square of a number is 41.
- Determine the number.
 - Is there more than one possible number? If so, determine the other number and explain why there is more than one.
16. Two less than a number is squared and its result is tripled. The difference between this and 7 is 41.
- Write the algebraic equation.
 - Determine both possible answers that satisfy the criteria.

LESSON

12.3 The Null Factor Law (extending)

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- recall the Null Factor Law
- solve quadratic equations using the Null Factor Law.



eles-4931

12.3.1 Using the Null Factor Law to solve equations

- The **Null Factor Law** states that if the product of two or more factors is zero, then at least one of the factors must be zero.

The Null Factor Law

If $a \times b = 0$ then:

$a = 0$ or $b = 0$ or both a and b equal 0.

For example, if $(x - 5)(x - 2) = 0$ then

$$(x - 5) = 0 \text{ or } (x - 2) = 0 \text{ or both } x - 5 = x - 2 = 0.$$

$$\text{If } x - 5 = 0 \quad \text{If } x - 2 = 0$$

$$x = 5 \quad x = 2$$

Both $x = 5$ and $x = 2$ make the equation true.

- This can be checked by substituting the values into the original equation.

For example:

If $x = 5$, then:

$$\begin{aligned}(x - 5)(x - 2) &= (5 - 5)(5 - 2) \\ &= 0 \times 3 \\ &= 0\end{aligned}$$

If $x = 2$, then:

$$\begin{aligned}(x - 5)(x - 2) &= (2 - 5)(2 - 2) \\ &= -3 \times 0 \\ &= 0\end{aligned}$$

Note: Quadratic equations can have a maximum of two solutions. This is known as the **Fundamental Theorem of Algebra**.

WORKED EXAMPLE 4 Using the Null Factor Law to solve quadratics

Solve each of the following quadratic equations.

a. $(x - 2)(2x + 1) = 0$

b. $(4 - 3x)(6 + 11x) = 0$

c. $x(x - 3) = 0$

d. $(x - 1)^2 = 0$

THINK

- a. 1. The product of 2 factors is 0, so apply the Null Factor Law.
2. One of the factors must equal zero.
3. Solve the equations.
4. Write the solutions.
- b. 1. The product of two factors is 0, so apply the Null Factor Law.
2. One of the factors must equal zero.
3. Solve the equations.
4. Write the solutions.
- c. 1. The product of two factors is 0, so apply the Null Factor Law.
2. One of the factors must equal zero.
3. Solve the equations.
4. Write the solutions.
- d. 1. Write the equation.
2. Rewrite the squared factors as the product of two factors.

WRITE

a. $(x - 2)(2x + 1) = 0$

Either $x - 2 = 0$ or $2x + 1 = 0$

$$\begin{aligned}x &= 2 & 2x &= -1 \\ & & &= -\frac{1}{2}\end{aligned}$$

$$x = -\frac{1}{2}, x = 2$$

b. $(4 - 3x)(6 + 11x) = 0$

Either $4 - 3x = 0$ or $6 + 11x = 0$

$$\begin{aligned}-3x &= -4 & 11x &= -6 \\ x &= \frac{4}{3} & x &= \frac{-6}{11}\end{aligned}$$

$$x = \frac{4}{3}, -\frac{6}{11}$$

c. $x(x - 3) = 0$

Either $x = 0$ or $x - 3 = 0$

$$x = 3$$

$$x = 0, 3$$

d. $(x - 1)^2 = 0$

$$(x - 1)(x - 1) = 0$$

- Apply the Null Factor Law.
- Solve the equations.
The two solutions are the same.
- Write the solution.

$$\text{Either } x - 1 = 0 \quad \text{or} \quad x - 1 = 0$$


$$x = 1 \qquad \qquad \qquad x = 1$$

$$x = 1$$

TI | THINK

- a–c.** On a Calculator page, press:
- Menu
 - 3. Algebra
 - 1. Solve
- Complete the entry lines as:
- solve
 $((x-2)(2x+1)=0, x)$
 solve
 $((4-3x)(6+11x)=0, x)$
 solve
 $((x-1)^2=0, x)$
 Press ENTER after each entry.

DISPLAY/WRITE




- a.** $x = -\frac{1}{2}, x = 2$
b. $x = -\frac{6}{11}, x = \frac{4}{3}$
c. $x = 1$

CASIO | THINK

- a–c.** On a Main screen, complete the entry lines as:
- solve
 $((x-2)(2x+1)=0)$
 solve
 $((4-3x)(6+11x)=0)$
 solve
 $((x-1)^2=0)$
 Press EXE after each entry.

DISPLAY/WRITE



- a.** $x = -\frac{1}{2}, x = 2$
b. $x = -\frac{6}{11}, x = \frac{4}{3}$
c. $x = 1$

on Resources

- eWorkbook** Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)
- Video eLesson** The Null Factor Law (eles-2312)
- Interactivities** Individual pathway interactivity: The Null Factor Law (int-8350)
The Null Factor Law (int-6095)

Exercise 12.3 The Null Factor Law (extending)

learn on

12.3 Quick quiz **on**

12.3 Exercise

Individual pathways

PRACTISE

1, 6, 9, 12

CONSOLIDATE

2, 4, 7, 10, 13

MASTER

3, 5, 8, 11, 14

Fluency

- 1. WE4** Solve each of the following quadratic equations.

a. $(x-2)(x+3)=0$

b. $(2x+4)(x-3)=0$

c. $(x+2)(x-3)=0$

d. $(2x+5)(4x+3)=0$

e. $(x+4)(2x+1)=0$

2. Solve each of the following quadratic equations.

a. $(2x - 1)(x + 30) = 0$

b. $(2x + 1)(3 - x) = 0$

c. $(1 - x)(3x - 1) = 0$

d. $x(x - 2) = 0$

e. $\left(x + \frac{1}{3}\right)\left(2x - \frac{1}{2}\right) = 0$

3. Solve each of the following quadratic equations.

a. $(5x - 1.5)(x + 2.3) = 0$

b. $\left(2x + \frac{1}{3}\right)\left(2x - \frac{1}{3}\right) = 0$

c. $(x - 2)^2 = 0$

d. $x(4x - 15) = 0$

e. $(x + 4)^2 = 0$

4. The Null Factor Law can be extended to products of more than two factors. Use this to determine all the solutions to the following equations.

a. $(x - 2)(x + 2)(x + 3) = 0$

b. $(x + 2)(x + 2)(2x - 5) = 0$

c. $(x + 2)(x + 2)(x + 4) = 0$

d. $x(x + 2)(3x + 12) = 0$

5. The Null Factor Law can be extended to products of more than two factors. Use this to determine all the solutions to the following equations.

a. $(2x - 2.2)(x + 2.4)(x + 2.6) = 0$

b. $(2x + 6)\left(x + \frac{1}{2}\right)(9x - 15) = 0$

c. $3(x - 3)^2 = 0$

d. $(x + 1)(x - 2)^2 = 0$

Understanding

6. **MC** Select the solutions to $(2x - 4)(x + 7) = 0$.

A. $x = 4, x = 7$

B. $x = 4, x = -7$

C. $x = 2, x = 7$

D. $x = 2, x = -7$

E. $x = -4, x = 7$

7. **MC** The Null Factor Law cannot be applied to the equation $x\left(x + \frac{1}{2}\right)\left(\frac{x}{2} - 1\right) = 1$ because:

A. there are more than two factors

B. the right-hand side equals 1

C. the first factor is a simple x -term

D. the third term has x in a fraction

E. there are three terms on the left-hand side

8. Rewrite the following equations so that the Null Factor Law can be used. Then solve the resulting equation.

a. $x^2 + 10x = 0$

b. $2x^2 - 14x = 0$

c. $25x^2 - 40x = 0$

Reasoning

9. Explain why $x^2 - x - 6 = 0$ has the same solution as $-2x^2 + 2x + 12 = 0$.



10. a. Explain why $x = 2$ is not a solution of the equation $(2x - 2)(x + 2) = 0$.

b. Determine the solutions to the equation.

11. Explain what is the maximum number of solutions a quadratic such as $x^2 - x - 56 = 0$ can have.

Problem solving

12. A bridge is constructed with a supporting structure in the shape of a parabola, as shown in the diagram. The origin $(0, 0)$ is at the left-hand edge of the bridge, which is 100 m long.
- Identify the maximum height of the bridge support.
 - If the equation of the support is $y = ax(b - x)$, determine the values of a and b . (*Hint: Let $y = 0$.*)
 - Calculate the height of the support when $x = 62$.
13. Consider a ball thrown upwards so that it reaches a height h of metres after t seconds. The expression $20t - 4t^2$ represents the height of the ball, in metres, after t seconds.



- Factorise the expression for height.
 - Calculate the height of the ball after:
 - 1 second
 - 5 seconds.
 - Explain if factorising the expression made it easier to evaluate.
14. The product of 7 more than a number and 14 more than that same number is 330. Determine the possible values of the number.

LESSON

12.4 Solving quadratic equations with two terms (extending)

LEARNING INTENTION

At the end of this lesson you should be able to:

- solve quadratic equations with two terms using the Null Factor Law.



eles-4932

12.4.1 Solving quadratic equations of the form $ax^2 + c = 0$

- In section 12.2.2, we saw how to solve equations in the form $ax^2 + c = 0$ using algebra. In most cases, we can use the Null Factor Law to solve them instead.
- To use the Null Factor Law the quadratic needs to be factorised and the RHS of the equation must be 0.

WORKED EXAMPLE 5 Using the Null Factor Law to solve $ax^2 + c = 0$

Solve each of the following quadratic equations.

a. $x^2 - 1 = 0$

b. $2x^2 - 18 = 0$

THINK

WRITE

a. 1. Write the equation: $x^2 - 1$ is the difference of two squares.

a. $x^2 - 1 = 0$

2. Factorise the left-hand side using the difference of two squares rule.

$$(x + 1)(x - 1) = 0$$

3. Apply the Null Factor Law.

$$\begin{aligned}x + 1 = 0 & \quad x - 1 = 0 \\x = -1 & \quad x = 1\end{aligned}$$

4. State the solutions.

$$x = 1, x = -1$$

(This can be abbreviated to $x = \pm 1$.)

b. 1. Write the equation.

b. $2x^2 - 18 = 0$

2. Take out the common factor.

$$2(x^2 - 9) = 0$$

3. Divide both sides of the equation by 2.

$$x^2 - 9 = 0$$

4. Factorise the left-hand side using the difference of two squares rule.

$$(x + 3)(x - 3) = 0$$

5. Apply the Null Factor Law.

$$\begin{aligned}x + 3 = 0 & \quad x - 3 = 0 \\x = -3 & \quad x = 3\end{aligned}$$

6. State the solutions.

$$x = \pm 3$$



eles-4933

12.4.2 Solving quadratic equations of the form $ax^2 + bx = 0$

- The equation $ax^2 + bx = 0$ can be solved easily using the Null Factor Law.
- To solve an equation of this form, factorise the LHS to $x(ax + b) = 0$ and then apply the Null Factor Law.
- Since both terms on the left-hand side involve the variable x , one solution will always be $x = 0$.

WORKED EXAMPLE 6 Solving equations in the form $ax^2 + bx = 0$

Solve each of the following equations.

a. $x^2 + 4x = 0$

b. $-2x^2 - 4x = 0$

THINK

WRITE

a. 1. Write the equation.

a. $x^2 + 4x = 0$

2. Factorise by taking out a common factor of x .

$$x(x + 4) = 0$$

3. Apply the Null Factor Law.

$$\begin{aligned}x = 0 & \quad x + 4 = 0 \\x = 0 & \quad x = -4\end{aligned}$$

4. Write the solutions.

$$x = -4, \quad x = 0$$

b. 1. Write the equation.

b. $-2x^2 - 4x = 0$

2. Factorise by taking out the common factor of $-2x$.

$$-2x(x + 2) = 0$$

3. Apply the Null Factor Law.

$$\begin{array}{l} -2x = 0 \\ x = 0 \end{array} \quad \begin{array}{l} x + 2 = 0 \\ x = -2 \end{array}$$

4. Write the solutions.

$$x = -2, \quad x = 0$$

WORKED EXAMPLE 7 Solving word problems using Null Factor Law

If the square of a number is multiplied by 5, the answer is 45. Calculate the number.

THINK

1. Define the number.
2. Write an equation that can be used to determine the number.
3. Transpose to make the right-hand side equal to zero.
Solve the equation.

WRITE

Let x be the number.

$$5x^2 = 45$$

$$5x^2 - 45 = 0$$

$$5(x^2 - 9) = 0$$

$$x^2 - 9 = 0$$

$$x^2 - 3^2 = 0$$

$$(x + 3)(x - 3) = 0$$

$$x + 3 = 0 \quad \text{or} \quad x - 3 = 0$$

$$x = -3 \quad \text{or} \quad x = 3$$

The number is either 3 or -3 .

4. Write the answer in a sentence.

on Resources



eWorkbook Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivity Individual pathway interactivity: Solving quadratic equations with two terms (int-8351)

Exercise 12.4 Solving quadratic equations with two terms (extending)

learn **on**

12.4 Quick quiz **on**

12.4 Exercise

Individual pathways

■ PRACTISE

1, 4, 7, 9, 12, 16

■ CONSOLIDATE

2, 5, 8, 10, 13, 17

■ MASTER

3, 6, 11, 14, 15, 18

Fluency

1. **WE5** Solve each of the following quadratic equations using the Null Factor Law.

a. $x^2 - 9 = 0$

b. $x^2 - 16 = 0$

c. $2x^2 - 18 = 0$

d. $2x^2 - 50 = 0$

e. $100 - x^2 = 0$

2. Solve each of the following quadratic equations using the Null Factor Law.

a. $49 - x^2 = 0$

b. $3x^2 - 27 = 0$

c. $5x^2 - 20 = 0$

d. $x^2 + 6 = 0$

e. $2x^2 + 18 = 0$

3. Solve each of the following quadratic equations using the Null Factor Law.

a. $-x^2 + 9 = 0$

b. $-3x^2 + 48 = 0$

c. $-4x^2 + 100 = 0$

d. $x^2 = 0$

e. $-x^2 = 0$

4. **WE6** Solve each of the following equations.

a. $x^2 + 6x = 0$

b. $x^2 - 8x = 0$

c. $x^2 + 9x = 0$

d. $x^2 - 11x = 0$

e. $2x^2 - 12x = 0$

5. Solve each of the following equations.

a. $2x^2 - 15x = 0$

b. $3x^2 - 2x = 0$

c. $4x^2 + 7x = 0$

d. $2x^2 - 5x = 0$

e. $x^2 + x = 0$

6. Solve each of the following equations.

a. $4x^2 - x = 0$

b. $-x^2 - 5x = 0$

c. $-2x^2 - 24x = 0$

d. $-x^2 + 18x = 0$

e. $x^2 - 2.5x = 0$

7. **MC** Select the solutions to $4x^2 - 36 = 0$.

A. $x = 3$ and $x = -3$

B. $x = 9$ and $x = -9$

C. $x = 1$ and $x = -1$

D. $x = 2$ and $x = -2$

E. $x = 6$ and $x = -6$

8. **MC** Select the solutions to $x^2 - 5x = 0$.

A. $x = 1$ and $x = 5$

B. $x = 0$ and $x = -5$

C. $x = 0$ and $x = 5$

D. $x = -1$ and $x = 5$

E. $x = -5$ and $x = 5$

Understanding

9. **WE7** If the square of a number is multiplied by 2, the answer is 32. Calculate the number.

10. A garden has two vegetable plots. One plot is a square; the other plot is a rectangle with one side 3 m shorter than the side of the square and the other side 4 m longer than the side of the square. Both plots have the same area. Sketch a diagram and determine the dimensions of each plot.

11. The square of a number is equal to 10 times the same number. Calculate the number.

Reasoning

12. Explain why the equation $x^2 + 9 = 0$ cannot be solved using the Null Factor Law.

13. Solve the equation $m^2x^2 - n^2 = 0$ using the Null Factor Law.

14. Explain why the equation $x^2 + bx = 0$ can only be solved using the Null Factor Law.

15. A quadratic expression may be written as $ax^2 + bx + c$. Using examples, explain why b and c may take any values, but a cannot equal zero.

Problem solving

16. The square of a number which is then tripled, is equal to 24 times the same number. Calculate the number.

17. The sum of 6 times the square of a number and 72 times the same number is equal to zero. Determine the value of the number.

18. A rectangular horse paddock has a width of x and a length that is 30 metres longer than its width. The area of the paddock is $50x$ square metres.

a. Write the algebraic equation for the area described.

b. Factorise the equation.

c. Solve the equation.

d. Can you use all your solutions? Explain why or why not.

LESSON

12.5 Factorising and solving monic quadratic equations (extending)

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- factorise monic quadratic equations
- solve monic quadratic equations using factorisation and the Null Factor Law.

12.5.1 Quadratic trinomials



eles-4934

- A **quadratic trinomial** is an expression of the form $ax^2 + bx + c$, where a , b and c are constants.
 - For the time being we will look only at quadratic trinomials for which $a = 1$; that is, trinomials of the form $x^2 + bx + c$.
- The area model of binomial expansion can be reversed to determine a pattern for factorising a general quadratic expression.

For example:

$$\begin{aligned}(x+f)(x+h) &= x^2 + fx + hx + fh \\ &= x^2 + (f+h)x + fh\end{aligned}$$

$$\begin{aligned}(x+4)(x+3) &= x^2 + 4x + 3x + 12 \\ &= x^2 + 7x + 12\end{aligned}$$



Factorising monic quadratics

To factorise a general quadratic, look for factors of c that sum to b .

$$x^2 + bx + c = (x + f)(x + h)$$

Factors of c that sum to b

For example, $x^2 + 7x + 12 = (x + 3)(x + 4)$.

$$3 + 4 = 7$$

$$3 \times 4 = 12$$

WORKED EXAMPLE 8 Factorising quadratic expressions

Factorise the following quadratic expressions.

a. $x^2 + 5x + 6$

b. $x^2 + 10x + 24$

THINK

a. 1. The general quadratic expression has the pattern $x^2 + 5x + 6 = (x+f)(x+h)$. f and h are a factor pair of 6 that add to 5.

Calculate the sums of factor pairs of 6. The factors of 6 that add to 5 are 2 and 3, as shown in blue.

2. Substitute the values of f and h into the expression in its factorised form.

b. 1. The general quadratic expression has the pattern $x^2 + 10x + 24 = (x+f)(x+h)$, where f and h are a factor pair of 24 that add to 10.

Calculate the sums of factor pairs of 24. The factors of 24 that add to 10 are 4 and 6, as shown in blue.

2. Substitute the values of f and h into the expression in its factorised form.

WRITE

a.

Factors of 6	Sum of factors
1 and 6	7
2 and 3	5

$$(x^2 + 5x + 6) = (x + 2)(x + 3)$$

b.

Factors of 24	Sum of factors
1 and 24	25
2 and 12	14
3 and 8	11
4 and 6	10

$$x^2 + 10x + 24 = (x + 4)(x + 6)$$

WORKED EXAMPLE 9 Factorising quadratic expressions with negative terms

Factorise the following expressions.

a. $x^2 - 9x + 18$

b. $x^2 + 6x - 16$

THINK

a. 1. The general quadratic expression has the pattern $x^2 - 9x + 18 = (x+f)(x+h)$, where f and h are a factor pair of 18 that add to -9 .

Calculate the sums of factor pairs of 18. As shown in blue, -3 and -6 are factors of 18 that add to -9 .

2. Substitute the values of f and h into the expression in its factorised form.

b. 1. The general quadratic expression has the pattern $x^2 + 6x - 16 = (x+f)(x+h)$, where f and h are a factor pair of -16 that add to 6.

Calculate the sums of factor pairs of -16 . As shown in blue, -2 and 8 are factors of -16 that add to 6.

2. Substitute the values of f and h into the expression in its factorised form.

WRITE

a.

Factors of 18	Sum of factors
1 and 18	19
-1 and -18	-19
2 and 9	11
-2 and -9	-11
3 and 6	9
-3 and -6	-9

$$x^2 - 9x + 18 = (x - 3)(x - 6)$$

b.

Factors of -16	Sum of factors
1 and -16	-15
-1 and 16	15
2 and -8	-6
-2 and 8	6

$$x^2 + 6x - 16 = (x - 2)(x + 8)$$

12.5.2 Solving quadratic equations of the form $x^2 + bx + c = 0$

- If the expression $x^2 + bx + c$ can be factorised, then the quadratic equation $x^2 + bx + c = 0$ can be solved using the Null Factor Law.

WORKED EXAMPLE 10 Solving monic quadratic equations

Solve the following quadratic equations.

a. $x^2 - 5x - 6 = 0$

b. $x^2 + 14x = 15$

THINK

- a. 1. Write the equation.
 2. Factorise $x^2 - 5x - 6$.
 3. Apply the Null Factor Law.
 4. Solve the equations.
 5. Write the solutions.
- b. 1. Write the equation and transpose so that the right-hand side equals 0.
 2. Factorise $x^2 - 14x - 15$.
 3. Apply the Null Factor Law.
 4. Solve the linear equations and write the solutions.

WRITE


a. $x^2 - 5x - 6 = 0$
 $(x - 6)(x + 1) = 0$
 $x - 6 = 0 \quad x + 1 = 0$
 $x = 6 \quad x = -1$
 $x = -1, \quad x = 6$

b. $x^2 + 14x = 15$
 $x^2 + 14x - 15 = 0$
 $(x + 15)(x - 1) = 0$
 $x + 15 = 0 \quad x - 1 = 0$
 $x = -15 \quad x = 1$
 $x = -15, \quad x = 1$

TI | THINK

- a, b. In a new problem on a Calculator page, complete the entry lines as:
 solve
 $(x^2 - 5x - 6 = 0, x)$
 solve
 $(x^2 + 14x = 15, x)$
 Press ENTER
 after each entry.
 Press ENTER
 after each entry.

DISPLAY/WRITE




- a. $x = -1, x = 6$
 b. $x = -15, x = 1$

CASIO | THINK

- a, b. On the Main screen, complete the entry lines as:
 solve
 $(x^2 - 5x - 6 = 0)$
 solve $(x^2 + 14x = 15)$
 Press EXE after each entry.

DISPLAY/WRITE



- a. $x = -1, x = 6$
 b. $x = -15, x = 1$



eWorkbook Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivities Individual pathway interactivity: Factorising and solving monic quadratics (int-8352)

Factorising monic quadratics (int-6092)

Exercise 12.5 Factorising and solving monic quadratic equations (extending)

learn**on**

12.5 Quick quiz **on**

12.5 Exercise

Individual pathways

PRACTISE

1, 4, 7, 10, 13, 18, 22, 25

CONSOLIDATE

2, 5, 8, 11, 14, 16, 19, 20, 23, 26

MASTER

3, 6, 9, 12, 15, 17, 21, 24, 27, 28

Fluency

1. Expand each of the following.

a. $(x + 4)(x + 2)$

b. $(x - 2)(x - 4)$

c. $(x - 4)(x - 5)$

d. $(x + 4)(x + 5)$

e. $(m + 1)(m + 5)$

f. $(m - 1)(m - 5)$

2. Expand each of the following.

a. $(t + 8)(t + 11)$

b. $(t - 10)(t - 20)$

c. $(x + 2)(x - 3)$

d. $(x + 3)(x - 2)$

e. $(v + 5)(v - 8)$

f. $(v - 5)(v + 8)$

3. Expand each of the following.

a. $(x + 7)(x - 2)$

b. $(t + 1)(t - 12)$

c. $(n + 15)(n - 2)$

d. $(a + 3)(a - 6)$

e. $(z + 4)(z + 4)$

f. $(z + 11)(z - 8)$

4. Make a complete systematic list of pairs of positive integers that add up to:

a. 5

b. 6

c. 12

d. 20.

5. Make a complete systematic list of pairs of negative integers that add up to:

a. -5

b. -7

c. -8

d. -11.

6. Make a systematic list of 5 pairs of integers with opposite signs whose sum is:

a. 6

b. -6

c. 3

d. -3

e. -8

f. 14.

7. **WE8** Factorise each of the following.

a. $x^2 + 4x + 3$

b. $x^2 - 4x + 3$

c. $x^2 + 12x + 11$

d. $x^2 - 12x + 11$

e. $a^2 + 6a + 5$

8. Factorise each of the following.

a. $a^2 - 6a + 5$

b. $x^2 - 7x + 12$

c. $x^2 - 7x + 10$

d. $n^2 + 8n + 16$

e. $n^2 + 10n + 16$

9. Factorise each of the following.

a. $y^2 - 12y + 27$

b. $x^2 - 13x + 42$

c. $t^2 - 8t + 12$

d. $t^2 + 11t + 18$

e. $u^2 + 5u + 6$

10. **WE9** Factorise each of the following.

a. $x^2 + 3x - 18$

b. $x^2 - 3x - 18$

c. $x^2 - 2x - 15$

d. $x^2 + 2x - 15$

e. $n^2 - 13n - 14$

11. Factorise each of the following.

a. $n^2 + 2n - 35$

b. $v^2 + 5v - 6$

c. $v^2 - 5v - 6$

d. $t^2 + 4t - 12$

e. $t^2 - 5t - 14$

12. Factorise each of the following.

a. $x^2 - x - 20$

b. $x^2 + x - 20$

c. $n^2 + n - 90$

d. $n^2 - 3n - 70$

e. $x^2 - 4x - 5$

13. **WE10** Solve each of the following quadratic equations.

a. $x^2 - 6x + 8 = 0$

b. $x^2 + 6x + 8 = 0$

c. $x^2 + 6x + 5 = 0$

d. $x^2 + x - 6 = 0$

e. $x^2 + 2x - 15 = 0$

14. Solve each of the following quadratic equations.

a. $x^2 + 4x + 4 = 0$

b. $x^2 + 2x - 24 = 0$

c. $x^2 - 5x - 24 = 0$

d. $x^2 - x - 12 = 0$

e. $x^2 + 13x + 12 = 0$

15. Solve each of the following quadratic equations.

a. $x^2 - 10x = 11$

b. $x^2 + x = 20$

c. $x^2 + 29x = -100$

d. $x^2 - 15x = -50$

e. $0 = x^2 - 2x - 8$

16. **MC** Select the solutions to the quadratic equation $x^2 + 2x - 8 = 0$.

A. $x = 4$ and $x = -2$

B. $x = -4$ and $x = 2$

C. $x = -4$ and $x = 4$

D. $x = -2$ and $x = 2$

E. $x = -4$ and $x = -2$

17. **MC** Select the solutions to the quadratic equation $x^2 - 7x - 8 = 0$.

A. $x = 1$ and $x = 8$

B. $x = -1$ and $x = -8$

C. $x = -8$ and $x = 1$

D. $x = -1$ and $x = 8$

E. $x = -1$ and $x = 1$

Understanding

18. Consider the quadratic trinomial $x^2 + 7x + c$ where c is a positive integer.

a. Factorise the expression if $c = 6$.

b. Determine what other positive whole number values c can take if the expression is to be factorised. Factorise the expression for each of these values.

19. Consider the quadratic trinomial $x^2 + 3x + c$ where c is a negative integer.

a. Factorise the expression if $c = -4$.

b. Determine three more values of c for which the expression can be factorised, and factorise each one.

20. **MC** When factorised, $x^2 - 3x - 18$ is equal to:

A. $(x - 3)(x - 6)$


B. $(x - 3)(x + 6)$

C. $(x + 3)(x + 6)$

D. $(x + 3)(x - 6)$


E. $(x + 2)(x - 9)$

21. The rectangle shown has an area of 45 cm^2 . By solving a quadratic equation, determine the dimensions of the rectangle.






Reasoning

22. Show that $x^2 + 8x + 10$ has no factors, if only whole numbers can be used.
23. Show that $x^2 + 12x + 24$ has no factors, if only whole numbers can be used.
24. The number of diagonals in a polygon is given by the formula $D = \frac{n}{2}(n - 3)$, where n is the number of sides in the polygon.
- Determine how many diagonals there are in:
 - a triangle
 - a square
 - a decagon.
 - Determine what type of polygon has:
 - 20 diagonals
 - 170 diagonals.



Problem solving

25. a. Write algebraic expressions for each of the parts below. Your friend picks a number, n .
- He adds 1 to it.
 - He adds 7 to it.
 - He multiplies these two new numbers together.
 - He then divides the product in part **iii** by the product of 2 more than the original number, n , and 1 more than the original number, n .
- b. The result of the division is 2. What number, n , did he begin with?
26. Rectangular floor mats have an area of $x^2 + 2x - 15$.
- If the length of the mat is $(x + 5)$ cm, develop an expression for the width.
 - If the length of the mat is 70 cm, calculate the width.
 - If the width of the mat is 1 m, calculate the length.
27. A girl, her brother and her teacher are shown below. The product of the teacher's and brother's ages is 256.



- Write an algebraic expression in terms of g to represent the teacher's age.
- Use the algebraic expressions to write an expression for the product of the brother's and teacher's ages and expand this product.
- Write an equation using your answer to part **b** for the product of their ages. *Hint:* An equation has an 'is equal to' part to it.
- Rearrange your answer to part **c** by collecting like terms and making it equal to zero.
- Determine the value of g by solving the equation in part **d** using the Null Factor Law.
Note: Discard any value that does not make sense.
- State the ages of the girl, her brother and the teacher.

28. A particular rectangle has an area of 120 m^2 .
- Generate a list of possible whole number dimensions for the rectangle.
 - The area of the rectangle can also be expressed as $A = x^2 + 2x - 48$. Determine algebraic expressions for the length and the width of the rectangle in terms of x .
 - Use your list of possible dimensions and your algebraic expressions for width and length to determine the dimensions of the rectangle. *Hint:* The value of x in the length and width must be the same.

LESSON

12.6 Graphs of quadratic functions

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- identify the features of a parabolic graph
- plot the graph of a quadratic equation using a table of values
- use a parabolic graph to determine values.

12.6.1 Graphs of quadratic functions

eles-4936

- When quadratic functions are graphed, they produce curved lines called parabolas.
- The graph shown is a typical parabola with features as listed below.



- The dotted line is the **axis of symmetry**; the parabola is the same on either side of this line.
- The **turning point** is the point where the graph changes direction. It is located on the axis of symmetry.


- The turning point is a local minimum if it is the lowest point on the graph, and a local maximum if it is the highest point on the graph.
- The **x -intercept(s)** is where the graph crosses (or sometimes just touches) the x -axis. Not all parabolas have **x -intercepts** (roots).
- The y -intercept is where the graph crosses the y -axis. All parabolas have one y -intercept.

WORKED EXAMPLE 11 Describing features of parabolas


For each of the following graphs:

- state the equation of the axis of symmetry
- state the coordinates of the turning point
- indicate whether it is a maximum or a minimum turning point.

a.



b.



THINK

- State the equation of the vertical line that cuts the parabola in half.
 - State the turning point.
 - Determine the nature of the turning point by observing whether it is the highest or lowest point of the graph.
- State the equation of the vertical line that cuts the parabola in half.
 - State the turning point.
 - Determine the nature of the turning point by observing whether it is the highest or lowest point of the graph.


WRITE

- Axis of symmetry is $x = 1$.
 - The turning point is at $(1, -4)$.
 - Minimum turning point.
- Axis of symmetry is $x = -2$.
 - The turning point is at $(-2, 3)$.
 - Maximum turning point.


The x - and y -intercepts

- The x -intercept (roots) is where the graph crosses (or just touches) the x -axis.
- The y -intercept is where the graph crosses the y -axis. All parabolas have one y -intercept.
- When sketching a parabola, the x -intercepts (if any) and the y -intercept should always be marked on the graph, with their respective coordinates.


Quadratic graphs can have two solutions (or roots), one solution (or root) or no solutions, depending on the position of the quadratic graph. Examples of these three situations are shown.



Two solutions (roots)



One solution (root)





No solution

WORKED EXAMPLE 12 Determining axis intercepts for a parabolic graph

For each of the following graphs, state:

- the equation of the axis of symmetry
- the coordinates of the turning point and whether the point is a maximum or a minimum
- the x - and y -intercepts.



THINK

- a.
- State the equation of the vertical line that cuts the parabola in half.
 - State the turning point and its nature; that is, determine whether it is the highest or lowest point of the graph.
 - Observe where the parabola crosses the x -axis. In this case, the graph touches the x -axis when $x = 2$, so there is only one x -intercept.

Observe where the parabola crosses the y -axis.

- b.
- State the equation of the vertical line that cuts the parabola in half.
 - State the turning point and its nature; that is, determine whether it is the highest or lowest point of the graph.
 - Observe where the parabola crosses the x -axis.

Observe where the parabola crosses the y -axis.

WRITE

- a.
- Axis of symmetry is $x = 2$.
 - Maximum turning point is at $(2, 0)$.
 - The x -intercept is 2. It occurs at the point $(2, 0)$.

The y -intercept is -4 . It occurs at the point $(0, -4)$.

- b.
- Axis of symmetry is $x = -1$.
 - Minimum turning point is at $(-1, -1)$.
 - The x -intercepts are -2 and 0 . They occur at the points $(-2, 0)$ and $(0, 0)$. The y -intercept is 0 . It occurs at the point $(0, 0)$.

12.6.2 Plotting points to graph quadratic functions

- If there is a rule connecting y and x , a table of values can be used to determine actual coordinates.
- When drawing straight line graphs, a minimum of two points is required. For parabolas there is *no minimum* number of points, but between 6 and 12 points is a reasonable number.
- The more points used, the ‘smoother’ the parabola will appear. The points should be joined with a smooth curve, not ruled.
- Ensure that points plotted include (or are near) the main features of the parabola, namely the axis of symmetry, the turning point and the x - and y -intercepts.
- Occasionally a list of x -values will be provided and the corresponding y -values can be calculated. In the following example, the set of x -values is specified as $-4 \leq x \leq 2$.

WORKED EXAMPLE 13 Plotting points to graph quadratic functions

Plot the graph of $y = x^2 + 2x - 3$, $-4 \leq x \leq 2$ and, hence, state:

- the equation of the axis of symmetry
- the coordinates of the turning point and whether it is a maximum or a minimum
- the x - and y -intercepts.

THINK


- Write the equation.
- Complete a table of values by substituting into the equation each integer value of x from -4 to 2 .
For example, when $x = -4$, $y = (-4)^2 + 2 \times (-4) - 3 = 5$.
- List the coordinates of the points.
- Draw and label a set of axes, plot the points listed and join the points to form a smooth curve.

WRITE/DRAW

$$y = x^2 + 2x - 3$$

x	-4	-3	-2	-1	0	1	2
y	5	0	-3	-4	-3	0	5

$(-4, 5)$, $(-3, 0)$, $(-2, -3)$, $(-1, -4)$, $(0, -3)$, $(1, 0)$, $(2, 5)$



- Determine the equation of the line that divides the parabola exactly into two halves.
 - Determine the point where the graph turns or changes direction, and decide whether it is the highest or lowest point of the graph. State the coordinates of this point.
1. State the x -coordinates of the points where the graph crosses the x -axis.
 2. State the y -coordinate of the point where the graph crosses the y -axis.


- Axis of symmetry is $x = -1$.
- Minimum turning point is at $(-1, -4)$.

- The x -intercepts are at -3 and 1 . They occur at the points $(-3, 0)$ and $(1, 0)$.
The y -intercept is at -3 . It occurs at the point $(0, -3)$.


TI | THINK

- a. In a new problem, on a Graphs page, complete the function entry line as:
 $f1(x) = x^2 + 2x - 3$
 $-4 \leq x \leq 2$
 Then press ENTER.

DISPLAY/WRITE



- b. To determine the equation of the axis of symmetry, we need to know the x -coordinate of the turning point. Press:
- MENU
 - 6: Analyze Graph
 - 2: Minimum.
- Press ENTER to fix the lower bound, then drag the vertical line to the right to locate an upper bound and press ENTER. State the coordinates of the turning point and whether it is a maximum or a minimum.




The axis of symmetry is $x = -1$.
 The turning point occurs at $(-1, -4)$ and it is a minimum point.


CASIO | THINK

- a. 1. On the Graph & Table screen, complete the function entry line as:
 $y1 = x^2 + 2x - 3$
 $-4 \leq x \leq 2$
 Then press EXE. Press the icon.

DISPLAY/WRITE



2. To determine the equation of the axis of symmetry, we need to know the x -coordinate of the turning point. Press the icon.



The axis of symmetry is $x = -1$

- b. State the coordinates of the turning point and whether it is a maximum or a minimum.


- b. The turning point occurs at $(-1, -4)$ and it is a minimum point.

- A rule connecting x and y will be occasionally provided. From this rule, pairs of x - and y -values can be calculated. In the following example, the rule is given as $h = -5x^2 + 25x$.
- Graphs can be drawn using a graphics calculator, a graphing program on your computer or by hand.

WORKED EXAMPLE 14 Using graphs of quadratic functions

Rudie, the cannonball chicken, was fired out of a cannon. His path could be traced by the equation $h = -5x^2 + 25x$, where h is Rudie's height, in metres, above the ground and x is the horizontal distance, in metres, from the cannon.

Plot the graph for $0 \leq x \leq 5$ and use it to determine the maximum height of Rudie's path.



THINK

- Write the equation.
- Complete a table of values by substituting into the equation each integer value of x from 0 to 5.
For example, when $x = 0$,
 $h = -5 \times 0^2 + 25 \times 0 = 0$.
- List the coordinates of the points.
- As a parabola is symmetrical, the greatest value of h must be greater than 30 and occurs when x lies between 2 and 3, so determine the value of h when $x = 2.5$.
- Draw and label a set of axes, plot the points from the table and join the points to form a smooth curve.

- The maximum height is the value of h at the highest point of the graph.
- Write the answer in a sentence.


WRITE/DRAW

$$h = -5x^2 + 25x$$

x	1	2	3	4	5
h	0	20	30	30	20

$(0, 0), (1, 20), (2, 30), (3, 30), (4, 20), (5, 0)$

$$\begin{aligned} \text{When } x = 2.5, h &= 5 \times (2.5)^2 + 25 \times 2.5 \\ &= 31.25 \end{aligned}$$



$$h = 31.25$$

The maximum height of Rudie's path is 31.25 metres.

Resources



eWorkbook

Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivities

Individual pathway interactivity: Graphs of quadratic functions (int-8353)
The y -intercept (int-3837)

Individual pathways

PRACTISE

1, 4, 5, 9, 10, 11, 16, 19

CONSOLIDATE

2, 6, 8, 12, 14, 17, 20







MASTER

3, 7, 13, 15, 18, 21, 22

Fluency







1. **WE11** For each of the graphs below:

- i. state the equation of the axis of symmetry
- ii. give the coordinates of the turning point
- iii. indicate whether it is a minimum or maximum turning point.









2. For each of the following graphs, state:

- i. the equation of the axis of symmetry
- ii. the coordinates of the turning point
- iii. whether the turning point is a maximum or minimum.



3. **WE12** For each of the following graphs, state:

- the equation of the axis of symmetry
- the coordinates of the turning point and whether the point is a maximum or a minimum
- the x - and y -intercepts.



4. **MC** a. Identify the axis of symmetry for the graph shown.

- A. $x = 0$ B. $x = -2$ C. $x = -4$
 D. $y = 0$ E. the y -axis

b. Identify the coordinates of the turning point for the graph.


- A. $(0, 0)$ B. $(-2, 4)$ C. $(-2, -4)$
 D. $(2, -4)$ E. $(2, 4)$

c. Identify the y -intercept.

- A. 0 B. -2 C. -4
 D. 2 E. 4

d. Identify the x -intercepts.

- A. 0 and 4 B. 0 and 2 C. 0 and -2
 D. 0 and -4 E. -2 and -4



For questions 5 to 7, plot the graph of the quadratic equation and hence state:

- the equation of the axis of symmetry
- the coordinates of the turning point and whether it is a maximum or a minimum
- the x - and y -intercepts.

5. **WE13** a. $y = x^2 + 8x + 15, -7 \leq x \leq 0$

b. $y = x^2 - 1, -3 \leq x \leq 3$

6. a. $y = x^2 - 4x, -1 \leq x \leq 5$

b. $y = x^2 - 2x + 3, -2 \leq x \leq 4$

7. a. $y = x^2 + 12x + 35, -9 \leq x \leq 0$

b. $y = -x^2 + 4x + 5, -2 \leq x \leq 6$

8. Consider the equations for $-3 \leq x \leq 3$:

i. $y = x^2 + 2$

ii. $y = x^2 + 3$

- Make a table of values and plot the points on the same set of axes.
- State the equation of the axis of symmetry for each equation.
- State the x -intercepts for each equation.

Understanding

9. Consider the table of values below.

x	-4	-3	-2	-1	0	1	2	3	4
y	12	5	0	-3	-4	-3	0	5	12

- a. Plot these points on graph paper. State the shape of the graph.
b. Locate the axis of symmetry.
c. Locate the y -intercept.
d. Locate the x -intercept(s).
10. Consider the function $y = x^2 + x$. Complete this table of values for the function.

x	-6	-4	-2	0	2	4	6	8
y				0				

11. **MC** Select which of the following rules is not a parabola.
A. $y = -2x^2$ B. $y = 2x^2 - x$ C. $y = -2 \div x^2$ D. $y = -2 + x^2$ E. $y = 3x^2$
12. Consider the graph of $y = -x^2$.
a. State the turning point of this graph.
b. State whether the turning point is a maximum or a minimum.
13. Given the following information, make a sketch of the graph involved.
a. Maximum turning point = $(-2, -2)$, y -intercept = $(0, -6)$.
b. Minimum turning point = $(-3, -2)$, x -intercept $(1, 0)$, and $(-7, 0)$.
14. Sketch a graph where the turning point and the x -intercept are the same. Suggest a possible equation.
15. Sketch a graph where the turning point and the y -intercept are the same. Suggest a possible equation.

Reasoning


16. On a set of axes, sketch a parabola that has no x -intercepts and has an axis of symmetry $x = -2$. Explain if the parabola has a maximum or minimum turning point, or both.
17. If the axis of symmetry of a parabola is $x = -4$ and one of the x -intercepts is $(10, 0)$, show the other x -intercept is $(-18, 0)$.
18. If the x -intercepts of a parabola are $(-2, 0)$ and $(5, 0)$, show that the axis of symmetry is $x = 1.5$.

Problem solving

19. Consider the parabola given by the rule $y = x^2$ and the straight line given by $y = x$. Show that the two graphs meet at $(0, 0)$ and $(1, 1)$.
20. a. The axis of symmetry of a parabola is $x = -4$. If one x -intercept is -10 , determine the other x -intercept.
b. Suggest a possible equation for the parabola.
21. **WE14** A missile was fired from a boat during a test. The missile's path could be traced by the equation $h = -\frac{1}{2}x^2 + x$, where h is the missile's height above the ground, in kilometres, and x is the horizontal distance from the boat, in kilometres.
Plot the graph for $0 \leq x \leq 2$ and use it to evaluate the maximum height of the missile's path, in metres.

22. SpaceCorp sent a lander to Mars to measure the temperature change over a period of time. The results were plotted on a set of axes, as shown. From the graph it can be seen that the temperature change follows the quadratic rule $T = -h^2 + 22h - 21$, where T is the temperature in degrees Celsius, and the time elapsed, h , is in hours.

- State the initial recorded temperature on Mars.
- State when the temperature measured 0°C .
- Identify when the highest temperature was recorded.
- State the highest temperature recorded.



LESSON

12.7 Sketching parabolas of the form $y = ax^2$

LEARNING INTENTIONS


At the end of this lesson you should be able to:

- sketch parabolas for quadratic equations of the form $y = ax^2$
- describe the features of a parabola of the form $y = ax^2$
- describe the effect of changing the value of a on parabolas of the form $y = ax^2$.

12.7.1 The graph of the quadratic function $y = x^2$

eles-4938

- The simplest parabola, $y = x^2$, is shown.




- Both the x - and y -axes are clearly indicated, along with their scales.
- The turning point $(0, 0)$ is indicated.
- The x - and y -intercepts are indicated. For this graph they are all $(0, 0)$.
- This is an example of a parabola that just touches (does not cross) the x -axis at $(0, 0)$.

12.7.2 Parabolas of the form $y = ax^2$, where $a > 0$

eles-4939

- A coefficient in front of the x^2 term affects the **dilation** of the graph, making it wider or narrower than the graph of $y = x^2$.
- If $a > 1$ then the graph becomes narrower, whereas if $0 < a < 1$, the graph becomes wider.
- The graph shows the effect of varying the coefficient a .



Notice that $y = 3x^2$ is narrower than $y = x^2$ and $y = \frac{1}{4}x^2$ is wider than $y = x^2$.

Parabolas of the form $y = ax^2$, where $a > 0$

For all parabolas of the form $y = ax^2$, where $a > 0$:

- the axis of symmetry is $x = 0$
- the turning point is $(0, 0)$
- the x -intercept is $(0, 0)$
- the y -intercept is $(0, 0)$
- the shape of the parabola is always upright, or a U shape (U).

WORKED EXAMPLE 15 Sketching dilation of parabolas with $a > 0$

On the same set of axes, sketch the graph of $y = x^2$ and $y = 3x^2$, marking the coordinates of the turning point and the intercepts. State which graph is narrower.

THINK

1. Write the equation of the first graph.
2. State its axis of symmetry.
3. State the coordinates of the turning point.
4. State the intercepts.
5. Calculate the coordinates of one other point.
6. Write the equation of the second graph.
7. State its axis of symmetry.
8. State the coordinates of the turning point.
9. State the intercepts.
10. Calculate the coordinates of one other point.

WRITE/DRAW

$$y = x^2$$

The axis of symmetry is $x = 0$.

The turning point is $(0, 0)$.

The x -intercept is $(0, 0)$ and the y -intercept is also $(0, 0)$.

When $x = 1$, $y = 1$; $(1, 1)$

$$y = 3x^2$$


The axis of symmetry is $x = 0$.

The turning point is $(0, 0)$.

The x -intercept is 0 and the y -intercept is 0 .

When $x = 1$, $y = 3$; $(1, 3)$

11. Sketch the graphs, labelling the turning point.



12. State the graph that is narrower.

The graph of $y = 3x^2$ is narrower.


12.7.3 Parabolas of the form $y = ax^2$, where $a < 0$

eles-4940

- When $a < 0$, the graph is inverted; that is, it is \cap shaped.
- A coefficient in front of the x^2 term affects the dilation of the graph, making it wider or narrower than the graph of $y = x^2$.
- If $-1 < a < 0$, the graph is wider than $y = -x^2$.
- If $a < -1$, the graph is narrower than $y = -x^2$.

- The graph shows the effect of varying the coefficient a .

Notice that $y = -3x^2$ is narrower than $y = -x^2$ and $y = -\frac{1}{4}x^2$ is wider than $y = -x^2$.



Parabolas of the form $y = ax^2$, where $a < 0$

For all parabolas of the form $y = ax^2$, where $a < 0$:

- the axis of symmetry is $x = 0$
- the turning point is $(0, 0)$
- the x -intercept is $(0, 0)$
- the y -intercept is $(0, 0)$
- the shape of the parabola is always inverted or an upside-down U shape (\cap).

WORKED EXAMPLE 16 Sketching dilation of parabolas with $a < 0$

On the same set of axes sketch the graphs of $y = -x^2$ and $y = -2x^2$, marking the coordinates of the turning point and the intercept. State which graph is narrower.

THINK

1. Write the equation of the first graph.
2. State its axis of symmetry.
3. State the coordinates of the turning point.
4. State the intercepts.

WRITE/DRAW


$$y = -x^2$$

The axis of symmetry is $x = 0$.

The turning point is $(0, 0)$.

The x -intercept is $(0, 0)$ and the y -intercept is also $(0, 0)$.

5. Calculate the coordinates of one other point. When $x = 1$, $y = -1$; $(1, -1)$
6. Write the equation of the second graph. $y = -2x^2$
7. State its axis of symmetry. The axis of symmetry is $x = 0$.
8. State the coordinates of the turning point. The turning point is $(0, 0)$.
9. State the intercepts. The x -intercept is $(0, 0)$ and the y -intercept is also $(0, 0)$.
10. Calculate the coordinates of one other point. When $x = 1$; $y = -2$; $(1, -2)$
11. Sketch the two graphs on a single set of axes, labelling the turning point (as well as intercepts and maximum).



12. State which graph is narrower.

The graph of $y = -2x^2$ is narrower.

on Resources



eWorkbook Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivities Individual pathway interactivity: Sketching parabolas of the form $y = ax^2$ (int-8354)
Dilation of parabolas (int-6096)

Exercise 12.7 Sketching parabolas of the form $y = ax^2$

learn on

12.7 Quick quiz on

12.7 Exercise

Individual pathways

■ PRACTISE

1, 5, 7, 11, 14

■ CONSOLIDATE

2, 4, 8, 12, 15

■ MASTER

3, 6, 9, 10, 13, 16

Fluency

- WE15** On the same set of axes, sketch the graph of $y = x^2$ and $y = 4x^2$, marking the coordinates of the turning point and the intercepts. State which graph is narrower.
- On the same set of axes, sketch the graph of $y = x^2$ and $y = \frac{1}{2}x^2$, marking the coordinates of the turning point and the intercepts. State which graph is narrower.
- Sketch the graph of the following table. State the equation of the graph.

x	-4	-3	-2	-1	0	1	2	3	4
y	4	2.25	1	0.25	0	0.25	1	2.25	4

4. **WE16** a. Using the same set of axes, sketch the graphs of $y = -x^2$ and $y = -0.5x^2$, marking the coordinates of the turning point and the intercepts. State which graph is narrower.
- b. Using the same set of axes, sketch the graph of $y = -5x^2$, marking the coordinates of the turning point and the intercepts. State which graph is the narrowest. State which graph is the narrowest.
5. **MC** a. The graph of $y = -3x^2$ is:
- A. wider than $y = x^2$
 - B. narrower than $y = x^2$
 - C. the same width as $y = x^2$
 - D. a reflection of $y = x^2$ in the x -axis.
 - E. a reflection of $y = x^2$ in the y -axis.
- b. The graph of $y = \frac{1}{3}x^2$ is:
- A. wider than $y = x^2$
 - B. narrower than $y = x^2$
 - C. the same width as $y = x^2$
 - D. a reflection of $y = x^2$ in the x -axis.
 - E. a reflection of $y = x^2$ in the y -axis.
- c. The graph of $y = \frac{1}{2}x^2$ is:
- A. wider than $y = \frac{1}{4}x^2$
 - B. narrower than $y = \frac{1}{4}x^2$
 - C. the same width as $y = \frac{1}{4}x^2$
 - D. a reflection of $y = \frac{1}{4}x^2$ in the x -axis.
 - E. a reflection of $y = x^2$ in the y -axis.

6. Match each of the following parabolas with the appropriate equation from the list.

i. $y = 3x^2$







ii. $y = -x^2$

iii. $y = 4x^2$

iv. $y = \frac{1}{2}x^2$

v. $y = -4x^2$

vi. $y = -2x^2$



Understanding

7. Write an equation for a parabola that has a minimum turning point and is narrower than $y = x^2$.
8. Write an equation for a parabola that has a maximum turning point and is wider than $y = x^2$.

LESSON

12.8 Sketching parabolas of the form $y = (x - h)^2 + k$

LEARNING INTENTIONS

At the end of this lesson you should be able to:


- determine the axes of symmetry, y -intercepts and turning points for graphs of the form $y = ax^2 + c$ and $y = (x - h)^2 + k$
- sketch graphs of the form $y = ax^2 + c$, $y = (x - h)^2$ and $y = (x - h)^2 + k$
- describe the transformations that transform the graph of $y = x^2$ into $y = (x - h)^2 + k$.



eles-4941

12.8.1 Sketching parabolas of the form $y = ax^2 + c$

- A parabola with the equation $y = x^2 + c$ is simply the graph of $y = x^2$ moved vertically by c units.
- If $c > 0$, the graph of $y = x^2$ moves up by c units.
- If $c < 0$, the graph of $y = x^2$ moves down by c units.
- The turning point of such a parabola is $(0, c)$.



WORKED EXAMPLE 17 Sketching parabolas of the form $y = x^2 + c$ and $y = -x^2 + c$

For each part of the question, sketch the graph of $y = x^2$, then, on the same axes, sketch the given graph, clearly labelling the turning point.


a. $y = x^2 + 2$

b. $y = -x^2 - 3$

THINK

- a. 1. Sketch the graph of $y = x^2$ by drawing a set of labelled axes, marking the turning point $(0, 0)$ and noting that it is symmetrical about the y -axis.


WRITE/DRAW




2. Determine the turning point of $y = x^2 + 2$ by adding 2 to the y -coordinate of the turning point of $y = x^2$.

The turning point of $y = x^2 + 2$ is $(0, 2)$.

3. Using the same axes as for the graph of $y = x^2$, sketch the graph of $y = x^2 + 2$, marking the turning point and making sure that it is the same width as the graph of $y = x^2$. (The coefficient of x^2 is the same for both graphs.)




- b. 1. Sketch the graph of $y = x^2$ by drawing a set of labelled axes, marking the turning point $(0, 0)$ and noting that it is symmetrical about the y -axis.



2. Determine the turning point of $y = -x^2 - 3$ by subtracting 3 from the y -coordinate of the turning point of $y = x^2$.
3. Using the same axes as for the graph of $y = x^2$, sketch the graph of $y = -x^2 - 3$, marking the turning point, inverting the graph and making sure that the graph is the same width as the graph of $y = x^2$.


The turning point of $y = -x^2 - 3$ is $(-0, -3)$.



12.8.2 Sketching parabolas of the form $y = (x - h)^2$

eles-4942

- Subtracting a constant value h from x in $y = x^2$ **translates** the parabola h units to the right. The parabola's shape is otherwise unchanged.
 - The equation is $y = (x - h)^2$, where h is a positive quantity.
 - The x -intercept occurs when $x = h$.
- Adding a constant value h to x in $y = x^2$ translates the parabola h units to the left; otherwise the parabola's shape is unchanged.
 - The equation is $y = (x + h)^2$, where h is a positive quantity.
 - The x -intercept occurs when $x = -h$.
- The y -intercept occurs when $x = 0$, and is always at $y = h^2$.



WORKED EXAMPLE 18 Sketching parabolas of the form $y = (x - h)^2$

On clearly labelled axes, sketch the graph of $y = (x - 2)^2$, marking the turning point and y-intercept. State whether the turning point is a maximum or a minimum.

THINK

1. Write the equation.
2. State the axis of symmetry ($x = h$ where h is 2).
3. State the turning point, which has been moved to the right.
4. The sign in front of the bracket is positive so the parabola is upright.
5. Determine the y-intercept by substituting $x = 0$ into the equation.
6. Draw a clearly labelled set of axes, mark the turning point and y-intercept and draw the graph of $y = (x - 2)^2$. Note that the sign in the brackets is negative so the graph moves 2 units to the right.

WRITE/DRAW

$$y = (x - 2)^2$$

The axis of symmetry is $x = 2$.


The turning point is $(2, 0)$.

Minimum turning point.

y-intercept: when $x = 0$,

$$\begin{aligned} y &= (0 - 2)^2 \\ &= 4 \end{aligned}$$

The y-intercept is 4.



WORKED EXAMPLE 19 Sketching parabolas of the form $y = -(x + h)^2$

Sketch the graph of $y = -(x + 1)^2$, labelling the turning point, stating whether it is a maximum or minimum, and determining the y-intercept and the equation of the axis of symmetry.

THINK

1. Write the equation.
2. State the turning point, which has been moved to the left.
3. The sign in front of the bracket is negative so the parabola is inverted.
4. Determine the y-intercept by substituting $x = 0$ into the equation.
5. State the axis of symmetry ($x = h$, where h is -1).

WRITE/DRAW

$$y = -(x + 1)^2$$

The turning point is $(-1, 0)$.

Maximum turning point.


y-intercept: when $x = 0$,

$$\begin{aligned} y &= -(0 + 1)^2 \\ &= -1 \end{aligned}$$

The y-intercept is -1 .

The axis of symmetry is $x = -1$.

6. Draw a clearly labelled set of axes, mark the turning point and y-intercept and draw the graph of $y = -(x + 1)^2$.




12.8.3 Sketching parabolas of the form $y = (x - h)^2 + k$

- The equation $y = (x - h)^2 + k$ combines a vertical translation of k and a horizontal translation of h together.
- The equation $y = (x - h)^2 + k$ is called **turning point form** because the turning point is given by the coordinates (h, k) .

Sketching the graph of $y = (x - h)^2 + k$

1. Identify the values of h and k .
2. Mark the following:
 - the turning point at (h, k)
 - y-intercept at $(0, h^2 + k)$
 - axis of symmetry at $x = h$.
3. Calculate the positions of the x-intercepts by setting $y = 0$ and solving for x .
Mark their positions on the x-axis.
4. Connect the points you have marked with a single neat curve to form a parabola.



WORKED EXAMPLE 20 Sketching graphs of the form $y = (x + h)^2 - k$

Sketch the graph of $y = (x + 2)^2 - 1$, marking the turning point and the y-intercept, and indicate the type of turning point.

THINK

1. Write the equation.
2. State the turning point. As the equation is in the form $y = (x - h)^2 + k$, the turning point is (h, k) .
3. There is no sign outside the brackets, so the parabola is upright.
4. Determine the y-intercept by substituting $x = 0$ into the equation.

WRITE/DRAW

$$y = (x + 2)^2 - 1$$


The turning point is $(-2, -1)$.

Minimum turning point.

y-intercept: when $x = 0$,

$$y = (0 + 2)^2 - 1 = 3$$

5. Draw clearly labelled axes, mark the coordinates of the turning point, the y-intercept and draw a symmetrical graph.



- Note that in the previous example the graph has shifted 2 units left ($h = 2$) and 1 unit down ($k = -1$).
- The graph has the same shape as $y = x^2$.

WORKED EXAMPLE 21 Sketching variations of the $y = x^2$ graph

On the same set of axes, sketch the graphs of each of the following, clearly marking the coordinates of the turning point and the y-intercept:


- $y = x^2$
- $y = (x - 2)^2$
- $y = (x - 2)^2 + 1$.

State the changes that are made from **a** to **b** and from **a** to **c**.

THINK

- Sketch the graph of $y = x^2$, marking the coordinates of the turning point and the y-intercept.

WRITE/DRAW




1. Write the equation.
2. Determine the coordinates of the turning point.
3. Calculate the y-intercept.

- $y = (x - 2)^2$
The turning point is $(2, 0)$.
y-intercept: when $x = 0$,

$$y = (0 - 2)^2 = 4$$

4. On the same set of axes, sketch the graph of $y = (x - 2)^2$, marking the coordinates of the turning point and the y-intercept.



- c. 1. Write the equation.
2. Determine the coordinates of the turning point.
3. Calculate the y-intercept.


c. $y = (x - 2)^2 + 1$

The turning point is $(2, 1)$.

y-intercept: when $x = 0$,

$$\begin{aligned} y &= (0 - 2)^2 + 1 \\ &= 4 + 1 \\ &= 5 \end{aligned}$$

4. On the same set of axes, sketch the graph of $y = (x - 2)^2 + 1$, marking the coordinates of the turning point and the y-intercept.



5. State how $y = x^2$ is changed to form $y = (x - 2)^2$.

If $y = x^2$ is moved 2 units to the right, the resulting graph is $y = (x - 2)^2$.


State how $y = x^2$ is changed to form $y = (x - 2)^2 + 1$.

If $y = x^2$ is moved 2 units to the right and 1 unit up, the resulting graph is $y = (x - 2)^2 + 1$.

TI | THINK

- a-c. In a new problem, on a Graphs page, complete the function entry lines as:
- $$f1(x) = x^2$$
- $$f2(x) = (x - 2)^2$$
- $$f3(x) = (x - 2)^2 + 1$$
- Press ENTER after each entry.

DISPLAY/WRITE




- a-c. The graph of $y = x^2$ has a turning point at $(0, 0)$. If this graph is moved 2 units to the right, it is the graph of $y = (x - 2)^2$, which has a turning point at $(2, 0)$. If this graph is moved 1 unit up, it is the graph of $y = (x - 2)^2 + 1$, which has a turning point at $(2, 1)$.

CASIO | THINK

- a-c. On the Graph & Table screen, complete the function entry lines as:
- $$y1 = x^2$$
- $$y2 = (x - 2)^2$$
- $$y3 = (x - 2)^2 + 1$$
- Press EXE after each entry. Press the Graph icon to view the graph.

DISPLAY/WRITE




- a-c. If $y = (x - h)^2 + k$, then h controls the horizontal translation of the graphs and k controls the vertical translations.

Digital technology

Using Desmos


In many cases it can be very helpful to use graphing software. One program that can be accessed free of charge is Desmos.

- Go to www.desmos.com.
- Under **Maths Tools**, select **graphing calculator**.
- You will see the following screen:




- Enter one of following equations into the box at top left, then click the + icon to create a new box, and repeat for the remaining equations: $y = x^2$; $y = 2x^2$; $y = -2x^2$.

- You will see the three parabolas appear. Note that the colours of the parabolas match the coloured circles in front of each of your equations in the panel.



- Clear parabolas from the graph space by clicking \times at right of each equation box.
- Now try the equation $y = (x - 3)^2 - 2$.
- Clicking on the grey circles at intersection points on the graph will display their coordinates. Below we can see the coordinates for the turning point, the x -intercepts and the y -intercept.



Resources

- eWorkbook** Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)
- Interactivities** Individual pathway interactivity: Sketching parabolas of the form $y = (x - h)^2 + k$ (int-8355)
 - Horizontal translations of parabolas (int-6054)
 - Vertical translations of parabolas (int-6055)
 - Turning point form (int-6099)

Exercise 12.8 Sketching parabolas of the form $y = (x - h)^2 + k$ learnon

12.8 Quick quiz **on**

12.8 Exercise

Individual pathways

PRACTISE

1, 4, 9, 10, 13, 16, 17, 20, 24, 27, 31, 36

CONSOLIDATE

2, 5, 7, 11, 14, 18, 21, 23, 26, 29, 32, 34, 37, 38

MASTER

3, 6, 8, 12, 15, 19, 22, 25, 28, 30, 33, 35, 39, 40

Fluency

1. **WE17** For each part of the question, sketch the graph of $y = x^2$; then, on the same axes, sketch the given graph, clearly labelling the turning point.

a. $y = x^2 + 1$

b. $y = x^2 - 4$

2. For each part of the question, sketch the graph of $y = x^2$; then, on the same axes, sketch the given graph, clearly labelling the turning point.

a. $y = x^2 + 4$

b. $y = -x^2 - 1$

3. For each part of the question, sketch the graph of $y = x^2$; then, on the same axes, sketch the given graph, clearly labelling the turning point.

a. $y = x^2 - 1$

b. $y = -x^2 + 1$

For questions 4 to 6, sketch the following graphs on clearly labelled axes, marking the turning point. State whether the turning point is a maximum or a minimum.

4. a. $y = x^2 + 2$

b. $y = -x^2 + 3$

5. a. $y = x^2 - 5$

b. $y = -x^2 + 4$

6. a. $y = -x^2 - 3$

b. $y = x^2 - \frac{1}{2}$

7. Sketch the following graphs, indicating the turning point and estimating the x -intercepts.

a. $y = 4 - x^2$

b. $y = -4 - x^2$

c. $y = 1 - x^2$

8. a. Does the turning point change if there is a negative number in front of the x^2 term in the equation $y = x^2$?

b. Explain how a negative coefficient of x^2 affects the graph.

9. **MC** a. The turning point for the graph of the equation $y = -x^2 + 8$ is:

A. (0, 0)

B. (-1, 8)

C. (0, 8)

D. (0, -8)

E. (1, 8)

b. The turning point of the graph of the equation $y = x^2 - 16$ is:

A. (1, -16)

B. (-16, 1)

C. (0, 0)

D. (0, -16)

E. (-1, 16)

c. The graph of $y = x^2 - 7$ moves the graph of $y = x^2$ in the following way:

A. Up 1

B. Down 1

C. Up 7

D. Down 7

E. Left 7

d. The y -intercept of the graph of $y = -x^2 - 6$ is:

A. 1

B. -1

C. 6

D. -6

E. -7

WE18 For questions 10–12, sketch the graphs of each of the following on clearly labelled axes, marking the turning point and y -intercept. State whether the turning point is a maximum or a minimum.

10. a. $y = (x - 1)^2$

b. $y = (x + 2)^2$

11. a. $y = (x - 3)^2$

b. $y = (x + 4)^2$

12. a. $y = (x - 5)^2$

b. $y = (x + 6)^2$

WE19 For questions 13–15, sketch the graphs of each of the following on clearly labelled axes, marking the turning point and y-intercept. State whether the turning point is a maximum or a minimum.

13. a. $y = -(x - 1)^2$

b. $y = -(x + 2)^2$

14. a. $y = -(x - 3)^2$

b. $y = -(x + 4)^2$

15. a. $y = -(x - 5)^2$

b. $y = -(x + 6)^2$

16. **MC** a. The axis of symmetry for the graph $y = (x + 3)^2$ is:

A. $y = 0$

B. $x = 3$

C. $x = -3$

D. $x = 2$

E. $x = 0$

b. The turning point of the graph $y = (x + 3)^2$ is:

A. (0, 0)

B. (1, 3)

C. (1, -3)

D. (-3, 0)

E. (3, 0)

WE20 For questions 17–19, sketch the graph of each of the following, marking the turning point, the type of turning point and the y-intercept.

17. a. $y = (x - 1)^2 + 1$

b. $y = (x + 2)^2 - 1$

18. a. $y = (x - 3)^2 + 2$

b. $y = (x + 3)^2 - 2$

19. a. $y = -(x + 2)^2 - 1$

b. $y = -(x - 1)^2 + 2$

20. **WE21** On the same set of axes, sketch the following graphs, clearly marking the coordinates of the turning points and the y-intercepts. State the changes that are made from a to b and from a to c.

a. $y = x^2$

b. $y = (x - 1)^2$

c. $y = (x - 1)^2 + 3$

21. On the same set of axes, sketch the following graphs, clearly marking the coordinates of the turning points and the y-intercepts. State the changes that are made from a to b and from a to c.

a. $y = x^2$

b. $y = (x + 3)^2$

c. $y = (x + 3)^2 + 2$

22. On the same set of axes, sketch the following graphs, clearly marking the coordinates of the turning points and the y-intercepts. State the changes that are made from a to b and from a to c.

a. $y = x^2$

b. $y = (x - 4)^2$

c. $y = (x - 1)^2 - 1$

23. **MC** a. For the graph of $y = (x + 5)^2 - 2$, the coordinates of the turning point are:

A. (5, -2)

C. (-2, -5)

E. (-5, 2)

B. (-2, 5)

D. (-5, -2)

b. For the graph of $y = -(x - 3)^2 + 7$, the axis of symmetry is:

A. $x = 3$

C. $x = -3$

E. $x = -7$

B. $x = 7$

D. $y = 3$

c. For the graph of $y = (x - 1)^2 - 4$, the y-intercept is:

A. 1

C. -3

E. -4

B. -1

D. 4

Understanding

24. Match each of the following parabolas with the appropriate equation from the list.

i. $y = x^2 - 3$







ii. $y = x^2 + 3$

iii. $y = 3 - x^2$

iv. $y = x^2 + 2$

v. $y = -x^2 + 2$

vi. $y = -x^2 - 2$









25. The vertical cross-section through the top of the mountain called the Devil's Tower can be approximated by the graph $y = -x^2 + 5$.
- Sketch the graph.
 - If the x -axis represents the sea level, and both x and y are in kilometres, calculate the maximum height of the mountain.

26. Match each of the following parabolas with the appropriate equation from the list.

i. $y = (x - 2)^2$
iv. $y = -(x + 2)^2$

ii. $y = x^2$
v. $y = (x - 3)^2$

iii. $y = (x + 3)^2$
vi. $y = -(x - 2)^2$



27. Determine the equation of the parabola that is of the form $y = (x - h)^2$ and passes through:
- (3, 1)
 - (-1, 9).

28. State the differences, if any, between the following pairs of parabolas.

- $y = (1 - x)^2$ and $y = (x - 1)^2$
- $y = (x - 2)^2$ and $y = (x + 2)^2$
- $y = -(x - 4)^2$ and $y = (4 - x)^2$


29. Write the equations for the following transformations on $y = x^2$.
- Reflection in the x -axis and translation of 2 units to the left.
 - Translation of 1 unit down and 3 units to the right.
 - Translation of 1 unit left and 2 units up.
 - Reflection in the x -axis, translation of 4 units to the right and 6 units up.
30. **MC** A parabola has an equation of $y = x^2 - 6x + 14$. Its turning point form is:
- $y = (x - 6)^2 + 14$
 - $y = (x - 14)^2 - 6$
 - $y = (x - 3)^2 + 5$
 - $y = (x - 5)^2 + 3$
 - $y = (x + 3)^2 + 14$

Reasoning

31. Explain why the equation presented in the form $y = (x - h)^2 + k$ is known as 'turning point form'.
32. Show that the equation of the parabola that is of the form $y = x^2 + c$ and passes through:
- $(2, 1)$ is $y = x^2 - 3$
 - $(-3, -1)$ is $y = x^2 - 10$.

33. The figure below shows the span of the Gateshead Millennium Bridge in England.

A set of axes has been superimposed onto the photo. Use the coordinates $(200, 0)$ and $(0, -75)$ to show that a possible equation is $y = -0.001\,875(x - 200)^2$.



34. Sketch $y = x^2 - 2$ and $y = -x^2 + 2$ on the same set of axes. Use algebra to explain where they intersect.
35. **a.** Describe how to transform the parabola $y = (x - 3)^2$ to obtain the parabolas $y = (x - 3)^2 - 2$ and $y = (x - 3)^2 + 1$.
- b.** Another parabola is created by moving $y = (x - 3)^2$ so that its turning point is $(5, -4)$. Write an equation for this parabola.

Problem solving

36. Nikki wanted to keep a carnivorous plant, so after school she recorded the temperature on her windowsill for 8 hours every day for several months. One summer evening the temperature followed the relationship $t = (h - 5)^2 + 15$, where t is the temperature in degrees Celsius, h hours after 4 pm.
- Calculate the temperature on the windowsill at 4 pm.
 - Show that the minimum temperature reached during the 8-hour period is 15°C .
 - Calculate the number of hours it took for the windowsill to reach the minimum temperature.
 - Sketch a graph of the relationship between the temperature and the number of hours after recording began. Mark the turning point and the t -intercept on the graph.
37. For the equation $y = x^2 - 5$, calculate the exact value of the y -coordinate when the x -coordinate is $2 + \sqrt{5}$.
38. A rocket is shot in the air from a given point $(0, 0)$. The rocket follows a parabolic trajectory. It reaches a maximum height of 400 m and lands 300 m away from the launching point.
- Calculate how far horizontally from the launching point the rocket is when it reaches its maximum height.
 - State the equation of the path of the rocket.
39. The path of a ball rolling off the end of a table follows a parabolic curve and can be modelled by the equation $y = ax^2 + c$. A student rolls a ball off a tabletop that is 128 cm above the floor, and the ball lands 80 cm horizontally away from the desk.
- If the student sets a cup 78 cm above the floor to catch the ball during its fall, determine where the cup should be placed.
40. Use the three points $(-2, 1)$, $(3, 1)$ and $(0, 7)$ to determine a , h and k in the equation $y = a(x - h)^2 + k$.

LESSON

12.9 Sketching parabolas of the form $y = (x + a)(x + b)$ (extending)

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- identify the x - and y -intercepts, the axis of symmetry and the turning point of parabolas of the form $y = (x + a)(x + b)$
- factorise quadratic equations of the form $y = x^2 + bx + c$ to sketch them using their x -intercepts.



eles-4944

12.9.1 Sketching parabolas of the form $y = (x + a)(x + b)$

- The equation $y = (x + a)(x + b)$ consists of a pair of linear factors $(x + a)$ and $(x + b)$ multiplied together.

Determining important points to sketch the $y = (x + a)(x + b)$ parabola

- The x -intercepts are found by setting each factor to 0:


$$(x + a) = 0, x = -a$$

$$(x + b) = 0, x = -b$$

- The y -intercept can be found by letting $x = 0$ in the original equation:

$$y = (0 + a)(0 + b) = ab$$

- The axis of symmetry is halfway between the x -intercepts $-a$ and $-b$; that is: $x = -\frac{a + b}{2}$.
- Substitute the x -value of the axis of symmetry into $y = (x + a)(x + b)$ to determine the y -coordinate of the turning point.



WORKED EXAMPLE 22 Sketching parabolas in intercept form

Sketch the graph of $y = (x - 4)(x + 2)$ by first determining the x - and y -intercepts and then the turning point.

THINK

1. Write the equation.
2. Determine the x -intercepts.
3. Determine the y -intercept.

WRITE/DRAW

$$y = (x - 4)(x + 2)$$

x -intercepts: when $y = 0$,

$$(x - 4)(x + 2) = 0$$

$$x - 4 = 0 \quad \text{or} \quad x + 2 = 0$$

$$x = 4 \quad \quad \quad x = -2$$

The x -intercepts are -2 and 4 .

y -intercept: when $x = 0$,

$$y = (0 - 4)(0 + 2)$$

$$= -4 \times 2$$

$$= -8$$


The y -intercept is -8 .

- Determine the x -value of the turning point by averaging the values of the two x -intercepts.
- Determine the y -value of the turning point by substituting the x -value of the turning point into the equation of the graph.
- State the coordinates of the turning point.
- Sketch the graph.

At the turning point, $x = \frac{4 + (-2)}{2} = 1$.

When $x = 1$,
 $y = (1 - 4)(1 + 2)$
 $= -3 \times 3$
 $= -9$

The turning point is $(1, -9)$.



- Sometimes, it is possible to rearrange equations in the general form ($y = x^2 + bx + c$) so that they are in intercept or factor form $y = (x + a)(x + b)$.

WORKED EXAMPLE 23 Sketching parabolas of the form $y = x^2 + bx + c$

Sketch the graph of $y = x^2 + 6x + 8$.

THINK

- Write the equation.
- Factorise the expression on the right-hand side of the equation, $x^2 + 6x + 8$. Remember that to factorise an equation, you should determine the factors of c that add up to b .
- Determine the two x -intercepts.
- Determine the y -intercept.
- Determine the x -value of the turning point.

WRITE/DRAW

$$y = x^2 + 6x + 8$$

$$= (x + 2)(x + 4)$$

x -intercepts: when $y = 0$,
 $(x + 2)(x + 4) = 0$
 $x + 2 = 0$ or $x + 4 = 0$
 $x = -2$ $x = -4$
 The x -intercepts are -4 and -2 .

y -intercept: when $x = 0$,
 $y = 0 + 0 + 8$
 $= 8$
 The y -intercept is 8 .

At the turning point, $x = \frac{-2 + (-4)}{2}$
 $= -3$

6. Determine the y -value of the turning point.


When $x = -3$

$$\begin{aligned}y &= (-3 + 2)(-3 + 4) \\ &= -1 \times 1 \\ &= -1\end{aligned}$$

7. State the turning point.

The turning point is $(-3, -1)$.

8. Sketch the graph.



on Resources



eWorkbook Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivity Individual pathway interactivity: Sketching parabolas of the form $y = (x + a)(x + b)$ (int-8356)

Parabolas of the form $y = (x + a)(x + b)$ (int-6100)

Exercise 12.9 Sketching parabolas of the form $y = (x + a)(x + b)$ (extending)

learn on

12.9 Quick quiz on

12.9 Exercise

Individual pathways

PRACTISE

1, 4, 7, 10, 11, 15, 18

CONSOLIDATE

2, 5, 8, 12, 14, 16, 19

MASTER

3, 6, 9, 13, 17, 20

Fluency

WE22 For questions 1–3, sketch the graph of each of the following by first determining the x - and y -intercepts and then the turning point.

1. a. $y = (x + 2)(x + 6)$

b. $y = (x - 3)(x - 5)$

2. a. $y = (x - 3)(x + 1)$

b. $y = (x + 4)(x - 6)$

3. a. $y = (x - 5)(x + 1)$

b. $y = (x + 1)(x - 2)$

WE23 For questions 4–9, sketch the graph of each of the following.

4. a. $y = x^2 + 8x + 12$

b. $y = x^2 + 8x + 15$

5. a. $y = x^2 + 6x + 5$

b. $y = x^2 + 10x + 16$

6. a. $y = x^2 + 2x - 8$

b. $y = x^2 + 4x - 5$

7. a. $y = x^2 - 4x - 12$

b. $y = x^2 - 6x - 7$

8. a. $y = x^2 - 10x + 24$

b. $y = x^2 - 6x + 5$

9. a. $y = x^2 - 5x + 6$

b. $y = x^2 - 3x - 10$

Understanding

10. Sketch the graph of a parabola whose x -intercepts are at $x = -4$ and $x = 6$ and whose y -intercept is at $y = 2$.
11. State a possible equation of a graph with an axis of symmetry at $x = 4$ and one intercept at $x = 6$.
12. Out on the cricket field, Michael Clarke chases the ball. He picks it up, runs 4 metres towards the stumps (which are 25 metres away from where he picks up the ball). He then throws the ball, which follows the path described by the quadratic equation $y = -0.1(x - m)(x - n)$, where m and n are positive integers. The ball lands 1 metre from the stumps, where another player quickly scoops it up and removes the bails. Take the origin as the point where Michael picked up the ball initially.
- Calculate how far the ball has travelled horizontally while in flight.
 - Calculate the values of m and n .
 - Calculate the x -intercepts.
 - Calculate the axis of symmetry.
 - Calculate the turning point.
 - Identify the highest point reached by the ball.
 - Determine how far horizontally the ball has travelled when it reaches its highest point.
 - Sketch the flight of the ball, showing all of the relevant details on the graph.

13. McDonald's golden arches were designed by Jim Schindler in 1962. The two arches each approximate the shape of a parabola. A large McDonald's sign stands on the roof of a shopping centre. The shape of one of the parabolas can be modelled by the quadratic function $y = -x^2 + 8x - 7$, where both x and y are measured in metres and the bottom of the outer edges of the arches touches the x -axis. The complete sign is supported by a beam underneath the arches. Determine the minimum length required for this beam.




14. For the equation $y = (x - 3)(x - 9)$:
- determine the turning point
 - rewrite the equation in turning point form
 - expand both forms of the equation, showing that they are equivalent
 - sketch the equation showing the x -intercepts and turning point.

Reasoning

15. The dimensions of a rectangular backyard can be given by $(x + 2)$ m and $(x - 4)$ m. Determine the value of x if the yard's area is 91 m^2 .

16. The height of an object, $h(t)$, thrown into the air is determined by the formula $h(t) = -8t^2 + 128t$, where t is time in seconds and h is height in metres.
- Explain if the graph of the formula has a maximum or a minimum turning point.
 - Calculate the maximum height of the object and the time that this height is reached.

17. Daniel is in a car on a roller-coaster ride as shown in the following graph, where the height, h , is in metres above the ground and time for the ride, t , is in minutes.



- The ride can be represented by three separate equations. Show that the first section of the ride is $h = 2t$ from $t = 0$ to $t = 6$. The middle section is a parabola and the final section is a straight line. Determine the other two equations for the ride.
- State the required domain (set of possible t -values) for each section of the ride.

Problem solving

18. On a set of axes, sketch several parabolas of the form $y = -x^2 + bx$. What do you notice about the turning points of each graph sketched?
19. Determine a quadratic equation given the x -intercepts of the parabola are $-\frac{1}{2}$ and $\frac{3}{4}$.
Explain if there is more than one possible equation.
20. Determine the equation of the parabola that passes through the points $(-1, -2)$, $(1, -4)$ and $(3, 10)$.

LESSON

12.10 Applications (extending)

LEARNING INTENTION

At the end of this lesson you should be able to:

- determine solutions to practical problems using your knowledge of quadratic equations.

12.10.1 Applications of quadratic equations

eles-4945

- Quadratic graphs and equations can be used to solve practical problems in science and engineering.
- They are often involved in problems where a maximum or minimum needs to be found. This will usually occur at the turning point of the parabola.
- When working with physical phenomena, ensure that the solution satisfies any physical constraints of the problem. For example, if measurements of length or time are involved, there can be no negative solutions.

WORKED EXAMPLE 24 Applying quadratic equations to word problems

A flare is fired from a yacht in distress off the coast of Brisbane.

The flare's height, h metres above the horizon t seconds after firing, is given by $h = -2t^2 + 18t + 20$.

- Calculate when the flare will fall into the ocean.
- Calculate the height of the flare after 2 seconds.
- Determine at what other time the flare will be at the same height.
- Calculate how long the flare is above the 'lowest visible height' of 56 m.

THINK

- Substitute $h = 0$ into the equation and solve for t using the Null Factor Law. Since t cannot equal -1 seconds, select the appropriate solution.
- Substitute $t = 2$ into the equation and solve for h .
 - Write the answer in a sentence.
- Substitute $h = 48$ into the equation, rearrange, factorise and solve for t .
- Substitute $h = 56$ into the equation, rearrange, factorise and solve for t .
 - Write the answer in a sentence.

WRITE

a. $0 = -2t^2 + 18t + 20$
 $= -2(t^2 - 9t - 10)$
 $= -2(t - 10)(t + 1)$
 $t = 10$ or -1
 $t = 10$ seconds

b. $h = -2(2)^2 + 18(2) + 20$
 $= -8 + 36 + 20$
 $= 48$

The flare is 48 metres high after 2 seconds.

c. $48 = -2t^2 + 18t + 20$
 $0 = -2t^2 + 18t - 28$
 $= -2(t^2 - 9t + 14)$
 $= -2(t - 7)(t - 2)$
 $t = 7$ seconds


d. $56 = -2t^2 + 18t + 20$
 $= -2t^2 + 18t - 36$
 $= -2(t^2 - 9t + 18)$
 $= -2(t - 6)(t - 3)$
 $t = 3$ seconds and 6 seconds

The flare is above 56 m for 3 seconds (between 3 and 6 seconds).

TI | THINK

a-d. On a Calculator page, complete the entry lines as:
 solve
 $(0 = -2t^2 + 18t + 20, t)$
 $-2t^2 + 18t + 20|t=2$
 solve
 $(48 = -2t^2 + 18t + 20, t)$
 solve
 $(56 = -2t^2 + 18t + 20, t)$
 Press ENTER after each entry.

DISPLAY/WRITE




- a.** $t = 10$ seconds
- b.** The flare is 48 metres high after 2 seconds.
- c.** $t = 7$ seconds
- d.** The flare is above 56 metres for 3 seconds.

CASIO | THINK

a-d. On a Main screen, complete the entry lines as:
 solve
 $(0 = 2t^2 + 18t + 20, t)$
 $2t^2 + 18t + 20|t=2$
 solve
 $(48 = 2t^2 + 18t + 20, t)$
 solve
 $(56 = 2t^2 + 18t + 20, t)$
 Press EXE after each entry.

DISPLAY/WRITE



- a.** $t = 10$ seconds
- b.** The flare is 48 metres high after 2 seconds.
- c.** $t = 7$ seconds
- d.** The flare is above 56 metres for 3 seconds.

on Resources

- eWorkbook** Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)
- Interactivity** Individual pathway interactivity: Applications (int-8357)

Exercise 12.10 Applications (extending)

learn on

12.10 Quick quiz **on**

12.10 Exercise

Individual pathways

PRACTISE

1, 4, 7, 9, 12

CONSOLIDATE

2, 5, 8, 10, 13, 14

MASTER

3, 6, 11, 15, 16

Fluency

- 1.** A spurt of water emerging from an outlet just below the surface of an ornamental fountain follows a parabolic path described by the equation $h = -x^2 + 8x$, where h is the height of the water and x is the horizontal distance from the outlet in metres.
 - a.** Sketch the graph of $h = -x^2 + 8x$.
 - b.** State the maximum height of the water above the surface of the fountain.
- 2.** The position d metres below the starting point of a ball when it is dropped from a great height is given by the equation $d = 250 - 4.9t^2$. If it falls for 1 second, it drops 4.9 m, giving a value of $d = 245.1$ m.
 - a.** Calculate how far it has dropped after:
 - i.** 2 seconds
 - ii.** 7 seconds.
 - b.** Sketch a graph of this relation.


3. A car travels along a highway for a number of minutes according to the relationship $P = 20t^2 + 20t - 120$, where P is the distance from home in metres and t is time in minutes.
- State the distance from home when $t = 0$.
 - Calculate how long it takes the car to reach home.

Understanding

4. **WE24** A rocket fired from Earth travels in a parabolic path. The equation for the path is $h = -0.05d^2 + 4d$, where h is the height in km above the surface of the earth and d is the horizontal distance travelled in km.




- Calculate the height of the rocket after:
 - 30 km
 - 60 km.
 - Calculate how far away the rocket lands.
 - Determine the maximum height of the rocket and how far it travelled before reaching this height.
 - Sketch the path of the rocket.
5. The height of a golf ball hit from the top of a hill is given by the quadratic rule $h = -t^2 + 5t + 14$, where h is in metres and t in seconds.
- Identify the height the golf ball was hit from.
 - Calculate the height of the ball after 2 seconds.
 - Calculate when the golf ball hits the ground.
 - Calculate the maximum height of the ball.
 - Sketch the graph of the flight of the ball.
6. Cave Ltd manufactures teddy bears. The daily profit, $\$P$, is given by the rule $P = -n^2 + 70n - 1200$, where n is the number of teddy bears produced.
- If they produce 40 bears a day, calculate the profit.
 - Sketch the graph of P for appropriate values of n .
 - Determine how many bears they need to produce before they start making a profit.
 - Determine the maximum profit and how many teddy bears they need to manufacture to make this amount.
7. A school playground is to be mulched and the grounds keeper needs to mark out a rectangular perimeter of 80 m.



- Write an expression for the perimeter of the playground in terms of x and y .
- Show that $y = 40 - x$.
- Write an expression for the area to be mulched, A , in terms of x only.
- Sketch the graph of A against x for suitable values of x .
- Determine the maximum area of the playground.
- Determine the dimensions of the playground for this maximum area.

8. A woman wished to build a fence around part of her backyard, as shown in the diagram.



The fence will have one side abutting the wall of the house. She has enough fencing material for 55 m of fence.

- Show that $y = 55 - 2x$.
- Write an expression for the area enclosed by the fence in terms of x alone.
- Determine the dimensions of the fence such that the area is a maximum and calculate that area.

Reasoning

9. A basketball thrown from the edge of the court to the goal shooter is described by the formula $h = -t^2 + 6t + 1$, where h is the height of the basketball in metres after t seconds.
- Identify the height the ball was thrown from.
 - Calculate the height of the basketball after 2 s.
 - Show that the ball was first at a height of 6 m above the ground at 1 second.
 - Determine the time interval during which the ball was above a height of 9 m.
 - Plot the path of the basketball.
 - Calculate the maximum height of the basketball during its flight.
 - Determine how long the ball was in flight if it was caught at a height of 1 m above the ground on its downward path.



10. Jack and his dog were playing outside. Jack was throwing a stick in the air for his dog to catch. The height of the stick (in metres) followed the equation $h = 20t - 5t^2$.
- The graph of h is a parabola. State if the parabola is upright or inverted.
 - Factorise the expression on the right-hand side of the equation.
 - Determine the t -intercepts. These will be the two points at which the stick is on the ground, once at take-off and once at landing. Calculate how long the stick remained in the air. (For simplicity, assume that Jack throws the stick from ground level.)
 - Justify whether the parabola has a maximum or minimum turning point, and determine its coordinates.
 - Calculate the maximum height reached by the stick.
 - Use the information you have found to produce a sketch of the path of the stick.
11. An engineer wishes to build a footbridge in the shape of an inverted parabola across a 40-m wide river. The bridge will be symmetrical and the greatest difference between the lowest part and highest part will be 4 m. Taking the origin as one side, show that the equation that models this footbridge is $y = -0.01x(x - 40)$. Explain what x and y are, and state the domain (the possible values that x can take).

Problem solving

12. A car engine spark plug produces a spark of electricity. The size of the spark depends on how far apart the terminals are.

The percentage performance, Z , of a certain brand is thought to be $Z = -400(g - 0.5)^2 + 100$, where g is the distance between the terminals.

- Sketch the graph of Z .
 - Identify when the performance is greatest.
 - From your graph, determine the values of g for which the percentage performance is greater than 50%.
13. The monthly profit or loss, p (in thousands of dollars), for a new brand of soft drink is given by $p = -2(x - 7.5)^2 + 40.5$, where x is the number (integer) of months after its introduction (when $x = 0$).
- Determine the month when the greatest profit was made.
 - Determine the months when the company made a profit.
14. NASA uses a parabolic flight path to simulate zero gravity and the gravity experienced on the moon. Use the internet to investigate the flight path and create a mathematical model to represent the flight path.
15. An arch bridge is modelled in the shape of a parabolic arch.



The arch span is 50 m wide and the maximum height of the arch above water level is 5.5 m. A floating platform 35-m wide is towed under the bridge.

Determine the greatest height of the deck above water level if the platform is to be towed under the bridge with at least a 35-cm clearance on either side.

16. First Nations Australians have been hunting for thousands of years, using a variety of hunting weapons. One of these hunting weapons is a spear, usually made from Tecoma vine and it is often used with a spear thrower, commonly known as a Woomera or Miru to help launch the spear. If a First Nations Australian launched a spear and it followed a parabolic path modelled by the equation $h = -0.025x^2 + x$, where h is the height of the spear in metres and x is the horizontal distance of the spear in metres.
- Determine the height of the spear after it has travelled 10 m horizontally.
 - Calculate how far the spear travelled horizontally.
 - Determine the maximum height the spear reached.

LESSON

12.11 Review

12.11.1 Topic summary

QUADRATIC EQUATIONS AND GRAPHS

Quadratic equations

- Quadratic equations are equations in which the largest power of x is 2. They contain an x^2 term.
- The general form of a quadratic equation is $y = ax^2 + bx + c$, where $a \neq 0$.
- Quadratic equations for which $a = 1$ are called monic quadratic equations.
- The largest number of solutions a quadratic equation can have is 2.

Axis of symmetry

- Every parabola has an axis of symmetry. This is the vertical line that passes through the parabola's turning point.

Turning point


- The turning point is the point at which a parabola changes direction.
- If $a > 0$ the turning point is the lowest point on the graph. This point is called the local minimum.
- If $a < 0$ the turning point is the highest point on the graph. This point is called the local maximum.

Null Factor Law

- If $a \times b = 0$, then:
 $a = 0$ or $b = 0$ or $a = b = 0$

Turning point form

- The turning point form of a parabola is $y = (x - h)^2 + k$.
- The turning point of a parabola is (h, k) .
- A basic parabola is translated h units horizontally and k units vertically.




Axis intercepts

- All parabolas have a single y -intercept. The location of this y -intercept can be found by letting $x = 0$.
- Parabolas can have 0, 1 or 2 x -intercepts.
- x -intercepts can be found by letting $y = 0$ and solving for x . If the equation can't be solved, there is no x -intercept.


Parabolas of the form $y = ax^2$

- Parabolas of the form $y = ax^2$ have a turning point at the origin.
- If $a > 0$ the graph is \cup -shaped and the turning point is a local minimum.
- If $a < 0$ the graph is \cap -shaped and the turning point is a local maximum.
- Larger values of a create narrower graphs. Smaller values of a create wider graphs.



Factor form




- Parabolas written in the form $y = (x + a)(x + b)$ have x -intercepts at $x = -a$ and $x = -b$.
- The y -intercept is $y = ab$.
- The axis of symmetry will be halfway between the x -intercepts.



12.11.2 Success criteria

Tick a column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

Lesson	Success criteria			
12.2	I can identify equations that are quadratic.			
	I can rearrange quadratic equations into the general form $y = ax^2 + bx + c$.			
	I can solve quadratic equations in the form $ax^2 + c = 0$.			
12.3	I can recall the Null Factor Law.			
	I can solve quadratic equations using the Null Factor Law.			
12.4	I can solve quadratic equations with two terms using the Null Factor Law.			
12.5	I can factorise monic quadratic equations.			
	I can solve monic quadratic equations using factorisation and the Null Factor Law.			
12.6	I can identify the features of a parabolic graph.			
	I can plot the graph of a quadratic equation using a table of values.			
	I can use a parabolic graph to determine values.			
12.7	I can sketch parabolas for quadratic equations of the form $y = ax^2$.			
	I can describe the features of a parabola of the form $y = ax^2$.			
	I can describe the effect of changing the value of a on parabolas of the form $y = ax^2$.			
12.8	I can determine the axes of symmetry, y -intercepts and turning points for graphs of the forms $y = ax^2 + c$ and $y = (x - h)^2 + k$.			
	I can sketch graphs of the form $y = ax^2 + c$, $y = (x + h)^2$ and $y = (x + h)^2 + k$.			
	I can describe the transformations that transform the graph of $y = x^2$ into $y = (x - h)^2 + k$.			
12.9	I can identify the x - and y -intercepts, the axis of symmetry and the turning point of a parabolas of the form $y = (x + a)(x + b)$.			
	I can factorise quadratic equations of the form $y = x^2 + bx + c$ to sketch them using their x -intercepts.			
12.10	I can determine solutions to practical problems using my knowledge of quadratic equations.			

12.11.3 Project


Constructing a parabola

The word *parabola* comes from the Greek language and means ‘thrown’, because it is the path followed by a projectile in flight. Notice that the water streams shown in the photo are moving in the path of a parabola. This investigation explores the technique of folding paper to display the shape of a parabola. The instructions are given below. Take care with each step to ensure your finished product is a well-constructed parabola that can be used in later parts of this investigation.



Forming a parabola by folding paper

- Take a sheet of A4 paper. Cut it into two pieces by dividing the longer side into two. Only one of the halves is required for this investigation.
- Along one of the longer sides of your piece of paper, mark points that are equally spaced 1 cm apart. Start with the first point being on the very edge of the paper.
- Turn over the piece of paper and mark a point, X, 3 cm above the centre of the edge that has the markings on the reverse side.
- Fold the paper so that the first point you marked on the edge touches point X. Make a sharp crease and open the paper flat.
- Fold the paper again so that the second mark touches the point X. Crease and unfold again.
- Repeat this process until all the marks have been folded to touch point X.
- With the paper flat and the point X facing up, you should notice the shape of a parabola appearing in the creases.



1. Trace the curve with a pencil.

The point X is called the focus of the parabola. Consider the parabola to represent a mirror. Rays of light from the focus would hit the mirror (parabola) and be reflected. The angle at which each ray hits the mirror is the same size as the angle at which it is reflected.

2. Using your curve traced from your folding activity, accurately draw a series of lines to represent rays of light from the point X to the parabola (mirror). Use a protractor to carefully measure the angle each line makes with the mirror and draw the path of these rays after reflection in the mirror.



3. Draw a diagram to describe your finding from question 2. Provide a brief comment on your description.

4. Retrace your parabola onto another sheet of paper. Take a point other than the focus and repeat the process of reflection of rays of light from this point by the parabolic mirror.

5. Draw a diagram to describe your finding from question 4. Provide a brief comment on your description.

6. Give examples of where these systems could be used in society.



eWorkbook Topic 12 Workbook (worksheets, code puzzle and project) (ewbk-2012)



Interactivities Crossword (int-0708)

Sudoku puzzle (int-3215)

Exercise 12.11 Review questions

Fluency

- MC** Identify which of the following is *not* a quadratic equation.

A. $x^2 - 1 = 0$ B. $x^2 - 1 + 2x = 0$ C. $x^2 - \frac{1}{2} = 0$ D. $x^2 - \frac{1}{x} = 0$ E. $x^2 + 4 = 0$
- MC** Identify which of the following is in general form.

A. $x^2 + 1 = 0$ B. $x^2 - 1 = 2$ C. $2x^2 = x^2 + 3$ D. $x^2 = x^2 + 1$ E. $x^2 + 5 = x$
- MC** The Null Factor Law cannot be applied to the equation $x(x + 3)(x - 2) = 1$ because:

A. the first factor is a simple x term. B. there are more than two factors.
 C. the right-hand side equals 1. D. the second term is positive.
 E. there is no x^2 term.
- MC** If the solutions to the quadratic equation $(x - 3)(x - b) = 0$ are 3 and -5 , then b is equal to:

A. 5 B. -5 C. 3 D. -3 E. 1
- MC** Select the solutions to $3x^2 - 27 = 0$.

A. $x = 3$ and $x = -3$ B. $x = 9$ and $x = -9$ C. $x = 1$ and $x = -1$
 D. $x = 2$ and $x = -2$ E. There are no solutions.
- Calculate the solutions to the quadratic equation $(x - 3)(2x + 8) = 0$.
- Calculate the solutions to the equation $(4 - x)(2x - 7) = 0$.
- Factorise each of the following quadratic trinomials.

a. $c^2 + 5c + 4$ b. $p^2 + 10p - 24$ c. $y^2 - 10y + 24$
 d. $x^2 + 3x + 2$ e. $m^2 - 7m + 10$ f. $m^2 + 24m + 44$
- Solve the following equations, identifying those with no real solutions.

a. $x^2 + 11x + 10 = 0$ b. $3x^2 + 6x = 0$ c. $-2x^2 - 1 = 0$
- Determine the solutions to the following equations.


a. $(x + 2)^2 - 16 = 0$ b. $4(x - 3)^2 - 36 = 0$ c. $(x + 1)^2 = 25$
- MC** a. The graph of $y = -4x^2$ is

A. wider than $y = x^2$. B. narrower than $y = x^2$.
 C. the same width as $y = x^2$. D. a reflection of $y = x^2$ in the x -axis.
 E. a reflection of $y = x^2$ in the y -axis.

- b. The graph of $y = \frac{1}{2}x^2$ is:
- A. wider than $y = x^2$.
 - B. narrower than $y = x^2$.
 - C. the same width as $y = x^2$.
 - D. a reflection of $y = x^2$ in the x -axis.
 - E. a reflection of $y = x^2$ in the y -axis.
- c. Compared to the graph of $y = x^2$, the graph of $y = -2x^2$ is:
- A. half as wide.
 - B. twice as wide.
 - C. moved 2 units to the right.
 - D. moved 2 units up.
 - E. moved 2 units to the left.
12. **MC** a. The turning point for the graph of the equation $y = -x^2 + 6$ is:
- A. (0, 0)
 - B. (-1, 6)
 - C. (0, 6)
 - D. (0, -6)
 - E. (1, -6)
- b. The turning point of the graph of the equation $y = x^2 - 12$ is:
- A. (1, -12)
 - B. (-12, 1)
 - C. (0, 12)
 - D. (0, -12)
 - E. (-1, 12)
- c. The graph of $y = x^2 - 4$ moves the graph of $y = x^2$ in the following way:
- A. Up 4
 - B. Down 4
 - C. No change
 - D. Left 4 units
 - E. Right 4 units
- d. The y -intercept of the graph of $y = -x^2 - 3$ is:
- A. 1
 - B. -1
 - C. 3
 - D. -3
 - E. -4
13. **MC** a. The axis of symmetry for the graph $y = (x + 1)^2$ is:
- A. $y = 0$
 - B. $x = 1$
 - C. $x = -1$
 - D. $x = 2$
 - E. $y = -1$
- b. The turning point of the graph $y = (x + 1)^2$ is:
- A. (0, 0)
 - B. (1, 1)
 - C. (1, 1)
 - D. (-1, 0)
 - E. (1, 0)
- c. Compared to the graph of $y = x^2$, the graph of $y = (x - 2)^2$ is:
- A. twice as wide.
 - B. moved 2 units to the left.
 - C. moved 2 units to the right.
 - D. moved 2 units up.
 - E. moved 2 units down.
14. **MC** a. For the graph of $y = (x + 6)^2 - 1$, the coordinates of the turning point are:
- A. (6, -1)
 - B. (-1, 6)
 - C. (-1, -6)
 - D. (-6, -1)
 - E. (-6, 1)
- b. For the graph of $y = -(x - 4)^2 + 5$, the axis of symmetry is:
- A. $x = 4$
 - B. $x = 5$
 - C. $x = -4$
 - D. $y = -5$
 - E. $y = 5$
- c. For the graph of $y = (x - 2)^2 - 3$, the y -intercept is:
- A. -5
 - B. 1
 - C. -4
 - D. -3
 - E. -1
15. **MC** To change the graph of $y = x^2$ to make the graph of $y = 2(x - 1)^2 - 3$, we need to:
- A. move it 1 unit to the right and 3 units down.
 - B. make it narrower by a factor of 3, move it 1 unit to the left and 3 units down.
 - C. make it wider by a factor of 2, move it 1 unit to the right and 3 units down.
 - D. make it narrower by a factor of 2, move it 1 unit to the right and 3 units down.
 - E. reflect the graph in the x -axis, and move it 1 unit to the right and 3 units up.
16. **MC** Human cannonball Stephanie Smith was fired from a cannon. Her flight path is a parabola. If c is her height above the net and b is her horizontal distance from the cannon, identify the rule that describes her flight.
- A. $y = x^2 + c$
 - B. $y = bx^2 + c$
 - C. $y = (x - b)^2 + c$
 - D. $y = -(x - b)^2 + c$
 - E. $y = -bx^2 + c$


17. **MC** Identify the rule for the curve shown.

- A. $y = -(x+1)^2 - 2$ B. $y = (x+1)^2 + 2$
 C. $y = (x+1)^2 - 2$ D. $y = -(x+2)^2 - 1$
 E. $y = -(x+1)^2 + 2$



18. **MC** Identify the equation describing the sketch in the diagram shown.







- A. $y = (x-1)(x-3)$ B. $y = (x+1)(x+3)$ C. $y = (x-1)(x+3)$
 D. $y = (x+1)(x-3)$ E. $y = (x-3)^2$



19. **MC** Select the x -intercepts of the graph $y = x^2 - x - 6$.

- A. -2 and 3 B. -3 and 2 C. 2 and 3
 D. -2 and -3 E. -2 and 2

20. For each of the following graphs, state the equation of the axis of symmetry, the coordinates of the turning point, whether the point is a maximum or a minimum, and the x - and y -intercepts.



21. Draw a table of values, use it to plot the graph of each of the following equations, and hence state the equation of the axis of symmetry, the coordinates of the turning point and whether it is a maximum or a minimum, and the x - and y -intercepts.

- a. $y = x^2 - 4x, -2 \leq x \leq 6$ b. $y = -x^2 - 2x + 8, -5 \leq x \leq 3$
 c. $y = 2x^2 - 4x + 4, -2 \leq x \leq 3$ d. $y = -x^2 + 6x - 5, 0 \leq x \leq 6$

22. Sketch each of the following graphs, labelling the axis of symmetry, the turning point and the intercepts, and stating the type of turning point.

- a. $y = 2x^2$ b. $y = \frac{1}{2}x^2$ c. $y = -4x^2$ d. $y = -\frac{1}{3}x^2$

23. Sketch each of the following graphs, labelling the axis of symmetry, the turning point and the y -intercept, and stating the nature of the turning point.

- a. $y = x^2 + 2$ b. $y = x^2 - 4$ c. $y = x^2 + 5$ d. $y = x^2 - 3$

24. Sketch each of the following graphs, labelling the turning point and the y -intercept, and stating the type of turning point.

a. $y = (x + 1)^2$

b. $y = -(x - 2)^2$

c. $y = (x - 5)^2$

d. $y = -(x + 2)^2$

25. Sketch each of the following graphs, labelling the turning point and the intercepts, and stating the nature of the turning point.

a. $y = (x - 1)^2 + 1$

b. $y = (x + 2)^2 + 3$

c. $y = (x - 3)^2 - 2$

d. $y = -(x + 1)^2 - 4$

Problem solving

26. The distance travelled by a motorbike is given by the formula $d = 18t + 2t^2$, where t is the time in seconds and d is the distance in metres.

Calculate how long it would take the motorbike to travel a distance of 180 m.

27. When 20 is subtracted from the square of a certain number, the result is 8 times that number. Determine the number, which is negative.

28. A ball is thrown from the balcony of an apartment building. The ball is h metres above the ground when it is a horizontal distance of x metres from the building. The path of the ball follows the rule

$$h = -x^2 + 3x + 28.$$

Determine how far from the building the ball will land.


29. David owned land in the shape of a square with side length p .

He decided to sell part of this land by reducing it by 50 metres in one direction and 90 metres in the other.

a. Write an expression in terms of p for the area of the original land.

b. Write an expression in terms of p , in its simplest form, for the area of land David has remaining after he sells the section of land shown.

c. Write an expression in terms of p for the area of land he sold.




30. A new computer monitor is made up of a rectangular screen surrounded by a hard plastic frame in which speakers can be inserted.

a. Write expressions for the length and width of the screen in terms of x .

b. Write an expression for the area of the screen, using expanded form.

c. Calculate the area of the screen if $x = 30$ cm.



31. An astronaut needs to find a positive number such that twice the number plus its square gives 35.

a. Write this as an equation.

b. Solve the equation to find the answer for her.

32. A window cleaner was 3 metres off the ground, cleaning the windows of a high-rise building, when he dropped his bucket. The height, h m, of the bucket above the ground t seconds after it is dropped is given by the equation $h = 3 - t^2 - 2t$.
- Identify the value of h when the bucket hits the ground.
 - Rewrite the equation, replacing h with the value it takes when the bucket hits the ground.
 - Solve the equation.
 - Calculate the time it takes for the bucket to reach the ground.



33. A railway bridge has an arch below it that can be modelled by the equation $H = -(x + 1)^2 + 16$, where H is the height of the bridge, in metres.
- Sketch a graph of the bridge and, hence, calculate its maximum height.

34. Sketch the graph of each of the following by determining the x - and y -intercepts and the turning point.
- $y = (x + 2)(x - 2)$
 - $y = (x + 3)(x - 7)$
 - $y = x^2 - 2x - 15$
 - $y = x^2 + 3x + 2$

35. A valley in the countryside has a river running through the centre of it. A vertical cross-section of the valley can be modelled by the equation $H = \frac{1}{10}x^2 - 6x + 100$, where H is the height above sea level, in metres, and x is the horizontal distance, in metres.
- If two villages are situated in the valley, one a horizontal distance of 10 metres from the river and the other 20 metres from the river, determine the height of each village above sea level.

36. Children playing on a cricket pitch throw a ball that follows the path $y = \frac{1}{120}(x - 16)^2$, where y is the height, in yards, above the ground and x is the horizontal distance from the stumps at the end from which the ball is thrown. If the stumps are 22 yards away and 0.5 yards high, explain if the ball will hit the stumps.
- Determine by sketching the graph. (As a cricket pitch is 22 yards long, a yard is a convenient unit to use in this example. A yard is about 91.5 cm.)

37. A toy rocket is thrown up into the air from a balcony. The path the rocket takes is given by the equation $h = 49 - (t - 2)^2$ where h is the height above the ground in metres and t is the time in seconds.
- Calculate the height of the balcony.
 - Calculate when the rocket lands on the ground.
 - Calculate the maximum height of the rocket.
 - Sketch a graph of the rockets path for $0 \leq t \leq 9$.



38. A piece of wire 1 m long is bent into the shape of a rectangle.
- If w cm is the width and l cm is the length, write a rule connecting w and l .
 - If the area of the rectangle is A cm², write an equation connecting l in terms of A .
 - Sketch the graph of A versus l .
 - Calculate the maximum area enclosed by the wire.

39. The organising committee of the Greensborough Tennis Club found that the profit, $\$P$, from a Saturday morning barbeque depended on the number of sausages sold, n .

The estimated profit is given by

$$P = -n^2 + 50n - 400.$$

- a. Calculate the profit if the number of sausages sold is 35.
- b. Sketch the graph of P versus n for $0 \leq n \leq 50$.
- c. Calculate the maximum profit possible and the number of sausages that need to be sold to make this amount.
- d. Determine the minimum number of sausages that would need to be sold to avoid a loss.
40. In a demonstration using the width of a street, a skateboarder performs an ollie over a distance of 2 m. He remains on the ground for $\frac{1}{4}$ m before he begins the ollie, the path of which can be described by the quadratic equation $y = -2(x - p)(x - q)$ (where all dimensions are in metres), and lands with 35 cm to spare before hitting the opposite edge of the curb. Take the origin as the start of his run.
- a. Calculate how far he has travelled horizontally.
- b. Determine the values of p and q .
- c. State the axis of symmetry of the ollie.
- d. Determine the highest point reached by the skateboarder.
- e. Sketch the path of the skateboarder, showing all of the relevant details on the graph.



To test your understanding and knowledge of this topic, go to your learnON title at www.jacplus.com.au and complete the **post-test**.

Answers

Topic 12 Quadratic equations and graphs

12.1 Pre-test

1. A
2. False
3. $(m-2)(m+4)$
4. $2(w-3)(w-2)$
5. E
6. $(1, -3)$
7. C
8. D
9. C
10. D
11. E
12. $(0, -9)$
13. B
14. C
15. True

12.2 Quadratic equations (extending)

1. a. $a = 1, b = 2, c = -1$
b. $a = 1, b = 1, c = -10$
c. $a = 4, b = 5, c = -30$
2. a. $a = 8, b = -11, c = 2$
b. $a = 2, b = 1, c = -60$
c. $a = 1, b = 0, c = -17$
3. D
4. C
5. a. $x = 4, x = -4$
b. There is no solution.
c. $x = 0$
6. a. $x = 1, x = -1$
b. There is no solution.
c. $x = 2, x = -2$
7. a. Y b. N c. Y
8. a. Y b. N c. N
9. a. N b. N c. Y
10. a. Y b. N
11. When $x = 2$:
 $(x+4)^2 = x^2 + 8x + 16$
 $= 2^2 + 8 \times 2 + 16$
 $= 4 + 16 + 16$
 $= 36$
 $x^2 + 16 = 2^2 + 16$
 $= 4 + 16$
 $= 20$
The two expressions are not equal.
12. a. Sample responses can be found in the worked solutions in the online resources.
b. 3.41 cm
c. x is a length, and lengths do not have negative values.

13. If $x^2 + 7x + 4 = 7x$, then $x^2 + 4 = 0$, which means that $x^2 = -4$ and, finally, $x = \sqrt{-4}$.
As the square root of a negative number is not a real number, x has no solutions in the real number range.

14. a. Side length_a = $x + 3$
Side length_b = $x + 5$
Side length_c = $x + 8$
Side length_d = $x - 3$
Side length_e = $x + 6$
b. I = $(x-3)^2$
II = $(x+3)^2$
III = $(x+5)^2$
IV = $(x+6)^2$
V = $(x+8)^2$
c. Area of square I = 4 cm^2
Area of square II = 64 cm^2
Area of square III = 100 cm^2
Area of square IV = 121 cm^2
Area of square V = 169 cm^2
15. a. 6 or -6
b. There are two possible numbers, as the square root of a positive number, in this case 36, has two possible solutions, one positive and one negative.
16. a. $3(x-2)^2 - 7 = 41$
b. -2 and 6

12.3 The Null Factor Law (extending)

1. a. $x = -3, x = 2$ b. $x = -2, x = 3$
c. $x = -2, x = 3$ d. $x = -2\frac{1}{2}, x = -\frac{3}{4}$
e. $x = -4, x = -\frac{1}{2}$
2. a. $x = \frac{1}{2}, x = -30$ b. $x = -\frac{1}{2}, x = 3$
c. $x = 1, x = \frac{1}{3}$ d. $x = 0, x = 2$
e. $x = -\frac{1}{3}, x = \frac{1}{4}$
3. a. $x = -2.3, x = 0.3$ b. $x = -\frac{1}{6}, x = \frac{1}{6}$
c. $x = 2$ d. $x = 0, x = \frac{15}{4}$
e. $x = -4$
4. a. $x = 2, x = -2, x = -3$
b. $x = -2, x = 2.5$
c. $x = -2, x = -4$
d. $x = 0, x = -2, x = -4$
5. a. $x = 1.1, x = -2.4, x = -2.6$
b. $x = -3, x = -\frac{1}{2}, x = 1\frac{2}{3}$
c. $x = 3$
d. $x = -1, x = 2$

6. D
7. B
8. a. $x(x + 10) = 0$; $x = 0$ or $x = -10$
 b. $2x(x - 7) = 0$; $x = 0$ or $x = 7$
 c. $5x(5x - 8) = 0$; $x = 0$ or $x = \frac{8}{5}$
9. By dividing both sides of the equation $-2x^2 + 2x + 12 = 0$ by -2 , we get $(x - 3)(x + 2) = 0$. Since the equations are equivalent, they have the same solution(s).
10. a. When $x = 2$, the first bracket equals 2 and the second bracket equals 4; therefore, the product is 8.
 b. -2 and 1
11. A quadratic can have a maximum of two solutions, because a quadratic can at most be factorised into two separate pairs of brackets, each of which represents one solution.
12. a. 10 metres
 b. $y = \frac{1}{250}x(100 - x)$; $a = \frac{1}{250}$, $b = 100$
 c. 9.424 m
13. a. $h = -4t(t - 5)$
 b. i. 16 m
 ii. 0 m
 c. Sample responses can be found in the worked solutions in the online resources.
14. -29 or 8

12.4 Solving quadratic equations with two terms (extending)

1. a. $x = -3, x = 3$ b. $x = -4, x = 4$
 c. $x = -3, x = 3$ d. $x = -5, x = 5$
 e. $x = -10, x = 10$
2. a. $x = -7, x = 7$ b. $x = -3, x = 3$
 c. $x = -2, x = 2$ d. No real solutions
 e. No real solutions
3. a. $x = -3, x = 3$ b. $x = -4, x = 4$
 c. $x = -5, x = 5$ d. $x = 0$
 e. $x = 0$
4. a. $x = 0, x = -6$ b. $x = 0, x = 8$
 c. $x = 0, x = -9$ d. $x = 0, x = 11$
 e. $x = 0, x = 6$
5. a. $x = 0, x = 7.5$ b. $x = 0, x = \frac{2}{3}$
 c. $x = 0, x = -1\frac{3}{4}$ d. $x = 0, x = 2\frac{1}{2}$
 e. $x = 0, x = -1$
6. a. $x = 0, x = \frac{1}{4}$ b. $x = 0, x = -5$
 c. $x = 0, x = -12$ d. $x = 0, x = 18$
 e. $x = 0, x = 2.5$
7. A
8. C
9. 4 or -4
10. The square plot is $12\text{ m} \times 12\text{ m}$; the rectangular plot is $16\text{ m} \times 9\text{ m}$.

11. The number is 0 or 10.
12. $x^2 + 9$ cannot be factorised.
13. $x = \pm \frac{n}{m}$
14. x cannot be isolated, so the only way to solve the equation will be to factorise it and use the Null Factor Law.
15. If $a = 0$, then the expression is not quadratic.
16. 0 or 8
17. -12 or 0
18. a. $x(x + 30) = 50x$
 b. $x(x - 20) = 0$
 c. $x = 0$ or $x = 20$
 d. No, x cannot be 0, because the width has to have a positive value.

12.5 Factorising and solving monic quadratic equations (extending)

1. a. $x^2 + 6x + 8$ b. $x^2 - 6x + 8$
 c. $x^2 - 9x + 20$ d. $x^2 + 9x + 20$
 e. $m^2 + 6m + 5$ f. $m^2 - 6m + 5$
2. a. $t^2 + 19t + 88$ b. $t^2 - 30t + 200$
 c. $x^2 - x - 6$ d. $x^2 + x - 6$
 e. $v^2 - 3v - 40$ f. $v^2 + 3v - 40$
3. a. $x^2 + 5x - 14$ b. $t^2 - 11t - 12$
 c. $n^2 + 13n - 30$ d. $a^2 - 3a - 18$
 e. $z^2 + 8z + 16$ f. $z^2 + 3z - 88$
4. a. 1, 4; 2, 3 b. 1, 5; 2, 4; 3, 3
 c. 1, 11; 2, 10... 6, 6 d. 1, 19; 2, 18... 10, 10
5. a. $-1, -4; -2, -3$
 b. $-1, -6; -2, -5; -3, -4$
 c. $-1, -7; -2, -6; -3, -5; -4, -4$
 d. $-1, -10; -2, -9... -5, -6$
6. a. 7 and -1 ; 8 and -2 ; 9 and -3 ; 10 and -4 ; 11 and -5
 b. 1 and -7 ; 2 and -8 ; 3 and -9 ; 4 and -10 ; 5 and -11
 c. 4 and -1 ; 5 and -2 ; 6 and -3 ; 7 and -4 ; 8 and -5
 d. 1 and -4 ; 2 and -5 ; 3 and -6 ; 4 and -7 ; 5 and -8
 e. 1 and -9 ; 2 and -10 ; 3 and -11 ; 4 and -12 ; 5 and -13
 f. 15 and -1 ; 16 and -2 ; 17 and -3 ; 18 and -4 ; 19 and -5
7. a. $(x + 1)(x + 3)$ b. $(x - 1)(x - 3)$
 c. $(x + 1)(x + 11)$ d. $(x - 1)(x - 11)$
 e. $(a + 5)(a + 1)$
8. a. $(a - 5)(a - 1)$ b. $(x - 4)(x - 3)$
 c. $(x - 5)(x - 2)$ d. $(n + 4)^2$
 e. $(n + 8)(n + 2)$
9. a. $(y - 9)(y - 3)$ b. $(x - 7)(x - 6)$
 c. $(t - 6)(t - 2)$ d. $(t + 9)(t + 2)$
 e. $(u + 2)(u + 3)$
10. a. $(x + 6)(x - 3)$ b. $(x - 6)(x + 3)$
 c. $(x - 5)(x + 3)$ d. $(x + 5)(x - 3)$
 e. $(n - 14)(n + 1)$
11. a. $(n + 7)(n - 5)$ b. $(v + 6)(v - 1)$
 c. $(v - 6)(v + 1)$ d. $(t + 6)(t - 2)$
 e. $(t - 7)(t + 2)$

12. a. $(x-5)(x+4)$
 c. $(n+10)(n-9)$
 e. $(x-5)(x+1)$
13. a. $x=2, x=4$
 c. $x=-1, x=-5$
 e. $x=3, x=-5$
14. a. $x=-2$
 c. $x=8, x=-3$
 e. $x=-12, x=-1$
15. a. $x=11, x=-1$
 c. $x=-25, x=-4$
 e. $x=4, x=-2$
16. B
17. D
18. a. $(x+1)(x+6)$
 b. $x^2 + 7x + 10 = (x+2)(x+5)$,
 $x^2 + 7x + 12 = (x+3)(x+4)$
19. a. $(x-1)(x+4)$
 b. Sample responses can be found in the worked solutions in the online resources.
20. D
21. 9 m by 5 m
22. Sample responses can be found in the worked solutions in the online resources.
23. Sample responses can be found in the worked solutions in the online resources.
24. a. i. 0
 ii. 2
 iii. 35
 b. i. Octagon
 ii. Icosahedron
25. a. i. $n+1$
 iii. $(n+1)(n+7)$
 b. 3
26. a. $w = (x-3)$ cm
 b. $w = 62$ cm
 c. $l = 108$ cm
27. a. $g+18$
 b. $g^2 + 12g - 108$
 c. $g^2 + 12g - 108 = 256$
 d. $g^2 + 12g - 364 = 0$
 e. $g = 14$
 f. The girl is 14 years old, her brother is 8 years old and her teacher is 32 years old.
28. a. 1×120 m
 2×60 m
 3×40 m
 4×30 m
 5×24 m
 6×20 m
 8×15 m
 10×12 m
 b. $l = x + 8, w = x - 6$
 c. 6×20 m

12.6 Graphs of quadratic functions


1. a. i. Axis of symmetry: $x = 0$,
 ii. turning point $(0, 0)$,
 iii. minimum
 b. i. Axis of symmetry: $x = 0$,
 ii. turning point $(0, -3)$,
 iii. minimum
 c. i. Axis of symmetry: $x = -1$,
 ii. turning point $(-1, -2)$,
 iii. minimum
 d. i. Axis of symmetry: $x = 0$,
 ii. turning point $(0, 0)$,
 iii. maximum
 e. i. Axis of symmetry: $x = 0$,
 ii. turning point $(0, 2)$,
 iii. maximum
 f. i. Axis of symmetry: $x = 2$,
 ii. turning point $(2, -1)$,
 iii. maximum
2. a. i. $x = 0$,
 ii. $(0, 1)$,
 iii. minimum
 b. i. $x = 1$,
 ii. $(1, -3)$,
 iii. minimum
 c. i. $x = -2$,
 ii. $(-2, 2)$,
 iii. maximum
 d. i. $x = -1$,
 ii. $(-1, -2)$,
 iii. maximum
 e. i. $x = 2$,
 ii. $(2, 2)$,
 iii. minimum
 f. i. $x = 0$,
 ii. $(0, 1)$,
 iii. maximum
3. a. i. $x = 0$;
 ii. TP $(0, -1)$, minimum;
 iii. x -intercepts are -1 and 1 , y -intercept is -1 .
 b. i. $x = 0$;
 ii. TP $(0, 1)$, maximum;
 iii. x -intercepts are -1 and 1 , y -intercept is 1 .
 c. i. $x = 1$;
 ii. TP $(1, -4)$, minimum;
 iii. x -intercepts are -1 and 3 , y -intercept is -3 .
 d. i. $x = -2$;
 ii. TP $(-2, 1)$, maximum;
 iii. x -intercepts are -3 and -1 , y -intercept is -3 .

- e. i. $x = -\frac{1}{2}$;
 ii. TP $(-\frac{1}{2}, -1)$, minimum;
 iii. x-intercepts are $-1\frac{1}{2}$ and $\frac{1}{2}$, y-intercept is $-\frac{3}{4}$.


- f. i. $x = \frac{1}{2}$;
 ii. TP $(\frac{1}{2}, 2)$, minimum;
 iii. no x-intercepts, y-intercept is $2\frac{1}{4}$.

4. a. B b. C c. A d. D


5. a. $y = x^2 + 8x + 15, -7 \leq x \leq 0$




- i. $x = -4$
 ii. $(-4, -1)$, minimum
 iii. x-intercepts are -5 and -3 , y-intercept is 15 .
 b. $y = x^2 - 1, -3 \leq x \leq 3$




- i. $x = 0$
 ii. $(0, -1)$, minimum
 iii. x-intercepts are -1 and 1 , y-intercept is -1 .
 6. a. $y = x^2 - 4x, -1 \leq x \leq 5$




- i. $x = 2$
 ii. $(2, -4)$, minimum
 iii. x-intercepts are 0 and 4 , y-intercept is 0 .
 b. $y = x^2 - 2x + 3, -2 \leq x \leq 4$



- i. $x = 1$
 ii. $(1, 2)$, minimum
 iii. No x-intercepts, y-intercept is 3 .
 7. a. $y = x^2 + 12x + 35, -9 \leq x \leq 0$




- i. $x = -6$
 ii. $(-6, -1)$, minimum
 iii. x-intercepts are -7 and -5 , y-intercept is 35 .
 b. $y = -x^2 + 4x + 5, -2 \leq x \leq 6$



- i. $x = 2$
 ii. $(2, 9)$, maximum
 iii. x-intercepts are -1 and 5 , y-intercept is 5 .

8.


x	-3	-2	-1	0	1	2	3
$y = x^2 + 2$	11	6	3	2	3	6	12
$y = x^2 + 3$	12	7	4	3	4	7	12



b. Axis of symmetry: $x = 0$ for both equations.

c. No x -intercepts for either equation.

9. a.



The shape is a parabola.

b. $x = 0$

c. $y = -4$

d. $x = -2, x = 2$

10.


x	-6	-4	-2	0	2	4	6	8
y	30	12	2	0	6	20	42	72

11. C


12. a. $(0, 0)$

b. Maximum

13. a.



b.



14. Answers will vary but must be of the form $y = a(x - h)^2$.

15. Answers will vary but must be of the form $y = ax^2 + c$.

16. Answers will vary. If the parabola is upright, it has a minimum turning point. If the parabola is inverted, it has a maximum turning point.

17. Axis of symmetry crosses halfway between the x -intercepts.

Let the unknown x -intercept be at p . Then,


$$-4 = \frac{10 + p}{2};$$

$$-8 = 10 + p;$$

$$p = -18$$

Therefore the other x -intercept is at $(-18, 0)$.


18. Let $(a, 0)$ be the point at which the axis of symmetry crosses the x -axis.
As a will lie halfway between the x -intercepts $(-2, 0)$ and $(5, 0)$;
- $$a = \frac{-2 + 5}{2} = \frac{3}{2} = 1.5$$
- Therefore, the axis of symmetry has the equation $x = 1.5$.



By inspection or algebra, the graphs meet at $(0, 0)$ and again at $(1, 1)$.

20. a. $\frac{x_2 + 10}{2} = -4$
 $x_2 = 2$


b. Answers will vary but must be of the form $y = a(x - 2)(x + 10)$.




The maximum height is 500 m.

22. a. -21°C b. 1 hour and at 21 hours
c. $t = 11$ hours d. 100°C


12.7 Sketching parabolas of the form $y = ax^2$



$y = 4x^2$ is narrower.
Turning point for each is at $(0, 0)$; x -int and y -int is 0 for both.



$y = x^2$ is narrower.
Turning point for each is at $(0, 0)$; x -int and y -int is 0 for both.



$y = 0.25x^2$

4. a, b. See the figure at the foot of the page.*
Looking at the graphs, $y = -5x^2$ is the narrowest.

5. a. B b. A c. B



6. a. iii b. vi c. i
d. ii e. iv f. v

7. Sample responses can be found in the worked solutions in the online resources; for example, $y = 2x^2$.

8. Sample responses can be found in the worked solutions in the online resources; for example, $y = -0.5x^2$.

9. a. $y = 3x^2$ b. $y = -x^2$

10. a. -350 b. -350 c. -31.5
d. -7.875 e. -16.94



12. $y = \frac{2}{3}x^2$

13. a. Sample responses can be found in the worked solutions in the online resources.

b. 6


14. a. 400 watts

b. 625 watts

15. $k = 1$


16. $x = y^2$

*4. a, b.




12.8 Sketching parabolas of the form $y = (x - h)^2 + k$


1. a.




b.




2. a.




b.




3. a.



b.




4. a.




Minimum


b.




Maximum




Minimum



Maximum




Maximum




Minimum

7. a. x-intercepts: $(-2, 0)$, $(2, 0)$



b. No x-intercepts



c. x-intercepts: $(-1, 0), (1, 0)$
See the figure at the foot of the page.*


8. a. No

b. A negative sign inverts the graph.

9. a. C b. D c. D


d. D

10. a. Minimum turning point $(1, 0)$




y-intercept $(0, 1)$

b. Minimum turning point $(-2, 0)$




y-intercept $(0, 4)$

*7. c.




11. a. Minimum turning point (3, 0)




y-intercept (0, 9)

b. Minimum turning point (-4, 0)




y-intercept (0, 16)

12. a. Minimum turning point (5, 0)




y-intercept (0, 25)

b. Minimum turning point (-6, 0)




y-intercept (0, 36)

13. a. Maximum turning point (1, 0)




y-intercept (0, -1)

b. Maximum turning point (-2, 0)




y-intercept (0, -4)

14. a. Maximum turning point (3, 0)




y-intercept (0, -9)

b. Maximum turning point (-4, 0)




y-intercept (0, -16)

15. a. Maximum turning point (5, 0)



y-intercept (0, -25)

b. Maximum turning point $(-6, 0)$




y-intercept $(0, -36)$

16. a. C


b. D

17. a.




Minimum

b.




Minimum

18. a.




Minimum

b.




Minimum

19. a.




Maximum

b.



Maximum


20. a.



b. 1 right

c. 1 right, 3 up


21. a.



b. 3 left

c. 3 left, 2 up

22. a.



b. 4 right

c. 4 right, 1 down

23. a. D

b. A

c. C

24. a. iii

b. iv

c. v

d. i


e. vi

f. ii

25. a. See figure at the foot of the page.*

b. 5 km above sea level.

*25.



26. a. vi b. ii c. iii
 d. i e. iv f. v

27. a. $y = (x - 2)^2$ or $y = (x - 4)^2$
 b. $y = (x - 2)^2$ or $y = (x + 4)^2$

28. a. No difference.
 b. 2nd graph is translated 4 units to the left of the 1st graph.
 c. 2nd graph is 'inverted' and is the mirror image of the 1st graph in the x -axis.

29. a. $y = -(x + 2)^2$ b. $y = (x - 3)^2 - 1$
 c. $y = (x + 1)^2 + 2$ d. $y = -(x - 4)^2 + 6$


30. C

31. The graph is a basic parabola that has been translated h units horizontally and k units vertically. Therefore the turning point becomes (h, k) . This form allows the turning point to be determined very quickly and easily.

32. Students should substitute the points into the given equations and show that the equations are true.

33. Students should substitute the points into the given equations and show that the equations are true.

34. Intersections at $(-\sqrt{2}, 0)$ and $(\sqrt{2}, 0)$



35. a. $y = (x - 3)^2 - 2$ is obtained by translating $y = (x - 3)^2$ vertically 2 units down.
 $y = (x - 3)^2 + 1$ is obtained by translating $y = (x - 3)^2$ vertically 1 unit up.


b. $y = (x - 5)^2 - 4$

36. a. 40°C

b. Sample responses can be found in the worked solutions in the online resources.

c. 5 hours

d. $t = (h - 5)^2 + 15$



37. $y = 4 + 4\sqrt{5}$

38. a. 150 metres


b. $y = \frac{-4}{225}(x - 150)^2 + 400$

39. 50 cm horizontally away from the desk.


40. $y = -\left(x - \frac{1}{2}\right)^2 + \frac{29}{4}$

12.9 Sketching parabolas of the form $y = (x + a)(x + b)$ (extending)


1. a.




b.




2. a.




b.




3. a.




b.




4. a.




b.












5. a.



b.







11. $y = (x - 2)(x - 6)$

12. a. 20 m

b. m4, n24

c. (4, 0), (24, 0)


d. $x = 14$

e. (14, 10)

f. 10 m

g. 10 m

h.



13. 12 m

14. a. (6, -9)


b. $y = (x - 6)^2 - 9$

c. Intercept form expanded:

$$y = x^2 - 12x + 27$$

Turning point form expanded:

$$y = x^2 - 12x + 36 - 9 = x^2 - 12x + 27$$



15. $x = 11$

16. a. Since the coefficient of t^2 is negative, the turning point is a maximum.

b. Maximum height = 512 m at $t = 8$ seconds

17. a. Sample responses can be found in the worked solutions in the online resources.

b. $0 \leq t \leq 6, 6 \leq t \leq 12, 12 \leq t \leq 21$

18. The general turning point is $\left(\frac{b}{2}, \frac{b^2}{4}\right)$.

19. Answers will vary but must be of the general form


$$y = a\left(x + \frac{1}{2}\right) - \left(x - \frac{3}{4}\right)$$

20. $y = 2x^2 - x - 5$

12.10 Applications (extending)


1. a.

b. 16 m



2. a. i. 19.6 m

ii. 240.1 m




3. a. 120 metres b. 2 minutes

4. a. i. 75 km

ii. 60 km

b. 80 km

c. 80 km high after travelling 40 km horizontally




5. a. 14 m


b. 20 m

c. 7 s

d. 20.25 m



6. a. No profit




c. 30 bears

d. \$25; 35 bears

7. a. $2(x + y) = 80$

b. Sample responses can be found in the worked solutions in the online resources.

c. $A = x(40 - x)$



e. 400 m^2

f. $x = 20 \text{ m}; y = 20 \text{ m}$

8. a. Perimeter $-2x + y = 55 \text{ m}$, so $y = 55 - 2x$

b. $55x - 2x^2$


c. $x = 13.75 \text{ m}, y = 27.5 \text{ m}$, area = 378.125 m^2 .

9. a. 1 m

b. 9 m

c. Sample responses can be found in the worked solutions in the online resources.

d. From 2 to 4 seconds



f. 10 m

g. 6 s


10. a. Inverted

b. $h = 5t(4 - t)$

c. $t = 0, t = 4$; 4 seconds

d. Maximum turning point at (2, 20)


e. 20 m



11. Let x = horizontal distance, y = vertical distance across the bridge ($0 \leq x \leq 40$)

$$y = -0.01x(x - 40)$$

12. a.



b. $g = 0.5$

c. $0.15 \leq g \leq 0.85$

13. a. 7th to 8th month b. 3rd and 12th months

14. Sample responses can be found in the worked solutions in the online resources.

15. The height of the platform deck above the water is 2.7 m.

16. a. 7.5 m b. 40 m c. 10 m

Project

1–5. Students need to construct a parabola by folding paper with the directions given in this project. Draw incident and reflected rays of light and use a protractor to carefully measure the angle each line makes with the mirror. Also comment on the diagram.


6. Parabolas could be seen in real life in, for example, water shot by the fountain in a parabolic path, a ball thrown into the air, bridges, headlights, satellite dishes or telescopes.

12.11 Review questions

1. D
2. A
3. C
4. B
5. A
6. $x = -4, x = 3$
7. $x = 3.5, x = 4$
8. a. $(c + 1)(c + 4)$ b. $(p - 2)(p + 12)$
- c. $(y - 6)(y - 4)$ d. $(x + 1)(x + 2)$
- e. $(m - 2)(m - 5)$ f. $(m + 22)(m + 2)$
9. a. $x = -10, x = -1$
- b. $x = -2, x = 0$
- c. No real solutions
10. a. $x = -6, x = 2$
- b. $x = 0, x = 6$
- c. $x = -6, x = 4$
11. a. B b. A c. A
12. a. C b. D c. B d. D
13. a. C b. D c. C
14. a. D b. A c. B
15. D
16. D
17. B
18. C
19. A
20. a. $x = 0$, TP (0, -4), minimum, x -intercepts are -2 and 2, y -intercept is -4.
- b. $x = 0$, TP (0, 9), maximum, x -intercepts are 3 and -3, y -intercept is 9.
- c. $x = 1$, TP (1, 0), minimum, x -intercepts are 1, y -intercept is 1.
- d. $x = -3$, TP (-3, 0), maximum, x -intercepts are -3, y -intercept is -9.
- e. $x = -1$, TP (-1, 1), minimum, no real x -intercepts, y -intercept is 2.
- f. $x = -1$, TP (-1, -4), minimum, x -intercepts are 1 and -3, y -intercept is -3.

21. a. $y = x^2 - 4x, -2 \leq x \leq 6$


x	-2	-1	0	1	2	3	4	5	6
y	12	5	0	-3	-4	-3	0	5	12



$x = 2$, TP $(2, -4)$, minimum, x -intercepts are 0 and 4, y -intercept is 0.

b. $y = -x^2 - 2x + 8, -5 \leq x \leq 3$

x	-5	-4	-3	-2	-1	0	1	2	3
y	-7	0	5	8	9	8	5	0	-7



$x = -1$, TP $(-1, 9)$, maximum, x -intercepts are -4 and 2, y -intercept is 8.

c. $y = 2x^2 - 4x + 4, -2 \leq x \leq 3$


x	-2	-1	0	1	2	3
y	20	10	4	2	4	10

See the figure at the foot of the page.*

$x = 1$, TP $(1, 2)$, minimum, no x -intercepts, y -intercept is 4.


d. $y = -x^2 + 6x - 5, 0 \leq x \leq 6$

x	0	1	2	3	4	5	6
y	-5	0	3	4	3	0	-5




$x = 3$, TP $(3, 4)$, maximum, x -intercepts are 1 and 5, y -intercept is -5.


22. a.




Minimum

*21. c.







Minimum




Minimum




Maximum




Minimum




Maximum




Minimum




Minimum




Minimum



Maximum




Minimum




Maximum


25. a.




Minimum



Minimum



Minimum



Maximum

26. 6 s

27. $x = -2$

28. 7 metres

29. a. p^2

b. $p^2 - 140p + 4500$

c. $140p - 4500$

30. a. Length = $(x-4)$ cm, width = $(x-6)$ cm

b. $(x^2 - 10x + 24)$ cm²

c. 624 cm²

31. a. $x^2 + 2x = 35$



b. 5

32. a. $h = 0$


b. $0 = 3 - t^2 - 2t$

c. $t = -3$ or $t = 1$


d. 1 second




Minimum



Minimum




Minimum



Minimum

35. 50 metres, 20 metres




Yes, the ball will hit the stumps because they are higher than 0.3 yards.


37. a. 45 m

b. 9 s

c. 49 m




38. a. $w + l = 50$
 b. $A = l(50 - l)$




d. 625 cm^2

39. a. \$125



- c. \$225 for 25 sausages
 d. 10 or 40 sausages

40. a. 2.6 m
 b. $p = 0.25, q = 2.25$
 c. $x = 1.25$
 d. 2 m



Semester review 2

The learnON platform is a powerful tool that enables students to complete revision independently and allows teachers to set mixed and spaced practice with ease.

Student self-study

Review the **Course Content** to determine which topics and subtopics you studied throughout the year. Notice the green bubbles showing which elements were covered.

The screenshot shows the 'learnON' interface for 'Jacaranda Maths Quest 7 AC 5e'. The 'Course Content' sidebar on the left lists various topics. The main content area displays the '1 Positive integers' section, which is highlighted with a red box. This section includes subtopics 1.1 Overview, 1.2 Place value, 1.3 Strategies for adding and subtracting positive integers, 1.4 Algorithms for adding and subtracting positive integers, 1.5 Multiplying positive integers, 1.6 Dividing positive integers, and 1.7 Rounding and estimating. Each subtopic has a green bubble next to it, indicating it was covered. The '1.2 Place value' and '1.3 Strategies for adding and subtracting positive integers' subtopics are also highlighted with a red box.

Review your results in **My Reports** and highlight the areas where you may need additional practice.

The screenshot shows the 'learnON' interface for 'Jacaranda Maths Quest 8 AC Third Edition'. The 'My Reports' section is displayed, showing 'Results by course'. The '1. Choose assignment type:' section is set to 'PRACTICE'. The '2. Filter further:' section is set to 'No filter'. The 'SHOW REPORT' button is visible. The 'Results by course' section shows a table of scores for 'Number and Algebra' and 'Linear equations'. The table is filtered to show 'PRACTICE' assignments. The 'Number and Algebra' row shows scores of 33% and 0% for two different assignments, with a red box highlighting these scores.



Course	Number and Algebra	Linear equations
Holiday Brighton	33% 0%	100% 20%

Use these and other tools to help identify areas of strengths and weakness and target those areas for improvement.


Teachers

It is possible to set questions that span multiple topics. These assignments can be given to individual students, to groups or to the whole class in a few easy steps.

Go to **Menu** and select **Assignments** and then **Create Assignment**. You can select questions from one or many topics simply by ticking the boxes as shown below.



Once your selections are made, you can assign to your whole class or subsets of your class, with individualised start and finish times. You can also share with other teachers.



More instructions and helpful hints are available at www.jacplus.com.au.

LESSON

13.1 Overview

13.1.1 Why learn this?

Computer programming is the process of creating instructions for a computer to follow to perform given tasks. In this topic the computer language JavaScript is used in most of the examples and questions. JavaScript has been chosen for its popularity and availability on nearly all web browsers.

A computer is often required to sort lists of data in ascending or descending order. These lists may be very long, but a program can loop through thousands of pieces of data every second. This topic will explore generating many random numbers to estimate probabilities, introduce how a program can process and organise large amounts of data, and explore various algorithms to sort data.

There are many areas of commerce where computer programmers are employed to facilitate the operations of a business. Website designers and developers are in high demand for their skills as more and more businesses operate online. Computers, phone apps and other digital devices are developed using coding. Learning to give detailed step-by-step instructions to a computer is also the basis of website development.



Hey students! Bring these pages to life online



Watch videos



Engage with interactivities



Answer questions and check solutions

Find all this and MORE in jacPLUS



Reading content and rich media, including interactivities and videos for every concept



Extra learning resources

Differentiated question sets

Questions with immediate feedback, and fully worked solutions to help students get unstuck

- MC** The value "1.25" is classified as:
A. Boolean B. Number C. String D. Array E. Decimal
- MC** The most suitable comment to the statement of the form // variable = value, for:

```
var myWeight = 52 + 15 + 18;
```


A. `var Weight = 52 + 15 + 18; // myWeight = 85`
B. `var myWeight = 52 + 15 + 18; // myWeight = 52`
C. `var myWeight = 52 + 15 + 18; // myWeight = 85`
D. `var myWeight = 52 + 15 + 18; // myWeight = 80`
E. `var Weight = 52 + 15 + 18; // Weight = 85`
- Determine the output to the console.

```
console.log((23%4)*8)
```
- MC** What is `[1, 6, 12]` classified as?
A. Number B. Boolean C. Array D. String E. Integer
- MC** Evaluate the following JavaScript manually:
`Array.isArray("[12]")`
A. "number" B. Array C. true D. "string" E. false
- MC** Determine what `[32, "square", "23"].length` evaluates to.
A. 3 B. square C. 23 D. 1 E. 32
- MC** Select which of the following JavaScript expressions is false.
A. `!true` B. `4.56 !== 5.46` C. `5.78 === 5.78`
D. `3.99 <= 4.01` E. `!false`
- MC** Select which of the following JavaScript expressions is false.
A. `true&&true` B. `false&&true` C. `false||true`
D. `true||true` E. `true||false`
- Manually simulate the running of the following program:

```
var value = 8;  
if (value%3 > 0) {  
    value = value + 3;  
}
```


State the final value stored.
- Determine how many times the loop is executed in the following program.

```
var i = 1;  
while (i <= 23) {  
    console.log(i);  
    i = i + 1;  
}
```

11. State how many times the loop code is executed in the following program.

```
var sum = 0;
var numbers = [2, 4, 6, 8, 10];
var j = 0;
while (j < numbers.length) {
    var number = numbers[j];
    sum = sum + number;
    j = j + 1;
}
console.log(sum);
```

12. Determine the output to the console of the following.

```
var sum = 0;
var numbers = [2, 4, 6, 8, 10];
var j = 0;
while (j < numbers.length) {
    var number = numbers[j];
    sum = sum + number;
    j = j + 1;
}
console.log(sum);
```

13. Determine the output to the console of the following program.

```
var values = [12, 9, 16, 21, 7, 20, 13];
var min = values[0];
var max = values[0];
var i = 0;
while (i < values.length) {
    var item = values[i];
    if (item > max) {
        max = item;
    }
    if (item < min) {
        min = item;
    }
    i = i + 1;
}
var result = max - min;
console.log(result);
```

14. **MC** By manually simulating running the following program, the final values stored in the array are:

```
var seed = 1;
function random () {
    seed = (7*seed)%5;
    return seed;
}
var array = [random(), random()];
```

A. [2, 2]

B. [2, 4]

C. [1, 2, 3]

D. [1, 2]

E. [1, 7, 5]

15. **MC** Given the function definition shown below, determine what is returned by the function call:

```
function swapItems (array, index1, index2) {
    var swap = array [index1];
    array [index1] = array [index2];
    array [index2] = swap;
}

function bubbleScan (array) {
    var swapped = false;
    var i = 1;
    while (i < array.length) {
        if (array[i - 1] > array[i]) {
            swapItems (array, (i - 1), i);
            swapped = true;
        }
        i = i + 1;
    }
    return swapped;
}

bubbleScan ([4, 12, 15, 13, 20, 33])
```

- A. false
B. bubbleScan
C. [13]
D. [12, 13, 15]
E. true

LESSON

13.2 Programs

LEARNING INTENTIONS

At the end of this lesson you should be able to:

- understand the types of values used in computer programming
- write numerical expressions in JavaScript and evaluating the output
- assign variables and insert comments.

13.2.1 Values

- Three of the most common types of values used in computer programming are:
 - **numbers** (numerical values): for example 87, -365, 1.5, -8.14213
 - **strings** (any text surrounded in quotation marks " "): for example "A name", "3.14", "State: Victoria", "29#\$\$3", ""
 - **Booleans** (used for logic, this type only has two possible values): either `false` or `true`.
- In JavaScript the expression `typeof value` returns either "number", "string" or "boolean".

WORKED EXAMPLE 1 Identifying types of values

Answer these questions for each of the following values.

- i. Classify the value as a number, a string or Boolean.
- ii. Apply the expression `typeof value` to each value. Write the result of the expression `typeof value`.
 - a. "Katherine"
 - b. 3.14159265
 - c. true
 - d. "false"

THINK

- a. i. "Katherine" is in quotation marks, so this value is a string.
 - ii. 1. Apply the expression `typeof value` to the value "Katherine".
2. `typeof "Katherine"` is "string".
- b. i. 3.14159265 is a numerical value and is not in quotation marks, so this value is a number.
 - ii. 1. Apply the expression `typeof value` to the value 3.14159265.
2. `typeof 3.14159265` is "number".
- c. i. true is a Boolean value.
 - ii. 1. Apply the expression `typeof value` to the value true.
2. `typeof true` is "boolean".
- d. i. "false" is in quotation marks, so it is a string.
 - ii. 1. Apply the expression `typeof value` to the value "false".
2. `typeof "false"` is "string".

WRITE

- a. i. String
 - ii. `typeof "Katherine"`
"string"
- b. i. Number
 - ii. `typeof 3.14159265`
"number"
- c. i. Boolean
 - ii. `typeof true`
"boolean"
- d. i. String
 - ii. `typeof "false"`
"string"

13.2.2 Variables

- Computer languages use memory locations to store values. These named containers are called **variables**. There are complex rules as to what is a valid variable name. For simplicity, this topic will restrict the variable names to simple rules.
- Variable names:
 - must not start with a number
 - can only contain upper and lower case letters, numbers, and the underscore character (`_`), and cannot contain spaces
 - cannot be JavaScript keywords. The following are JavaScript keywords which should not be used as variable names.

abstract, arguments, Array, boolean, break, byte, case, catch, char, class, const, continue, Date, debugger, default, delete, do, double, else, enum, eval, export, extends, final, finally, float, for, function, goto, hasOwnProperty, if, implements, import, in, Infinity, instanceof, int, interface, isFinite, isNaN, isPrototypeOf, length, let, long, Math, name, NaN, native, new, null, Number, Object, package, private, protected, prototype, public, return, short, static, String, super, switch, synchronized, this, throw, throws, toString, transient, true, try, typeof, undefined, valueOf, var, void, volatile, while, with, yield, false

WORKED EXAMPLE 2 Checking the validity of a variable names

Identify whether each of the following is a valid variable name.

- a. `variableName` b. `100_hundred` c. `base^power`
 d. `number_1` e. `while`

THINK

- a. `variableName` is valid because it only uses letters.
 b. `100_hundred` is invalid because it uses a number as the first character.
 c. `base^power` is invalid because it uses a `^` character.
 d. `number_1` is valid as it only uses letters, numbers and the `_` character. It also does not start with a number.
 e. `while` is invalid because "`while`" is a JavaScript keyword.

WRITE

- a. True
 b. False
 c. False
 d. True
 e. False

13.2.3 Numerical expressions

- In JavaScript, numerical expressions involving numbers, brackets, and plus and minus signs evaluate normally.
- Multiplication uses the character `*`.
- Division uses the character `/`.
- Fractions can be evaluated using the division character, but the numerator and denominator expressions must be put in brackets.

$$\bullet \frac{3+1}{31+1+32} = (3+1) / (31+1+32)$$

- The `%` symbol is used to find remainder after a division. For example, `76%7` evaluates to `6`, as 7 divides into 76 ten times with a 6 remainder.

b. 1. Identify the variable name, which in this case is `topic`.

2. Identify the value, which in this case is `"Programming"`.

3. Apply the JavaScript structure to write the statement.

b.

```
var topic = "Programming";
```

c. 1. Identify the variable name, which in this case is `seconds`.

2. Identify the mathematical expression, which in this case is $60 \times 60 \times 24 \times 7$.

3. Convert the mathematical expression into a JavaScript expression: `60*60*24*7`.


4. Apply the JavaScript structure to write the statement.

c.

```
var seconds = 60*60*24*7;
```

13.2.5 Reassigning

- A variable can change value and type as the program runs through a sequence of statements in order. When a variable is reassigned, the statement does not require JavaScript keyword `var`. JavaScript uses the following sequence to assign and reassign a value or expression to a variable. The value of the variable changes with each assignment.



WORKED EXAMPLE 5 Simulating programs manually

Manually simulate running the following programs. For each program:

- identify the final type of the variable `a`
- determine the final value stored in the variable `a`.

- ```
var a = 123.3;
a = "123.3";
```
- ```
var a = (2+3+15)/(20-7-3);  
a = a / (40*7/140);
```
- ```
var a = "String";
var b = 122;
a = b;
b = "String";
```

#### THINK

- Write the initial assignment.
  - Write the reassignment.
- `"123.3"` is in quotation marks, so this value is a string.
  - The last value assigned to `a` was `"123.3"`.

#### WRITE

- `a = 123.3`
  - `a = "123.3"`
- String
  - `"123.3"`

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>b.</b></p> <ol style="list-style-type: none"> <li>1. Write the initial assignment.</li> <li>2. Evaluate the expression.</li> <li>3. Write the reassignment.</li> <li>4. On the right side of the assignment, substitute 2 for <b>a</b>.</li> <li>5. Substitute 2 for <math>40 * 7 / 140</math>.</li> <li>6. Substitute 1 for <math>2 / 2</math>.</li> </ol> <p><b>i.</b> 1 is a numerical value not in quotation marks, so this value is a number.</p> <p><b>ii.</b> The last value assigned to <b>a</b> was 1.</p> | <p><b>b.</b> <math>a = (2+3+15) / (20-7-3)</math></p> <p><math>a = 2</math></p> <p><math>a = a / (40 * 7 / 140)</math></p> <p><math>a = 2 / (40 * 7 / 140)</math></p> <p><math>a = 2 / 2</math></p> <p><math>a = 1</math></p> <p><b>i.</b> Number</p> <p><b>ii.</b> 1</p> |
| <p><b>c.</b></p> <ol style="list-style-type: none"> <li>1. Write the initial assignment to <b>a</b>.</li> <li>2. Write the initial assignment to <b>b</b>.</li> <li>3. Write the reassignment to <b>a</b>.</li> <li>4. Substitute 122 for <b>b</b>.</li> <li>5. Write the reassignment to <b>b</b>.</li> </ol> <p><b>i.</b> 122 is a numerical value not in quotation marks, so this value is a number.</p> <p><b>ii.</b> The last value assigned to <b>a</b> was 122.</p>                                                | <p><b>c.</b> <math>a = \text{"String"}</math></p> <p><math>b = 122</math></p> <p><math>a = b</math></p> <p><math>a = 122</math></p> <p><math>b = \text{"String"}</math></p> <p><b>i.</b> Number</p> <p><b>ii.</b> 122</p>                                                 |

## 13.2.6 Comments

- Someone reading a code may know the computer language in which the code is written, but that does not necessarily mean they will be able to understand it.
- Comments are added to codes to give hints about function of the program. The comments are completely ignored by the computer, so they are only for our benefit.
- JavaScript comments start with `//`. All the text after `//` on a line is ignored by the computer.

For example, consider the following line:

```
var PI = 3.1415926536; // A 10 digit approximation to π
```

The first part of the line, `var PI = 3.1415926536;`, is the actual JavaScript code to be run.

The second part (until the end of the line), `// A 10 digit approximation to  $\pi$ ,` is ignored by the computer and is only there for our own reference. The text could be changed and would make no difference to the running of the program.

### WORKED EXAMPLE 6 Adding comments to statements

**Add a comment to each statement of the form `// variable=value` where `variable` is the variable name and `value` is the evaluated value.**

**a.** `var pi = 3+1/10+3/100;`

**b.** `var myAge = 21+22;`

**c.** `var sum = 1+2+3+4;`

**THINK**

- a. 1. The variable name is `pi`.
2. The calculated value is  $3+1/10+4/100 = 3.14$ .
3. Append the comment.
- b. 1. The variable name is `myAge`.
2. The calculated value is  $21+22 = 43$ .
3. Append the comment.
- c. 1. The variable name is `sum`.
2. The calculated value is  $1+2+3+4 = 10$ .
3. Append the comment.

**WRITE**

- a. 


```
var pi=3+1/10+4/100;//pi=3.14
```
- b. 

```
var myAge=21+22; // myAge=43
```
- c. 

```
var sum=1+2+3+4; // sum=10
```

## 13.2.7 Output

- The **console** is a special region in a web browser for monitoring the running of JavaScript programs. Most web browsers, including Chrome, Firefox, Safari, Internet Explorer, Microsoft Edge and Opera, allow you to activate the console through the menu options.
- In order to see the result of an expression or value, the JavaScript `console.log` function can be used. This function outputs results to the console.



Note that the quotation marks at the start and end of strings are not shown in the output.

### WORKED EXAMPLE 7 Determining the output of statements

Determine the output to the console after each of the following statements runs.

- a. `console.log(((12%5)+12)/(17%10));`
- b. `console.log("Some information");`

**THINK**

- a. 1. Show the original statement.
2.  $(12\%5)$  simplifies to 2.  
 $(17\%10)$  simplifies to 7.

**WRITE**

- a. 

```
console.log(((12%5)+12)/(17%10));
```

```
console.log((2+12)/7);
```

3.  $(2+12)$  simplifies to 14.
  4.  $14/7$  simplifies to 2.
  5. Write the output to console.
- b.**
1. Show the original statement.
  2. Write the output to console. Note that the quotation marks in strings are not shown in the output.

```
console.log(14/7);
console.log(2);
2
b. console.log("Some information");
Some information
```

## on Resources

 **Interactivity** CodeBlocks Activity 1 (int-6573)

## Exercise 13.2 Programs

learn **on**

13.2 Quick quiz **on**

13.2 Exercise

### Individual pathways

**PRACTISE**

1, 2, 4, 5, 6

**CONSOLIDATE**

3, 7, 8, 9, 10, 14, 16

**MASTER**

11, 12, 13, 15, 17

### Fluency

1. **WE1** Answer these questions for each of the following values.
  - i. Classify the value as a number, a string or Boolean.
  - ii. Apply the expression `typeof value` to each value. Write the result of the expression `typeof value`.
 

|                             |                                      |
|-----------------------------|--------------------------------------|
| a. <code>-0.69314718</code> | b. <code>false</code>                |
| c. <code>"true"</code>      | d. <code>1000000</code>              |
| e. <code>true</code>        | f. <code>"Yet another string"</code> |
| g. <code>18</code>          |                                      |
2. Answer these questions for each of the following values.
  - i. Classify the value as a number, a string or Boolean.
  - ii. Apply the expression `typeof value` to each value. Write the result of the expression `typeof value`.
 

|                            |                                 |                        |
|----------------------------|---------------------------------|------------------------|
| a. <code>"false"</code>    | b. <code>"False"</code>         | c. <code>"-10"</code>  |
| d. <code>2.71828183</code> | e. <code>"What is this?"</code> | f. <code>"1283"</code> |
3. Answer these questions for each of the following expressions and values.
  - i. Classify the value as a number, string or Boolean.
  - ii. Apply the expression `typeof expression` to each expression or value. Write the result of the expression `typeof expression`.
 

|                          |                        |
|--------------------------|------------------------|
| a. <code>10/2</code>     | b. <code>"True"</code> |
| c. <code>10*10*10</code> | d. <code>99%5</code>   |



4. **WE2** Identify whether each of the following is a valid variable name.
- a. `4_four`                      b. `value_122`                      c. `divide/this`  
d. `value_-10`                      e. `for`
5. Identify whether of the following is a valid variable name.
- a. `someName`                      b. `throw`                      c. `some_name`  
d. `21_twenty_one`                      e. `multiply*this`
6. **WE5** Manually simulate running the following programs. For each program:
- identify the final type of the variable `result`
  - determine the final value stored in the variable `result`.
    - `var result = "Was a string, now a number.";  
result = 12`
    - `var result = 12;  
var x = 145;  
result = x;`
    - `var result = "Was a string. Now what?";  
result = false`
    - `var result = (3+15)/(20-2*7);  
result = result / (158%5);`
    - `var result = (1+2+3+4+5)/(1+1+1);  
result = result / (125/(5*5));  
result = true;`
    - `var result = "result is a string";  
var FALSE = true;  
result = FALSE;  
FALSE = "true";`
7. **WE7** Determine the output to the console after each of the following statements runs.
- `console.log("A simple string");`
  - `console.log("Some data");`
  - `console.log(111%(23%12));`
  - `console.log(true);`
  - `console.log("True");`
  - `console.log("12");`

### Understanding

8. **WE3** Write the following as JavaScript expressions.
- $\frac{12-6}{5-2}$                       b. The remainder from the division  $16 \div 5$
  - $12 + 12 - (-13 - 14 - 1)$                       d.  $12 \div (4 \times 3)$
  - $12 + (6 - 4 - 1)$
9. Write the following as JavaScript expressions.
- $112 \times 2.5$                       b.  $160 \div (4 \times 4 \times 8 \div 2)$
  - The remainder from the division  $111 \div 14$                       d.  $24 \times 2.5 \times 2.5$

10. **WE4** Write JavaScript statements to make the following variable assignments.
- Assign "JavaScript" to the variable `computerLanguage`.
  - Assign the value `true` to the variable `event`.
  - Assign the value `false` to the variable `finished`.
  - Assign the expression `1024*1024` to the variable `megabyte`.
  - Assign "Mathematics" to the variable `subject`.
  - Assign the mathematical expression  $60 \times 24 \times 7$  to the variable `minutes`.
  - Assign the value `1000000` to the variable `million`.
11. **WE6** Add a comment to each statement of the form `// variable=value` where `variable` is the variable name and `value` is the evaluated value.
- `var lotsOfFactors = 2*3*5*7*11*13;`
  - `var e = 2+7/10+1/100+8/1000+3/10000;`
  - `var product = 1*2*3*4;`
  - `var yourAge = 225/15;`
12. Manually simulate running the following program.
- ```
var final = 12;
final = final % 5;
final = 160 / final;
final = final * final;
final = final / 100;
final = final / 32;
final = final * final + final;
```
- Identify the final type of the variable `final`.
 - Determine the final value stored in the variable `final`.
13. Manually simulate running the following program.
- ```
var w = "xyz";
var x = 1;
var y = 1;
var z = x+y;
y = z*(x+x+x);
x = y+z+x;
z = (y+z)*x;
x = x+z+y*x;
z = w;
```
- Identify the final type of the variable `x`.
  - Identify the final type of the variable `y`.
  - Identify the final type of the variable `z`.
  - Determine the final value stored in the variable `x`.
  - Determine the final value stored in the variable `y`.
  - Determine the final value stored in the variable `z`.

## Reasoning


14. Write a program to:
- store the value `2` in the variable `inLitres`
  - convert `inLitres` to millilitres and store into the variable `inMillilitres`.
15. Write a program to:
- store the value `100` in the variable `crossSectionalArea`
  - store the value `12` in the variable `height`
  - calculate the volume of a prism given the `crossSectionalArea` and `height`, and store the result into the variable `prismVolume`.

## Problem solving

16. Write a program to calculate the **area** of a sector with **radius** equal to 10 and **angle** equal to 45 degrees.

The start of the program is shown below.


```
var radius = 10;
var angle = 45;
```



17. Write a program to calculate the **perimeter** of a sector with **radius** equal to 19 and **angle** equal to 120 degrees.

The start of the program is shown below.

```
var radius = 19;
var angle = 120;
```



# LESSON

## 13.3 Arrays

### LEARNING INTENTIONS

At the end of this lesson you should be able to:

- classify values as an array, number, string or Boolean
  - determine the JavaScript index of a value in an array
  - evaluate the JavaScript expression from a given array.
- 
- Computers are very good at repeating the same program many times extremely fast. However, it is redundant to apply the same program on the same data many times, so it is important to apply the program on a list of many values. This list is called an **array**.



### 13.3.1 List of values

- A variable can contain an array of values. An array takes the form `[value0, value1, value2, ...]` where `value0`, `value1`, `value2`, `...` are the different values of the array. An array can have zero, one or more values, and the values can be any type (i.e. number, string or Boolean).

- An array with many values can be split over multiple lines. For example:

```
var array = [
 value0,
 value1,
 value2,
 ...
];
```

## WORKED EXAMPLE 8 Classifying values

Classify each of the following as an array, number, string or Boolean.

- |              |                           |
|--------------|---------------------------|
| a. "Array"   | b. 3.4                    |
| c. [2,3,4]   | d. [ ]                    |
| e. ["Array"] | f. [true,false]           |
| g. false     | h. [1,true,"string",[ ] ] |

### THINK

- "Array" is in quotation marks, so this value is a string.
- 3.4 is a numerical value not in quotation marks, so this value is a number.
- [2,3,4] is a list of 3 numbers, so this is an array.
- [ ] is an empty list, so this is an array.
- ["Array"] is a list of 1 string, so this is an array.
- [true,false] is a list of 2 Booleans, so this is an array.
- false is a Boolean value.
- [1,true,"string",[ ] ] is a list of 4 different types of values, so this is an array.

### WRITE

- String
- Number
- Array
- Array
- Array
- Array
- Boolean
- Array

## 13.3.2 Testing for arrays

The function `Array.isArray(value)` can be used to test if `value` is an array. The function returns `true` if `value` is an array; otherwise, the function returns `false`.

## WORKED EXAMPLE 9 Evaluating JavaScript expressions manually

Evaluate the following JavaScript expressions manually.

- |                                     |                                         |
|-------------------------------------|-----------------------------------------|
| a. <code>Array.isArray([ ])</code>  | b. <code>Array.isArray("[3]")</code>    |
| c. <code>Array.isArray(true)</code> | d. <code>Array.isArray("array")</code>  |
| e. <code>Array.isArray([2])</code>  | f. <code>Array.isArray([2,"45"])</code> |

### THINK

- [ ] is an empty list, so this is an array.
  - The function `Array.isArray` returns `true`.
- "[3]" is in quotation marks, so this value is a string, even though the contents looks like an array.
  - The function `Array.isArray` returns `false`.
- `true` is a Boolean value.
  - The function `Array.isArray` returns `false`.
- "array" is in quotation marks, so this value is a string.
  - The function `Array.isArray` returns `false`.
- [2] is a list of 1 number, so this is an array.
  - The function `Array.isArray` returns `true`.
- [2,"45"] is a list of 2 different types of values, so this is an array.
  - The function `Array.isArray` returns `true`.

### WRITE

- `true`
  - `false`
- `false`
  - `false`
- `false`
  - `false`
- `true`
  - `true`
- `true`
  - `true`

### 13.3.3 Length of an array

The expression `array.length` will return the number of values in the `array`.

#### WORKED EXAMPLE 10 Evaluating JavaScript expressions manually

Evaluate the following JavaScript expressions manually.

- `[] .length`
- `[3] .length`
- `[99, "99"] .length`
- `[3.1416, "pi", true, [2, 3]] .length`

#### THINK

- `[]` is an empty list, so it has a length of 0.
- `[3]` is a list of 1 number.
- `[99, "99"]` is a list of 2 different types of values.
- `[3.1416, "pi", true, [2, 3]]` is a list of 4 different types of values.

*Note:* Do not count the number of values in the `[2, 3]` array, as this is counted as one value in the longer array.

#### WRITE

- 0
- 1
- 2
- 4

### 13.3.4 Array indexes

- When we isolate values in a list, we usually count the item at the start of the list as the first term. JavaScript arrays are slightly different: the values inside an array are accessed using an integer **index** starting at 0. That is, the first value in an `array` (of length `N`) is accessed with index 0, the second value is accessed with index 1, the third value is accessed with index 2 and the last value is accessed with index `N-1`.

|                          |                |                      |                |                       |                |                      |                     |                     |                 |
|--------------------------|----------------|----------------------|----------------|-----------------------|----------------|----------------------|---------------------|---------------------|-----------------|
| <code>var array =</code> | <code>[</code> | <code>"first"</code> | <code>,</code> | <code>"second"</code> | <code>,</code> | <code>"third"</code> | <code>, ...,</code> | <code>"last"</code> | <code>];</code> |
| <b>Location</b>          |                | 1st                  |                | 2nd                   |                | 3rd                  | ...                 | Nth                 |                 |
| <b>JavaScript index</b>  |                | 0                    |                | 1                     |                | 2                    |                     | <code>N-1</code>    |                 |

#### WORKED EXAMPLE 11 Determining the JavaScript index of values in an array

Identify the JavaScript index of the following values in the array

`["first", "second", "third", "fourth", "second last", "last"]`.

- `"first"`
- `"third"`
- `"last"`
- `"second last"`

#### THINK

- `"first"` is the 1st value, so its index is 0.
- `"third"` is the 3rd value, so its index is 2.
- `"last"` is the 6th value, so its index is 5.
- `"second last"` is the 5th value, so its index is 4.

#### WRITE

- 0
- 2
- 5
- 4

### 13.3.5 Accessing values in an array

- A value inside an `array` is accessed by its `index` using the expression `array[index]`. This expression returns the value with that index.

#### WORKED EXAMPLE 12 Evaluating JavaScript expressions manually from a given array

Given the array assignment

```
var array = ["a string", true, 4, 5, "false"];
```

evaluate the following JavaScript expressions manually.

- a. `array[1]`
- b. `array[3]`
- c. `array[2]`
- d. `array[array[2]]`
- e. `[1, 2, 5][2]`

#### THINK

- a. `array[1]` has an index of 1, which is the 2nd value in the array.
- b. `array[3]` has an index of 3, which is the 4th value in the array.
- c. `array[2]` has an index of 2, which is the 3rd value in the array.
- d. 1. `array[2]` has an index of 2, which is the 3rd value in the array.  
2. The 3rd value is 4.  
3. Write the original expression.  
4. Substitute 4 for `array[2]`.  
5. `array[4]` has an index of 4, which is the 5th value in the array.
- e. 1. Indexes can be applied directly to an array. `[1, 2, 5][2]` has an index of 2, which is the 3rd value in the array `[1, 2, 5]`.  
2. The 3rd value in the array `[1, 2, 5]` is 5.

#### WRITE

- a. `true`
- b. `5`
- c. `4`
- d.  

```
array[2]=4
array[array[2]]
= array[4]
= "false"
```
- e.  

```
[1, 2, 5][2] = 5
```

#### Resources

 **Interactivity** CodeBlocks Activity 2 (int-6574)

## Individual pathways

## ■ PRACTISE

1, 3, 5, 6, 7

## ■ CONSOLIDATE

2, 4, 8, 10, 12, 14, 16

## ■ MASTER

9, 11, 13, 15, 17

## Fluency

- WE8** Classify each of the following as an array, number, string or Boolean.
  - $3+0.1+0.04$
  - `[2.1, 3.2, 4.3, 5.4, 6.5, 7.6]`
  - `[true, false, true, true]`
  - `false`
  - $1237.3+1.2$
  - `["a", "b", "c", "d"]`
- Classify each of the following as an array, number, string or Boolean.
  - `["Many", "strings", "in", "an", "array"]`
  - `[1.3/2, false, "one type of many", []]`
  - `"Anthony"`
  - `[]`
  - `["A single string in an array"]`
  - `"Is this an array, number or string?"`
- WE9** Evaluate the following JavaScript expressions manually.
  - `Array.isArray(["true, false, true, true"])`
  - `Array.isArray([true, "true", "True", "TRUE"])`
  - `Array.isArray([3.14159, 2.718282])`
  - `Array.isArray(true)`
  - `Array.isArray([2.2, -13])`
  - `Array.isArray([])`
- Evaluate the following JavaScript expressions manually.
  - `Array.isArray("[2.1, 3.4, 5]")`
  - `Array.isArray(false)`
  - `Array.isArray("[true, false, true, true]")`
  - `Array.isArray([2, "45", true, false, [1, 23, 4]])`
  - `Array.isArray("Strings are like arrays")`
- WE10** Evaluate the following JavaScript expressions manually.
  - `[true, "true", "True", "TRUE"].length`
  - `[2.718282, "e", true, [2, 7, 1, 8, 2, 8, 2].length].length`
  - `[2, 3, 4, 5].length`
  - `[3, [2, 3], 4].length`
- WE11** Identify the JavaScript index of the following values in the array `[3, 1, 4, 1, 5, 9, 2, 6, 5, 3, 5, 8, 9, 7, 9, 3]`.
 

|      |      |      |      |      |
|------|------|------|------|------|
| a. 2 | b. 4 | c. 6 | d. 7 | e. 8 |
|------|------|------|------|------|

7. **WE12** Given the array assignment
- ```
var listOfItems = [true,5,4,false,2.3,"some string",
[2,31]];
```
- evaluate the following JavaScript expressions manually.
- `listOfItems[2]`
 - `listOfItems[0]`
 - `listOfItems[1+3]`
 - `listOfItems[listOfItems[1]]`

Understanding

8. Evaluate the following JavaScript expressions manually.
- `[2,3,4,5][0]`
 - `Array.isArray([3,2][1])`
 - `[2.718282,"e",true,[2,7,1,8,2,8,2].length][1+1]`
 - `typeof Array.isArray(["[false,2,3]"])`
 - `Array.isArray([false,"false","False","FALSE"].length)`
9. Evaluate the following JavaScript expressions manually.
- `typeof Array.isArray([3,[2,4]][1])`
 - `Array.isArray([2.2,[-13]][1].length)`
 - `[Array.isArray([[]].length),2]`
 - `[3,[2,3],4][2]`
10. Manually simulate running the following program.
- ```
var value = 12;
value = [value,value,value*2];
value = [value.length,value[0],value[1]*value[2]];
value = value.length;
value = [value, value*value, value*value*3];
```
- Identify the final type of the variable `value`.
  - Determine the final value stored in the variable `value`.
11. Manually simulate running the following program.
- ```
var pi = 3.14;
var pi100 = pi*100;
var d0 = pi100%10;
var d1 = ((pi100-d0)/10)%10;
var d2 = ((pi100-10*d1-d0)/100)%10;
var digits = [d2,d1,d0];
```
- Identify the final type of the variable `digits`.
 - Determine the final value stored in the variable `digits`.

12. Manually simulate running the following program.

```
var values = [10,2,3,5,6];
var total = 0;
total = total + values[0];
total = total + values[1];
total = total + values[2];
total = total + values[3];
total = total + values[4];
var mean = total/values.length;
```

- Identify the final type of the variable `mean`.
 - Determine the final value stored in the variable `mean`.
13. Manually simulate running the following program.
- ```
var numbers = [1,2,3,4,5,6];
var factorial = 1;
factorial = factorial*numbers[0];
factorial = factorial*numbers[1];
factorial = factorial*numbers[2];
factorial = factorial*numbers[3];
factorial = factorial*numbers[4];
```
- Identify the final type of the variable `factorial`.
  - Determine the final value stored in the variable `factorial`.

### Reasoning

14. Write a program to:
- store the values 2, 3, 5, 7, 11 and 13 in the array `primes`
  - multiply all the values and store into the variable `primorial`.
15. Write a program to total all the prime numbers between 2 and 20.

### Problem solving

16. Manually determine all of the possible unique combinations of rearranging the items in the array `[1,2,3]`. Store this list as an array of arrays in the variable `combinations`.
17. Let a three-item array represent three coin tosses, where one side of the coin shows `true` and the other `false`. For example, the throw below would be represented as `[true,true,false]`.



Determine all of the possible unique combinations of tossing the three coins. Store this list as an array of arrays in the variable `sampleSpace`.

# LESSON

## 13.4 Loops

### LEARNING INTENTIONS

At the end of this lesson you should be able to:

- evaluate JavaScript expressions from a given statement
- understand, count and evaluate loops.

### 13.4.1 Loops and Boolean expressions

- A similar task can be repeated many times in sequence. For example, the following program calculates  $4 + 4 + 4 + 4 + 4$  and outputs the sum.

```
var sum = 0;
sum = sum + 4;
sum = sum + 4;
sum = sum + 4;
sum = sum + 4;
sum = sum + 4;
console.log(sum);
```



- Loops can execute a task many times without needing multiple copies of the same code.
- Two numerical values can be compared with each other by using different combinations of symbols as shown in the following table. The comparisons return Boolean results as shown.

| Expression             | Description                                                                                                     |
|------------------------|-----------------------------------------------------------------------------------------------------------------|
| <code>!x</code>        | Opposite of <code>x</code> , where <code>x</code> is a Boolean.                                                 |
| <code>x1===x2</code>   | <code>true</code> if <code>x1</code> equals <code>x2</code> , otherwise <code>false</code>                      |
| <code>x1!==x2</code>   | <code>true</code> if <code>x1</code> does not equal <code>x2</code> , otherwise <code>false</code>              |
| <code>x1&gt;x2</code>  | <code>true</code> if <code>x1</code> is greater than <code>x2</code> , otherwise <code>false</code>             |
| <code>x1&gt;=x2</code> | <code>true</code> if <code>x1</code> is greater than or equal to <code>x2</code> , otherwise <code>false</code> |
| <code>x1&lt;x2</code>  | <code>true</code> if <code>x1</code> is less than <code>x2</code> , otherwise <code>false</code>                |
| <code>x1&lt;=x2</code> | <code>true</code> if <code>x1</code> is less than or equal to <code>x2</code> , otherwise <code>false</code>    |

### WORKED EXAMPLE 13 Evaluating JavaScript expressions manually

Evaluate the following JavaScript expressions manually.

- |                               |                               |                             |
|-------------------------------|-------------------------------|-----------------------------|
| a. <code>!false</code>        | b. <code>!true</code>         | c. <code>3.24===3.42</code> |
| d. <code>2.4!==4.2</code>     | e. <code>6.4&gt;-3.2</code>   | f. <code>9.9&gt;=9.9</code> |
| g. <code>3.005&lt;3.01</code> | h. <code>1.99&lt;=1.01</code> |                             |

#### THINK

- The opposite of `false` is `true`.
- The opposite of `true` is `false`.
- The statement "`3.24` equals `3.42`" is `false`.

#### WRITE

- `true`
- `false`
- `false`

- |                                                                 |          |
|-----------------------------------------------------------------|----------|
| d. The statement "2.4 does not equal 4.2" is true.              | d. true  |
| e. The statement "6.4 is greater than -3.2" is true.            | e. true  |
| f. The statement "9.9 is greater than or equal to 9.9" is true. | f. true  |
| g. The statement "3.005 is less than 3.01" is true.             | g. true  |
| h. The statement "1.99 is less than or equal to 1.01" is false. | h. false |

- The statement `boolean1 && boolean2` returns `true` if `boolean1` and `boolean2` are both `true`; otherwise, the statement returns `false`.

### WORKED EXAMPLE 14 Evaluating JavaScript && expressions manually

Evaluate the following JavaScript expressions manually.

- `false && false`
- `true && false`
- `false && true`
- `true && true`

#### THINK

- At least one Boolean is `false`.
- At least one Boolean is `false`.
- At least one Boolean is `false`.
- Both of the Booleans are `true`.

#### WRITE

- `false`
- `false`
- `false`
- `true`

- The statement `boolean1 || boolean2` returns `true` if either `boolean1` or `boolean2` are `true`; otherwise, the statement returns `false`.

### WORKED EXAMPLE 15 Evaluating JavaScript || expressions manually

Evaluate the following JavaScript expressions manually.

- `false || false`
- `true || false`
- `false || true`
- `true || true`

#### THINK

- Both of the Booleans are `false`.
- At least one Boolean is `true`.
- At least one Boolean is `true`.
- At least one Boolean is `true`.

#### WRITE

- `false`
- `true`
- `true`
- `true`



### 13.4.3 While loops

- In JavaScript, a `while` loop is used to repeat a section of code {`statement 1`, `statement 2`, ..., `statement n`} while `repeatDecision` is `true`.

#### Using `while` loops to repeat sections of code

```
while (repeatDecision) {
 statement 1
 statement 2
 statement n
}
```

- It is extremely important that at least one statement in the loop eventually changes the `repeatDecision` expression to `false`. If `repeatDecision` never changes to `false`, the program will never finish (without intervention). This is referred to as an **infinite loop**.

#### WORKED EXAMPLE 17 Checking for infinite loops

Each of the following causes an infinite loop. True or False?

- |                                                                                                  |                                                                                                                  |
|--------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| <b>a.</b> <pre>while(true) {<br/>    console.log("For ever");<br/>}</pre>                        | <b>b.</b> <pre>while(false) {<br/>    console.log("Never");<br/>}</pre>                                          |
| <b>c.</b> <pre>var y=1;<br/>while(y&lt;=10) {<br/>    console.log(y);<br/>    y=y+1;<br/>}</pre> | <b>d.</b> <pre>var z = 1;<br/>var y = 1;<br/>while(z&lt;=5) {<br/>    console.log(y);<br/>    y=2*y;<br/>}</pre> |

#### THINK

- |                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                   |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>a.</b> 1. <code>repeatDecision</code> is <code>true</code> .<br>2. <code>repeatDecision</code> never changes to <code>false</code> and causes an infinite loop.                                                                                                   | <b>b.</b> 1. <code>repeatDecision</code> is <code>false</code> .<br>2. <code>repeatDecision</code> starts <code>false</code> , so the loop is never executed.                                                                                                                     |
| <b>c.</b> 1. <code>repeatDecision</code> is <code>y&lt;=10</code> .<br>2. <code>Y</code> is incremented each time the loop is executed with <code>y=y+1</code> ;<br>3. After 10 executions, <code>y=11</code> and <code>y&lt;=10</code> becomes <code>false</code> . | <b>d.</b> 1. <code>repeatDecision</code> is <code>z&lt;=5</code> .<br>2. <code>z</code> is initialised at 1 and never changes inside the loop.<br>3. <code>repeatDecision</code> is <code>z&lt;=5</code> and never changes to <code>false</code> , so it causes an infinite loop. |

#### WRITE

- |                    |                    |
|--------------------|--------------------|
| <b>a.</b><br>True  | <b>b.</b><br>False |
| <b>c.</b><br>False | <b>d.</b><br>True  |

## 13.4.4 Loop a number of times

The following structure is one way to repeat a section of code { `statement 1`, `statement 2`, ..., `statement n` } a fixed `number` of times. The variable `i` is an index used to track the number of times through the loop.

### Using `while` to repeat code a fixed number of times

```
var i=1;
while (i <= number) {
 statement 1
 statement 2
 statement n
 i=i+1;
}
```

### WORKED EXAMPLE 18 Counting loop code execution in programs

State how many times the loop code is executed in each of the following programs.

- a. 

```
var i=1;
while (i<=42) {
 console.log(i);
 i=i+1;
}
```
- b. 

```
var a=1;
var i=1;
while (i<=1000) {
 a=a+i;
 i=i+1;
}
```
- c. 

```
var j=1;
var b=1;
while (j<=5) {
 b=b*j;
 j=j+1;
}
```

#### THINK

- a. The number is 42.
- b. The number is 1000.
- c. 1. The change of the index from `i` to `j` is irrelevant to the loop count. Any variable can be used as an index as long as it is changed at every location in the loop structure (i.e. at initialisation, in the decision and when it is increased by 1).
2. The number is 5.

#### WRITE

- a. 42
- b. 1000
- c.
- 5

- The previous program can be restructured to calculate  $4 + 4 + 4 + 4 + 4$  and output the total. Notice the use of comments after each line to explain what the code is doing.

```

var total = 0; // Initialise the running total
var i = 1; // i tracks number from 1 to 5
while (i <= 5) { // Check i does not exceed 5
 total = total + 4; // Add 4 to the running total
 i = i + 1; // Increment i
}
console.log(total); // Output the total

```

### WORKED EXAMPLE 19 Evaluating loops and determining the stored value

For each of the following programs:

- state how many times the loop is executed
- determine the final value stored in the variable  $a$ .

- ```

var a=1;
var i=1;
while (i<=4) {
    a=a*i;
    i=i+1;
}

```
- ```

var a=0;
var i=1;
while (i<=3) {
 a=a+i;
 i=i+1;
}

```
- ```

var a=0;
var i=1;
while (i<=24334) {
    a=a+7;
    i=i+1;
}

```

THINK

- The number is 4.
 - Write the first assignment involving a .
 - The statement is $a = a * i$.
 - Repeat for all values of i .
 - Note the pattern.
 - Evaluate a .

WRITE

- ```

4
a=1

a=a*1
a=a*2
a=a*3
a=a*4
a=1*1*2*3*4
a=24

```

- b. i. The number is 3.  
 ii. 1. Write the first assignment involving a.  
 2. The statement is  $a = a + i$ ;  
 3. Repeat for all values of i.

4. Note the pattern.  
 5. Evaluate a.

- c. i. The number is 24334.  
 ii. 1. Write the first assignment involving a.

2. The statement is  $a = a + 7$ ;  
 3. Repeat for 24334.

4. Note the pattern.  
 5. Note the simplification as a multiplication.  
 6. Evaluate a.

- b. 3  
 $a = 0$

$a = a + 1$   
 $a = a + 2$   
 $a = a + 3$   
 $a = 0 + 1 + 2 + 3$   
 $a = 6$

- c. 24334  
 $a = 0$

$a = a + 7$ ; // i=1  
 $a = a + 7$ ; // i=2  
 ...  
 $a = a + 7$ ; // i=24334  
 $a = 0 + 7 + 7 + \dots + 7$   
 $a = 7 * 24334$   
 $a = 170338$

## on Resources

 **Interactivity** CodeBlocks Activity 3 (int-6575)

## Exercise 13.4 Loops

learn on

13.4 Quick quiz **on**

13.4 Exercise

### Individual pathways

**PRACTISE**

1, 2, 4, 5, 6, 7

**CONSOLIDATE**

3, 8, 9, 10, 12, 14, 16

**MASTER**

11, 13, 15, 17

### Fluency

1. **WE13** Evaluate the following JavaScript expressions manually.

a.  $16.41 > -13.12$

b.  $-999 > = -999$

c.  $4.06 < 4.1$

d.  $!true$

e.  $113.3 === 113.30$

f.  $2.41 !== 412.2$

g.  $101.99 < = 102.01$

2. Evaluate the following JavaScript expressions manually.

a.  $12 === 21$

b.  $13.40 === 13.400$

c.  $21.41 !== -42212.22$

d.  $-1.1 > -1.12$

e.  $!false$

f.  $123.3 === 213.2$

g.  $-4 < = -4$



3. **WE14** Evaluate the following JavaScript expressions manually.

- a. `false&&true`
- b. `true&&false`
- c. `true&&true`
- d. `false&&false`

4. **WE15** Evaluate the following JavaScript expressions manually.

- a. `true||true`
- b. `false||true`
- c. `false||false`
- d. `true||false`

5. **WE16** Manually simulate running the following programs. For each program, determine the final value stored in `x`.

- a. 

```
var x=153;
if ((x%7)===6) {
 x = x*x;
}
```
- b. 

```
var x=false&&true;
if (x===false) {
 x = 100;
}
```

6. **WE17** Each of the following causes an infinite loop. True or False?

- a. 

```
while(false) {
 var i = 10;
 console.log(i);
}
```
- b. 

```
while(true) {
 console.log("Too many");
}
```



7. **WE18** State how many times the loop code is executed in each of the following programs.

- a. 

```
var i=1;
while (i<=99) {
 console.log(2*i);
 i=i+1;
}
```
- b. 

```
var u=1;
var i=1;
while (i<=100) {
 u=u*i;
 i=i+1;
}
```
- c. 

```
var k=1;
var b=1000;
while (k<=5) {
 b=b-k;
 k=k+1;
}
```
- d. 

```
while(false) {
 console.log("This sentence is ignored.");
}
```

```
e. var i=1;
 while (i<=900) {
 console.log(2*i);
 i=i+1;
 }
```

8. **WE19** For each of the following programs:

- state how many times the loop is executed.
- determine the final value stored in the variable **h**.

```
a. var h=0;
 var i=1;
 while (i<=14) {
 h=h+i;
 i=i+1;
 }
```

```
b. var h=1;
 var k=1;
 while (k<=9) {
 h=h*k;
 k=k+1;
 }
```

```
c. var h=0;
 var index=1;
 while (index <=5) {
 h=h*3;
 index=index+1;
 }
```

## Understanding

9. Consider the following program.

```
var total=0;
var i=1;
var j=1;
while (j<=5) {
 total=total+2*i;
 j=j+1;
}
```

- Determine the final value stored in the variable **total**.
- Determine the final value stored in the variable **j**.
- Determine the final value stored in the variable **i**.

10. Each of the following causes an infinite loop. True or False?

```
a. var repeat=true;
 var i=1;
 while(repeat) {
 repeat = (i%31)!=0;
 i=i+1;
 }
```

```

b. var repeat=true;
 var i=1;
 while(repeat) {
 repeat = (i%31) !==0;
 i=(i%30)+1;
 }

```

11. State how many times the loop code is executed in each of the following programs. Keep in mind the loop indexes are not always increased by one each time.

```

a. var max=1000;
 var m=1;
 while (m<=max) {
 console.log(m*m);
 m=m+1;
 }

```

```

b. var a=0;
 var b=100;
 while (a<=b) {
 a=a+2;
 b=b+1;
 }

```

```

c. var k=0;
 while (k<100) {
 k=k+1;
 }

```

```

d. var j=0;
 while (j<100) {
 j=j+2;
 }

```

12. Determine the final value stored in the variable `total` by the following program.

```

var total=0;
var x=1;
while (x<=100) {
 total = total+20;
 x=x+1;
}

```

13. Determine the output of the following program.

```

var total = 0;
var a=1;
while (a<=10) {
 total = total + a*a;
 a=a+1;
}
console.log(total);

```

### Reasoning

14. Write a program to multiply all of the integers between 1 and 10, and store the product into the variable `factorial_10`.
15. Write a program to add all of the integers between 1 and 1 000 000, and store the sum into the variable `total`.

## Problem solving

16. Given two variables `value1` and `value2` assigned to 362 and 232 respectively, assign the first positive integer to `product` where `value1` and `value2` divide exactly. Output the value of `product` to the console. The start and end of the program is shown below. Fill in the middle to find the value `product`.

```
var value1 = 362;
var value2 = 232;
...
console.log(product);
```

17. Given an `array` of integers `[362, 232, 4, 8, 5]`, assign the first positive integer to `product` where all the values in the `array` divide into `product` exactly. Write a program to find and output the value `product`. The start and end of the program is shown below. Fill in the middle to find the value `product`.

```
var array = [362, 232, 4, 8, 5];
...
console.log(product);
```

# LESSON

## 13.5 Set structures

### LEARNING INTENTIONS

At the end of this lesson you should be able to:

- represent data as an array
- determine the output to the generated console
- understand the concept of the mean of the values in an array.

### 13.5.1 Array set structure

- An array can be used to store a set of values. This set structure is useful to represent a list of values where the order might be important and values might be repeated in the list.

### WORKED EXAMPLE 20 Representing data as an array of values

For each of the following situations, represent the data as an array of values.

- Tim finished last in a race between friends. Jane was first to cross the line, closely followed by Wayne. Grace also finished the race.
- The list of prime numbers from 1 to 21
- The first 8 digits of  $\pi$  after the decimal place

#### THINK

- The friends finished the race in the order Jane, Wayne, Grace and Tim.

#### WRITE

- `["Jane", "Wayne", "Grace", "Tim"]`

- b. The primes between 1 and 21 are 2, 3, 5, 7, 11, 13, 17 and 19.      b. [2,3,5,7,11,13,17,19]
- c.  $\pi = 3.141\,592\,65$ , correct to 8 decimal places.      c. [1,4,1,5,9,2,6,5]  
The 8 digits after the decimal place are 1, 4, 1, 5, 9, 2, 6 and 5.

## 13.5.2 Loop over an array

The following structure is one way to repeat a section of code

```
{statement 1, statement 2, ..., statement n}
```

for each `item` in an `array`. The variable `i` is the array index and is also used to track the current loop iteration.

### Using loops over arrays to repeat a section of code

```
var array = [...];
var i=0;
while (i < array.length) {
 var item = array[i];
 statement 1
 statement 2
 statement n
 i=i+1;
}
```

## WORKED EXAMPLE 21 Counting loop code execution in programs

Determine how many times the loop code is executed in each of the following programs.

- a. 

```
var friends = ["Jane","Wayne","Grace","Tim"];
var places = ["1st","2nd","3rd","Last"];
var i=0;
while (i < friends.length) {
 var item = friends[i];
 var place = places[i];
 console.log(place+" : "+item);
 i=i+1;
}
```
- b. 

```
var sum = 0;
var primes = [2,3,5,7,11,13,17,19];
var j=0;
while (j < primes.length) {
 var prime = primes[j];
 sum = sum + prime;
 j=j+1;
}
console.log(sum);
```

```

c. var pi = 3;
 var place = 1;
 var digits = [1,4,1,5,9,2,6,5];
 var k=0;
 while (k < digits.length) {
 var item = digits[k];
 place = place / 10;
 pi = pi + item * place;
 k=k+1;
 }
 console.log(pi);

```

**THINK**

- a. 1. The code follows the structure to loop over the array `friends=["Jane","Wayne","Grace","Tim"]`.
- 2. The length of the `friends` array is 4.
- b. 1. The code follows the structure to loop over the array `primes=[2,3,5,7,11,13,17,19]`.
- 2. The length of the `primes` array is 8.
- c. 1. The code follows the structure to loop over the array `digits=[1,4,1,5,9,2,6,5]`.
- 2. The length of the `digits` array is 8.

**WRITE**

- a. 4
- b. 8
- c. 8

### 13.5.3 Sum

- There are various reasons for looping over an array. One reason is to sum all the values. The following structure is one way to sum all the values in an `array`.

**Determining the sum of an array**

```

var array = [...];
var total = 0;
var i=0;
while (i < array.length) {
 var item = array [i];
 total = total + item;
 i=i+1;
}
console.log(total);

```

1. First the program initialises the running `total` with the value `0`.
2. Then the `while` loop adds each `item` to a running `total` one at a time.
3. Finally, the program outputs the `total` to the console.

## WORKED EXAMPLE 22 Determining the output to the generated console

Determine the output to the console generated by the following program.

```
var sum = 0;
var primes = [2,3,5,7,11,13];
var j=0;
while (j < primes.length) {
 var item = primes[j];
 sum = sum + item;
 j=j+1;
}
console.log(sum);
```

### THINK

1. The code follows the structure to find the sum of all the items in the array `primes=[2,3,5,7,11,13]`.
2. Write the sum of all items in the array `primes` and assign to `sum`.
3. Evaluate the sum of items.
4. Write the `total` to the console.

### WRITE

```
sum=2+3+5+7+11+13
```

```
sum=41
```

```
41
```

## 13.5.4 Mean

- The mean or average of a set of values is the sum of all the values divided by the number of values. Finding the **mean** from an **array** is only a little more work after calculating the **total**. Building on the sum structure, the following structure is one way to find the **mean** of the values in an **array**.

### Determining the mean of an array

```
var array = [...];
var total = 0;
var i=0;
while (i < array.length) {
 var item = array[i];
 total = total + item;
 i=i+1;
}
var mean = total / array.length;
console.log(mean);
```

1. First the program initialises the running `total` with the value 0.
2. The `while` loop adds each `item` to a running `total` one at a time.
3. The `mean` is calculated by dividing the `total` with the number of items.
4. Finally, the program outputs the `mean` to the console.

## WORKED EXAMPLE 23 Determining the mean

Determine the output to the console generated by the following program.

```
var total = 0;
var digits = [3,1,4,1,5,9,2,6,5,4];
var k=0;
while (k < digits.length) {
 var item = digits[k];
 total = total + item;
 k=k+1;
}
var mean = total / digits.length;
console.log(mean);
```

### THINK

1. The code follows the structure to find the mean of all the items in the array `digits=[3,1,4,1,5,9,2,6,5,4]`.
2. Write the sum of all items in the array `digits` and assign to `total`.
3. Evaluate the sum of the items.
4. Write the assignment to calculate the `mean`.
5. Substitute 40 for `total` and 10 for `digits.length`.
6. Evaluate the `mean`.
7. Write the `mean` to the console.

### WRITE

```
total=3+1+4+1+5+9+2+6+5+4
total=40
mean=total/digits.length
mean=40/10

mean=4
4
```

## Exercise 13.5 Set structures

learnon

13.5 Quick quiz **on**

13.5 Exercise

### Individual pathways

#### ■ PRACTISE

1, 2, 4, 5, 6

#### ■ CONSOLIDATE

3, 7, 8, 11, 13

#### ■ MASTER

9, 10, 12, 14

### Fluency

1. **WE21** Determine how many times the loop code is executed in each of the following programs.

```
a. var e = 2;
 var base = 1;
 var eDigits = [7,1,8,2,8,1,8,2,8,4,5,9];
 var m=0;
```



```

while (m < eDigits.length) {
 var item = eDigits[m];
 base = base / 10;
 e = e + item * base;
 m=m+1;
}
console.log(e);
b. var d = ["Sun", "Mon", "Tues", "Wednes", "Thurs", "Fri",
"Saturn"];
var j=0;
while (j < d.length) {
 var start = d[j];
 console.log(start+"day");
 j=j+1;
}
c. var primorial = 1;
var primes = [2,3,5,7,11,13,17,19,23,29];
var j=0;
while (j < primes.length) {
 var prime = primes[j];
 primorial = primorial * prime;
 j=j+1;
}
console.log(primorial);

```

2. **WE22** Determine the output to the console generated by the program below, to 12 decimal places.

```

var e = 2;
var base = 1;
var eDigits = [7,1,8,2,8,1,8,2,8,4,5,9];
var m=0;
while (m < eDigits.length) {
 var item = eDigits[m];
 base = base / 10;
 e = e + item * base;
 m=m+1;
}
console.log(e);

```

3. Determine the output to the console generated by the program below.

```

var primorial = 1;
var primes = [2,3,5,7,11,13,17,19,23,29];
var j=0;
while (j < primes.length) {
 var prime = primes[j];
 primorial = primorial * prime;
 j=j+1;
}
console.log(primorial);

```



4. **WE23** Determine the output to the console generated by the program below.

```
var sum = 0;
var eDigits = [2,7,1,8,2,8,1,8,2,8,4,5,9];
var i=0;
while (i < eDigits.length) {
 var digit = eDigits[i];
 sum = sum + digit;
 i=i+1;
}
var mean = sum / eDigits.length;
console.log(mean);
```

### Understanding

5. **WE20** For each of the following situations, represent the data as an array of values.
- The days of the week starting at "Monday".
  - The months of the year.
  - The lowercase letters of alphabet.
  - The number of days in each month during a leap year.

6. Determine the output to the console generated by the program below.

```
var isPrime = true;
var testPrime=31;
var j = 2;
while (j < testPrime) {
 var notMultiple = testPrime%j !== 0;
 isPrime = isPrime && notMultiple;
 j = j+1;
}
console.log(isPrime);
```

7. Determine the output to the console generated by the program below.

```
var primorial = 1;
var testPrime=2;
while (testPrime <= 30) {
 var isPrime = true;
 var j = 2;
 while (j < testPrime) {
 var notMultiple = testPrime%j !== 0;
 isPrime = isPrime && notMultiple;
 j = j+1;
 }
 if (isPrime) {
 primorial = primorial * testPrime;
 }
 testPrime = testPrime+1;
}
console.log(primorial);
```



8. Consider the program below.

```
var values = [1.2,3,5,6,2,2,1];
var result = values[0];
var i = 0;
while (i < values.length) {
 var item = values[i];
 if (item > result) {
 result = item;
 }
 i = i + 1;
}
console.log(result);
```

- Determine the output to the console generated by the program.
- Explain the purpose of the program.

9. Consider the program below.

```
var values = [2,13,2,1,1,8];
var min = values[0];
var max = values[0];
var i = 0;
while (i < values.length) {
 var item = values[i];
 if (item > max) {
 max = item;
 }
 if (item < min) {
 min = item;
 }
 i = i + 1;
}
var result = max - min;
console.log(result);
```

- Determine the output to the console generated by the program.
- Explain the purpose of the program.

10. Consider that the function `Math.sqrt(value)` calculates the square root of `value`. Also consider the program below. Determine the output to the console generated by the program.

```
var scores = [10, 16, 6, 12];
var n = scores.length;
var sum = 0;
var i = 0;
while (i < n) {
 var x = scores[i];
 sum = sum + x;
 i = i + 1;
}
var mean = sum/n;
var sumSquares = 0;
var i = 0;
```





## Swapping the contents of variables

- Imagine you are holding an item in each of your hands. Each hand can hold only one of each item. In order to swap the items, you need to place one item down in a temporary location.

| Step | Diagram | Explanation                                                       | Swapping program                                  |
|------|---------|-------------------------------------------------------------------|---------------------------------------------------|
| 1.   |         | Start with green in the left hand and red in the right.           | <pre>var left = "green"; var right = "red";</pre> |
| 2.   |         | Move the contents in the right hand to the swap location.         | <pre>var swap = right;</pre>                      |
| 3.   |         | Move the contents in the left hand to the right hand.             | <pre>right = left;</pre>                          |
| 4.   |         | Finally, move the contents in the swap location to the left hand. | <pre>left = swap;</pre>                           |

## WORKED EXAMPLE 24 Identifying if programs swap the contents of variables

Each of the following programs swaps the contents of `variable1` and `variable2`. True or False?

a. `var variable1 = 12;`  
`var variable2 = 3;`  
`var swap = variable1;`  
`variable1 = variable2;`  
`variable2 = swap;`

b. `var variable1 = "blue";`  
`var variable2 = "yellow";`  
`var swap = variable1;`  
`variable2 = variable1;`  
`variable1 = swap;`

c. `var variable1 = true;`  
`var variable2 = false;`  
`var swap = variable2;`  
`variable2 = variable1;`  
`variable1 = swap;`

### THINK

- a. 1. Write the first assignment.  
2. Write the second assignment.  
3. Write the third assignment.  
4. Substitute 12 for `variable1`.  
5. Write the fourth assignment.  
6. Substitute 3 for `variable2`.  
7. Write the fifth assignment.  
8. Substitute 12 for `swap`.  
9. `variable1` and `variable2` have successfully swapped.
- b. 1. Write the first assignment.  
2. Write the second assignment.  
3. Write the third assignment.  
4. Substitute "blue" for `variable1`.  
5. Write the fourth assignment.  
6. Substitute "blue" for `variable1`.  
7. Write the fifth assignment.  
8. Substitute "blue" for `swap`.  
9. `variable1` and `variable2` both ended up with the same value, so they did not successfully swap.
- c. 1. Write the first assignment.  
2. Write the second assignment.  
3. Write the third assignment.  
4. Substitute `false` for `variable2`.  
5. Write the fourth assignment.  
6. Substitute `true` for `variable1`.  
7. Write the fifth assignment.  
8. Substitute `false` for `swap`.  
9. `variable1` and `variable2` have successfully swapped.

### WRITE

a. `variable1 = 12`  
`variable2 = 3`  
`swap = variable1`  
`swap = 12`  
`variable1 = variable2`  
`variable1 = 3`  
`variable2 = swap`  
`variable2 = 12`  
True

b. `variable1 = "blue"`  
`variable2 = "yellow"`  
`swap = variable1`  
`swap = "blue"`  
`variable2 = variable1`  
`variable1 = "blue"`  
`variable1 = swap`  
`variable2 = "blue"`  
False

c. `variable1 = true`  
`variable2 = false`  
`swap = variable2`  
`swap = false`  
`variable2 = variable1`  
`variable2 = true`  
`variable1 = swap`  
`variable2 = false`  
True

## WORKED EXAMPLE 25 Completing the program to swap the contents of variables

Complete the rest of the program to swap the contents of the two variables.

```
var containerA = "Cream";
var containerB = "Jam";
```

### THINK

1. Write the start of the program as provided.
2. Store the contents of `containerA` in a temporary variable `swap`.
3. Overwrite the contents of `containerA` with the value in `containerB`.
4. Overwrite the contents of `containerB` with the contents of the temporary variable `swap`.

### WRITE

```
var containerA = "Cream";
var containerB = "Jam";
var swap = containerA;

containerA = containerB;

containerB = swap;
```

## 13.6.2 Swapping two array items

- To sort an array, you must be able to swap values that are out of place. The following structure is one way to `swap` the two items in an `array` with the indexes `index1` and `index2`.

### Swapping two item arrays

```
var array = [...];
var swap = array[index1];
array[index1] = array[index2];
array[index2] = swap;
```

## WORKED EXAMPLE 26 Completing the program to swap items in an array

Complete the rest of the program to swap the contents of the 3rd and 6th item in the array.

```
var array = [15,11,2,4,5,9,11];
```

### THINK

1. Rewrite the start of the program.
2. Store the array index 2 of the 3rd item in the variable `index1`.
3. Store the array index 5 of the 6th item in the variable `index2`.
4. Store the contents of `array[index1]` in a temporary variable `swap`.
5. Overwrite the contents of `array[index1]` with the value in `array[index2]`.
6. Overwrite the contents of `array[index2]` with the contents of the temporary variable `swap`.

### WRITE

```
var array=[15,11,2,4,5,9,11];
var index1=2;

var index2=5;

var swap=array[index1];

array[index1]=array[index2];

array[index2]=swap;
```

## 13.6.3 Defining a function

- In more complex programs, it is useful to wrap a section of code in a JavaScript function. This gives the section of code `{statement 1, statement 2, ..., statement n}` a descriptive **name**. A function with a **name** can be given zero, one or more inputs, `x1, x2, ...`.

### Defining a function to wrap a section of code

```
function name(x1,x2,...) {
 statement 1
 statement 2
 statement n
 return output;
}
var store = name(v1,v2,...);
```

- Once a function is defined, it can be called with the statement `name(v1,v2,...);`. The value `v1` is assigned to the input `x1`, the value `v2` is assigned to the input `x2`, and so on.
- In JavaScript, a function **name** can **return** an internal value **output** to be **stored**.

### WORKED EXAMPLE 27 Determining the output of a given console

Determine the output to the console of each of the following programs.

- a. 

```
function outputSquared(r) {
 console.log(r*r);
}
outputSquared(-11);
```
- b. 

```
function hexagonPerimeter(side) {
 console.log(6*side);
}
hexagonPerimeter(8.5);
```
- c. 

```
function noInputs() {
 return "No inputs";
}
console.log(noInputs());
```

#### THINK

- a. 1. The input `r` equals `-11`.  
2. Write the output statement.  
3. Substitute `-11` for `r`.  
4. Evaluate the expression.  
5. Write the output to the console.
- b. 1. The input `side` equals `8.5`.  
2. Write the output statement.  
3. Substitute `8.5` for `side`.  
4. Evaluate the expression.  
5. Write the output to the console.

#### WRITE

- a. 

```
r=-11
console.log(r*r)
console.log(-11*-11)
console.log(121)
121
```
- b. 

```
side=8.5
console.log(6*side)
console.log(6*8.5)
console.log(51)
51
```



- c. 1. The function has no inputs.
2. Write the return expression.
3. Write the output statement.
4. Substitute "No inputs" for `noInputs()`.
5. Write the output to the console.

```
c. noInputs()="No inputs"
 console.log(noInputs());
 console.log("No inputs");
 No inputs
```

### 13.6.4 The swapping function

- A sorting algorithm requires swapping two array items. It is useful to define the swapping algorithm as a reusable function. The function `swapItems` has three inputs `array`, `index1` and `index2`, where `array` is the list of values, and `index1` and `index2` are the indexes of the items to be swapped.

#### Defining swapping algorithms as functions

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
var anArray = ["1","3","2"];
swapItems(anArray, 1, 2);
console.log(anArray);
```

- Once this function is defined, it can be called with the statement `swapItems(array, index1, index2)`.

#### WORKED EXAMPLE 28 Determining the output using `swapItems`

Given the swapping function shown below, determine the output to the console of each of the following programs.

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
```

- a. 

```
var digits = [1,2,3,4,5,6,7,8];
swapItems(digits, 2, 7);
console.log(digits[2]);
```
- b. 

```
var a = ["1st","2nd","last","3rd"];
swapItems(a, 2, 3);
console.log(a[a.length-1]);
```

**THINK**

- a. 1. Write the first assignment.
2. Calling `swapItems` with array `digits` and indexes 2 and 7 will swap the 3rd and 8th items.
3. Write the output statement.
4. `digits[2]` is the 3rd item in the `digits` array.
5. Substitute 8 for `digits[2]`.
6. Write the output to the console.
- b. 1. Write the first assignment.
2. Calling `swapItems` with array `a` and indexes 2 and 3 will swap the 3rd and 4th items.
3. Write the output statement.
4. Substitute 4 for `a.length`
5. Substitute 3 for `4-1`.
6. `a[3]` is the 4th item in the `a` array.
7. Substitute "last" for `a[3]`.
8. Write the output to the console.

**WRITE**

```
a. digits = [1,2,3,4,5,6,7,8]
 digits = [1,2,8,4,5,6,7,3]

 console.log(digits[2])

 console.log(8)
 8
b. a = ["1st","2nd","last","3rd"];
 a = ["1st","2nd","3rd","last"]

 console.log(a[a.length-1]);
 console.log(a[4-1]);
 console.log(a[3]);

 console.log("last")
 last
```

## 13.6.5 Decisions

- While sorting an array, you must make many decisions to swap two values in the array if they are out of place. In JavaScript, the `if` structure can be used to make a decision to swap items if `decision` is `true`.

**Using if structures to make decisions**

```
if (decision) {
 // swap
}
```

**WORKED EXAMPLE 29** Determining the final value using `if`

Manually simulate running the following programs. For each program, determine the final value stored in the array.

```
a. var primes=[2,3,5,13,11,7];
 var i=0;
 var j=1;
 if (primes[i]>primes[j]) {
 var swap=primes[i];
 primes[i]=primes[j];
 primes[j]=swap;
 }
```

```

b. var digits=[2,3,4,9,8,7];
 var i=3;
 var j=4;
 if (digits[i]>digits[j]) {
 var swap=digits[i];
 digits[i]=digits[j];
 digits[j]=swap;
 }

```

#### THINK

- a. 1. Write the first assignment.
2. Write the second assignment.
3. Write the third assignment.
4. Write the expression for the if `decision`.
5. Substitute 0 for `i` and 1 for `j`.
6. Substitute 2 for `primes[0]` and 3 for `primes[1]`.
7. Substitute `false` for `2>3`.
8. `decision` is `false`, so ignore the statement inside the `{}` block.
9. The array `primes` remains unchanged.
- b. 1. Write the first assignment.
2. Write the second assignment.
3. Write the third assignment.
4. Write the expression for the if `decision`.
5. Substitute 3 for `i` and 4 for `j`.
6. Substitute 9 for `digits[3]` and 8 for `digits[4]`.
7. Substitute `true` for `9>8`.
8. `decision` is `true`, so run the statements inside the `{}` block.
9. The code inside the `{}` block swaps the `digits[3]` and `digits[4]` items.

#### WRITE

```

a. primes=[2,3,5,13,11,7]
 i=0
 j=1

 decision=primes[i]>primes[j]
 decision=primes[0]>primes[1]
 decision=2>3

 decision=false

 primes=[2,3,5,13,11,7]
b. digits=[2,3,4,9,8,7]
 i=3
 j=4

 decision=digits[i]>digits[j]
 decision=digits[3]>digits[4]
 decision=9>8

 decision=true

 digits=[2,3,4,8,9,7]

```

- In JavaScript, the `if else` structure is used to execute one of the two different sections of code.
  1. `{statement 1, statement 2, ..., statement n}` if `decision` is `true`.
  2. `{statement A, statement B, ..., statement Z}` if `decision` is `false`.

### Using if else structures

```
if (decision) {
 statement 1
 statement 2
 statement n
} else {
 statement A
 statement B
 statement Z
}
```

### WORKED EXAMPLE 30 Determining the final value using if else

Manually simulate running the following programs. For each program, determine the final value stored in the variable `mean`.

a. 

```
var mean=0;
var values = [4,10];
if (values.length > 1) {
 mean = (values[0]+values[1])/2;
} else {
 mean = values[0];
}
```

b. 

```
var mean=0;
var values = [99];
if (values.length > 1) {
 mean = (values[0]+values[1])/2;
} else {
 mean = values[0];
}
```

#### THINK

- a. 1. Write the first assignment.
2. Write the second assignment.
3. Write the `decision`.
4. Substitute 2 for `values.length`.
5. Substitute `true` for `2 > 1`.
6. `decision` is `true`, so run the statements inside the first `{}` block
7. Substitute 4 for `values[0]` and 10 for `values[1]`.
8. Substitute 7 for  $(4+10)/2$ .
- b. 1. Write the first assignment.
2. Write the second assignment.
3. Write the `decision`.

#### WRITE

a. 

```
mean=0
values = [4,10]
decision = values.length > 1
= 2 > 1
= true
mean =
(values[0]+values[1])/2
mean = (4+10)/2

mean = 7
```

b. 

```
mean = 0
values = [99]
decision = values.length > 1
```

- |                                                                                                                |                               |
|----------------------------------------------------------------------------------------------------------------|-------------------------------|
| 4. Substitute 1 for <code>values.length</code> .                                                               | <code>= 1 &gt; 1</code>       |
| 5. Substitute <code>false</code> for <code>1 &gt; 1</code> .                                                   | <code>= false</code>          |
| 6. <code>decision</code> is <code>false</code> , so run the statements inside the second <code>{}</code> block | <code>mean = values[0]</code> |
| 7. Substitute 99 for <code>values[0]</code> .                                                                  | <code>mean = 99</code>        |

### 13.6.6 Bubble sort

- One of the simplest sorting algorithms is the **bubble sort**. The algorithm scans an array, comparing adjacent values. If any adjacent values are out of place, they are swapped. This scanning process is repeated until no swaps are made during a complete scan.



| Values |   |   |   |   | Scan | Array           | Swap?         | Action | After swap      |
|--------|---|---|---|---|------|-----------------|---------------|--------|-----------------|
| 9      | 1 | 5 | 2 | 3 | 1    | [9, 1, 5, 2, 3] | 9 > 1 = true  | Swap   | [1, 9, 5, 2, 3] |
| 1      | 9 | 5 | 2 | 3 | 1    | [1, 9, 5, 2, 3] | 9 > 5 = true  | Swap   | [1, 5, 9, 2, 3] |
| 1      | 5 | 9 | 2 | 3 | 1    | [1, 5, 9, 2, 3] | 9 > 2 = true  | Swap   | [1, 5, 2, 9, 3] |
| 1      | 5 | 2 | 9 | 3 | 1    | [1, 5, 2, 9, 3] | 9 > 3 = true  | Swap   | [1, 5, 2, 3, 9] |
| 1      | 5 | 2 | 3 | 9 | 2    | [1, 5, 2, 3, 9] | 1 > 5 = false |        |                 |
| 1      | 5 | 2 | 3 | 9 | 2    | [1, 5, 2, 3, 9] | 5 > 2 = true  | Swap   | [1, 2, 5, 3, 9] |
| 1      | 2 | 5 | 3 | 9 | 2    | [1, 2, 5, 3, 9] | 5 > 3 = true  | Swap   | [1, 2, 3, 5, 9] |
| 1      | 2 | 3 | 5 | 9 | 2    | [1, 2, 3, 5, 9] | 5 > 9 = false |        |                 |
| 1      | 2 | 3 | 5 | 9 | 3    | [1, 2, 3, 5, 9] | 1 > 2 = false |        |                 |
| 1      | 2 | 3 | 5 | 9 | 3    | [1, 2, 3, 5, 9] | 2 > 3 = false |        |                 |
| 1      | 2 | 3 | 5 | 9 | 3    | [1, 2, 3, 5, 9] | 3 > 5 = false |        |                 |
| 1      | 2 | 3 | 5 | 9 | 3    | [1, 2, 3, 5, 9] | 5 > 9 = false |        |                 |

- The function `bubbleScan` has one input called `array`, where `array` is the list of values to be scanned in sequence.
- The `bubbleScan` swaps any adjacent items not in order and stops after one pass.
- The `bubbleScan` returns `true` if any values were swapped (the array was modified) during the scan; otherwise, it returns `false`.
- The function `bubbleScan` uses the previously defined `swapItems` function.

### Using `bubbleScan` sorting algorithm

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
function bubbleScan(array) {
 var swapped = false;
 var i=1; // start at the second item
 while (i < array.length) {
 // compare adjacent items
 if (array[i-1]>array[i]) {
 // swap if items are out of place
 swapItems(array, i-1, i);
 swapped = true;
 }
 i=i+1;
 }
 return swapped;
}
var anArray = [9,1,5,2,3];
console.log(bubbleScan(anArray));
console.log(anArray);
```

- Once the function is defined, it can be called with the statement `bubbleScan(array)`.

### WORKED EXAMPLE 31 Determining the final output using `bubbleScan`

Given the function definitions shown below, determine what is returned from each of the following function calls.

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
function bubbleScan(array) {
 var swapped = false;
 var i=1;
 while (i < array.length) {
 if (array[i-1]>array[i]) {
 swapItems(array, i-1, i);
 swapped = true;
 }
 i=i+1;
 }
```

```

 }
 return swapped;
}

```

- a. `bubbleScan([2,1])`
- b. `bubbleScan([1,2,3,4,5,7,6])`
- c. `bubbleScan([])`
- d. `bubbleScan([1,2,3,4,5,7])`
- e. `bubbleScan([99])`

#### THINK

- a. `[2,1]` is not sorted, so `bubbleScan` returns `true`.
- b. `[1,2,3,4,5,7,6]` is not sorted (7 and 6 are out of place), so `bubbleScan` returns `true`.
- c. `[]` is empty but is still considered sorted, so `bubbleScan` returns `false`.
- d. `[1,2,3,4,5,7]` is sorted, so `bubbleScan` returns `false`.
- e. `[99]` is sorted, so `bubbleScan` returns `false`.

#### WRITE

- a. `true`
- b. `true`
- c. `false`
- d. `false`
- e. `false`

- The function `bubbleSort` has one input `array`, where `array` is the list of values to be sorted. The function `bubbleSort` uses the previously defined `bubbleScan` and `swapItems` functions.
- The `bubbleScan` is called many times until the `array` is no longer modified.

### Using `bubbleSort` sorting algorithm

```

function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}

function bubbleScan(array) {
 var swapped = false;
 var i=1; // start at the second item
 while (i < array.length) {
 // compare adjacent items
 if (array[i-1]>array[i]) {
 // swap if items are out of place
 swapItems(array, i-1, i);
 swapped = true;
 }
 i=i+1;
 }
 return swapped;
}

function bubbleSort(array) {
 // Scan the array until there are no more changes
 // bubbleScan(array) is now used as the decision to
 // loop.
 // bubbleScan(array) is called many times each time

```

```

// changing the array.
// Eventually the array becomes sorted and
// bubbleScan(array) returns false.
while(bubbleScan(array)) {
// Does nothing here
}
return array;
}
console.log(bubbleSort([9,1,5,2,3]));

```

- Once the function is defined, it can be called with the statement `bubbleSort(array)`. The original `array` is sorted in place and returned once the `bubbleScan` detects no more changes.

### WORKED EXAMPLE 32 Determining the output using `bubbleSort`

Given the function definitions shown below, determine what is returned from each of the following function calls.

```

function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
function bubbleScan(array) {
 var swapped = false;
 var i=1;
 while (i < array.length) {
 if (array[i-1]>array[i]) {
 swapItems(array, i-1, i);
 swapped = true;
 }
 i=i+1;
 }
 return swapped;
}
function bubbleSort(array) {
 while(bubbleScan(array)) {
 }
 return array;
}

```

- a. `bubbleSort([2,1])`
- c. `bubbleSort([])`
- e. `bubbleSort([99])`

- b. `bubbleSort([1,2,3,4,5,7,6])`
- d. `bubbleSort([1,2,3,4,5,7])`

#### THINK

- a. `[2,1]` when sorted is `[1,2]`.
- b. `[1,2,3,4,5,7,6]` when sorted is `[1,2,3,4,5,6,7]`.

#### WRITE

- a. `[1,2]`
- b. `[1,2,3,4,5,6,7]`



- c. `[]` is empty and already sorted.
- d. `[1, 2, 3, 4, 5, 7]` is already sorted.
- e. `[99]` is already sorted.
- c. `[]`
- d. `[1, 2, 3, 4, 5, 7]`
- e. `[99]`

### 13.6.7 Selection sort

- Another sorting algorithm is called **selection sort**. The algorithm swaps the minimum value with the item at the start of the array. This minimum value is locked in place and the process is repeated on the rest of the array. The second smallest value is now found and locked into place. The process is repeated on the remainder of the array until the entire array is sorted.

| Stage         | Values |   |   |   |   | Subset array                 | Minimum |
|---------------|--------|---|---|---|---|------------------------------|---------|
| Initial order | 9      | 1 | 5 | 2 | 3 | <code>[9, 1, 5, 2, 3]</code> | 1       |
| 1st pass      | 1      | 9 | 5 | 2 | 3 | <code>[9, 5, 2, 3]</code>    | 2       |
| 2nd pass      | 1      | 2 | 5 | 9 | 3 | <code>[3, 1, 2]</code>       | 3       |
| 3rd pass      | 1      | 2 | 3 | 9 | 5 | <code>[2, 1]</code>          | 5       |
| 4th pass      | 1      | 2 | 3 | 5 | 9 | <code>[1]</code>             | 9       |
| Sorted        | 1      | 2 | 3 | 5 | 9 | <code>[]</code>              |         |

### 13.6.8 Median

- The **median** is the middle item of a sorted array.
- If `array.length` is odd, then the median is the middle item.

$$\text{index} = (\text{array.length} - 1) / 2$$

For example, an odd length array `[12, 15, 23, 86, 101]` of length 5:

$$\text{index} = (\text{array.length} - 1) / 2$$

$$\text{index} = (5 - 1) / 2$$

$$\text{index} = 2$$

|       |    |    |           |    |     |
|-------|----|----|-----------|----|-----|
| Value | 12 | 15 | 23        | 86 | 101 |
| index | 0  | 1  | index = 2 | 3  | 4   |

- If `array.length` is even, then the median is the average of the two middle items.

$$\text{index1} = \text{array.length} / 2 - 1$$

$$\text{index2} = \text{array.length} / 2$$

For example, an even length array `[8, 15, 16, 18, 100, 201]` of length 6:

$$\text{index1} = \text{array.length} / 2 - 1$$

$$\text{index1} = 6 / 2 - 1$$

$$\text{index1} = 2$$

$$\text{index2} = \text{array.length} / 2$$

$$\text{index2} = 6 / 2$$

$$\text{index2} = 3$$

|       |   |    |            |            |     |     |
|-------|---|----|------------|------------|-----|-----|
| Value | 8 | 15 | 16         | 18         | 100 | 201 |
| index | 0 | 1  | index1 = 2 | index2 = 3 | 4   | 5   |

### WORKED EXAMPLE 33 Using `array.length` to determine the median

Given the function definition shown below, determine what is returned from each of the following function calls.

```
function isOdd(items) {
 return (items%2)===1;
}
function middleValue(array) {
 var middle = 0;
 if (isOdd(array.length)) {
 var index = (array.length-1)/2;
 middle = array[index];
 } else {
 var index1 = array.length/2 - 1;
 var index2 = array.length/2;
 middle = (array[index1]+array[index2])/2;
 }
 return middle;
}
```

a. `isOdd(3)`

b. `middleValue([3,6,9])`

#### THINK

- a.
  1. The input `items` equals 3.
  2. Write the output statement.
  3. Substitute 3 for `value`.
  4. Substitute 1 for  $3\%2$ , as 2 divides into 3 with 1 remainder.
  5. Substitute `true` for  $1===1$ .
- b.
  1. Write the function call.
  2. Write the first assignment in the function.
  3. Write the decision in the function.
    4. Substitute `[3,6,9]` for `array`.
    5. Substitute 3 for `[3,6,9].length`.
    6. Substitute `true` for `isOdd(3)`.
    7. `decision` is `true`, so run the statements inside first `{ }` block.
    8. The first assignment is in the first `{ }` block.
    9. Substitute `[3,6,9]` for `array`.
    10. Substitute 3 for `[3,6,9].length`.
    11. Substitute 1 for  $(3-1)/2$ .
    12. The second assignment is in the first `{ }` block.
    13. Substitute 1 for `index`.
    14. Substitute 6 for `array[1]`.
    15. Write the return statement.
    16. Substitute 6 for `middle`.

#### WRITE

```
a. items = 3
isOdd(3) = (items%2)===1
isOdd(3) = (3%2)===1
isOdd(3) = 1===1

isOdd(3) = true
middleValue([3,6,9])
middle = 0
decision =
isOdd(array.length)
= isOdd([3,6,9].length)
= isOdd(3)
= true

index = (array.length-1)/2
index = ([3,6,9].length-1)/2
index = (3-1)/2
index = 1
middle = array[index]

middle = array[1]
middle = 6
middleValue([3,6,9]) = middle
middleValue([3,6,9]) = 6
```

## Individual pathways

**PRACTISE**

1, 2, 3, 4, 5, 11

**CONSOLIDATE**

6, 7, 8, 9, 10, 12

**MASTER**

13, 14, 15, 16

## Fluency

1. **WE24** Each of the following programs swap the contents of `x1` and `x2`. True or False?

a. `var x1 = true;  
var x2 = false;  
var a = x2;  
x2 = x1;  
x1 = a;`

b. `var x1 = -12;  
var x2 = 32;  
var temporary = x1;  
x1 = x2;  
x2 = temporary;`

c. `var x1 = "Monday";  
var x2 = "Tuesday";  
var temp = x1;  
x2 = x1;  
x1 = temp;`

2. **WE27** Determine the output to the console of each the following programs.

a. `function alwaysDoingTheSameThing() {  
 return "This text again!!!!";  
}  
console.log(alwaysDoingTheSameThing());`

b. `function outputCubed(x) {  
 var squared = x*x;  
 console.log(x*squared);  
}  
outputCubed(-5);`

c. `function octagonPerimeter(sideLength) {  
 var sides = 8;  
 var perimeter = sides * sideLength;  
 console.log(perimeter);  
}  
octagonPerimeter(1.25);`

d. `function regularPolygonPerimeter(sides, sideLength)  
{  
 var perimeter = sides * sideLength;  
 return perimeter;  
}  
console.log(regularPolygonPerimeter(200,0.1));`

3. **WE28** Given the swapping function shown below, determine the output to the console of each of the following programs.

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
```

- a. 

```
var items = [true, false, "true", "false", 1, 2];
swapItems(items, 4, 5);
console.log(items[5]);
```
- b. 

```
var months = [
 "", "January", "February", "March", "April", "September", "June",
 "July", "August", "May", "October", "November",
 "December"
];
swapItems(months, 9, 5);
console.log(months[9]);
```

4. **WE30** Manually simulate running the following programs. For each program, determine the final value stored in the variable `changing`.

- a. 

```
var changing=0;
var scores = [2,6,73,2];
if (scores.length === 3) {
 changing = scores[0]+scores[1]+scores[2];
} else {
 changing = scores[0];
}
```
- b. 

```
var changing=0;
var values = [237.2,237.1];
if (values[0]>values[1]) {
 changing = values[0];
} else {
 changing = values[1];
}
```
- c. 

```
var changing=0;
var values = [1,6,2];
if (values[0]>values[1]) {
 changing = values[0];
} else {
 changing = values[1];
}
if (changing < values[2]) {
 changing = values[2];
}
```

5. **WE31** Given the function definitions shown below, determine what is returned from each of the following function calls.

```
function swapItems(array, i, j) {
 var temporary = array[j];
 array[j] = array[i];
 array[i] = temporary;
}
function scan(array) {
 var hasSwapped = false;
 var i=1;
 while (i < array.length) {
 if (array[i-1]>array[i]) {
 swapItems(array, i-1, i);
 hasSwapped = true;
 }
 i=i+1;
 }
 return hasSwapped;
}
```

- a. scan([10,200,30])
- b. scan([1.5,2.4,3.3,4.2,5.1])
- c. scan([-3.2])
- d. scan([10,20,30,40,50,70,60])
- e. scan([2.2,2.1])
- f. scan([])

6. **WE33** Given the function definition shown below, determine what is returned from each of the following function calls?

```
function isEven(items) {
 return (items%2)===0;
}
function middleValue(array) {
 var middle = 0;
 if (isEven(array.length)) {
 var i1 = array.length/2 - 1;
 var i2 = array.length/2;
 middle = (array[i1]+array[i2])/2;
 } else {
 var i = (array.length-1)/2;
 middle = array[i];
 }
 return middle;
}
```

- a. isEven(3)
- b. middleValue([3,6,9])
- c. isEven([2,3,5,7,11,13,17,19,23,29].length)
- d. middleValue([2,3,5,7,11,13,17,19,23,29])

### Understanding

7. **WE25** Complete the rest of the program to swap the contents of the two variables.

```
var valueA = 1.414213562373095;
var valueB = 3.141592653589793;
```

8. **WE26** Complete the rest of the program to swap the contents of the 3rd and 4th item in the array.  
`var array=["Mon","Tue","Thu","Wed","Fri","Sat","Sun"];`
9. **WE29** Manually simulate running the programs. For each program, determine the final value stored in the array.

a. `var squares=[1,4,9,16,100,25,36,49,64,81];  
var i=4;  
var j=9;  
if (squares[i]>squares[j]) {  
var temporary=squares[i];  
squares[i]=squares[j];  
squares[j]=temporary;  
}`

b. `var scores=[1.2,1.3,1.6,1.5];  
var a=2;  
var b=3;  
if (scores[a]>scores[b]) {  
var swap=scores[a];  
scores[a]=scores[b];  
scores[b]=swap;  
}`

10. **WE32** Given the function definitions shown below, determine what is returned from each of the following function calls.

```
function swapItems(array, i1, i2) {
var temp = array[i1];
array[i1] = array[i2];
array[i2] = temp;
}
```

```
function bubbleScan(array) {
var swapDone = false;
var i=1;
while (i < array.length) {
if (array[i-1]>array[i]) {
swapItems(array, i-1, i);
swapDone = true;
}
i=i+1;
}
return swapDone;
}
```

```
function sort(array) {
while(bubbleScan(array)) {
}
return array;
}
```

- a. `sort([10,200,30])`  
b. `sort([4.2,1.5,5.1,2.4,3.3])`  
c. `sort([-3.2])`  
d. `sort([10,50,70,20,30,40,60])`  
e. `sort([2.1,2.2])`  
f. `sort([])`

11. Given the function definitions shown below, what is returned from each of the following function calls?

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}

function selectionScan(startIndex, array) {
 var minIndex = startIndex;
 var i = startIndex;
 while (i < array.length) {
 if (array[i] < array[minIndex]) {
 minIndex = i;
 }
 i=i+1;
 }
 swapItems(array, startIndex, minIndex);
 return array;
}
```

- a. selectionScan(0, [1.5,2.4,3.3,4.2,0.1])
- b. selectionScan(0, [-3.3])
- c. selectionScan(0, [2.2,2.1])
- d. selectionScan(0, [-233.2, -22])
- e. selectionScan(1, [2.1,2.2])
- f. selectionScan(0, [10,200,3])
- g. selectionScan(1, [3,200,10])
- h. selectionScan(2, [3,10,200])

### Reasoning

12. The `selectionScan` function scans for the minimum value on the subset of the `array`. The subset starts from the `startIndex` until the end of the `array`.

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}

function selectionScan(startIndex, array) {
 var minIndex = startIndex;
 var i = startIndex;
 while (i < array.length) {
 if (array[i] < array[minIndex]) {
 minIndex = i;
 }
 i=i+1;
 }
 swapItems(array, startIndex, minIndex);
 return array;
}
```

Write a function `selectionSort` that has one input called `array`, where `array` is the list of values to be sorted using the selection sort algorithm in ascending order. The function `selectionSort` can use the previously defined `selectionScan` and `swapItems` functions.



13. Rewrite the following functions so the function `bubbleSort` now sorts the array in descending order.

```
function swapItems(array, index1, index2) {
 var swap = array[index1];
 array[index1] = array[index2];
 array[index2] = swap;
}
function bubbleScan(array) {
 var swapped = false;
 var i=1;
 while (i < array.length) {
 if (array[i-1]>array[i]) {
 swapItems(array, i-1, i);
 swapped = true;
 }
 i=i+1;
 }
 return swapped;
}
function bubbleSort(array) {
 while(bubbleScan(array)) {
 }
 return array;
}
```

### Problem solving

14. Write a function `findMedian` to find and return the median of an unsorted `array`.
15. Write two functions, `findLowerQuartile` and `findUpperQuartile`, to find the 1st quartile and 3rd quartile respectively of a sorted `array`.
16. Write a function `findIQR` to find the interquartile range (IQR) of an unsorted `array`.

## LESSON

### 13.7 The Monte Carlo method

#### LEARNING INTENTIONS

At the end of this lesson you should be able to:

- understand the Monte Carlo method to solve difficult problems
- generate random integers and random samples
- test a sample against the event in probability.

#### 13.7.1 Pseudo-random numbers

- A technique called the **Monte Carlo method** was developed to solve problems that are too difficult to analyse directly. The Monte Carlo method approximates the solutions using many random samples. The more random samples are generated, the higher is the accuracy of the final result.



- In order to generate random samples, a random number generator is required. Computers do not generate truly random numbers; instead, they use a mathematical algorithm to generate **pseudo-random** numbers that appear to be random. These algorithms require a **seed** value to get started. One method is to initialise the seed using the computer's internal clock. The following assignment stores the number of milliseconds since midnight on 1 January 1970.



```
var seed = new Date().getTime();
```

The seed is then updated and stored each time the pseudo-random generator is called.

### WORKED EXAMPLE 34 Using `seed` to generate random numbers

Manually simulate running the following program. Determine the final values stored in the `array`.

```
var seed = 1;
function random() {
 seed = (5*seed)%7;
 return seed;
}
var array = [random(),random(),random()];
```

#### THINK

- `random()` is called 3 times starting with the `seed` 1.
- Write the 1st `seed` update.
- Substitute 1st for `seed` only on the right-hand side.
- Substitute 5 for  $(5*1)\%7$ .
- Write the 2nd `seed` update.
- Substitute 5 for `seed` only on the right-hand side.
- Substitute 4 for  $(5*5)\%7$ .
- Write the 3rd `seed` update.
- Substitute 4 for `seed` only on the right-hand side.
- Substitute 6 for  $(5*4)\%7$ .
- Write the `array` assignment.
- `random()` returns a different `seed` value each time.

#### WRITE

```
seed = (5*seed)%7
seed = (5*1)%7
seed = 5
seed = (5*seed)%7
seed = (5*5)%7
seed = 4
seed = (5*4)%7
seed = (5*4)%7
seed = 6

array = [random(),random(),
random()]
= [5,4,6]
```


- JavaScript has a function called `Math.random` that generates pseudo-random numbers. Every time `Math.random()` is called, a different random number between 0 (inclusive) and 1 (exclusive) will be returned. The following statement stores a random number in the variable `sample`.

```
var sample = Math.random();
```

## 13.7.2 Scaled random numbers

- `Math.random()` returns a number between 0 (inclusive) and 1 (exclusive). Often a larger or smaller range of random numbers is required. The value returned from `Math.random()` can be multiplied by a scaling factor to adjust the range. The following statement stores a random number between 0 (inclusive) and `scale` (exclusive) in the variable `sample`.

```
var sample = scale*Math.random();
```



- The shaded region and blue lines in the diagram represent the values transitioning from the random number to the scaled random number. Numbers from 0 to 1 are scaled to numbers from 0 to `scale`. The transition is done in one step.

### WORKED EXAMPLE 35 Writing an expression to generate a random number

Write an expression to return a random number between 0 and:

a. 2

b. 2.5

c. -2.

#### THINK

- The `scale` factor is 2.
- The `scale` factor is 2.5.
- The `scale` factor is -2.


#### WRITE

- `2*Math.random()`
- `2.5*Math.random()`
- `-2*Math.random()`

## 13.7.3 Floor

- The expression `scale*Math.random()` returns a number between 0 and `scale`, but sometimes random integers are required. JavaScript has a function called `Math.floor` that rounds a `real` number down to the `integer` below. For example, this function will round 1.367 to 1.

```
var integer = Math.floor(real);
```



- The shaded region and blue lines in the diagram represent the values transitioning from the real number to the integer `Math.floor(real)`. The transition is done in one step. The diagram shows how values slightly larger than 1 are reduced to 1, values slightly larger than 2 are reduced to 2, and so on.

## WORKED EXAMPLE 36 Using `Math.floor` to round down

Evaluate the following JavaScript expressions manually.

- `Math.floor(2.3)`
- `Math.floor(501)`
- `Math.floor(-6)`
- `Math.floor(-1.4)`
- `Math.floor(0)`

### THINK

- Remove the digits after the decimal place to floor a positive number.
- No change is required to floor an integer.
- No change is required to floor an integer.
- Remove the digits after the decimal place and subtract 1 to floor a negative number.
- No change is required to floor an integer.


### WRITE

- `2 = 2`
- `501`
- `-6`
- `-1 - 1 = -2`
- `0`

## 13.7.4 Random integers

- By combining `Math.random` with `Math.floor`, we can generate a random integer between 0 and `scale-1` inclusive.


```
var integer = Math.floor(scale*Math.random());
```



- The shaded region and blue lines in the diagram above represent the values transitioning from `Math.random()` to `scale*Math.random()` to the random integer. The transition is done in two steps.

- Adding 1 to `Math.floor(scale*Math.random())` will generate a random integer between 1 and `scale` inclusive.

```
var integer = Math.floor(scale*Math.random()) + 1;
```



- The shaded region and blue lines in the diagram above represent the values transitioning in three steps from `Math.random()` to the random integer.

### WORKED EXAMPLE 37 Using `Math.floor` and `Math.random` to list possible outcomes

List all of the possible outcomes of the following expressions as ascending arrays of numbers.

- `Math.floor(6*Math.random())`
- `Math.floor(2*Math.random())`
- `Math.floor(4*Math.random())+1`
- `Math.floor(Math.random())`

#### THINK

- The `scale` factor is 6.
  - The expression randomly returns integers between 0 and 6-1.
- The `scale` factor is 2.
  - The expression randomly returns integers between 0 and 2-1.
- The `scale` factor is 4.
  - The expression adds 1, so it randomly returns integers between 1 and 4.
- The `scale` factor is hidden, but it is 1.
  - The expression randomly returns integers between 0 and 1-1.

#### WRITE

- `[0, 1, 2, 3, 4, 5]`
- `[0, 1]`
- `[1, 2, 3, 4]`
- `[0]`

## 13.7.5 Random index and random integer functions

- Generating a random integer between 0 and  $N-1$  can be wrapped into a function `index`.

Using `index` function to generate random numbers>

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
```

- Generating a random integer between 1 and  $N$  can be wrapped into a function `integer`.

Using `index` function to generate random integers between 1 and  $N$

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
```

### WORKED EXAMPLE 38 Using `index` function to list possible outcomes

Given the following function definitions, list all of the possible outcomes of the expressions below as ascending arrays of numbers.

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
```

- a. `integer(7)`      b. `integer(3)`      c. `index(4)`      d. `index(1)`

#### THINK

- The expression `integer(7)` randomly returns integers between 1 and 7.
- The expression `integer(3)` randomly returns integers between 1 and 3.
- The expression `index(4)` randomly returns integers between 0 and  $4-1$ .
- The expression `index(1)` randomly returns integers between 0 and  $1-1$ .

#### WRITE

- [1,2,3,4,5,6,7]
- [1,2,3]
- [0,1,2,3]
- [0]

## 13.7.6 Random selection function and random samples

- Randomly selecting an `item` in an `array` can be wrapped into a function `select`. First an index `i` is randomly selected between `0` and `array.length-1`. Then the `item` is selected from the `array` using the random index `i`.

### Using `select` to randomly select an item in an arrayUsing

```
function index(N) {
 return Math.floor(real);
 var real = Math.random()*N;
}
function select(array) {
 var i = index(array.length);
 var item = array[i];
 return item;
}
```

The `generate` function returns a `random` sample. The details of how the `random` sample is generated depend on the problem. If more than one random value is required, the `random` sample can become an array of values.

### Using `generate` to return a random sample

```
function generate() {
 ...
 return random;
}
```

## WORKED EXAMPLE 39 Generating a random sample

Given the following function definitions, define a `generate` function to return a random sample for each of the problems shown below.

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
function select(array) {
 var i = index(array.length);
 var item = array[i];
 return item;
}
```

- Select one of the strings "1st", "2nd" or "3rd".
- Show the results from throwing two fair six-sided dice.
- Show an index between 0 and 99.
- Show an hour integer between 1 and 12 and a minute integer between 0 and 59.

#### THINK

- Define the function `generate`.
  - Store the `list` of items.
  - select a random item.
  - Return the `random` item.
- Define the function `generate`.
  - Randomly generate the 1st die.
  - Randomly generate the 2nd die.
  - Create the `random` sample array.
  - Return the `random` item.
- Define the function `generate`.
  - Randomly generate an `index` between 0 and  $100-1$ .
  - Return the `random` item.
- Define the function `generate`.
  - Randomly generate the 1st integer between 1 and 12.
  - Randomly generate the 2nd integer between 0 and 59.
  - Create the `random` sample array.
  - Return the `random` item.

#### WRITE

```
a. function generate() {
 var list = ["1st","2nd","3rd"];
 var random = select(list);
 return random;
}

b. function generate() {
 var die1 = integer(6);
 var die2 = integer(6);
 var random = [die1,die2];
 return random;
}

c. function generate() {
 var random = index(100);

 return random;
}

d. function generate() {
 var hour = integer(12);

 var minute = index(60);

 var random = [hour,minute];
 return random;
}
```

### 13.7.7 Testing a sample

- For the Monte Carlo method, each random sample needs to be tested against the event. The `isEvent` function tests the sample and returns `true` if the `test` sample is the event; otherwise, it returns `false`. The details of how the `test` sample is checked against the event depend on the problem.

#### Using `isEvent` to test a sample

```
function isEvent(test) {
 ...
 return event;
}
```

## WORKED EXAMPLE 40 Defining the `isEvent` function to test the sample

For each of the following situations, define the `isEvent` function to return `true` if the `test` sample is the event, and otherwise return `false`.

- The three items in an array `test` sample add to 20.
- The sample `test` is even.

### THINK

1. Define the function `isEvent`.
  2. Add the three items in the array.
  3. Compare the sum with 20.
  4. Return the result of the test `event`.
1. Define the function `isEvent`.
  2. Check if `test` is even.
  3. Return the result of the test `event`.

### WRITE

```
a. function isEvent(test) {
 var t = test[0]+test[1]+test[2];
 var event = t === 20;
 return event;
}

b. function isEvent(test) {
 var event = test%2 === 0;
 return event;
}
```

## 13.7.8 Approximating probability

- For some events, the theoretical probability can be difficult to calculate. The Monte Carlo method can be used to simulate the event with many random samples. If a very large number of samples are used, the relative frequency of an event will approach the theoretical probability of the event.
- The follow program skeleton can be used to **estimate** the probability using a number of random **samples**.
- The `generate` function return a **random** sample. The details of how the **random** sample is generated depend on the problem.
- The `isEvent` function tests the sample and returns `true` if the `test` sample is the event; otherwise, it returns `false`. The details of how the `test` sample is checked against the event depend on the problem.

### Simulating the probability of events using the Monte Carlo method

```
function generate() {
 ...
 return random;
}
function isEvent(test) {
 ...
 return event;
}
function estimate(samples) {
 var events = 0;
 var i = 1;
 while(i <= samples) {
 if (isEvent(generate())) {
 events = events + 1;
 }
 }
}
```



```

 i = i + 1;
 }
 return events/samples;
}
var probability = estimate(totalSamples);

```

- Once the function is defined, it can be called with the statement `estimate(totalSamples)`.

### WORKED EXAMPLE 41 Estimating the probability of given outcomes

Create a program to estimate the probability that the results of three fair 6-sided dice sum to 10 using a million samples. Output the estimated probability to the console.

#### THINK

1. Selecting a random integer between 1 and 6 will be useful for simulating each die. Define the random generating `integer` and `index` functions.
2. Define the `generate` function to return an array of three integers, `d1`, `d2` and `d3`, each independently randomly selected between 1 and 6.
3. Define the `isEvent` function to test if the sum of three integers in an array `d` equals 10.
4. Define the function to `estimate` the probability using a number of random `samples`. The function `estimate` calls the already defined functions `generate` and `isEvent`.
5. Call the function `estimate` with 1000000 samples and store the result in `probability`.
6. Output `probability` to the console.

#### WRITE

```

function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
function generate() {
 var d1 = integer(6);
 var d2 = integer(6);
 var d3 = integer(6);
 return [d1, d2, d3];
}
function isEvent(d) {
 var sum = d[0]+d[1]+d[2];
 var event = sum === 10;
 return event;
}
function estimate(samples) {
 var events = 0;
 var i = 1;
 while(i <= samples) {
 if (isEvent(generate())) {
 events = events + 1;
 }
 i = i + 1;
 }
 return events/samples;
}

var probability=estimate
(1000000);

console.log(probability);

```

## Individual pathways

**PRACTISE**

1, 2, 4, 5, 7, 8

**CONSOLIDATE**

3, 6, 9, 11, 13, 15, 17

**MASTER**

10, 12, 14, 16, 18

## Fluency

1. **WE34** Manually simulate running the following program. Determine the final values stored in the `array`.

```
var seed = 1;
function random() {
 seed = (13*seed)%29;
 return seed;
}
var array = [random(),random(),random(),random(),random()];
```

2. **WE36** Evaluate the following JavaScript expressions manually.

- `Math.floor(-10.42342)`
- `Math.floor(-6)`
- `Math.floor(-1.422)`
- `Math.floor(1.45)`
- `Math.floor(267)`

3. Evaluate the following JavaScript expressions manually.

- `Math.floor(999)`
- `Math.floor(-612)`
- `Math.floor(0)`
- `Math.floor(3.45)`

4. **WE37** List all of the possible outcomes of the following expressions as ascending arrays of numbers.

- `Math.floor(8*Math.random())`
- `Math.floor(9*Math.random()+1)`
- `Math.floor(13*Math.random())`
- `Math.floor(4*Math.random()+1)`

5. **WE38** Given the following function definitions, list all of the possible outcomes of the expressions below as ascending arrays of numbers.

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
```

- |                          |                           |                            |
|--------------------------|---------------------------|----------------------------|
| a. <code>index(2)</code> | b. <code>index(11)</code> | c. <code>integer(6)</code> |
| d. <code>index(5)</code> | e. <code>index(1)</code>  |                            |




- a. A day integer between 1 and 365
  - b. One of the strings "January", "February", "March", "April", "May", "June", "July", "August", "September", "October", "November" and "December"
  - c. The results from throwing one fair twenty-sided die
  - d. One of the strings "Monday", "Tuesday", "Wednesday", "Thursday" or "Friday"
  - e. The results from throwing three fair six-sided dice
  - f. An index between 0 and 9
12. Given the following function definitions, define a `generate` function to return a `random` sample for each of the problems shown below.
- a. An integer between 100 and 200
  - b. Two independent real numbers between 0 and 1
  - c. One real number between 0 and 2
  - d. One real number between -1 and 1
  - e. Two independent real numbers between -1 and 1
13. **WE40** For each of the following situations, define the `isEvent` function to return `true` if the `test` sample is the event, and otherwise return `false`.
- a. The four items in an array `test` add to less than 10.
  - b. The two items in an array `test` have a difference of 5.
  - c. The sample `test` is odd.
  - d. The sum of the squares of two items in an array `test` is less than 1.
  - e. At least one of the two items in an array `test` is 6.
14. a. Define a `generate` function to return an array of two independent real numbers between -1 and 1.
- b. Define the `isEvent` function where the input `test` is an array of two numbers. The function `isEvent` returns `true` if `test[0]*test[0]+test[1]*test[1]` is less than 1; otherwise, it returns `false`.

### Reasoning

15. **WE41** Create a program to estimate the `probability` that four fair 6-sided dice sum to 12 using a thousand samples. Output the estimated probability to the console.
16. Create a program to estimate the probability that five fair 6-sided dice form a 'straight'. A 'straight' is where the five dice can be arranged to form either [1, 2, 3, 4, 5] or [2, 3, 4, 5, 6]. Use a million samples and output the estimated probability to the console.

### Problem solving

17. Create a Monte Carlo program to estimate the probability that any random point in a unit square is also inside the enclosed quarter circle that has a unit radius.
18. Create a Monte Carlo program to estimate the value of  $\pi$ .






# LESSON

## 13.8 Review


### 13.8.1 Success criteria

Tick a column to indicate that you have completed the lesson and how well you think you have understood it using the traffic light system.

(**Green:** I understand; **Yellow:** I can do it with help; **Red:** I do not understand)

| Lesson | Success criteria                                                         |  |  |  |
|--------|--------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| 13.2   | I can understand the types of values used in computer programming.       |                                                                                     |                                                                                     |                                                                                     |
|        | I can write numerical expressions in JavaScript and evaluate the output. |                                                                                     |                                                                                     |                                                                                     |
|        | I can assign variables and insert comments.                              |                                                                                     |                                                                                     |                                                                                     |
| 13.3   | I can classify values as an array, number, string or Boolean.            |                                                                                     |                                                                                     |                                                                                     |
|        | I can determine the JavaScript index of a value in an array.             |                                                                                     |                                                                                     |                                                                                     |
|        | I can evaluate the JavaScript expression from a given array.             |                                                                                     |                                                                                     |                                                                                     |
| 13.4   | I can evaluate JavaScript expressions from a given statement.            |                                                                                     |                                                                                     |                                                                                     |
|        | I can understand, count and evaluate loops.                              |                                                                                     |                                                                                     |                                                                                     |
| 13.5   | I can represent data as an array.                                        |                                                                                     |                                                                                     |                                                                                     |
|        | I can determine the output to the generated console.                     |                                                                                     |                                                                                     |                                                                                     |
|        | I can understand the concept of the mean of the values in an array.      |                                                                                     |                                                                                     |                                                                                     |
| 13.6   | I can swap the contents of variables and items in an array.              |                                                                                     |                                                                                     |                                                                                     |
|        | I can understand the structure used to make decisions to swap items.     |                                                                                     |                                                                                     |                                                                                     |
|        | I can understand sorting using the bubble sort or the selection sort.    |                                                                                     |                                                                                     |                                                                                     |
| 13.7   | I can understand the Monte Carlo method to solve difficult problems.     |                                                                                     |                                                                                     |                                                                                     |
|        | I can generate random integers and random samples.                       |                                                                                     |                                                                                     |                                                                                     |
|        | I can test a sample against the event in probability.                    |                                                                                     |                                                                                     |                                                                                     |

### Resources

 **Digital document** How to write your own program (doc-18769)

## Fluency

- Consider the following values and expressions.
  - Classify the value or expression result as a number, a string or Boolean.
  - Apply `typeof expression` to each value or expression. Write the result of `typeof expression`.
 

|                         |                         |                                |
|-------------------------|-------------------------|--------------------------------|
| a. <code>-0.125</code>  | b. <code>true</code>    | c. <code>"more strings"</code> |
| d. <code>"FALSE"</code> | e. <code>"False"</code> | f. <code>"True"</code>         |
- Consider the following values and expressions.
  - Classify the value or expression result as a number, a string or Boolean.
  - Apply `typeof expression` to each value or expression. Write the result of `typeof expression`.
 

|                       |                        |                       |
|-----------------------|------------------------|-----------------------|
| a. <code>100</code>   | b. <code>false</code>  | c. <code>124/4</code> |
| d. <code>5*5*5</code> | e. <code>949%25</code> |                       |
- Each of the following is a valid variable name. True or False?
 

|                                   |                                    |
|-----------------------------------|------------------------------------|
| a. <code>someRandomName</code>    | b. <code>41_forty_one</code>       |
| c. <code>and&amp;&amp;this</code> | d. <code>42_meaning_of_life</code> |
| e. <code>number_22</code>         | f. <code>remainder%this</code>     |
| g. <code>negative_-10</code>      | h. <code>function</code>           |
- Manually simulate running the following programs. For each program:
  - identify the final type of `variable`
  - determine the final value stored in `variable`.
    - `var variable = (1*2*3*4)/(1+1+1);  
variable = true;`
    - `var variable = 2;  
var x = false;  
variable = x;`
    - `var x = 1272.2/5;  
var t = 0;  
var variable = -(15%4)/(230-23332*7.2);  
variable = variable*x*t/23712.3;`
    - `var variable = "Am I a string, a Boolean or a number?";  
variable = true;  
variable = 12.23;`
    - `var variable = "I have changed a few times.";  
variable = 2.34;  
variable = true;  
variable = [true,false];  
variable = "String";`



12. Evaluate the following JavaScript expressions manually.
- |                                        |                                 |
|----------------------------------------|---------------------------------|
| a. <code>-34 &lt;= -34</code>          | b. <code>1.41 &gt; -1.12</code> |
| c. <code>true &amp;&amp; false</code>  | d. <code>!true</code>           |
| e. <code>false &amp;&amp; false</code> | f. <code>123 === 132</code>     |
13. Evaluate the following JavaScript expressions manually.
- |                                   |                                      |
|-----------------------------------|--------------------------------------|
| a. <code>143.3 === 143.30</code>  | b. <code>-49 &gt;= -49</code>        |
| c. <code>true    true</code>      | d. <code>40.08 &lt; 40.1</code>      |
| e. <code>71.99 &lt;= 72.01</code> | f. <code>true &amp;&amp; true</code> |
14. Evaluate the following JavaScript expressions manually.
- |                                       |                                   |
|---------------------------------------|-----------------------------------|
| a. <code>false    true</code>         | b. <code>false    false</code>    |
| c. <code>4.500 === 4.5000</code>      | d. <code>2.4221 !== 242.21</code> |
| e. <code>false &amp;&amp; true</code> | f. <code>true    false</code>     |
15. Manually simulate running the following programs. For each program, determine the final value stored in `t`.
- a. 

```
var t = 81;
 if (t === 9*9) {
 t = t/3;
 }
```
- b. 

```
var t = false || true;
 if (t) {
 t = 817;
 }
 if (t%2 === 1) {
 t = "false";
 }
```
16. Each of the following causes an infinite loop. True or False?
- a. 

```
var double=1;
 while(double===1025) {
 console.log(double);
 double=2*double;
 }
```
- b. 

```
var j = 0;
 while(false) {
 j = j + 1;
 console.log(i);
 }
```
- c. 

```
while(true) {
 console.log("Go for ever!!!!");
}
```
- d. 

```
var triple=1;
 while(triple!==243) {
 console.log(triple);
 triple=3*triple;
 }
```



17. For each the following programs:

- i. state how many times the loop is executed
- ii. determine the final value stored in the variable `w`.

a. 

```
var w = 200;
var p=1;
while (p<=9) {
 w=w-p;
 p=p+1;
}
```

b. 

```
var w=0;
var i=1;
while (i <= 13) {
 w=w+i;
 i=i+1;
}
```

c. 

```
var w=0;
var z=1;
while (z<=10) {
 console.log(5*z);
 z=z+1;
 w=w+2;
}
```

d. 

```
var w=2048;
var i=1;
while (i<=10) {
 w=w/2;
 i=i+1;
}
```

e. 

```
var y=1;
var w=1000;
while (y<=5) {
 w=w-y;
 y=y+1;
}
```

f. 

```
var w = "Some string";
while(false) {
 w = "Another string";
}
```

18. State how many times the loop code is executed in each of the following programs.

a. 

```
var number = 0;
var digits = [0,1,2,3,4,5,6,7,8,9];
var i=0;
while (i < digits.length) {
 var item = digits[i];
 number = number*10 + item;
 i=i+1;
}
console.log(number);
```

```
b. var weekend = ["Saturday", "Sunday"];
 var k=0;
 while (k < weekend.length) {
 var day = weekend[k];
 console.log(day);
 k=k+1;
 }
```

19. Each of the following programs swaps the contents of `var1` and `var2`. True or False?

```
a. var var1 = true;
 var var2 = false;
 var temporary = var2;
 var2 = var1;
 var1 = temporary;
b. var var1 = 3.14;
 var var2 = 3.1416;
 var temp = var1;
 var1 = var2;
 var1 = temp;
c. var var1 = "April";
 var var2 = "May";
 var t = var1;
 var2 = var1;
 var1 = t;
```

20. Determine the output to the console of each of the following programs.

```
a. function theSame() {
 return "Always this!!!";
 console.log(theSame());
}
b. function outputFloor(x) {
 var floor = Math.floor(x);
 console.log(floor);
}
outputFloor(2.34);
c. function circleArea(radius) {
 var pi = 3.14159;
 var area = pi * radius * radius;
 console.log(area);
}
circleArea(10);
d. function circumference(radius) {
 var pi = 3.14159;
 var perimeter = 2 * pi * radius;
 console.log(perimeter);
}
circumference(10);
```

21. Given the swapping function shown below, determine the output to the console of each of the following programs.

```
function swap(array, i, j) {
 var temporary = array[i];
 array[i] = array[j];
 array[j] = temporary;
}
```

- a. `var array = [5,1,2,3,4,0];  
swap(array, 0, 5);  
console.log(array[5]);`
- b. `var w =  
["Monday","Wednesday","Friday","Thursday","Tuesday"];  
var weekdays = w;  
swap(weekdays, 1, 2);  
console.log(weekdays[1]);  
swap(weekdays, 4, 1);`

22. Given the function definitions shown below, determine what is returned from the following function calls.

```
function swapItems(array, i, j) {
 var temporary = array[j];
 array[j] = array[i];
 array[i] = temporary;
}
```

```
function bubbleScan(array) {
 var swapped = false;
 var k=1;
 while (k < array.length) {
 if (array[k-1]>array[k]) {
 swapItems(array, k-1, k);
 swapped = true;
 }
 k=k+1;
 }
 return swapped;
}
```

- a. `bubbleScan([7,4,5,2,3,1])`  
b. `bubbleScan([12.2,12.1,12.3])`  
c. `bubbleScan([1,2,2,3])`  
d. `bubbleScan([10,20,30,40,50,60])`  
e. `bubbleScan([10,20,15])`  
f. `bubbleScan([])`

23. Evaluate the following JavaScript expressions manually.

- a. `Math.floor(44)`  
b. `Math.floor(-4.442)`  
c. `Math.floor(-6)`  
d. `Math.floor(13.415)`  
e. `Math.floor(0)`  
f. `Math.floor(-3.45)`  
g. `Math.floor(3.45)`

24. Given the function definitions shown below, list all the possible outcomes of each of the following expressions as ascending arrays of numbers.

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
```

- |                                    |                            |
|------------------------------------|----------------------------|
| a. <code>index(0)</code>           | b. <code>integer(0)</code> |
| c. <code>index(11)</code>          | d. <code>integer(5)</code> |
| e. <code>index(11)+20</code>       | f. <code>integer(9)</code> |
| g. <code>integer(4)*10 + 60</code> | h. <code>integer(1)</code> |

25. Given the function definitions shown below, list all the possible outcomes of each of the following expressions as ascending arrays of numbers.

```
function index(N) {
 var real = Math.random()*N;
 return Math.floor(real);
}
function integer(N) {
 return index(N)+1;
}
```

- a. `integer(4)*12`
- b. `Math.floor(7*Math.random())+1`
- c. `index(2)`
- d. `index(8)`
- e. `index(8)*100`
- f. `Math.floor(6*Math.random())`
- g. `index(10)`

26. Determine the output to the console generated by the program below.

```
var number = 0;
var digits = [0,1,2,3,4,5,6,7,8,9];
var i=0;
while (i < digits.length) {
 var item = digits[i];
 number = number*10 + item;
 i=i+1;
}
console.log(number);
```

27. Manually simulate running the following program. Determine the final value stored in the variable `final`.

```
var final=0;
var scores = [3,4];
if (scores.length === 3) {
 final = scores[0]+scores[1]+scores[2];
}
```

```
if (scores.length === 2) {
 final = scores[0]+scores[1];
}
if (scores.length === 1) {
 final = scores[0];
}
```

28. Manually simulate running the following program. Determine the final values stored in the `array`.

```
var seed = 1;
function random() {
 seed = (23*seed)%31;
 return seed;
}
var array = [random(),random(),random(),random()];
```

### Problem solving

29. Create a program to estimate the probability that the results of five fair 6-sided dice form exactly four of a kind. Use 100 000 samples and output the estimated probability to the console.



To test your understanding and knowledge of this topic, go to your learnON title at [www.jacplus.com.au](http://www.jacplus.com.au) and complete the **post-test**.

# Answers

## Topic 13 Algorithmic thinking

### 13.1 Pre-test

- C
- C
- 24
- C
- E
- A
- A
- B
- 11
- 23 times
- 5
- 30
- 14
- B
- E

### 13.2 Programs

- Number
    - "number"
  - Boolean
    - "boolean"
  - String
    - "string"
  - Number
    - "number"
  - Boolean
    - "boolean"
  - String
    - "string"
  - Number
    - "number"
- String
    - "string"
  - String
    - "string"
  - String
    - "string"
  - Number
    - "number"
  - String
    - "string"
  - String
    - "string"
- Number
    - "number"
  - String
    - "string"
  - Number
    - "number"
  - Number
    - "number"

- False
  - True
  - False
  - False
  - False
- True
  - False
  - True
  - False
  - False

- Number
    - 12
  - Number
    - 145
  - Boolean
    - false
  - Number
    - 1
  - Boolean
    - true
  - Boolean
    - true
- A simple string
  - Some data
  - 1
  - true
  - True
  - 12

- $(12-6)/(5-2)$
  - $16\%5$
  - $12+12-(-13-14-1)$
  - $12/(4*3)$
  - $12+(6-4-1)$
- $112*2.5$
  - $160/(4*4*8/2)$
  - $111\%14$
  - $24*2.5*2.5$

- ```
var computerLanguage = "JavaScript";
```
 - ```
var event = true;
```
  - ```
var finished = false;
```
 - ```
var megabyte = 1024*1024;
```
  - ```
var subject = "Mathematics";
```
 - ```
var minutes = 60*24*7;
```
  - ```
var million = 1000000;
```
- ```
var lotsOfFactors = 2*3*5*7*11*13; // lotsOfFactors=30030
```
  - ```
var e = 2+7/10+1/100+8/1000+3/10000; // e = 2.7183
```
 - ```
var product = 1*2*3*4; // product = 24
```
  - ```
var yourAge = 225/15; // yourAge = 15
```
- Number
 - 6
- Number
 - 135
 - Number
 - 6
 - String
 - "xyz"

- ```
var inLitres = 2; var inMillilitres = 1000*inLitres;
```
- ```
var crossSectionalArea = 100; var height = 12; var prismVolume = height*crossSectionalArea;
```

16. There are many possible programs to solve this problem; an example is shown. The value of `area` should be 39.27 (correct to 2 decimal places).

```
var radius = 10;
var angle = 45;
var pi = 3.141592653589793;
var area = pi*radius*radius *
angle/360;
```

17. There are many possible programs to solve this problem; an example is shown. The value of `perimeter` should be 77.79 (correct to 2 decimal places).

```
var radius = 19;
var angle = 120;
var pi = 3.141592653589793;
var arcLength = 2*pi*radius *
angle/360;
var perimeter = arcLength +
radius + radius;
```

13.3 Arrays

- | | | |
|------------|-----------|----------|
| a. Number | b. Array | c. Array |
| d. Boolean | e. Number | f. Array |
- | | | |
|----------|----------|-----------|
| a. Array | b. Array | c. String |
| d. Array | e. Array | f. String |
- | | | |
|----------|---------|---------|
| a. true | b. true | c. true |
| d. false | e. true | f. true |
- | | | |
|----------|----------|----------|
| a. false | b. false | c. false |
| d. true | e. false | |
- | | | | |
|------|------|------|------|
| a. 4 | b. 4 | c. 4 | d. 3 |
|------|------|------|------|
- | | | |
|-------|-------|------|
| a. 6 | b. 2 | c. 7 |
| d. 13 | e. 11 | |
- | | |
|--------|------------------|
| a. 4 | b. true |
| c. 2.3 | d. "some string" |
- | | |
|----------|--------------|
| a. 2 | b. false |
| c. true | d. "boolean" |
| e. false | |
- | | |
|--------------|----------|
| a. "boolean" | b. false |
| c. [true,2] | d. 4 |
- | | |
|----------|-------------|
| a. Array | b. [3,9,27] |
|----------|-------------|
- | | |
|----------|------------|
| a. Array | b. [3,1,4] |
|----------|------------|
- | | |
|-----------|--------|
| a. Number | b. 5.2 |
|-----------|--------|
- | | |
|-----------|--------|
| a. Number | b. 120 |
|-----------|--------|

14. There are many possible solutions to this problem; an example is shown. The value of `primorial` should be 30030.

```
var primes = [2,3,5,7,11,13];
var primorial = 1;
primorial = primorial*primes[0];
primorial = primorial*primes[1];
primorial = primorial*primes[2];
primorial = primorial*primes[3];
primorial = primorial*primes[4];
primorial = primorial*primes[5];
```

15. There are many possible solutions to this problem; an example is shown. The value of `total` should be 77.

```
var primes = [2,3,5,7,11,13,17,19];
var total = 0;
total = total + primes[0];
total = total + primes[1];
total = total + primes[2];
total = total + primes[3];
total = total + primes[4];
total = total + primes[5];
total = total + primes[6];
total = total + primes[7];
```

16. There are many solutions to this problem; an example is shown. The 6 arrays could be in any order and still be correct.

```
var combinations = [
[1,2,3],
[1,3,2],
[2,1,3],
[2,3,1],
[3,1,2],
[3,2,1]
];
```

17. There are many solutions to this problem; an example is shown. The 8 arrays could be in any order and still be correct.

```
var sampleSpace = [
[false,false,false],
[false,false,true],
[false,true,false],
[false,true,true],
[true,false,false],
[true,false,true],
[true,true,false],
[true,true,true]
];
```

13.4 Loops

- | | | |
|----------|---------|---------|
| a. true | b. true | c. true |
| d. false | e. true | f. true |
| g. true | | |
- | | | |
|----------|---------|----------|
| a. false | b. true | c. true |
| d. true | e. true | f. false |
| g. true | | |
- | | |
|----------|----------|
| a. false | b. false |
| c. true | d. false |
- | | |
|----------|---------|
| a. true | b. true |
| c. false | d. true |
- | | |
|----------|--------|
| a. 23409 | b. 100 |
|----------|--------|
- | | |
|----------|---------|
| a. False | b. True |
|----------|---------|
- | | | |
|-------|--------|------|
| a. 99 | b. 100 | c. 5 |
| d. 0 | e. 900 | |
- | | |
|----------|------------|
| a. i. 14 | ii. 105 |
| b. i. 9 | ii. 362880 |
| c. i. 5 | ii. 0 |

9. a. 10 b. 6 c. 1

10. a. False b. True

11. a. 1000 b. 101
 c. 100 d. 50

12. 2000

13. 385

14. There are many programs to solve this problem; an example is shown. The value of `factorial_10` should be 3628800.

```
var product = 1;
var a=1;
while (a<=10) {
  product = product*a;
  a=a+1;
}
var factorial_10 = product;
```

15. There are many programs to solve this problem; an example is shown. The value of `total` should be 500000500000.

```
var sum = 0;
var a=1;
while (a<=1000000) {
  sum = sum + a;
  a=a+1;
}
var total = sum;
```

16. There are many programs to solve this problem; an example is shown. The value of `product` should be 41992.

```
var value1 = 362;
var value2 = 232;
var product = 0;
var found = false;
while (!found) {
  product = product + 1;
  if ((product%value1 === 0) &&
      (product%value2 === 0)) {
    found = true;
  }
}
console.log(product);
```

17. There are many programs to solve this problem; an example is shown. The value of `product` should be 209960.

```
var array = [362,232,4,8,5];
var product = 0;
var found = false;
while (!found) {
  product = product + 1;
  found = true;
  var i = 0;
  while (i < array.length) {
    if (product%array[i] !== 0) {
      found = false;
    }
    i = i + 1;
  }
}
console.log(product);
```

13.5 Set structures

1. a. 12 b. 7 c. 10

2. 2.718281828459

3. 6469693230

4. 5

5. a. ["Monday","Tuesday","Wednesday",
 "Thursday","Friday",
 "Saturday","Sunday"]

b. ["January","February","March",
 "April","May","June","July",
 "August","September","October",
 "November","December"]

c. ["a","b","c","d","e","f","g","h",
 "i","j","k","l","m","n","o","p",
 "q","r","s","t","u","v","w","x",
 "y","z"]

d. [31,29,31,30,31,30,31,31,30,31,
 30,31]

6. true

7. 6469693230

8. a. 6

b. The purpose of the program is to scan the array and find the largest value. The largest value is then sent as output to the console.

9. a. 12

b. The purpose of the program is to scan the array and find the largest and smallest value. The difference of the largest and smallest value is computed to find the range of the values. The range is then sent as output to the console.

10. 4.1633

11. There are many possible programs to solve this problem; an example is shown. The value of `mean` should be 10.

```
var total = 0;
var digits = [10,17,6,3,26,23,  
1,23,4,5];
var k=0;
while (k < digits.length) {
  var item = digits[k];
  total = total + item;
  k=k+1;
}
var mean = total /
digits.length;
console.log(mean);
```

12. There are many possible programs to solve this problem; an example is shown. The value of `count` should be 168.

There are more efficient algorithms to count primes; this is a simple version that uses only the concepts shown in this topic.

```
var count = 0;
var i = 2;
while (i <= 1000) {
  var isPrime = true;
  var j = 2;
  while ((j < i) && isPrime) {
    var multiple = i%j === 0;
```



```

    isPrime = isPrime &&
    !multiple;
    j = j+1;
}
if (isPrime) {
    count = count + 1;
}
i = i+1;
}
console.log(count);

```

13. There are many possible programs to solve this problem; an example is shown. The value of `prime` should be 7919. There are more efficient algorithms to count primes; this is a simple version that uses only the concepts shown in this topic.

```

var count = 0;
var prime = 1;
while (count !== 1000) {
    prime = prime + 1;
    var isPrime = true;
    var j = 2;
    while ((j < prime) && isPrime) {
        var multiple = prime%j === 0;
        isPrime = isPrime &&
        !multiple;
        j = j+1;
    }
    if (isPrime) {
        count = count + 1;
    }
}
console.log(prime);

```

14. The program should output the following list in any order.

```

a=3 b=4 c=5
a=5 b=12 c=13
a=6 b=8 c=10
a=7 b=24 c=25
a=8 b=15 c=17
a=9 b=12 c=15
a=10 b=24 c=26
a=12 b=16 c=20
a=15 b=20 c=25
a=16 b=30 c=34
a=18 b=24 c=30
a=20 b=21 c=29
a=21 b=28 c=35

```

There are many possible programs to solve this problem; an example is shown. There are more efficient algorithms to find solutions to the equation; this is a simple version that uses only the concepts shown in this topic.

```

var a = 1;
while (a <= 30) {
    var b = a;
    while (b <= 30) {
        var c = b;
        while (c <= a+b) {
            if (a*a+b*b===c*c) {
                console.log("a="+a+" b="+b+"
                c="+c);
            }
            c = c+1;
        }
        b = b+1;
    }
    a = a+1;
}

```

```

}
b = b+1;
}
a = a+1;
}

```

13.6 Sorting algorithms

1. a. True b. True c. False

2. a. This text again!!!!
 b. -125
 c. 10
 d. 20

3. a. 1 b. September

4. a. 2 b. 237.2 c. 6

5. a. true b. false c. false
 d. true e. true f. false

6. a. false b. 6
 c. true d. 12

7. var valueA = 1.414213562373095;
 var valueB = 3.141592653589793;
 var swap = valueA;
 valueA = valueB;
 valueB = swap;

8. var array = ["Mon","Tue","Thu",
 "Wed","Fri","Sat","Sun"];
 var swap = array[2];
 array[2] = array[3];
 array[3] = swap;

9. a. [1, 4, 9, 16, 81, 25, 36, 49,
 64, 100]

- b. [1.2, 1.3, 1.5, 1.6]

10. a. [10,30,200]

- b. [1.5,2.4,3.3,4.2,5.1]

- c. [-3.2]

- d. [10,20,30,40,50,60,70]

- e. [2.1,2.2]

- f. []

11. a. [0.1,2.4,3.3,4.2,1.5]

- b. [-3.3]

- c. [2.1,2.2]

- d. [-233.2, -22]

- e. [2.1,2.2]

- f. [3,200,10]

- g. [3,10,200]

- h. [3,10,200]

12. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions.

```

function selectionSort(array) {
    var j = 0;
    while (j < array.length) {
        selectionScan(j, array);
        j = j+1;
    }
}

```

13. There are many possible programs to solve this problem; an example is shown. This is a simple change from the original version that only swaps the decision.

```

array[i-1]>array[i] to
array[i-1]< array[i].
function swapItems(array, index1,
index2) {
    var swap = array[index1];
    array[index1] = array[index2];
    array[index2] = swap;
}
function bubbleScan(array) {
    var swapped = false;
    var i=1;
    while (i < array.length) {
        if (array[i-1]< array[i]) {
            swapItems(array, i-1, i);
            swapped = true;
        }
        i=i+1;
    }
    return swapped;
}
function bubbleSort(array) {
    while(bubbleScan(array)) {
    }
    return array;
}

```

14. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions.

```

function swapItems(array, index1,
index2) {
    var swap = array[index1];
    array[index1] = array[index2];
    array[index2] = swap;
}
function bubbleScan(array) {
    var swapped = false;
    var i=1;
    while (i < array.length) {
        if (array[i-1]>array[i]){
            swapItems(array, i-1, i);
            swapped = true;
        }
        i=i+1;
    }
    return swapped;
}
function bubbleSort(array) {
    while(bubbleScan(array)) {
    }
    return array;
}
function isOdd(items) {
    return (items%2)===1;
}
function middleValue(array) {
    var middle = 0;
    if (isOdd(array.length)) {
        var index = (array.length-1)/2;

```

```

        middle = array[index];
    } else {
        var index1 = array.length/2 -
1;
        var index2 = array.length/2;
        middle = (array[index1]+
array[index2])/2;
    }
    return middle;
}
function findMedian(array) {
    bubbleSort(array);
    median = middleValue(array);
    return median;
}

```

15. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions.

```

function isOdd(items) {
    return (items%2)===1;
}
function middle(array, min, max)
{
    var range = max - min;
    var middle = 0;
    if (isOdd(range)) {
        var index = min + (range-1)/2;
        middle = array[index];
    } else {
        var index1 = min + range/2 - 1;
        var index2 = min + range/2;
        middle = (array[index1]+array
[index2])/2;
    }
    return middle;
}
function findLowerQuartile(array) {
    var half = 0;
    if (isOdd(array.length)) {
        half = (array.length-1)/2;
    } else {
        half = array.length/2;
    }
    var Q1 = middle(array, 0,
half);
    return Q1;
}
function findUpperQuartile(array)
{
    var half = 0;
    if (isOdd(array.length)) {
        half = (array.length-1)/2;
    } else {
        half = array.length/2;
    }
    var Q3 = middle(array,
array.length-half, array.length);
    return Q3;
}

```

16. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions.

```
function swapItems(array, index1,
index2) {
  var swap = array[index1];
  array[index1] = array[index2];
  array[index2] = swap;
}
function bubbleScan(array) {
  var swapped = false;
  var i=1;
  while (i < array.length) {
    if (array[i-1]>array[i]) {
      swapItems(array, i-1, i);
      swapped = true;
    }
    i=i+1;
  }
  return swapped;
}
function bubbleSort(array) {
  while(bubbleScan(array)) {
  }
  return array;
}
function isOdd(items) {
  return (items%2)===1;
}
function middle(array, min, max)
{
  var range = max - min;
  var middle = 0;
  if (isOdd(range)) {
    var index = min + (range-1)/2;
    middle = array[index];
  } else {
    var index1 = min + range/2 -
    1;
    var index2 = min + range/2;
    middle = (array[index1]+array
    [index2])/2;
  }
  return middle;
}
function findIQR(array) {
  bubbleSort(array);
  var half = 0;
  if (isOdd(array.length)) {
    half = (array.length-1)/2;
  } else {
    half = array.length/2;
  }
  var Q1 = middle(array, 0, half);
  var Q3 = middle(array,
  array.length-half, array.length);
  var IQR = Q3 - Q1;
  return IQR;
}
```

13.7 The Monte Carlo method

- [13,24,22,25,6]
- a. -11 b. -6 c. -2
d. 1 e. 267
- a. 999 b. -612 c. 0 d. 3
- a. [0,1,2,3,4,5,6,7]
b. [1,2,3,4,5,6,7,8,9]
c. [0,1,2,3,4,5,6,7,8,9,10,11,12]
d. [1,2,3,4]
- a. [0,1]
b. [0,1,2,3,4,5,6,7,8,9,10]
c. [1,2,3,4,5,6]
d. [0,1,2,3,4]
e. [0]
- a. [1]
b. [1,2,3,4,5,6,7,8,9,10]
c. [1]
d. [0]
- a. [10,11,12,13,14,15,16,17,18,19,20]
b. [0,10,20,30,40,50,60,70,80,90,100,110]
c. [11,22,33,44,55,66,77,88,99]
d. [600,700,800,900,1000]
- a. 100*Math.random()
b. 0.001*Math.random()
c. -0.1252*Math.random()
d. 3.14*Math.random()
e. -99*Math.random()
f. 10*Math.random()
- a. -120 b. -16 c. 17 d. 10 e. 57
- a. 444 b. -63 c. 0 d. -17 e. 9
- a. function generate() {
 var random = integer(365);
 return random;
}
b. function generate() {
 var months = [
 "January", "February", "March",
 "April", "May", "June", "July",
 "August", "September", "October",
 "November", "December"];
 var random = select(months)
 return random;
}
c. function generate() {
 var random = integer(20);
 return random;
}
d. function generate() {
 var weekdays = [
 "Monday", "Tuesday", "Wednesday",
 "Thursday", "Friday"];
 var random = select(weekdays)
 return random;
}

```

e. function generate() {
  var d1 = integer(6);
  var d2 = integer(6);
  var d3 = integer(6);
  var random = [d1,d2,d3];
  return random;
}
f. function generate() {
  var random = index(10);
  return random;
}
12. a. function generate() {
  var random = 100+index(101);
  return random;
}
b. function generate() {
  var real1 = Math.random();
  var real2 = Math.random();
  var random = [real1,real2];
  return random;
}
c. function generate() {
  var random = 2*Math.random();
  return random;
}
d. function generate() {
  var random = 2*Math.random()-1;
  return random;
}
e. function generate() {
  var real1 = 2*Math.random()-1;
  var real2 = 2*Math.random()-1;
  var random = [real1, real2];
  return random;
}
13. a. function isEvent(test) {
  var sum = test[0]+test[1]+
  test[2]+test[3];
  var event = sum < 10;
  return event;
}
b. function isEvent(test) {
  var diff1 = test[0] -test[1];
  var diff2 = test[1] -test[0];
  var event = (diff1 === 5) ||
  (diff2 === 5);
  return event;
}
c. function isEvent(test) {
  var event = (test%2)===1;
  return event;
}
d. function isEvent(test) {
  var square1 = test[0]*test[0];
  var square2 = test[1]*test[1];
  var sum = square1 + square2;
  var event = sum < 1;
  return event;
}
e. function isEvent(test) {
  var event = (test[0]===6) ||
  (test[1]===6);
  return event;
}
14. a. function generate() {
  var real1 = 2*Math.random()-1;
  var real2 = 2*Math.random()-1;
  return [real1, real2];
}
b. function isEvent(test) {
  var sum = test[0]*test[0] +
  test[1]*test[1];
  var event = sum < 1;
  return event;
}
15. There are many possible programs to solve this problem; an
example is shown. This is a simple version that uses only
the concepts shown in this topic and previously defined
functions. The probability should be approximately
0.095.
function index(N) {
  var real = Math.random()*N;
  return Math.floor(real);
}
function integer(N) {
  return index(N)+1;
}
function generate() {
  var d1 = integer(6);
  var d2 = integer(6);
  var d3 = integer(6);
  var d4 = integer(6);
  return [d1,d2,d3,d4];
}
function isEvent(test) {
  var sum = test[0]+test
  [1]+test[2]+test[3];
  var event = sum === 12;
  return event;
}
function estimate(samples) {
  var events = 0;
  var i = 1;
  while(i <= samples) {
    if (isEvent(generate())) {
      events = events + 1;
    }
    i = i + 1;
  }
  return events/samples;
}
var probability =
estimate(1000);
console.log(probability);
16. There are many possible programs to solve this problem; an
example is shown. This is a simple version that uses only
the concepts shown in this topic and previously defined
functions. The probability should be approximately
0.031.

```

```

function index(N) {
  var real = Math.random()*N;
  return Math.floor(real);
}
function integer(N) {
  return index(N)+1;
}
function generate() {
  var d1 = integer(6);
  var d2 = integer(6);
  var d3 = integer(6);
  var d4 = integer(6);
  var d5 = integer(6);
  return [d1,d2,d3,d4,d5];
}
function swapItems(array, index1,
index2) {
  var swap = array[index1];
  array[index1] = array[index2];
  array[index2] = swap;
}
function bubbleScan(array) {
  var swapped = false;
  var i=1;
  while (i < array.length) {
    if (array[i-1]>array[i]) {
      swapItems(array, i-1, i);
      swapped = true;
    }
    i=i+1;
  }
  return swapped;
}
function bubbleSort(array) {
  while(bubbleScan(array)) {
  }
  return array;
}
function isEvent(test) {
  bubbleSort(test);
  var straight1 =
test[0]===1 &&
test[1]===2 &&
test[2]===3 &&
test[3]===4 &&
test[4]===5;
  var straight2 =
test[0]===2 &&
test[1]===3 &&
test[2]===4 &&
test[3]===5 &&
test[4]===6;
  var event = straight1 ||
straight2;
  return event;
}
function estimate(samples) {
  var events = 0;
  var i = 1;
  while(i <= samples) {
    if (isEvent(generate())) {
      events = events + 1;

```

```

    }
    i = i + 1;
  }
  return events/samples;
}
var probability =
estimate(1000000);
console.log(probability);

```

17. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions. The `probability` should be approximately $\frac{\pi}{4} \approx 0.785$.

```

function generate() {
  var real1 = Math.random();
  var real2 = Math.random();
  return [real1, real2];
}
function isEvent(test) {
  var sum = test[0]*test[0] +
test[1]*test[1];
  var event = sum < 1;
  return event;
}
function estimate(samples) {
  var events = 0;
  var i = 1;
  while(i <= samples) {
    if (isEvent(generate())) {
      events = events + 1;
    }
    i = i + 1;
  }
  return events/samples;
}
var probability =
estimate(1000000);
console.log(probability);

```

18. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions. The `probability` should be approximately $\frac{\pi}{4} \approx 0.785$. This can be multiplied by 4 to estimate $\pi \approx 3.14$.

```

function generate() {
  var real1 = Math.random();
  var real2 = Math.random();
  return [real1, real2];
}
function isEvent(test) {
  var sum = test[0]*test[0] +
test[1]*test[1];
  var event = sum < 1;
  return event;
}
function estimate(samples) {
  var events = 0;
  var i = 1;
  while(i <= samples) {

```

```

    if (isEvent(generate())) {
        events = events + 1;
    }
    i = i + 1;
}
return events/samples;
}
var probability =
estimate(1000000);
var estimatePi = probability * 4;
console.log(estimatePi);

```

10. a. 12 b. 0 c. 6
 d. 8 e. 2
11. a. 5 b. 4
 c. "Eight" d. 8
12. a. true b. true c. false
 d. false e. false f. false
13. a. true b. true c. true
 d. true e. true f. true
14. a. true b. false c. true
 d. true e. false f. true
15. a. 27 b. "false"

13.8 Review questions

1. a. i. Number
 ii. "number"
- b. i. Boolean
 ii. "boolean"
- c. i. String
 ii. "string"
- d. i. String
 ii. "string"
- e. i. String
 ii. "string"
- f. i. String
 ii. "string"
2. a. i. Number b. i. Boolean
 ii. "number" ii. "boolean"
- c. i. Number d. i. Number
 ii. "number" ii. "number"
- e. i. Number
 ii. "number"
3. a. True b. False c. False
 d. False e. True f. False
 g. False h. False
4. a. i. Boolean
 ii. true
- b. i. Boolean
 ii. false
- c. i. Number
 ii. 0
- d. i. Number
 ii. 12.23
- e. i. String
 ii. "String"
5. a. Testing b. Results
 c. 0 d. false
6. a. Array b. String c. Boolean
 d. Array e. Array f. String
7. a. Number b. Array c. Array
 d. Number e. Array f. Array
8. a. false b. 5 c. true
 d. 4 e. false f. true
 g. false
9. a. true b. 4 c. false
 d. 5 e. true f. true
10. a. 12 b. 0 c. 6
 d. 8 e. 2
11. a. 5 b. 4
 c. "Eight" d. 8
12. a. true b. true c. false
 d. false e. false f. false
13. a. true b. true c. true
 d. true e. true f. true
14. a. true b. false c. true
 d. true e. false f. true
15. a. 27 b. "false"
16. a. False b. False
 c. True d. False
17. a. i. 9
 ii. 155
- b. i. 13
 ii. 91
- c. i. 10
 ii. 20
- d. i. 10
 ii. 2
- e. i. 5
 ii. 985
- f. i. 0
 ii. "Some string"
18. a. 10 b. 2
19. a. True b. True c. False
20. a. Always this!!!
 b. 2
 c. 314.159
 d. 62.8318
21. a. 5 b. Friday
22. a. true b. true c. false
 d. false e. true f. false
23. a. 44 b. -5 c. -6
 d. 13 e. 0 f. -4
 g. 3
24. a. [0]
 b. [1]
 c. [0,1,2,3,4,5,6,7,8,9,10]
 d. [1,2,3,4,5]
 e. [20,21,22,23,24,25,26,27,28,29,30]
 f. [1,2,3,4,5,6,7,8,9]
 g. [70,80,90,100]
 h. [1]
25. a. [12,24,36,48]
 b. [1,2,3,4,5,6,7]
 c. [0,1]
 d. [0,1,2,3,4,5,6,7]
 e. [0,100,200,300,400,500,600,700]
 f. [0,1,2,3,4,5]
 g. [0,1,2,3,4,5,6,7,8,9]
26. 123456789
27. 7
28. [23,2,15,4]

29. There are many possible programs to solve this problem; an example is shown. This is a simple version that uses only the concepts shown in this topic and previously defined functions. The `probability` should be approximately 0.019.

```
function index(N) {
  var real = Math.random()*N;
  return Math.floor(real);
}
function integer(N) {
  return index(N)+1;
}
function generate() {
  var d1 = integer(6);
  var d2 = integer(6);
  var d3 = integer(6);
  var d4 = integer(6);
  var d5 = integer(6);
  return [d1,d2,d3,d4,d5];
}
function isEvent(test) {
  var counts = [0,0,0,0,0,0];
  counts[test[0]] =
  counts[test[0]] + 1;
  counts[test[1]] =
  counts[test[1]] + 1;
  counts[test[2]] =
  counts[test[2]] + 1;
  counts[test[3]] =
  counts[test[3]] + 1;
  counts[test[4]] =
  counts[test[4]] + 1;
  var four1 = counts[1] === 4;
  var four2 = counts[2] === 4;
  var four3 = counts[3] === 4;
  var four4 = counts[4] === 4;
  var four5 = counts[5] === 4;
  var four6 = counts[6] === 4;
  var event = four1||four2||four3|
  |four4||four5||four6;
  return event;
}
function estimate(samples) {
  var events = 0;
  var i = 1;
  while(i <= samples) {
    if (isEvent(generate())) {
      events = events + 1;
    }
    i = i + 1;
  }
  return events/samples;
}
var probability =
estimate(100000);
console.log(probability);
```

GLOSSARY

- adjacent side** the side next to the reference angle in a right-angled triangle
- algorithm** a step-by-step set of tasks to solve a particular problem. A program is an implementation of an algorithm.
- altitude** *see* opposite side
- angle of depression** the angle measured down from a horizontal line (through the observation point) to the line of vision
- angle of elevation** the angle measured up from the horizontal line (through the observation point) to the line of vision
- area** the amount of surface enclosed by a shape; measured in square units, such as square metres (m^2) and square kilometres (km^2)
- array** (probability) a table that systematically displays all the outcomes of an experiment; (programming) a list of values. A JavaScript array takes the form $[x_0, x_1, x_2, \dots]$, where x_0, x_1, x_2 etc. are the different values of the array.
- asymptote** a line that a graph approaches but never touches
- axis of symmetry** the straight line that sits midway between two halves of a symmetrical graph, or between an object and its image. An object is reflected along an axis of symmetry (mirror).
- back-to-back stem-and-leaf plot** a method for comparing two data distributions by attaching two sets of 'leaves' to the same 'stem' in a stem-and-leaf plot; for example, to compare the pulse rate before and after exercise
- balanced** an equation whose left-hand side and right-hand side are equal
- bar chart (graph)** a graph drawn in a similar way to a column graph, with horizontal bars instead of vertical columns, with categories graphed on the vertical axis and the frequencies (numbers) on the horizontal axis
- base** *see* adjacent side
- bimodal distribution** a distribution of data that has two modes
- binomial** an expression containing two terms, for example $x + 3$ or $2y - z^2$
- Booleans** a JavaScript data type with two possible values: true or false. JavaScript Booleans are used to make logical decisions.
- bubble sort** a simple sorting algorithm. The algorithm scans an array, comparing adjacent values. If any adjacent values are out of place, they are swapped. This scanning process is repeated until no swaps are made during a complete scan.
- capacity** the maximum amount of fluid contained in an object, usually applied to the measurement of liquids and measured in units such as millilitres (mL), litres (L) and kilolitres (kL)
- Cartesian plane** an area that consists of two number lines at right angles to each other and includes all the space between the two number lines
- categorical data** data that can be organised into categories; for example, satisfaction rating or favourite colour. Categorical data can be either numerical or nominal.
- chance experiment** a process, such as rolling a die, that can be repeated many times
- circumference** the distance around the outside of a circle, given by the rule $2\pi r$ or πd , where r is the radius and d is the diameter of the circle
- coefficient** the number part of a term
- complement** (of a set) the complement of a set, A , written A' ; the set of elements that are in ξ but not in A
- compound interest** interest that is paid on the sum of the principal plus preceding interest over time
- congruent figures** identical figures with exactly the same shape and size
- console** a special region in a web browser for monitoring the running of JavaScript programs
- constant** a term with a fixed value
- constant of proportionality** the number multiplying x in the equation $y = kx$, which is equal to the gradient of the corresponding graph; also called the constant of variation
- constant rate** a rate of change that changes a quantity by the same amount for a given unit of time

continuous data data that has been collected by measuring; for example, the time it takes to get to school

cross-section the identical 'slice' produced when cuts are made across a prism parallel to its ends

cube a solid object with six faces, each of which is a square

cube root the inverse of cubing (raising to the power 3); a cube root is equivalent to raising a number to an index of $\frac{1}{3}$

cylinder a solid object with two identical flat circular ends and one curved side

dependent variable a variable with a value that changes in response to changes in the value of other variables

diameter a straight line passing through the centre of a circle from one side of the circumference to the other

difference of two squares an expression in which one perfect square is subtracted from another, for example $x^2 - 100$; this can be factorised by $(a + b)(a - b) = a^2 - b^2$

dilation (geometry) changing the size of an object (to larger or smaller) but not its shape; (parabolas) the coefficient of x^2 in the equation of a parabola making the graph wider or narrower than the graph of $y = x^2$

direct linear proportion describes a particular relationship between two variables or quantities; as one variable increases, so does the other variable

discrete data data that has been collected by counting amounts; for example, number of pens

Distributive Law the product of one number with the sum of two others equals the sum of the products of the first number with each of the others; for example, $4(6 + 2) = 4 \times 6 + 4 \times 2$. It is also applicable to algebra; for example, $3x(x + 4) = 3x^2 + 12x$.

dodecahedron a solid object with 12 faces, each of which is a regular polygon

dot plot a way of recording and organising data that displays each member of a sample as an individual dot and assigns each dot a value along an x -axis

edges a line between two vertices in a shape or solid

equally likely describes outcomes that have the same probability of occurring in an experiment

equation a mathematical statement that shows two equal expressions

Euler's formula a formula that shows the relationship between the number of edges, the number of faces and the number of vertices in any polyhedron: $F + V - 2 = E$

event set of favourable outcomes in each trial of a probability experiment

expanded an expression written without the brackets

expanding using the Distributive Law to remove the brackets from an expression

experimental probability probability determined by observing an experiment and gathering data, expressed by the formula: $\text{Pr}(\text{event}) = \frac{\text{number of ways the event can occur}}{\text{total number of equally likely outcomes}}$

explanatory variable *see* independent variable

expression group of terms separated by + or - signs, but which do not contain equals signs

faces plane shapes which are joined across edges to form a solid

factorised when an expression shows one number (or factor) multiplied by another

factorising writing a number or term as the product of a pair of its factors

favourable outcome any outcome of a trial that belongs to the event; for example, if A is the event of getting an even number when rolling a die, the favourable outcomes are 2, 4 and 6

First Index Law when terms with the same base are multiplied, the indices are added; therefore, $a^m \times a^n = a^{m+n}$

formula a mathematical rule, usually written using pronumerals

Fourth Index Law to remove brackets, multiply the indices inside the brackets by the index outside the brackets. Where no index is shown, assume that it is 1; therefore, $(a^m)^n = a^{mn}$.

frequency table a kind of table that is used to record and organise data

Fundamental Theorem of Algebra theorem that states that quadratic equations can have a maximum of two solutions

gradient (slope) (m) measure of how steep something is; the gradient of a straight line is given by $m = \frac{\text{rise}}{\text{run}}$ or

$$m = \frac{y_2 - y_1}{x_2 - x_1} \text{ and is constant anywhere along that line}$$

grouped data data presented using a frequency table to show a range instead of a value

hectare (ha) a unit of area equal to the space enclosed by a square with side lengths of 100 m (1 ha = 10 000 m²); often used to measure land area. There are 100 hectares in 1 square kilometre.

highest common factor (HCF) the largest of the set of factors common to two or more numbers; for example, the HCF of 16 and 24 is 8

histogram a column graph with no gaps between the columns and each column 'straddles' an x -axis score, starting and finishing halfway between scores. The x -axis scale is continuous and usually a half-interval is left before the first column and after the last column.

hypotenuse the longest side of a right-angled triangle, always opposite the right angle

icosahedron a solid object with 20 faces, each of which is an equilateral triangle

image an enlarged or reduced copy of a shape (the original shape is called the object)

independent variable a variable that does not change value in response to changes in the value of other variables

index (programming) an integer that points to a particular item in an array

index notation a short way of writing a number or variable when it is multiplied by itself repeatedly (for example, $6 \times 6 \times 6 \times 6 = 6^4$)

infinite loop when a program gets into a state where it will not finish without external intervention. The problem happens in the loop when it is impossible to change the terminating condition.

integer (Z) any number from the set of numbers that includes positive whole numbers, negative whole numbers and zero

interest a fee charged for the use of someone else's money, normally as a percentage of the amount borrowed

interquartile range (IQR) the difference between the third quartile, Q_3 and the first quartile, Q_1 , expressed as $IQR = Q_3 - Q_1$. It is the range of the middle half of the data.

intersection in probability, an intersection of two events (written as $A \cap B$) describes all outcomes in event A and in event B

inverse operation an operation that reverses the action of another; for example, $+$ is the inverse of $-$, and \times is the inverse of \div

irrational number (I) a number that cannot be written as a fraction. Examples of irrational numbers include surds, π and non-terminating, non-recurring decimals.

like terms terms containing exactly the same pronumeral part, including the power; for example, $3ab$ and $7ab$ are like terms, but $5a$ is not

linear equations equations in which the variable has an index (power) of 1

linear graph a relationship between two variables where a straight line is formed when the points are plotted on a graph

literal equation formulas are defined as literal equations because rearranging formulas does not require finding a value for the pronumeral (or pronumerals)

mean the measure of the centre (average) of a set of data, given by $\text{mean} = \frac{\text{sum of all scores}}{\text{number of scores}}$ or $\bar{x} = \frac{\sum x}{n}$. When data is presented in a frequency distribution table, $\bar{x} = \frac{\sum(f \times x)}{n}$.

median the measure of the centre (middle) of a set of data. The median is the middle score for an odd number of scores arranged in numerical order (ordered). For an even number of scores, the median is the average of the two middle scores when they are ordered. Its location is determined by the rule $\frac{n+1}{2}$, where n is the number of scores.

metric system the system of measurement based on the metre

midpoint the halfway point between the end points of a line segment

modal class a term used when analysing grouped data, given by the class interval with the highest frequency

mode the score that occurs most often. There may be no mode (for example, when all scores occur only once), one mode, or more than one mode (for example, when two or more scores occur equally frequently).

monomial an expression with one term

Monte Carlo method a method that approximates the solution to a problem using many random samples. The more random samples generated, the higher the accuracy of the final result.

multistep equation an equation that performs more than two operations on a pronumeral

natural numbers (N) positive whole numbers; 1, 2, 3, 4, ...

negatively skewed distribution describes a plot that has larger amounts of data as the values of the data increase

nominal data data in categories that have names, but cannot be ranked in a logical order; for example, brands of car or types of pet

Null Factor Law (NFL) law used to solve quadratic equations, which states that if $a \times b = 0$, then either $a = 0$ or $b = 0$ or both $a = 0$ and $b = 0$

numbers (programming) a JavaScript data type that represents numerical values

numerical data data that has been counted or measured; for example, number of siblings or heights of fences.

Numerical data can be either discrete and continuous.

object the original shape before it is copied and the copy is enlarged or reduced (the changed shape is called the image)

octahedron a solid object with 8 faces, each of which is an equilateral triangle

opposite side the side opposite the reference angle in a right-angled triangle

ordinal data data in categories that can be ranked in a logical order; for example, star ratings or race results

outcome the result obtained from a probability experiment

overtime the time in which a worker is working hours in addition to, or outside, their regular hours of employment

parabola the graph of a quadratic function, for example $y = x^2$ or $y = x^2 + 8$

parallel lines straight lines with the same gradient

perfect square a number or expression that is the result of the square of a whole number

perimeter the distance around the outside of a 2D shape

perpendicular lines straight lines that intersect at right angles to each other

pi (π) represents the ratio of the circumference of any circle to its diameter. The number represented by π is irrational, with an approximate value of $\frac{22}{7}$ and a decimal value of $\pi = 3.141\ 59 \dots$

pie chart (graph) a type of graph mostly used to represent categorical data. A circle is used to represent all the data, with each category being represented by a sector of the circle. The size of each sector is proportional to the size of that category compared to the total.

point of inflection where a curve changes from concave upward to concave downward (or vice versa)

polyhedron a three-dimensional solid with faces that are polygons, for example a tetrahedron

population the whole group from which a sample is drawn

positively skewed distribution describes a plot that has larger amounts of data as the values of the data decrease

principal the amount of money you start with when you put money into, or borrow from, a financial institution such as a bank or credit union

prism a solid shape with identical opposite ends joined by straight edges; a 3-dimensional figure with uniform cross-section

probability likelihood or chance of a particular event (result) occurring, given by the formula:

$$\text{Pr}(\text{event}) = \frac{\text{number of favourable outcomes}}{\text{number of possible outcomes}}$$
The probability of an event occurring ranges from 0 (impossible) to 1 (certain) inclusive.

pronumeral a letter or symbol that stands for a number

proportion equality of two or more ratios

pseudo-random appearing to be random. A pseudo-random number or sequence of numbers is generated by a mathematical algorithm. The algorithm is seeded with a different value each time so the same sequence is not generated again.

Pythagoras' theorem a theorem stating that in any right-angled triangle, the square on the hypotenuse is equal to the sum of the squares on the other two sides. This is often expressed as $c^2 = a^2 + b^2$.

Pythagorean triad a group of any three whole numbers that satisfy Pythagoras' theorem; for example, {5, 12, 13} and {7, 24, 25} are Pythagorean triads ($5^2 + 12^2 = 13^2$)

quadrant a quarter of a circle; one of the four regions of a Cartesian plane produced by the intersection of the x - and y -axes

quadratic equation an equation in which the term with the highest power is a squared term; for example, $x^2 + 2x - 7 = 0$

quadratic trinomial an expression of the form $ax^2 + bx + c$, where a , b and c are non-zero constants

radius (plural: radii) the distance from the centre of a circle to its circumference

range the difference between the maximum and minimum values in a data set

rate a ratio that compares the size of one quantity with that of another quantity; a method of measuring the change in one quantity in response to another

ratio comparison of two or more quantities of the same kind

rational number (Q) a number that can be expressed as a fraction with whole numbers both above and below the dividing sign

real number (R) any number from the set of all rational and irrational numbers

recurring decimal a decimal number with one or more digits repeated continuously; for example, 0.999 ...
A rational number can be expressed exactly by placing a dot or horizontal line over the repeating digits; for example, $8.343\ 434\ \dots = 8.\overline{34}$ or $8.\overline{34}$.

relative frequency the frequency of a particular score divided by the total sum of the frequencies; given by the rule: Relative frequency of a score = $\frac{\text{frequency of the score}}{\text{total sum of frequencies}}$

response variable *see* dependent variable

right-angled triangle a triangle with one 90° angle

salary a fixed annual amount of payment to an employee, usually paid in fortnightly or monthly instalments

sample part of a whole population

sample space the set of all possible outcomes from a probability experiment, written as ξ or S . A list of every possible outcome is written within a pair of curled brackets $\{ \}$.

scale factor the factor by which an object is enlarged or reduced compared to the image

scientific notation (standard form) the format used to express very large or very small numbers. To express a number in standard form, write it as a number between 1 and 10 multiplied by a power of 10; for example, 64 350 000 can be written as 6.435×10^7 in standard form.

Second Index Law when terms with the same base are divided, the indices are subtracted; therefore, $a^m \div a^n = a^{m-n}$

sector the shape created when a circle is cut by two radii

seed the starting number used in a pseudo-random number generator

selection sort a simple sorting algorithm. The selection sort algorithm swaps the minimum value with the item at the start of the array. This minimum value is locked in place and the process is repeated on the rest of the array until the array is sorted.

semicircle half of a circle

similar figures identical shapes with different sizes. Their corresponding angles are equal in size and their corresponding sides are in the same ratio, called a scale factor.

simple interest the interest accumulated when the interest payment in each period is a fixed fraction of the principal. The formula used to calculate simple interest is $I = \frac{Pin}{100}$, where I is the interest earned (in \$) when a principal of \$ P is invested at an interest rate of $i\%$ p.a. for a period of n years.

simple random sampling a type of probability sampling method in which each member of the population has an equal chance of selection

skewed distributions describes a non-symmetric data distribution, i.e. when more data values occur on one side of the peak than on the other

square root the inverse of squaring (raising to the power 2); a square root of a number is equivalent to raising that number to an index of $\frac{1}{2}$

standard form *see* scientific notation

stem-and-leaf plot format used to display organised data. Each piece of data in a stem-and-leaf plot is made up of two components, a 'stem' and a 'leaf'.

strings a JavaScript data type that represents text

subject (of a formula) the pronumeral or variable that is written by itself, usually on the left-hand side of the equation

surd any n th root of a number that results in an irrational number

symmetrical distribution a distribution that has a clear centre and an even spread on either side; also referred to as a bell-shaped or normal distribution

term part of an expression. Terms may contain one or more pronumerals, such as $6x$ or $3xy$, or they may consist of a number only.

terminate to end **terminating decimals** are decimals with a fixed number of decimal places, for example 0.6 and 2.54

tetrahedron a solid object with four faces, each of which is an equilateral triangle

Third Index Law any term (excluding 0) with an index of 0 is equal to 1; therefore, $a^0 = 1$

translate (quadratics) to move a parabola horizontally (left/right) or vertically (up/down)

tree diagram diagram used to list all possible outcomes of two or more events; the branches show the possible links between one outcome and the next outcome

trial one performance of an experiment to get a result; for example, each roll of a die is a trial

triangular prism a prism with a triangular base

trigonometry the branch of mathematics that deals with the relationship between the sides and the angles of a triangle

trinomial an algebraic expression containing three terms

turning point the point at which the graph of a quadratic function (parabola) changes direction (changes from going up to going down, or going down to going up)

turning point form the form of the equation of a graph that gives the coordinates of the turning point; for a parabola with the equation $y = (x - h)^2 + k$, the turning point is (h, k)

two-step equation an equation in which two operations have been performed on the pronumeral; for example, $2y + 4 = 12$, $6 - x = 8$, $\frac{x}{3} - 4 = 2$

two-way table a table listing all the possible outcomes of a probability experiment in a logical manner

undefined a numerical value that cannot be calculated

union in probability, the union of two events (written as $A \cup B$) is all possible outcomes in events A or in event B

variable (algebra) a letter or symbol in an equation or expression that may take many different values; (programming) a named container or memory location that holds a value

Venn diagram a series of circles, representing sets, within a rectangle, representing the universal set. They show the relationships between the sets.

vertex (geometry) a point or a corner of a shape where the straight edges meet; (quadratics) *see* turning point

volume amount of space a 3-dimensional object occupies; measured in cubic units, such as cubic centimetres (cm^3) and cubic metres (m^3)

wage a payment to an employee based on a fixed rate per hour

without replacement describes an experiment carried out with each event reducing the number of possible events; for example, taking an object permanently out of a bag before taking another object out of the bag. In this case, the number of the objects in the bag is reduced by one after every selection.

with replacement describes an experiment carried out without each event reducing the number of possible events; for example, taking an object out of a bag and then placing it back in the bag before taking another object out of the bag. In this case, the number of objects in the bag remains constant.

x-intercept the point(s) where a graph cuts the x -axis

y-intercept the point(s) where a graph cuts the y -axis. In the equation of a straight line, $y = mx + c$, the constant term, c , represents the y -intercept of that line.

INDEX

A

addition
 like terms 88–9
 of surds 49–51
adjacent side 274
algebra 72–144
 difference of two squares 104–5
 expanding brackets 95–104
 further expansions 109–12
 highest common binomial factor 118–22
 highest common factor 112–8
 in worded problems *see* worded problems
 perfect squares 105–7
 pronumeral in the denominator 153–4
 simplification 88–94
 substitution 75–7
 using pronumerals 74–83
algebraic expressions 74
 converting worded problems into 122–5
 simplification of 88–95
 worded problems into 83
algebraic terms
 division of 90–1
 multiplication of 89–90
angle of depression 299–301
angles of elevation 299–301
angle-angle-angle (AAA) condition of similarity 218
angle-side-angle (ASA) condition for congruence 205
angles, using inverse trigonometric ratios 292
applications
 of quadratic equations 760–2
arc length of sector 513
area
 converting units of 502
 definition 501
 of composite shapes 504–6
 of figures 503
 of quadrant 512
 of sector 512–3, 519
 of similar figures 229–46
 units of 229–30
 using formulas 502–4
array 597

ASA condition for congruence *see* angle-side-angle condition for congruence
asymptotes 376
Australian currency 479
average speed 426–8
axis of symmetry 372, 729

B

back-to-back stem-and-leaf plots 657–8
balanced, linear equation 150
bar charts 649–52
 displaying data in 649
 frequency table as 650
bimodal distributions 670
binomial
 definition 74, 98
 factors 98
 highest common factor 118
brackets, linear equations with 161–5

C

cancelling fractions 19–21
capacity 530–2
 of cylinder 534
Cartesian plane
 definition 325
 plotting linear graphs on 325–8
categorical data 630
chance experiment 565
circle 379–82
 circumference of 493
 equation 379
 graphing 381
circumference of circle 493–7
coefficients 74
collecting data 639–49
 dot plots 641–2
 frequency tables 639–41
common binomial factor 118
complement of an event 580–1
complementary events 581
composite number 16
composite shapes 504–6
 area of 504
 to solve problems 267–9
compound interest 469–78
 calculating 470

 definition 469
 formula for 473
concentric squares 58
congruence 192
 angle-side-angle condition for 205
 right angle-hypotenuse-side condition 205
 side-angle-side condition 205
 side-side-side condition 205
 symbol for 204
 testing triangles for 204–7
 triangles 206
congruent figures 204–12
constant 74
constant of proportionality 421–3
 as ratio 423
constant rates 429–30
 distance-time graph 430
continuous data 630
cosine ratio, unknown side length using 284
cross-section
 definition 519
 prism 532
cube, volume of 532–4
cube roots 41–2
cuboid, volume of 532–4
cylinders
 capacity of 534
 surface area of 525–7
 volume of 534–6

D

data
 as back-to-back stem-and-leaf plot 657
 categorical 630–1
 collection methods 633
 displaying *see* displaying data
 discrete 630
 graphical displays of 690
 grouped 668–70
 nominal 630
 numerical 630–1
 ordinal 630
 organising 639–49
 stem-and-leaf plots 642–5
 types of 630–2
decimal currency 479
decimal notation, converting scientific notation to 36–8

decimal numbers
 rounding 11–12
 in scientific notation 35
 in significant figures 12
 dependent variable 362, 422
 Desmos Graphing Calculator 356
 difference of two squares 104–5
 dilation of parabolas 739
 direct linear proportion 416–21
 checking for 417
 direct proportion
 relationship 422
 discrete data 630
 displaying data 649–64
 bar charts 649–52
 histograms 652–5
 misleading graphs 655–7
 pie charts 649–52
 distance-time graph 430
 distribution types 670–3
 and measures of centre 671
 Distributive Law 95–6
 definition 95
 division
 of algebraic terms 90–1
 of surds 46–8
 using indices 17–18
 dodecahedron 213
 dot plots 641–2

E
 edges 213
 equally likely outcomes 566
 equations
 circle 379
 linear *see* linear equations
 literal 175, 177
 multistep *see* multistep equations
 of straight line 339–48
 solving with brackets 161
 using inverse operations 152
 with pronumerals in
 denominator 154
 equivalent ratios method 154
 Euler's formula 213–4
 events, in probability 565
 expanding
 and simplifying 97–8
 binomial factors 98–100
 brackets 95–104
 Distributive Law 95–6
 multiple sets of brackets 109–11
 perfect squares 106
 using arrows 96
 using difference of two
 squares 104

experimental probability 571–3
 calculating 572
 expected number of results 573
 formula for 572
 relative frequency 571–2
 expressions 74–5
 coefficients of terms in 75
 using substitution 76
 extended form, recurring
 decimals in 6

F
 faces of polyhedra 213
 factorised 95
 factorising
 definition 112
 expressions 112–3
 by binomial common
 factor 118
 by grouping in pairs 118–20
 by highest common
 factor 113–5
 monic quadratics 723
 quadratic expressions 724
 terms 113
 favourable outcome 565
 Fifth Index Law 24–5
 financial mathematics
 compound interest 469–78
 penalty rates 455–61
 salaries and wages 451–5
 simple interest 461–9
 First Index Law
 and Third index laws 19
 expression using 17
 formulas
 Euler 213–4
 for compound interest 473
 for simple interest 462
 gradient 331
 rearranging 175–80
 substitution into 77–9
 to calculate area 502–4
 Fourth Index Law 24
 fractions
 as recurring decimals 6
 cancelling 19–21
 denominator in 5
 rational numbers as 5
 frequency tables 639–41
 as bar chart and pie chart 650
 with categories 640
 with values 640
 Fundamental Theorem of
 Algebra 716

G
 generating random numbers 634
 gradient 331–5
 formula 331
 line passing through two
 points 332–3
 of horizontal and vertical
 lines 333–5
 of straight line 333
 gradient-intercept method 350–2
 sketching graphs using 351
 graphing
 circle 381
 hyperbola 375
 parabola 372, 729–38
 parallel lines 356
 perpendicular lines 357
 with constant rates 430
 with technology 356
 with variable rates 431
 graphing calculator 750
 grouped data 668–70

H
 HCF *see* highest common factor
 hectare 502
 hexagonal prism 533
 highest common binomial factor
 118–22
 highest common factor (HCF) 112–8
 definition 113
 factorising expressions by 113–5
 of two terms 113
 histograms 652–5
 horizontal lines 352–3
 horizontal translations 374–5
 hyperbola 375–6
 hypotenuse 258

I
 icosahedron 213
 image 198
 independent variable 362, 422
 index laws
 cancelling fractions 19–21
 cube roots 41–2
 division using indices 17–8
 multiplication using indices 17
 negative indices 28–34
 raising powers 24–8
 review of 16–23
 scientific notation 34–41
 square roots 41
 surds 44–50
 zero index 18–9

- index notation 16
 - integers 4
 - interest
 - compound 469–77
 - simple 461–9
 - interquartile range (IQR) 680–3
 - intersection, of two events 587
 - inverse cosine values 292
 - inverse operations
 - definition 150
 - equations using 152
 - inverse tangent, word problems
 - using 294
 - inverse trigonometric ratios 291–3
 - angles using 292
 - IQR *see* interquartile range
 - irrational numbers 7–9
- L**
- left-skewed distributions 670
 - length
 - converting units of 492
 - of hypotenuse 259
 - of shorter side 260
 - length units 229
 - like terms 88
 - line segment, midpoint of 365–71
 - line(s)
 - gradient of 332
 - points on 328–9
 - rule of 340
 - y-intercepts of 335
 - linear equations 148–190
 - definition 150
 - multistep equations 151–3
 - rearranging formulas 175–80
 - solving problems with 169–75
 - solving 150–61
 - translating worded expressions
 - into 170
 - with brackets 161–5
 - with pronumerals on both sides 165–9
 - linear graphs 322–70
 - by plotting points 327
 - distance between two points 367–9
 - equation of straight line 339–48
 - features of 331–9
 - gradient 331–5
 - y-intercept 335–6
 - midpoint of line segment 365–71
 - plotting *see* plotting linear graphs
 - practical applications
 - of 360–5
 - sketching 348–55
 - technology and 356–60
 - literal equations 175–8
 - local minimum 729

M

 - Maths Tools 750
 - mean 664–8
 - measurement 488–555
 - area 501–12
 - area and perimeter of a sector 512–8
 - length 492
 - perimeter 493–7
 - surface area of a cylinder 525–9
 - surface area of prisms 519–25
 - volume of prisms and cylinders 530–42
 - measures of centre 664–80
 - grouped data 668–70
 - mean, median and mode 664–8
 - measures of spread 680–9
 - analysing data 683–4
 - range and interquartile range 680–3
 - median 664–8
 - metric system 492
 - midpoint of line segment 365–71
 - modal class 668
 - mode 664–8
 - modelling linear
 - relationships 361–2
 - monic quadratics
 - equations 723–9
 - monomial 74
 - multiple sets of brackets
 - expanding 109–11
 - simplifying 110
 - multiplication
 - of algebraic terms 89–90
 - of surds 46–8
 - using indices 17
 - multistep equations 151–3

N

 - natural numbers 4–5
 - definition 4
 - integers 4
 - rational numbers 4–5
 - negative indices 28–34
 - products of powers with 30
 - quotients of powers with 31
 - Seventh Index Law 28–32
 - negatively skewed distributions 670
 - nets 212
 - of octahedron 213
 - nominal data 630
 - non-linear relations
 - circle 379–82
 - cubic 376–7
 - exponential 377–9
 - hyperbola 375–6
 - parabola 372–5
 - non-right-angled triangles 268
 - Null Factor Law 715–9
 - word problems using 721
 - numbers
 - cube roots 41–2
 - irrational 7–9
 - rational 4–5
 - real *see* real numbers
 - rounding *see* rounding numbers
 - square roots 41
 - surds 44–56
 - numerical data 630

O

 - object 197
 - octahedron 213
 - Ohm's Law 78
 - opposite side 274
 - ordinal data 630
 - outcomes 565
 - expected number of 573
 - experimental probability of 572
 - number of 584–7
 - overtime 455–6
 - and penalty rates 455–6
 - calculating 455
 - definition 455
 - wages with 456

P

 - parabolas 372–5, 729–99
 - constructing 768
 - dilation of 739
 - features of 730
 - form $y = ax^2$ 739–41
 - form $y = ax^2 + c$ 744–5
 - form $y = (x + a)(x + b)$ 756–60
 - form $y = (x - h)^2$ 745–7
 - form $y = (x - h)^2 + k$ 744–52, 756
 - form $y = x^2 + bx + c$ 757
 - horizontal translations 374–5
 - in intercept form 756
 - reflection 375
 - sketching *see* sketching parabolas
 - vertical translations 373–4
 - parallel lines
 - definition 356
 - graphing 356
 - pay slips 456–8

- penalty rates 455–61
 - definition 455
 - overtime and 455–6
 - time sheets and pay slips 456–8
 - perfect squares 105–7
 - perimeter 493–7
 - circumference 493–7
 - of sector 512–5, 519
 - perpendicular lines 357
 - pi (π)
 - definition 8
 - value 2
 - with circles 493, 503
 - pie charts 649–52
 - displaying data in 649
 - frequency table as 650
 - plotting linear graphs 325–31
 - by plotting two points 327
 - from a rule 325–8
 - on a Cartesian plane 325–8
 - points on line 328–9
 - polyhedron 212–3
 - positive indices
 - converting scientific notation with 36
 - products of powers with 30
 - quotients of powers with 31
 - positively skewed distributions 670
 - prime factors 16
 - principal 461–2
 - prisms 519–22
 - and cylinders 530–43
 - base 532
 - cross-section 532
 - definition 519
 - rectangular 520
 - surface area of 519–25
 - triangular 519, 521
 - volume of 532–4
 - probability 558–623
 - compound events 597–609
 - definition 564
 - expected number of results 573–6
 - experimental 571–80
 - key terms of 565–6
 - language of 564–5
 - scale 564
 - theoretical 564–70
 - two-way tables 580–97
 - union of two events 589
 - Venn diagrams 580–97
 - problem solving, with linear equations 169–75
 - pronumerals 74–83
 - in congruent triangles 206
 - definition 74
 - in denominator, algebraic fractions with 153–4
 - proportion 414–25
 - direct linear 416–21
 - proportionality 421–6
 - constant of 421–3
 - ratio 423–4
 - Pythagoras 254
 - Pythagoras' theorem 258–67
 - angle of depression 299–301
 - angles of elevation 299–301
 - applications of 260–1, 267–74
 - distance between two points 367–8
 - definition 258
 - in 3D 269–71
 - introducing 258–9
 - problems using 261
 - Pythagorean triads 261–4
 - to non-right-angled triangles 268
 - trigonometric ratios 274–8, 282
 - unknown angles 291–9
 - unknown side lengths 259–60, 282–91
 - Pythagorean triads 261–263
- Q**
- quadrant 512
 - quadratic equations 708–28
 - applications of 760–2
 - definition 710
 - factorising 723–9
 - general form 711
 - monic quadratic equations 723–9
 - Null Factor Law 715–7
 - of form $ax^2 + bx = 0$ 720–1
 - of form $ax^2 + c = 0$ 711–4, 719–20
 - of form $x^2 + bx + c = 0$ 725–6
 - solutions to 711–5
 - to word problems 761
 - with two terms 719–23
 - quadratic graphs *see* sketching
 - parabolas
 - quadratic trinomials 723–4
 - quartile 680
- R**
- radius 493
 - raising powers 24–8
 - more index laws 24–6
 - range 680–3
 - rates 414, 426–35
 - average speed 426–8
 - calculating 427
 - constant 429–30
 - introduction to 426–9
 - penalty 455–61
 - reports 426
 - variable 431
 - rational numbers 4–5
 - definition 4
 - writing as fractions 5
 - ratios 195–7, 423–4
 - constant of proportionality as 423
 - simplifying 196
 - trigonometric 274–304
 - real numbers 4–11
 - irrational numbers *see* irrational numbers
 - natural numbers *see* natural numbers
 - recurring decimals 5–7
 - real-world problems 122
 - rearranging formulas 175–80
 - rectangular prism, surface
 - area of 520
 - recurring decimals 5–7
 - definition 5
 - fractions as 6
 - in extended form 6
 - writing 6
 - reflection 375
 - relative frequency 571–2
 - RHS condition for congruence *see*
 - right angle-hypotenuse-side condition for congruence
 - right angle-hypotenuse-side (RHS) condition
 - for congruence 205
 - for similarity 218
 - right-angled triangle 254–319
 - right-skewed distributions 670
 - rounding numbers 11–5
 - decimals 11–2
 - to significant figures 12–3
- S**
- salaries 451–5
 - definition 451
 - employees 451–3
 - sample space 565
 - sampling 632–5
 - generating random numbers 634
 - SAS condition *see* side-angle-side condition
 - scale 197–200
 - scale factors 197–231
 - scientific notation 34–41
 - converting 36–8
 - decimal numbers in 35

- scientific notation (*Cont.*)
 definition 34
 with negative indices 36
 with positive indices 36
- Second Index Law 17–18
 simplifying using 18
 with non-cancelling coefficients 18
- sectors 512
 arc length of 513
 area of 512–3
 definition 512
 perimeter of 513–4
- semicircle 512
- Seventh Index Law 28–32
- side-angle-side (SAS) condition
 for congruence 205
 for similarity 218–9
- side-side-side (SSS) condition
 for congruence 205
 for similarity 218
- significant figures 11–6
 in decimal number 12
 rounding to 12–3
- similar figures 217–29
 area of 229–46
 definition 217
 surface area of 231–2
 volume of 229–35
- similarity 192
 angle-angle-angle condition 218
 side-angle-side condition 218–9
 side-side-side condition 218
 testing triangles for 218
- simple interest 461–9
 calculating 463
 formula for 462
 interest 461–2
 on item 464
 principal 461–2
- Sixth Index Law 24–5
- sketching linear graphs 348–56
 gradient–intercept method 350–2
 horizontal and vertical lines 352–3
 using technology 356
 x - and y -intercept method 348–50
- sketching parabolas 738–44
 form $y = ax^2 + c$ 744–5
 form $y = (x + a)(x + b)$ 756–60
 form $y = (x - h)^2$ 745–7
 form $y = (x - h)^2 + k$ 744–52, 756
 form $y = x^2 + bx + c$ 757
 in intercept form 756
- skewed distributions 670
- speed 426
 average 426–8
- square roots 41
- SSS condition *see* side-side-side condition
- standard form 34
- statistics 626–706
 collecting data 630–9
 data *see* data
 displaying data *see* displaying data
 measures of centre 664–80
 measures of spread 680–9
 organising data 639–48
 sampling 630–9
 science of 626
- stem plot *see* stem-and-leaf plots
- stem-and-leaf plots 642–5
 back-to-back 657–8
 histogram 655
 measures of centre 672
- straight line
 equation of 339–48
 general equation of 339–40
 gradient of 333
- subject of a formula 175
- substitution
 expression using 76
 into formulas 77–9
 to check equations 150
 values into terms 75
- subtraction
 like terms 88–9
 of surds 49–51
- surds 44–55
 addition and subtraction of 49–51
 comparing 49
 definition 44
 entire 49
 expressions 50
 identifying 44–6
 multiplying and dividing 46–8
 simplifying 48
- surface area (SA)
 of cylinder 525–7
 of prisms 519–22
 of rectangular prism 520
 of triangular prism 521
- symmetrical distributions 670
- T**
- table of values 360–1
- tangent ratio, unknown side length using 283
- technology, graphing with 356
- terminate 5
- terms
 definition 74
 factorising 113
 in expression 75
 substituting values into 75
- testing triangles, for congruence 204–7
- tetrahedron 213
- Texas Instruments 356
- theoretical probability 564–7
- Third Index Law 18, 19
 First Index Law and 19
- time of an investment 463
- time sheets 456–8
- transcendental number 8
- tree diagram 597
- trial 565
- triangles
 congruence 204–7
 similar 218–21
- triangular prism 519
 surface area of 521
- trigonometric ratios 274–304
 inverse 291–3
 side lengths using 283–6
 using calculator 282
 values of 282–3
 worded problems using 285
- trigonometry 274–314
 angles of elevation and depression 299–304
 trigonometric ratios 274–81
 unknown side lengths 282–90
 unknown angles 290–9
- trinomial 74
- turning point 729
- turning point form 747
- two-step equation 151
- two-step experiments
 language of 604
 probabilities 601
 sample space of 597–600
 with replacement 601–2
 without replacement 601–5
- two-way tables 580–4, 597
 probability 587
 Venn diagrams to 586
- types of distributions 670–2
- U**
- undefined gradient 352
- union 587–8
- units of length 229, 492
- units of measurements, converting between 230

unknown angles 291–9
 calculating 292–3
 when two sides are known 293
unknown side lengths 259–60
 calculating 282–91
 in composite shapes 267
 using cosine ratio 284
 using tangent ratio 283

V

value of the investment
 with compound interest 471
 with simple interest 461
values
 frequency tables with 640
 in proportion 197
 of trigonometric ratios 282–3
 table of 360–1
variable rates 431–2
variables 74, 175, 360
Venn diagrams 580–96
 completing 588
 definition 581
 intersection of events 581–3
 to two-way table 586
 union of events 587–91
vertex 213, 372
 in polyhedron 213
 of a parabola 372

vertical lines 352–3
vertical translations of parabolas
 373–4
volume
 and capacity 530–2
 of composite solids 535
 of cube/cuboid 532–4
 of cylinder 534–6
 of prisms 530–4, 543
 of similar figures 229–35
 units of 230–1

W

wages 451–5
 definition 451
 employees 451–3
 with overtime 456
with replacement, experiment 601–2
without replacement,
 experiment 602–5
worded problems
 algebra in 83–8
 into algebraic expressions,
 converting 84
 key language used in 83
 quadratic equations to 761
 solving 84–131
 using inverse tangent 294
 using Null Factor Law 721

 using trigonometric ratios 285
writing
 decimal numbers in scientific
 notation 35
 entire surds 49
 fractions as recurring decimals 6
 ratios 195
 real number to decimal places 12
 recurring decimals 6
 scientific notation, number in 35
 with positive indices 30

X

x-intercept
 in parabolas 730–1
 method to sketch linear graphs
 348–50

Y

y-intercept
 in parabolas 730–1
 in the equation of a linear graph
 339–42
 method to sketch linear graphs
 348–50
 of linear graphs 335

Z

zero index 18–9