

Trial Examination 2008

VCE Biology Unit 3

Written Examination

Question and Answer Booklet

Reading time: 15 minutes Writing time: 1 hour 30 minutes

Student's Name:		
eacher's Name:		

Structure of Booklet

Section	Number of questions	Number of questions to be answered	Number of marks
А	25	25	25
В	8	8	50
			Total 75

Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers. Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape. No calculator is allowed in this examination.

Materials supplied

Question and answer booklet of 20 pages.

Answer sheet for multiple-choice questions.

Instructions

Write your name and teacher's name on this booklet and in the space provided on the answer sheet for multiple-choice questions. All written responses should be in English.

At the end of the examination

Place the answer sheet for multiple-choice questions inside the front cover of this booklet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Students are advised that this is a trial examination only and cannot in any way guarantee the content or the format of the 2008 VCE Biology Unit 3 Written

Neap Trial Exams are licensed to be photocopied and used only within the confines of the school purchasing them, for the purpose of examining that school's students only. They may not be placed on the school intranet or otherwise reproduced or distributed. The copyright of Neap Trial Exams remains with Neap. No Neap Trial Exam or any part thereof is to be issued or passed on by any person to any party inclusive of other schools, nonpractising teachers, coaching colleges, tutors, parents, students, publishing agencies or websites without the express written consent of Neap

SECTION A: MULTIPLE-CHOICE QUESTIONS

Instructions for Section A

Answer all questions in pencil on the answer sheet provided for multiple-choice questions.

Choose the response that is **correct** for the question.

A correct answer scores 1, an incorrect answer scores 0.

Marks will **not** be deducted for incorrect answers.

No marks will be given if more than one answer is completed for any question.

Question 1

Water serves as a solvent in biological stems.

With regard to this role, it is most important that water

- **A.** is non-polar.
- **B.** is polar.
- **C.** is organic.
- **D.** solidifies at room temperature.

Question 2

The chemical bonds that link amino acids together in a protein are called

- **A.** hydrogen bonds.
- **B.** peptide bonds.
- **C.** glycosidic bonds.
- **D.** phosphodiester bonds.

Question 3

Examples of lipids include

- **A.** phospholipids and antibodies.
- **B.** glucose and cellulose.
- C. DNA and water.
- **D.** triglycerides and steroids.

Question 4

Consider the following chemical reaction.

$$ATP \rightarrow ADP + P_i$$

The reaction could be described as

- A. endergonic.
- **B.** anabolic.
- C. catabolic.
- **D.** uphill.

Use the following information to answer Questions 5, 6 and 7.

Consider the following DNA template strand.

AGCTGGACCA

Question 5

This strand of DNA contains the nucleotide called adenosine.

Adenosine is composed of three components, namely

- **A.** a purine, a ribose and a phosphate.
- **B.** a pyrimidine, a deoxyribose and a phosphate.
- **C.** a purine, a nitrogenous base and a phosphate.
- **D.** a nitrogenous base, a phosphate and a deoxyribose.

Ouestion 6

The mRNA sequence complementary to the DNA sequence is

- A. UCGACCUGGU.
- **B.** TCGACCTGGT.
- C. AGCUCCAGGA.
- **D.** UGCTCCUCCU.

Ouestion 7

The process that uses a DNA template to create an mRNA sequence is called

- A. transcription.
- **B.** DNA replication.
- **C.** translation.
- **D.** protein synthesis.

Ouestion 8

Despite the limited number of different amino acids, many different types of proteins exist.

This is because

- **A.** the size of any given amino acid can vary.
- **B.** the chemical composition of a given amino acid can vary.
- C. the sequence and number of amino acids in a given protein is unique.
- **D.** the same amino acid can have many different properties.


Question 9

Consider a cell in a multicellular organism.

The proteome of the cell refers to

- **A.** all the genes and gene products in the cell.
- **B.** all the proteins within the cell at any moment in time.
- **C.** all the DNA in the cell and its associated functions.
- **D.** all the proteins within the cell throughout the cell's entire period of existence.

The graph below shows the effect of environmental factors on the rate of photosynthesis by barley seedlings.




The graph shows that

- **A.** carbon dioxide concentration is the factor limiting the rate of photosynthesis at point X.
- **B.** the rate of photosynthesis increases when the temperature is raised from 20°C to 30°C. This is because the enzymes controlling the Calvin cycle are nearer to their optimal temperature.
- **C.** the rate of photosynthesis levels off at point Y because chlorophyll molecules have been denatured by the high temperature
- **D.** water availability was the variable limiting the rate of photosynthesis at point Y.

Question 11

The graph below compares the absorption spectrum of chlorophyll *a* with the rate of photosynthesis of a plant upon exposure to a range of different light wavelengths.



From this graph, it can be concluded that chlorophyll *a* is not the only pigment involved in photosynthesis. Evidence that supports this conclusion includes the fact that

- **A.** the rate of photosynthesis remains high when the plant is exposed to light wavelengths between 450 nm and 650 nm. The percentage of light absorbed by chlorophyll *a* over these wavelengths is low.
- **B.** the rate of photosynthesis is low when the plant is exposed to light wavelengths such as 450 nm. Absorption of light by chlorophyll *a* is higher at these wavelengths.
- C. the rate of photosynthesis and the percentage of light absorbed by chlorophyll a are equal at 700 nm.
- **D.** between wavelengths 450 nm and 650 nm, the percentage of light absorbed by chlorophyll *a* is constant.

Use the following information to answer Questions 12 and 13.

This flowchart below shows the enzyme-controlled conversion of the amino acid threonine into isoleucine.

$$\begin{array}{c} \mathsf{COOH} \\ \mathsf{HCNH}_2 \\ \mathsf{HCOH} \\ \mathsf{CH}_3 \end{array} \xrightarrow{ \begin{array}{c} \mathsf{enzyme} \\ \mathsf{enzyme} \end{array}} \begin{array}{c} \mathsf{enzyme} \\ \mathsf{enzyme} \end{array} \xrightarrow{ \begin{array}{c} \mathsf{enzyme} \\ \mathsf{enzyme} \end{array}} \begin{array}{c} \mathsf{enzyme} \\ \mathsf{HCNH}_2 \\ \mathsf{HCOH} \\ \mathsf{CH}_3 \end{array} \xrightarrow{ \begin{array}{c} \mathsf{COOH} \\ \mathsf{HCNH}_2 \\ \mathsf{CH}_3 \end{array}} \xrightarrow{ \begin{array}{c} \mathsf{CNOH}_2 \\ \mathsf{CH}_2 \\ \mathsf{CH}_3 \end{array}}$$
 threonine

Question 12

The process labelled 'X' is best described as

- **A.** non-competitive inhibition.
- **B.** denaturation.
- **C.** feedback inhibition.
- **D.** catalysis.


Question 13

To convert threonine to isoleucine in a laboratory, the minimum materials you would need to place into a 'cell-like' solution would be

- **A.** enzymes 1-5.
- **B.** threonine and enzymes 1–5.
- C. intermediates A–D and enzymes 1–5.
- **D.** threonine and intermediates A–D.

Use the following information to answer Questions 14 and 15.

A scientist researching male infertility develops a drug that she thinks might increase sperm production in humans. The control of sperm production is shown in the diagram below.



Question 14

Some volunteer male subjects take the drug.

An effective drug may

- **A.** lower the levels of gonadotrophin-releasing hormone (GnRH) in the blood.
- **B.** increase testis size.
- **C.** increase testosterone production, leading to a reduction in sperm formation.
- **D.** increase the levels of luteinizing hormone (LH) in the blood.

Question 15

In the context of the diagram above, FSH is acting as a(n)

- A. neurotransmitter.
- **B.** pheromone.
- C. hormone.
- **D.** exocrine gland.

Question 16

In the mammalian circulatory system, excess fluid remaining in tissue spaces (interstitial fluid) is

- **A.** used to form urine.
- **B.** removed in the form of sweat.
- **C.** drained away by the lymphatic system.
- **D.** absorbed by fat cells.

A store placed a large basket of fruit containing oranges and unripe bananas by the cash register. There were too many bananas to fit in the basket, so they put the extra bananas in the back room. The next day, the bananas in the basket had ripened, but the ones in the back room had not.

The most likely explanation for this is that

- **A.** the heat in the store accelerated the ripening process.
- **B.** the oranges released ethylene into the air, triggering the bananas to ripen.
- C. the oranges released gibberellins into the air, triggering the bananas to ripen.
- **D.** auxin in the oranges diffused into the bananas where the two fruits touched each other, triggering the bananas to ripen.

Question 18

Phytochrome-deficient mutants of the plant Arabidopsis thaliana (mouse-ear cress) are unable to

- **A.** control their flowering by detecting changes in day length.
- **B.** carry out photosynthesis, due to an inability to absorb light.
- C. produce antibiotics that kill bacterial and fungal pathogens.
- **D.** seal wounds caused by grazing insect pests.

Question 19

Most T cells are tolerant to self antigens and become active only when receptors on their membranes are stimulated by non-self antigens. Non-self antigens are recognised by these receptors only when they are exposed on the surface of an antigen-presenting cell (APC).

An example of a cell which can function as an APC is

- **A.** a macrophage.
- **B.** a B cell.
- **C.** a dendritic cell.
- **D.** all of the above.

Ouestion 20

An experiment was conducted to investigate the role of the pineal gland in controlling reproductive behaviour in mammals. The pineal gland was surgically removed from one group of adult male Syrian hamsters (the *pinx* group). The operation was then faked (incisions made but glands not removed) on another group (the *sham* group).

At the beginning of the study all hamsters had large testes. Each of these two groups was then subdivided and placed under either a long day photoperiod (sixteen hours of light followed by eight hours of darkness) or short day photoperiod (one hour of light followed by twenty-three hours of darkness). After six weeks they weighed the testes of the hamsters and reported the following results.

	Long day, Pinx	Long day, Sham	Short day, Pinx	Short day, Sham
Average testis weight (mg)	2100	2088	2077	493

The best conclusion that can be drawn from these data is that

- **A.** the pineal gland secretes hormones that increase testis size in hamsters.
- **B.** testis size in hamsters decreases as daylength increases.
- C. short days are a stimulus for the pineal gland to cause a reduction in testicular size.
- **D.** removal of the pineal gland leads to reduced testis size in Syrian hamsters.

Consider the following statements regarding avian (bird) influenza.


- 1. The bacterium that causes bird flu is resistant to most antibiotics.
- 2. The pathogen that causes bird flu may mutate so that it can spread from human to human.
- 3. The pathogen that causes bird flu is carried by wild birds and can be spread around the world as birds migrate.
- 4. The pathogen that causes bird flu can be spread by eating cooked chicken.
- 5. Existing flu vaccines will not work against the type of pathogen that causes bird flu.

The correct statements are

- **A.** 1, 2, 3, 4 and 5.
- **B.** 1 and 5 only.
- **C.** 2, 3 and 5 only.
- **D.** 2 and 3 only.

Question 22

The diagram below shows an event in the human immune response.



The diagram shows

- **A.** antigens (R) from a pathogen (P) causing a B cell (Q) to secrete antibodies (S).
- **B.** antigens (S) on a pathogen (Q) causing a B cell (P) to secrete antibodies (R).
- **C.** plasma cells (Q) forming antibodies (R) in the endoplasmic reticulum (S), and secreting them in response to a pathogen (P).
- **D.** a B cell (P) proliferating into a clone of plasma cells (Q) that secrete antibodies (R) formed in the endoplasmic reticulum (S).

Question 23

Individuals with HIV (Human Immunodeficiency Virus) sometimes contract a pneumonia infection that is rare in the rest of the population.

This is because people with HIV

- **A.** are unable to fight off these pneumonia-causing organisms.
- **B.** are more often exposed to these pneumonia-causing organisms.
- **C.** release pheromones that attract the pneumonia-causing organisms.
- **D.** release substances that increase the strength of the pneumonia-causing organisms.

A **nude mouse** is a mouse that has had its thymus gland removed. The outward appearance of the mouse is a lack of body hair, which gives it the 'nude' nickname.

Nude mice are commonly used in experiments on tissue grafting because

- **A.** they have B cells that produce antibodies. The antibodies can then be used to study the process of tissue rejection.
- **B.** they are much cheaper than ordinary laboratory mice.
- **C.** they have a much less effective immune system due to a greatly reduced number of T cells. Grafted tissue is therefore not rejected.
- **D.** the lack of body hair makes them less vulnerable to infections that would lead to an immune response and tissue rejection.

Question 25

Salicylic acid is a chemical found in most plants in elevated levels in response to microbial pathogens (it is also found in aspirin). The recent discovery of salicylic acid-binding protein 2 (SABP2) showed how increased levels of salicylic acid can initiate what is being called a type of innate immune response.

When stimulated by raised levels of salicylic acid, SABP2 could play a role in restricting infection within a plant by

- **A.** inducing infected areas of the plant to undergo programmed cell death.
- **B.** promoting localised cell division in the infected area.
- **C.** stimulating less salicylic acid production.
- **D.** changing its tertiary structure to no longer be sensitive to salicylic acid.

SECTION B: SHORT-ANSWER QUESTIONS

Instructions for Section B

Answer this section in **pen**.

Answer all questions in the spaces provided.

Question 1

Biomacromolecules play a vital role within cells. DNA is a relatively simple molecule, but it provides a blueprint for proteins that are extremely complex in structure. The flowchart below illustrates how these biomacromolecules relate to each other.

 $DNA \rightarrow mRNA \rightarrow protein$

are to explain how proteins fold to form a specific
are to explain how proteins fold to form a specific
are to explain how proteins fold to form a specific
are to explain how proteins fold to form a specific
_

Membranes have a very important role as a selective barrier to movement of chemicals across them.

a. Complete the table below by filling in the missing information.


Type of movement	Membrane component involved	Movement of solutes	Example
active transport		low to high	absorption of glucose into the blood
	protein channels	high to low	the movement of sodium into a cell
diffusion		high to low	the movement of oxygen into a cell

i.	Name this process.
ii.	Identify the sequence of organelles the protein would pass through from synthesis to secretion
	1 + 1 = 2 mark
	ry membrane is in direct contact with water on either side of it, yet it still provides a structure that as an effective barrier.
	a labelled diagram to show the structure of a membrane, and explain how it can remain stable in a watery environment.

2 marks Total 6 marks

b.

One of the first enzymes to be purified and studied (in the 1930s) was urease. Urea molecules will slowly break down to carbon dioxide and ammonia in the absence of a catalyst, but the reaction proceeds much more quickly in the presence of urease. This is shown in the figure below.



a.	Define the term enzyme.

1	mark

In the gra	ph, why is x greater that	ın y?		1 mark

2 marks

An experiment was performed to test the effect of pH on the activity of urease. Enzyme activity was determined by measuring the amount of urea broken down to ${\rm CO_2}$ and ${\rm NH_3}$ per unit time. The table below shows the results of the investigation.

pН	Enzyme activity (arbitrary units)
4.6	0.06
5.0	0.12
5.6	0.32
6.0	0.50
6.4	0.70
6.8	0.88
7.2	0.84
7.6	0.62
8.0	0.40
8.4	0.14

i.	With reference to the table, what is the most likely optimum pH value for ure	ase:
ii.	Explain why a change in pH affects the activity of an enzyme.	
State	e a factor that must be kept constant during the investigation described above.	1 + 2 = 3

1 mark Total 7 marks

	ts with isolated mitochondria can lead to a better understanding of how they operate.
i.	Describe the function of mitochondria.
ii.	Describe a structural feature of a mitochondrion and show how this feature assists the function of the organelle.
	1 + 1 = 2 marks
This	of the important enzymes that operate within mitochondria is pyruvate dehydrogenase (PDH). enzyme is synthesised in cytoplasmic ribosomes and is then transported to the mitochondrial ace for delivery into the mitochondrion. While it can cross the outer mitochondrial membrane,
	I is excluded from other organelles.
PDF	
PDF	I is excluded from other organelles.
PDF Description	I is excluded from other organelles. cribe a feature of the outer mitochondrial membrane that allows PDH to cross.
PDF Description	I is excluded from other organelles. cribe a feature of the outer mitochondrial membrane that allows PDH to cross. 1 mark DH is prevented from entering the mitochondrion, the Krebs cycle will not function. A build-up of
PDF Desc	I is excluded from other organelles. cribe a feature of the outer mitochondrial membrane that allows PDH to cross. 1 mark DH is prevented from entering the mitochondrion, the Krebs cycle will not function. A build-up of c acid in the cytoplasm will result.

1 + 1 + 1 = 3 marks Total 6 marks

Many diseases arise from errant molecular interactions or faults in molecular mechanisms at the cellular level. Finding molecules that prevent these interactions or rectify these faults can, in theory, be used to develop drugs to treat such diseases.

a. What name is given to this process of drug development?

1 mark

Apples and tea leaves are both rich in molecules called flavonoids that could form the basis for developing new drugs. Between 1985 and 1990, a study was conducted on 805 elderly Dutch men aged between 65 and 84 years who were not previously diagnosed with coronary heart disease. Their dietary intake of flavonoids was varied by controlling their consumption of tea and apples. Death rates and mortality due to coronary disease were also recorded for each group. The results are shown in the following table.

Number of men	268	268	269
Flavonoid intake (mg/day)	<19	19–30	>30
Deaths from coronary heart disease	22	11	10

							1 mai
How c	ould the concl	usion from par	t b be linked to	drug develop	oment?		
							1 mar
them. I Cells in	Large quantitie	s of a molecul petunia (a bed	ounts of flavond le called chalcor lding plant) can	narigenin are	found in the	peel of yellow t	omatoes
	chal	conarigenin cl	nalcone isomera	se kaempfero	ol (a flavonoi	d)	
. .	he what could	he done to ens	able yellow toma	atoes to make	e flavonoids		

1 mark

		-	knowledge of how reasure		vork to explain h	now kaempferol
						2 mark Total 6 mark
Question 6						
ear. Extensive he number of	e research is b people who ha	eing cond	reventable cause of lucted to try to find moking after having ebo (an injection of	a solution to this p been given an 'an	problem. The gra	ph below shows
	_	_				
		18 16 14 12				
		16 14 12 10 8 6 4				
	smokers who	16 14 12 10 8 6	vaccinated	smokers wh		
•	smokers who quit (%)	16	vaccinated smokers need to be kept congenerating the above	smokers what plates a plates stant between the	cebo	sound scientific

1 mark


The	vaccine promotes an antibody response against nicotine (one of the addictive agents in cigarette smoke). antibodies then bind to the nicotine which is then unable to enter brain cells to continue ing addiction.
c.	Identify one of the cell types involved in this process, and describe its role in promoting an antibody response.
d.	Draw and label a diagram of the interaction between nicotine and the antibody.
	2 marks Total 6 marks
The Thy	human thyroid gland is situated in the neck and secretes the hormones thyroxine and calcitonin. oxine causes tissues to generate heat by increasing the rate of aerobic respiration in target cells. itonin helps to regulate blood calcium levels. Thyroxine is lipophilic and calcitonin is water-soluble.
a.	Describe how thyroxine and calcitonin will differ in the way they deliver their signals to target cells.
b.	Explain how thyroxine may cause responses in a target cell.
	2 marks

glandi.	d and can result in hypothyroidism. What is an 'antigen'?
1.	——————————————————————————————————————
ii.	What type of disease does this kind of hypothyroidism represent?
	1 + 1 = 2 ma
often can b	umatoid arthritis (which can display similar symptoms to hypothyroidism) is a condition which
can l	n extremely painful due to inflammation of the joints. A corticosteroid (e.g. cortisone) injection be administered to reduce this inflammation. Cortisone causes vasoconstriction and has mistamine properties.
can l antih	be administered to reduce this inflammation. Cortisone causes vasoconstriction and has
can l antih	be administered to reduce this inflammation. Cortisone causes vasoconstriction and has nistamine properties.
can l antih	be administered to reduce this inflammation. Cortisone causes vasoconstriction and has nistamine properties.
can l antih	be administered to reduce this inflammation. Cortisone causes vasoconstriction and has nistamine properties.
can l antih	nistamine properties.

18 TEVBIOU3_QA_08.FM Copyright © 2008 Neap

The antibody microarray is attracting a great deal of interest because it can measure protein abundance. There are many applications of this technology, but cancer research is one of the largest and most promising. With increasing use of protein biomarkers in the blood for screening and treatment of cancer, it is important to have a good, quantitative method for measuring and comparing the amounts of particular proteins.

All antibody arrays begin with a number of antibodies bonded to a plate or slide. In a sandwich assay, immobilised antibodies fixed to the slide capture unlabelled proteins from blood serum. A second, labelled antibody is used to detect the bound protein. In the microarray shown below, each slide has sixteen antibodies in sixty-four wells with an identical antibody spotted at the bottom of each row of four wells.



KEY:

AS - alignment spots

PS - positive control spots

IL - interleukin (cytokine)


 $\textbf{GM-CSF}-granulocyte\ macrophage-colony$


stimulating factor

IFN – interferon


TNF - tumour necrosis factor

This sandwich microarray was used to test blood serum from an elderly female. The serum was passed over the microarray. The microarray was then rinsed before being washed with radioactively labelled antibodies. The microarray was rinsed again, then dried, and then placed under photographic film. The images below show the developed film. Dark spots indicate the presence of radioactive antibodies. The darker the spot, the more antibody is present and hence the more serum protein has been bound.






serum from a person without bowel cancer



serum from an elderly female suspected of having bowel cancer



a. What is the purpose of the alignment and positive control si		iat is the nurnose of the alignmen	ent and positive contro	SDOTS
--	--	------------------------------------	-------------------------	-------

2 marks

b. Does the elderly female appear to have bowel cancer? Use evidence from the microarrays to support your answer.

1 mark

1 + 1 = 2 marks Total 7 marks

END OF QUESTION AND ANSWER BOOKLET