

# MAV Mathematical Methods Examination 1

## Answers & Solutions

### Part I (Multiple-choice) Answers

- |       |       |       |       |       |
|-------|-------|-------|-------|-------|
| 1. E  | 2. D  | 3. D  | 4. C  | 5. D  |
| 6. D  | 7. B  | 8. C  | 9. B  | 10. D |
| 11. C | 12. E | 13. A | 14. A | 15. E |
| 16. B | 17. A | 18. D | 19. A | 20. A |
| 21. E | 22. C | 23. A | 24. E | 25. A |
| 26. D | 27. B |       |       |       |

#### Question 1 [E]

$$f(x) = 7 - 6\sin(4x)$$

$$\text{amplitude} = 6$$

$$\text{period} = \frac{2\pi}{4} = \frac{\pi}{2}$$

#### Question 2 [D]

$$\sin^2(2\theta) = \frac{3}{4}$$

$$\sin(2\theta) = \pm \frac{\sqrt{3}}{2}$$

$$2\theta = \frac{\pi}{3}, \frac{2\pi}{3} + 2n\pi (n = -2, -1, 0, 1)$$

$$\theta = -\frac{5\pi}{6}, -\frac{2\pi}{3}, -\frac{\pi}{3}, -\frac{\pi}{6}, \frac{\pi}{6}, \frac{\pi}{3}, \frac{2\pi}{3}, \frac{5\pi}{6}$$

#### Question 3 [D]

$$y = a \cos nx$$

$$\text{period} = 2\pi, \text{amplitude} = 1$$

$$\text{Double: period} = 4\pi \therefore n = \frac{1}{2}$$

$$\text{Amplitude} = 2$$

$$\Rightarrow y = 2\cos(0.5x) - 1$$

#### Question 4 [C]

Reflect  $y = \tan x$  in the  $x$ -axis. A or C

There has been a horizontal and vertical translation, hence C

#### Question 5 [D]

$$\begin{aligned} \text{Turning point: } x &= -\frac{b}{2a} \\ &= \frac{-6}{-6} = 1 \end{aligned}$$

Substitute  $x = 1$  into  $y = -3x^2 + 6x - 3a$

$$y = -3 + 6 - 3a$$

$$y = 3 - 3a$$

#### Question 6 [D]

$$\left(2x^2 - \frac{3}{x}\right)^6$$

$$\text{General term: } {}^6C_r (2x^2)^{6-r} \left(-\frac{3}{x}\right)^r$$

$$\therefore 12 - 2r - r = 0$$

$$\therefore r = 4$$

$$= {}^6C_4 (2x^2)^2 \left(-\frac{3}{x}\right)^4$$

$$= {}^6C_4 (2)^2 (-3)^4$$

$$= 15 \times 4 \times 81$$

$$= 4860$$

#### Question 7 [B]

$$2 \log_2 x - \log_2(x+4) = 1$$

$$\Rightarrow \log_2(x^2) - \log_2(x+4) = 1$$

$$\Rightarrow \log_2\left(\frac{x^2}{x+4}\right) = 1$$

$$\Rightarrow \frac{x^2}{x+4} = 2$$

$$\Rightarrow x^2 - 2x - 8 = 0$$

$$\Rightarrow (x-4)(x+2) = 0$$

$$\Rightarrow x = 4, -2$$

Note:  $-2$  is not possible

**Question 8** [C]

$$3^{2x} - 3^{x+1} = 54$$

$$\Rightarrow 3^{2x} - 3 \times 3^x - 54 = 0$$

Let  $a = 3^x$

$$a^2 - 3a - 54 = 0$$

$$(a - 9)(a + 6) = 0$$

$a = 9$  or  $a = -6$

$3^x = 9$  or  $3^x = -6$

$x = 2$

Note:  $3^x = -6$  is not possible.

**Question 9** [B]

$$3x - 5 \neq 0$$

Hence  $x = \frac{5}{3}$  is an asymptote.

The largest domain is  $\left(\frac{5}{3}, 0\right)$

$$b = \frac{5}{3} = 1\frac{2}{3}$$


**Question 10** [D]

$$y = -4(x + 1)^2$$

Inverse:  $x = -4(y + 1)^2$

$$y = -\sqrt{-\frac{x}{4}} - 1$$

$$= \frac{-\sqrt{-x}}{2} - 1$$



dom:  $(-\infty, 0]$

**Question 11** [C]

Note  $a$  and  $b$  are negative.

$y = \frac{1}{x}$  has been reflected in the  $x$ -axis

$y = -\frac{1}{x}$  and then translated  $-a$  units to the left and  $-b$  units down.

$$y = -\frac{1}{x-a} + b$$

**Question 12** [E]

The curve has a stationary point of inflection at  $(B, C)$ .

**Question 13** [A]

Translation 3 to right  $\therefore B = -3$

Substitute  $(7, -1) \Rightarrow -1 = A \log_e 4$

$$A = \frac{-1}{\log_e 4}$$

**Question 14** [A]

$$f(0) = -1$$

$$f(2) = e^2$$

$$\text{Gradient} = \frac{e^2 - -1}{2 - 0}$$

$$= \frac{e^2 + 1}{2}$$

**Question 15** [E]

When  $x = \pi, y = \frac{\pi}{-1} = -\pi$

From calculator: tangent is  $y = -x$

$\Rightarrow$  normal:  $y = x - 2\pi$

**Question 16** [B]

Chain rule:  $\frac{dy}{dx} = \frac{1}{\tan x} \times \sec^2 x$

$$= \frac{\cos x}{\sin x} \times \frac{\sec x}{\cos x}$$

$$= \frac{\sec x}{\sin x}$$

**Question 17** [A]

$$y = \left(\sqrt{x^2 + 1}\right)^3 = (x^2 + 1)^{\frac{3}{2}}$$

$$\frac{dy}{dx} = \frac{3}{2}(x^2 + 1)^{\frac{1}{2}} \times 2x$$

$$= 3x(x^2 + 1)^{\frac{1}{2}}$$

$$= 3xy^{\frac{1}{3}}$$

**Question 18** [D]

From calculator: note: point of inflection at  $x = 1$

**Question 19****[A]**

Use TABLE on calculator (or graphics calculator program).

$$A = \frac{1}{2} [f(0) + f(0.5) + f(1) + f(1.5) + f(2) + f(2.5)]$$

$$= 231.01 \text{ square units}$$

**Question 20****[A]**

$f(x)$  is gradient function of  $h(x)$

$$h'(x) = 0 \text{ when } x \approx -2.3$$

$$h'(x) < 0 \text{ when } x < -2.3$$

$$h'(x) > 0 \text{ when } x > -2.3$$

Either A or D

$$f(x) = (x+1)^3 + 2$$

$$h(x) = \int ((x+1)^3 + 2) dx$$

$$= \frac{(x+1)^4}{4} + 2x + c$$

Hence A

**Question 21****[E]**

$$\int \frac{e^{3x} + 1}{e^x} dx$$

$$= \int (e^{2x} + e^{-x}) dx$$

$$= \frac{e^{2x}}{2} - e^{-x} + c$$

**Question 22****[C]**

$$\int_1^a \frac{1}{(x-2)^3} dx = -\frac{1}{2}$$

$$\left[ \frac{1}{-2(x-2)^2} \right]_1^a = -\frac{1}{2}$$

$$\frac{1}{-2(a-2)^2} + \frac{1}{2} = -\frac{1}{2}$$

$$\frac{1}{-2(a-2)^2} = -1$$

$$2(a-2)^2 = 1$$


$$(a-2)^2 = \frac{1}{2}$$

$$a-2 = \pm \sqrt{\frac{1}{2}}$$

$$a = 2 \pm \sqrt{\frac{1}{2}}$$

$$= 2 \pm \frac{\sqrt{2}}{2}$$

cannot be  $2 + \frac{\sqrt{2}}{2}$

**Question 23****[A]**

From calculator:

$$\text{Area} = \left| \int_0^1 (2^x(x-1)^3) dx \right|$$

$$= 0.289$$

**Question 24****[E]**

$$10a = 1$$

$$a = 0.1$$

$$E(X) = 1 \times 0.2 + 2 \times 0.4 + 3 \times 0.3$$

$$= 1.9$$

$$\therefore E(2X - 1) = 2 \times 1.9 - 1$$

$$= 2.8$$


**Question 25**

$\Pr(\text{correct}) = 0.2$

$$\begin{aligned} \Pr(20 \text{ correct}) &= {}^{33}C_{20}(0.2)^{20}(0.8)^{13} \\ &= {}^{33}C_{13}(0.2)^{20}(0.8)^{13} \end{aligned}$$


**Question 26**

$$\Pr(\text{pair}) = \frac{\binom{b}{2}\binom{n}{0} + \binom{b}{0}\binom{n}{2}}{\binom{b+n}{2}}$$



$$\Pr(\text{pair}) = \frac{b(b-1) + n(n-1)}{(b+n)(b+n-1)}$$

**Question 27**



$\text{invNorm}(0.95) = 1.6449$

$$z = \frac{x - \mu}{\sigma}$$

$$1.6449 = \frac{6.5 - 6}{\sigma}$$

$\Rightarrow \sigma = 0.304$

[A]

[D]

[B]

**Part II Solutions**

**Question 1**

$4\cos^2x + 4\sin x = 1$

$\Rightarrow 4 - 4\sin^2x + 4\sin x = 1$

[M]

$\Rightarrow 4\sin^2x - 4\sin x - 3 = 0$

$\Rightarrow (2\sin x - 3)(2\sin x + 1) = 0$

[M]

$\Rightarrow \sin x = \frac{3}{2}$  (not possible) or  $\sin x = \frac{-1}{2}$

$\Rightarrow x = \frac{7\pi}{6}, \frac{11\pi}{6}$

[A]

**Question 2**

a. 
$$\frac{x^2 + 2x + 2}{(x+1)^2}$$

$$= \frac{x^2 + 2x + 1 - 1 + 2}{(x+1)^2}$$

$$= \frac{(x+1)^2 + 1}{(x+1)^2}$$

$$= 1 + \frac{1}{(x+1)^2}$$

$$= \frac{1}{(1+x)^2} + 1$$

[A][A]

b.  $f_1(x) = \frac{2}{x^2} + 2$

[A]

c.  $(2, \infty)$

[A]

**Question 3**

a.  $d = 1$  [A]

$$\frac{dy}{dx} = 3x^2 + 2bx + c$$

$$\Rightarrow 0 = 3 + 2b + c$$

$$2b + c = -3 \quad \dots\dots \textcircled{1}$$

$$y = x^3 + bx^2 + cx + 1$$

$$2 = 1 + b + c + 1$$

$$\therefore b + c = 0 \quad \dots\dots \textcircled{2}$$

Solve  $\textcircled{1}$  and  $\textcircled{2}$  simultaneously

$$b = -3, c = 3$$

$$\Rightarrow y = x^3 - 3x^2 + 3x + 1 \quad \text{[A][A]}$$

b.  $y = (x - 1)^3 + 2$  [A]

c. If  $y = 0$

$$\Rightarrow (x - 1)^3 + 2 = 0$$

$$\Rightarrow (x - 1)^3 = -2$$

$$\Rightarrow x - 1 = \sqrt[3]{-2}$$

$$\Rightarrow x = 1 + \sqrt[3]{-2} \quad \text{[M]}$$

**Question 4**

a.  $m = \frac{8}{4} = 2$  [A]

$$y = 2x^4 - 8 \quad \text{[A]}$$

b.  $y = 2(x^4 - 4)$

$$= 2(x^2 - 2)(x^2 + 2)$$

$$= 2(x + \sqrt{2})(x - \sqrt{2})(x^2 + 2)$$

When  $y = 0$ ,  $x = \pm\sqrt{2}$  [A][A]

c.  $A = \left| \int_{-\sqrt{2}}^{\sqrt{2}} (2x^4 - 8) dx \right|$  [M]

( = 18.10) for checking

$$= \left| \left[ \frac{2}{5} x^5 - 8x \right]_{-\sqrt{2}}^{\sqrt{2}} \right|$$

$$= \left| \left( \frac{2 \times 4\sqrt{2}}{5} - 8\sqrt{2} \right) - \left( \frac{2 \times -4\sqrt{2}}{5} + 8\sqrt{2} \right) \right|$$

$$= \frac{64}{5} \sqrt{2} \quad \text{[A]}$$

**Question 5**

Y	0	1	2	3	4
f(Y)	0	$\frac{1}{10}$	$\frac{2}{10}$	$\frac{3}{10}$	$\frac{4}{10}$

[A]

$$E(Y) = 3 ; \quad E(Y^2) = 10 \quad \text{[A]}$$

$$\begin{aligned} \text{Var}(Y) &= E(Y^2) - [E(Y)]^2 \\ &= 1 \Rightarrow \sigma = 1 \quad \text{[A]} \end{aligned}$$

**Question 6**

$$\Pr(X \geq 3) = 1 - \Pr(X \leq 2)$$

$$= 1 - \text{binomcdf}(10, 0.4, 2) \quad \text{[M]}$$

$$= 0.8327 \quad \text{[A]}$$