

'2016 Examination Package' - Trial Examination 3 of 5

<i>'</i>		<i>`\</i>
i		1
i		1
1		- 1
1		Į.
1	THIS BOX IS FOR ILLUSTRATIVE PURPOSES ONLY	ļ.
1		Į.
1		!
1		!
1		,
\ .		_/

	STUDENT	Γ NUMBE	R			_	Letter
Figures							
Words							

MATHEMATICAL METHODS

Units 3 & 4 – Written examination 1

(TSSM's 2013 trial exam updated for the current study design)

Reading time: 15 minutes Writing time: 1 hour

QUESTION & ANSWER BOOK

Structure of book

Number of questions	Number of questions to be answered	Number of marks
8	8	40
		Total 40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers
- Students are NOT permitted to bring into the examination room: notes of any kind, blank sheets of paper, white out liquid/tape or a calculator of any type.

Materials supplied

- Question and answer book of 9 pages.
- Working space is provided throughout the book.

Instructions

- Print your name in the space provided on the top of this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic communication devices into the examination room.

© TSSM 2013 Page 1 of 9

Instructions

Answer all questions in the spaces provided.

A decimal approximation will not be accepted if an exact answer is required to a question. In questions where more than one mark is available, appropriate working must be shown. Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Ouestion	1
Oueshon	1

$$Let f(x) = ke^x - 1 + ke^{-x}$$

				2 n
Find the solutic	on(s) to the equation	f(x) = 1 in terms of	f <i>k</i> .	

2 marks Total 4 marks

© TSSM 2013 Page 2 of 9

Question 2

Consider $f(x) = 2x\sqrt{1-x}$

a.	Find $f'(x)$.
	2 marks
b.	Hence, find the area between the curve $y = \frac{x}{\sqrt{1-x}}$, the x-axis and the lines $x = -2$ and $x = 0$.

3 marks Total 5 marks

TURN OVER

© TSSM 2013 Page 3 of 9

Question 3

Consider the function $g(x) = \sin(x) + \cos(x)$

a. Find the general solution for the equation g(2x) = 0.

2 marks

b. Hence, find all solutions to the equation g(2x)=0 for $x \in (-\pi,\pi)$.

2 marks

Total 4 marks

Question 4

a. Find the image, g(x), of the equation $f(x) = \frac{1}{2x}$ under the matrix transformation $T = \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix}$.

3 marks

© TSSM 2013 Page 4 of 9

).	State the transformations made to the function $f(x)$ to obtain $g(x)$.	
		2 marks
	Find the inverse, $g^{-1}(x)$, of the transformed function $g(x)$.	
		3 marks
		3 marks
l.	State the domain and range of $g^{-1}(x)$.	
		2 marks Total 10 marks
		TOTAL TO IIIAIK

TURN OVER

© TSSM 2013 Page 5 of 9

Question 5

A random variable X has probability density function given by

$$f(x) = \begin{cases} 0.3 p, & 0 \le x \le 4 \\ xp, & 4 \le x \le 5 \\ 0, & elsewhere \end{cases}$$

a.	Find the exact value of p .	
		3 marks
b.	Find the exact value of Pr $(1 < x < 2.5)$.	
		2 marks

Total 5 marks

© TSSM 2013 Page 6 of 9

Question 6

An approximate 95% confidence interval for the population proportion is given by the interval (r, s).

a.	Find an expression for the sample proportion \hat{p} in terms of r and s .
	2 marks
b.	Find an expression for the margin of error for this confidence interval in terms of r and s .
	2 marks
	Total 4 marks

TURN OVER

© TSSM 2013 Page 7 of 9

Question 7 Find the equation of the tangent(s) to the curve $y = -x^2 - 8x$ such that the tangent	gent(s) cross the
y axis at $(0,3)$.	<i>5</i> ()
	
	4 marks
	THAIRS
Question 8	
Consider the function $f(x) = \sin\left(\frac{4x}{3}\right)$	
Consider the ranction $I(x) = \sin\left(\frac{1}{3}\right)$	
a. Find the value of $\int_0^{\frac{\pi}{8}} f(x) dx$.	
a. Find the value of $\int_0^8 f(x) dx$.	

2 marks

© TSSM 2013 Page 8 of 9

b.	Find the average value of $f(x)$ over the interval $\left[0, \frac{\pi}{4}\right]$.
	2 marks
	Total 4 marks

END OF QUESTION AND ANSWER BOOK

© TSSM 2013 Page 9 of 9