

2015 Trial Examination

STUDENT NUMBER

Figures	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	Letter
Words	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

MATHEMATICAL METHODS (CAS)

Units 3 & 4 – Written examination 2

Reading time: 15 minutes

Writing time: 2 hours

QUESTION & ANSWER BOOK

Structure of book

<i>Section</i>	<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
1	22	22	22
2	5	5	58
			Total 80

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, one bound reference, one approved CAS calculator (memory DOES NOT need to be cleared) and, if desired, one scientific calculator.
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 21 pages including answer sheet for multiple-choice questions.

Instructions

- Print your name in the space provided on the top of this page and the multiple-choice answer sheet.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic communication devices into the examination room.

SECTION 1 – Multiple-choice questions**Instructions for Section 1**

Answer all questions on the answer sheet provided for multiple choice questions.

Choose the response that is **correct** for the question.

A correct answer scores 1, an incorrect answer scores 0.


Marks will **not** be deducted for incorrect answers.

No marks will be given if more than one answer is completed for any question.

Question 1

The x -value of the point of intersection of the graphs of $y = 3$ and $y = \frac{5}{2}x + 1$ is:

- A. 0.8
- B. 1.6
- C. 3
- D. 5
- E. 8

Question 2

The function that represents the graph above is:

- A. $y = x^3 - x^2$
- B. $y = 3x^3 - x^2$
- C. $y = 3x^2 - x^3$
- D. $y = x^2 - 3x^3$
- E. $y = 3x^4 - x^2$

Question 3

If $f(x) = x^2 + 2x + 5$ and $g(x) = x - 1$, then $f(g(x))$ is:

- A. $x^2 + 3x + 4$
- B. $x^2 - 4$
- C. $x^2 + 4x - 1$
- D. $x^2 - 1$
- E. $x^2 + 4$

Question 4

The function $f(x) = 2\sqrt{x} + 1$ is dilated by a factor of $\frac{1}{2}$ unit from the x -axis to transform to the function $g(x)$. The rule for $g(x)$ is:

- A. $g(x) = \sqrt{x} + 1$
- B. $g(x) = \sqrt{\frac{x}{2}} + 1$
- C. $g(x) = \sqrt{2x} + 1$
- D. $g(x) = \sqrt{x} + \frac{1}{2}$
- E. $g(x) = \sqrt{\frac{x}{2}} + \frac{1}{2}$

Question 5

The domain and range of the function $f(x) = \sqrt{4 - x}$ respectively are:

- A. $[4, \infty)$ and $[2, \infty)$
- B. $(-\infty, 4]$ and R
- C. $(-\infty, 4]$ and $[0, \infty)$
- D. $[4, \infty)$ and $(-\infty, 0]$
- E. R and $[0, \infty)$


SECTION 1 - continued

TURN OVER

Question 6

The average rate of change for the function $g(x) = \sqrt{2x}$ between $x = 2$ and $x = 8$ is:

- A. $\frac{1}{3}$
- B. $-\frac{3}{10}$
- C. 2
- D. 3
- E. 7

Question 7

The function that describes the graph of the hybrid function $y = f(x)$ shown above is:

- A. $f(x) = \begin{cases} \frac{1}{2}x, & -4 < x < 0 \\ x, & 0 \leq x \leq 4 \end{cases}$
- B. $f(x) = \begin{cases} -\frac{1}{2}x, & -4 \leq x < 0 \\ \frac{1}{2}x, & 0 \leq x \leq 4 \end{cases}$
- C. $f(x) = \begin{cases} -\frac{1}{2}x, & -4 \leq x < 0 \\ x, & 0 \leq x \leq 3 \end{cases}$
- D. $f(x) = \begin{cases} -\frac{1}{2}x, & -4 \leq x < 0 \\ x, & 0 < x < 4 \end{cases}$
- E. $f(x) = \begin{cases} -\frac{1}{2}x, & -4 \leq x < 0 \\ 2x, & 0 \leq x \leq 4 \end{cases}$

Question 8

The domain of the composite function $f \circ g$ where $f(x) = \frac{x}{x+6}$ and $g(x) = \frac{18}{x+2}$ is:

- A. R
- B. $R \setminus \{-5\}$
- C. $R \setminus \{-2, -5\}$
- D. $R \setminus \{-2, -6\}$
- E. $R \setminus \{0, -2, -5\}$

Question 9

Which of the following functions is one-to-one?

- A. $f: R \rightarrow R, f(x) = x^2$
- B. $f: [-1, \infty) \rightarrow R, f(x) = x^2$
- C. $f: [-1, \infty) \rightarrow R, f(x) = |x|$
- D. $f: (-\infty, 3] \rightarrow R, f(x) = 1 - x^2$
- E. $f: [1, \infty) \rightarrow R, f(x) = |x - 1|$

Question 10

The amplitude and period of $y = 2 \cos\left(\frac{x}{5}\right)$ respectively are:

- A. 5 and 4π
- B. 2 and $\frac{2\pi}{5}$
- C. $\frac{1}{2}$ and $\frac{2\pi}{5}$
- D. 2 and 10π
- E. $\frac{1}{2}$ and 10π

SECTION 1 - continued

TURN OVER

Question 11

The gradient of the tangent line to the graph of $y = -x^2 + 4\sqrt{x}$ at $x = 4$ is:

- A. -8
- B. -9
- C. -10
- D. -5
- E. -7

Question 12

Let $f(x) = x^3$

Let A_1 be the area bounded by the graphs of $y = -1$ and $y = f(x)$ between $x = -1$ and $x = 0$ and let A_2 be the area bounded by the graphs of $y = 1$ and $y = f(x)$ between $x = 0$ and $x = 1$. The total area of these two regions can be represented in the integral form as:

- A. $\int_{-1}^1 (1 - x^3) dx$
- B. $2 \int_0^1 (1 - x^3) dx$
- C. $2 \int_0^1 (1 + x^3) dx$
- D. $\int_{-1}^1 (1 + x^3) dx$
- E. $\int_0^1 (-1 - x^3) dx$

Question 13

The equation of the normal line to the graph of $y = 2x\sqrt{x^2 + 8} + 2$ at the point $(0, 2)$ is:

- A. $x - 4\sqrt{2}y + 8\sqrt{2} = 0$
- B. $x + 4\sqrt{2}y = 8\sqrt{2}$
- C. $4\sqrt{2}x + y = 2$
- D. $-4\sqrt{2}x + y = 2$
- E. $x + 4\sqrt{2}y = 2$

Question 14

Which of the following functions satisfy the condition $(f(x))^2 \times (f(y))^2 = f(2x + 2y)$?

- A. $f(x) = x + y$
- B. $f(x) = \sqrt{x}$
- C. $f(x) = e^x$
- D. $f(x) = \log_e x$
- E. $f(x) = \frac{1}{x}$

Question 15

The function $f(x) = x^3$ has an average value of 9 on the closed interval $[0, k]$.

The value of k is:

- A. $\sqrt[3]{36}$
- B. $\sqrt{3}$
- C. $\sqrt[3]{18}$
- D. 6
- E. 3

Question 16

A test consists of 10 multiple choice questions with five choices for each question. If you guess on each and every answer without even reading the questions, the probability of getting exactly six questions correct is:

- A. 0.999
- B. 0.006
- C. 0.088
- D. 0.12
- E. 0.046

SECTION 1 - continued

TURN OVER

Question 17

Heights of college women have a distribution that can be approximated by a normal curve with a mean of 165cm and a standard deviation equal to 7.62cm. The proportion of college women between 165 and 170 tall is closest to:

- A. 0.50
- B. 0.75
- C. 0.24
- D. 0.15
- E. 0.17

Question 18

The probability that a certain machine will produce a defective item is 0.20. If a random sample of 6 items is taken from the output of this machine, the probability that there will be 5 or more defectives in the sample is:

- A. 0.0016
- B. 0.0154
- C. 0.0015
- D. 0.2458
- E. 0.0001

Question 19

Tina's score on her midterm exam was at the 50th percentile. The grades were normally distributed and the exam average was 78 and the standard deviation was 6. Tina's score on the exam was:

- A. 90
- B. 50
- C. 84
- D. 78
- E. 72

Question 20

The maximum value of $f(x) = \sin(x) + \cos(2x)$ over $0 \leq x \leq 2\pi$ is:

- A. 0
- B. 1
- C. 1.125
- D. 2.5
- E. 2.7

Question 21

For the following discrete probability distribution, the value of $E(2X + 1)$ is closest to:

x	0	2	4	6
$\text{Pr}(x)$	0.15	0.20	k	0.45

- A. 7
- B. 8
- C. 9
- D. 12
- E. 13

Question 22

Two successive asymptotes of the function $f(x) = \tan(nx)$ have equations $x = 3$ and $x = 6$.
The value of n is:

- A. π
- B. $\frac{\pi}{3}$
- C. $\frac{\pi}{2}$
- D. $\frac{3\pi}{2}$
- E. $\frac{\pi}{6}$

END OF SECTION 1

TURN OVER

SECTION 2 – Analysis questions**Instructions for Section 2**

Answer **all** questions in the spaces provided.

A decimal approximation will not be accepted if an **exact** answer is required to a question.


In questions where more than one mark is available, appropriate working **must** be shown.

Where an instruction to **use calculus** is stated for a question, you must show an appropriate derivative or anti-derivative.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1 (12 marks)

An inverted cone of radius r cm and height h cm is shown below.



Let the slant height of the cone be l cm and the semi-vertical angle be α .

- a. Find the radius and the height of the cone in terms of l and α .

2 marks

- b. Show that the volume of the cone can be expressed in terms of l and α as

$$V = \frac{\pi}{3} l^3 \sin^2(\alpha) \cos(\alpha)$$

1 mark

- c. Find the stationary point(s) of $V(\alpha)$.

4 marks

- d. If the slant height of the cone is 6cm find the maximum volume of the cone.

2 marks

SECTION 2 - continued

TURN OVER

Let $\alpha = 30^\circ$. Water is poured into this conical flask at the rate of $7\text{cm}^3/\text{sec}$.

- e. Find the rate, correct to two decimal places, at which the slant height is changing when the radius is 4 cm.

3 marks

Question 2 (9 marks)

The population of foxes in a certain forest varies according to the rule

$$P(t) = 500 - 300\cos\left(\frac{\pi}{2.2}(t - 2.9)\right)$$

where $P(t)$ is the number of foxes t years after 2011.

- a. Find the period and amplitude of this function.

2 marks

- b. Find the maximum and minimum number of foxes in this forest.

2 marks

- c. After how many months is the population of foxes a maximum in the first 5 years?

2 marks

Foxes are declared a vulnerable species if their population drops below 300.

- d. Over which two time intervals will the population of foxes become vulnerable in the first 10 years? Give your answer correct to one decimal place.

3 marks


SECTION 2 - continued

TURN OVER

Question 3 (11 marks)

While smoking tobacco, the body absorbs many chemical compounds in addition to nicotine, including cyanide (which is highly toxic to humans).

The function $C(t) = 0.1 + 0.3t^{0.6}e^{-0.17t}$ is a reasonable model of the measured blood cyanide concentrations in $\mu\text{g/mL}$ after t minutes, which is shown in the figure below.



- a. Find the maximum concentration of blood cyanide, correct to 2 decimal places.

1 mark

- b. When, to the nearest tenth of a minute, is the concentration a maximum?

1 mark

- c. Find the concentration, correct to two decimal places, after 10 minutes.

2 marks

SECTION 2 - continued

- d. Find the value of the average rate of change of concentration over the time interval $\left[\frac{3}{2}, 5\right]$ correct to four decimal places.

2 marks

- e. Find the time interval, correct to 2 decimal places, when the rate of change of concentration is negative?

2 marks

SECTION 2 - continued

TURN OVER

Another function $C_1(t) = ate^{-bt}$ is used to measure the concentration of another substance in the blood in $\mu\text{g/mL}$ after t minutes.

- f. Find the values of a and b if the maximum amount of this substance in the blood was $120 \mu\text{g/mL}$ after 2 hours?

3 marks

Question 4 (11 marks)

Consider the function $f(x) = x^2 + bx + 3$, where $b > 0$ and $g(x) = x^2$.

- a. If the function $f(x)$ can be written as $f(x) = (x + 5)^2 - 22$, find the value of b .

2 marks

b. State the transformations that transformed the graph of $y = f(x)$ to $y = g(x)$.

2 marks

c. Explain why $f(g(x))$ exists and find the rule for $f(g(x))$.

3 marks

Let $h(x) = f(g(x))$.

d. Find the equation of the tangent to the graph of $y = h(x)$ at the point $x = k$.

2 marks

SECTION 2 - continued

TURN OVER

- e. Write down the integral that will find the area bounded by the graph of $y = h(x)$ and the x -axis between $x = 0$ and $x = 3$.

2 marks

Question 5 (15 marks)

The distance, in kilometres, that Jack walks each week follows a normal distribution with a mean of 7.5 km and standard deviation of 2.5km.

- a. Find the probability, correct to four decimal places, that in a particular week Jack walks
- i. less than 11km.

- ii. between 5.5km and 10.5km

2 marks

10% of the time Jack walks less than d km.
Find the value of d , correct to one decimal place.

2 marks

Jack observes his pattern of walking over a 6 week time period.

- b. Find, to 4 decimal places, the probability that in 4 out of 6 weeks, he walks for at least 6.8km.

3 marks

Julie also walks each week following the normal pattern with mean distance of 6.4km. She has 65% chance of walking at least 5km each week.

- c. Find, to 2 decimal places, the standard deviation of the distance she covers each week.

3 marks

SECTION 2 - continued

TURN OVER

Jeremy also decides to walk each week to keep fit. The chance of him walking a particular day when he walked the previous day is 0.45. If he does not walk on any day his chance of walking the following day is 0.67.

- d. If Jeremy walks on Sunday, find the probability that he walks on at least 2 out of the next 3 days.

3 marks

- e. Find, to 2 decimal places, the long-term chance of Jeremy going for a walk.

2 marks

END OF QUESTION AND ANSWER BOOK

MULTIPLE CHOICE ANSWER SHEET

Student Name: _____

Circle the letter that corresponds to each correct answer.

Question					
1	A	B	C	D	E
2	A	B	C	D	E
3	A	B	C	D	E
4	A	B	C	D	E
5	A	B	C	D	E
6	A	B	C	D	E
7	A	B	C	D	E
8	A	B	C	D	E
9	A	B	C	D	E
10	A	B	C	D	E
11	A	B	C	D	E
12	A	B	C	D	E
13	A	B	C	D	E
14	A	B	C	D	E
15	A	B	C	D	E
16	A	B	C	D	E
17	A	B	C	D	E
18	A	B	C	D	E
19	A	B	C	D	E
20	A	B	C	D	E
21	A	B	C	D	E
22	A	B	C	D	E