


MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

Question 1



correct shape [A1]

x-intercept: $y = 0$ $\frac{4}{x} - 2x^2 = 0$

$$4 - 2x^3 = 0$$

$$x^3 = 2$$

$$x = \sqrt[3]{2}$$

[A1]

Turning point: $\frac{dy}{dx} = 0$ $y = 4x^{-1} - 2x^2$

$$\frac{dy}{dx} = -4x^{-2} - 4x$$

$$-\frac{4}{x^2} - 4x = 0$$

$$4 + 4x^3 = 0$$

$$x^3 = -1$$

$$x = -1$$

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

When, $y = \frac{4}{-1} - 2(-1)^2$

Maximum at $(-1, -6)$

[A1]

Asymptotes: $x = 0, y = -2x^2$

[A2]

Total 5 marks

Question 2

a.

$$P(z) = z^2 + (1 + i\sqrt{2})z + 4i\sqrt{2} - 12$$

$$P(-4) = 16 - 4 - 4i\sqrt{2} + 4i\sqrt{2} - 12 = 0$$

[M1]

b.

$$z^2 + (1 + i\sqrt{2})z + 4i\sqrt{2} - 12 = 0$$

$$z^2 + (1 + i\sqrt{2})z + 4(i\sqrt{2} - 3) = (z + 4)(z + a + bi)$$

[M1]

$$= z^2 + az + biz + 4z + 4a + 4bi$$

$$= z^2 + (a + 4 + bi)z + 4a + 4bi$$

[A1]

Equating coefficients of z :

[M1]

$$1 + i\sqrt{2} = a + 4 + bi \quad \text{equating real components and equating imaginary components}$$

$$a + 4 = 1 \quad \text{and} \quad b = \sqrt{2}$$

$$a = -3$$

Therefore the other factor is $(z - 3 + i\sqrt{2})$, hence the other solution is $3 - i\sqrt{2}$

[A1]


Total 5 marks

Question 3

The three forces P , T , and $24g$ will be in equilibrium.


$$\cos \theta = \frac{2}{4}$$

$$\theta = 60^\circ$$



MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

Adding the forces head to tail.



[M1]

From the triangle of forces

$$\sin(30^\circ) = \frac{P}{T} \quad \text{and}$$

$$\cos(30^\circ) = \frac{24g}{T}$$

$$P = \frac{1}{2}T \quad \dots (1)$$

$$T = \frac{24g}{\cos(30^\circ)} = \frac{48g}{\sqrt{3}} = 16\sqrt{3}g \quad \dots (2) \quad \text{[A1]}$$

Substituting (2) into (1)

$$P = \frac{1}{2} \times 16\sqrt{3}g$$

$$P = 8\sqrt{3}g \text{ newtons}$$

[A1]

Total 3 marks

Question 4

$$\cot(2x) - \cot(x) = 2$$

$$\frac{\cos(2x)}{\sin(2x)} - \frac{\cos(x)}{\sin(x)} = 2$$

$$\frac{\sin(x)\cos(2x) - \cos(x)\sin(2x)}{\sin(2x)\sin(x)} = 2 \quad \text{[M1]}$$

$$\frac{\sin(x - 2x)}{\sin(2x)\sin(x)} = 2 \quad \text{[M1]}$$

$$\frac{\sin(-x)}{\sin(2x)\sin(x)} = 2$$

$$\frac{-\sin(x)}{\sin(2x)\sin(x)} = 2$$

$$\frac{-1}{\sin(2x)} = 2$$

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS


$$\sin(2x) = -\frac{1}{2} \quad \text{[A1]}$$

$$2x = -\frac{5\pi}{6}, \quad -\frac{\pi}{6}, \quad \frac{7\pi}{6}, \quad \frac{11\pi}{6}$$

$$x = -\frac{5\pi}{12}, \quad -\frac{\pi}{12}, \quad \frac{7\pi}{12}, \quad \frac{11\pi}{12} \quad \text{[A1]}$$

Total 4 marks

Question 5



There is a turning point on $y = f(x)$ at $x = a$ that touches the x -axis, so $y = F(x)$ will have a stationary point of inflexion at $x = a$. [A1]

There is a maximum turning point on $y = f(x)$ at $x = c$, so $y = F(x)$ will have a non-stationary point of inflexion at $x = c$. This is the point of steepest slope on $y = F(x)$ for $a \leq x \leq d$ [A1]

There is an x -intercept on $y = f(x)$ at $x = d$ cutting the x -axis from positive to negative, so $y = F(x)$ will have a local maximum at $x = d$. [A1]

Total 3 marks

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

Question 6

$$y = \cot(x)$$

$$y = \frac{\cos(x)}{\sin(x)}$$

$$\frac{dy}{dx} = \frac{\sin(x) \times (-\sin(x)) - \cos(x) \times \cos(x)}{\sin^2(x)}$$

[M1]

$$= \frac{-(\sin^2(x) + \cos^2(x))}{\sin^2(x)}$$

$$= \frac{-1}{\sin^2(x)}$$

$$= -\operatorname{cosec}^2(x)$$

[A1]

Total 2 marks

Question 7

Use implicit differentiation and apply the product rule.

$$5y = 2xy^2 - 3x$$

$$\frac{d}{dx}(5y) = \frac{d}{dx}(2xy^2) - \frac{d}{dx}(3x)$$

$$5 \frac{dy}{dx} = \left(y^2 \frac{d}{dx}(2x) + 2x \frac{d}{dx}(y^2) \right) - \frac{d}{dx}(3x)$$

$$5 \frac{dy}{dx} = \left(2 \times y^2 + 2x \times 2y \times \frac{dy}{dx} \right) - 3$$

[M1]

$$5 \frac{dy}{dx} = 2y^2 + 4xy \frac{dy}{dx} - 3$$

$$(5 - 4xy) \frac{dy}{dx} = 2y^2 - 3$$

$$\frac{dy}{dx} = \frac{2y^2 - 3}{5 - 4xy}$$

[A1]

Total 2 marks

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

Question 8

$$\int \frac{\sqrt{x}}{x+1} dx$$

Let $u = \sqrt{x} = x^{\frac{1}{2}}$ hence $x = u^2$

$$\begin{aligned} \frac{du}{dx} &= \frac{1}{2} x^{-\frac{1}{2}} \\ &= \frac{1}{2\sqrt{x}} \\ &= \frac{1}{2u} \end{aligned}$$

[M1]

$$\begin{aligned} &\int \frac{u}{u+1} 2u \frac{du}{dx} dx \\ &= \int \frac{2u^2}{u^2+1} du \end{aligned}$$

$$\begin{aligned} \frac{2u^2}{u^2+1} &= \frac{(u^2+1) + (u^2+1) - 2}{u^2+1} && \text{Alternatively, perform long division.} \\ &= 2 - \frac{2}{u^2+1} \end{aligned}$$

[M1, A1]

$$\begin{aligned} &= \int \left(2 - \frac{2}{u^2+1} \right) du \\ &= 2u - 2 \tan^{-1}(u) + c \\ &= 2\sqrt{x} - 2 \tan^{-1}(\sqrt{x}) + c \end{aligned}$$

[A1]

Total 4 marks

Question 9

$$\int \frac{1}{1+\cos(x)} dx$$

$$\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$$

$$= \int \frac{1}{2 \cos^2\left(\frac{x}{2}\right)} dx$$

$$2 \cos^2\left(\frac{x}{2}\right) = 1 + \cos(x)$$

[M1]

$$= \int \frac{1}{2} \sec^2\left(\frac{x}{2}\right) dx$$

[A1]

$$= \frac{1}{2} \times 2 \tan\left(\frac{x}{2}\right) + c$$

$$= \tan\left(\frac{x}{2}\right) + c$$

[A1]

Total 3 marks

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

Question 10

a.

Finding a relationship between the radius and height of the solution in the filter:

$$\tan(60^\circ) = \frac{r}{h} \Rightarrow r = \sqrt{3}h \dots (1)$$

Volume of cone: $V = \frac{1}{3}\pi r^2 h$

Substitute (1) $V = \frac{1}{3}\pi (\sqrt{3}h)^2 h$

$$V = \pi h^3 \dots (2) \quad \text{[A1]}$$

Related rate equation:

$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt} \quad \text{Given } \frac{dV}{dt} = -8\sqrt{h} \quad \text{and from (2)} \quad \frac{dV}{dh} = 3\pi h^2$$

$$-8\sqrt{h} = 3\pi h^2 \times \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{8}{3\pi h^{\frac{3}{2}}} \quad \text{[A1]}$$

$$\frac{dh}{dt} = -\frac{8}{3\pi\sqrt{h^3}} \quad \text{as required.}$$

b.

$$\frac{dt}{dh} = -\frac{3\pi\sqrt{h^3}}{8}$$

$$t = -\frac{3\pi}{8} \int h^{\frac{3}{2}} dt$$

$$t = -\frac{3\pi}{8} \times \frac{2}{5} h^{\frac{5}{2}} + c$$

$$t = -\frac{3\pi}{20} h^{\frac{5}{2}} + c \quad \text{[M1]}$$

When $t = 0$, $h = 4 \Rightarrow 0 = -\frac{3\pi}{20} \times (4)^{\frac{5}{2}} + c$

$$\Rightarrow c = \frac{3\pi}{20} \times 2^5$$

$$\Rightarrow c = \frac{24\pi}{5} \quad \text{[A1]}$$

Hence $t = -\frac{3\pi}{20} h^{\frac{5}{2}} + \frac{24\pi}{5}$

Solution will have passed through filter when $h = 0$

MAV Trial Examination Papers 2009
Specialist Examination 1
SOLUTIONS

$$t = -\frac{3\pi}{20}(0)^5 + \frac{24\pi}{5}$$

$$t = \frac{24\pi}{5} \text{ minutes}$$

The solution will have completely passed through the filter after $\frac{24\pi}{5}$ minutes.

[A1]

Total 5 marks

Question 11

a.

Finding an expression for v in terms of x .

$$a = -\frac{3\sqrt{x}}{8}$$

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -\frac{3\sqrt{x}}{8}$$

$$\frac{1}{2}v^2 = \int -\frac{3\sqrt{x}}{8} dx$$

[M1]

$$\frac{1}{2}v^2 = -\frac{3}{8} \int x^{\frac{1}{2}} dx$$

$$\frac{1}{2}v^2 = -\frac{3}{8} \times \frac{2}{3} x^{\frac{3}{2}} + c$$

Initially $x = 0$, $v = -2$

$$\frac{1}{2}(-2)^2 = -\frac{1}{4}(0)^{\frac{3}{2}} + c$$

$$\therefore c = 2$$

$$\frac{1}{2}v^2 = 2 - \frac{1}{4}x^{\frac{3}{2}}$$

[A1]

$$v^2 = 4 - \frac{1}{2}x^{\frac{3}{2}}$$

$$v = -\sqrt{4 - \frac{1}{2}x^{\frac{3}{2}}} \quad \text{Negative root since } v = -2 \text{ m/s when } x = 0$$

[A1]

b.

Acceleration valid for $4 - \frac{1}{2}x^{\frac{3}{2}} \geq 0$

$$8 - (\sqrt{x})^3 \geq 0$$

$$x \geq 0 \text{ and } \sqrt{x} \leq 2$$

$$\therefore 0 \leq x \leq 4$$

[A1]

Total 4 marks