

SPECIALIST MATHEMATICS UNIT 4
SAC 3: INTEGRAL CALCULUS AND APPLICATIONS TEST

NAME: _____

PAPER TWO: Technology Active

Time: 25 Minutes Total = 18 marks

SECTION A: MULTIPLE CHOICE

Please circle the correct answer.

Question 1

An antiderivative of $\frac{1}{x^2 - 2x + 2}$ is:

- A. $-(x^2 - 2x + 2)^{-2}$
- B. $\log_e(x^2 - 2x + 2)$
- C. $\log_e \left| \frac{x-2}{x+1} \right|$
- D. $\operatorname{arcsec}(x-1)$
- E. $\arctan(x-1)$

Question 2

A solid is constructed by rotating the function $y = 1 - \cos(2x)$, where $0 \leq x \leq \frac{\pi}{2}$, about the y-axis. The volume of this solid is:

- A. $\frac{\pi(\pi^2 - 4)}{4}$
- B. $\frac{-\pi(\pi^2 - 20)}{4}$
- C. $\frac{\pi^3}{2}$
- D. $\frac{-\pi(\pi^2 - 4\pi - 4)}{4}$
- E. $\frac{\pi(\pi^2 + 4)}{4}$

Question 3

The region enclosed by the graph of $y = x^2 + 1$ and the lines $y = 1$ and $y = 4$ is rotated about the y -axis to form a solid of revolution. The volume of the solid is given by

- A. $\pi \int_0^{\sqrt{3}} (x^2 + 1) dx$
- B. $\pi \int_1^4 (y-1) dx$
- C. $\pi \int_1^4 (x^2 + 1) dx$
- D. $\pi \int_1^4 (y-1) dy$
- E. $\pi \int_0^{\sqrt{3}} (y-1) dy$

Question 4

Using a suitable substitution, $\int_{-\frac{\pi}{6}}^{\frac{\pi}{3}} (\tan(x) \log_e(\sec(x))) dx$ can be expressed completely in terms of u as:

- A. $\int_{\frac{2}{\sqrt{3}}}^2 (\log_e(u)) du$
- B. $\int_{-\log\left(\frac{\sqrt{3}}{2}\right)}^{\log_e(2)} (u) du$
- C. $-\int_{-\frac{\sqrt{3}}{2}}^{\frac{1}{2}} (u) du$
- D. $\int_{-\frac{\pi}{6}}^{\frac{\pi}{3}} (\log_e(u)) du$
- E. $\int_{-\frac{\pi}{6}}^{\frac{\pi}{3}} (u) du$

Question 5

If the substitutions $u = \frac{x}{2}$ is made, the integral $\int_2^4 \frac{1 - \left(\frac{x}{2}\right)^2}{x} dx$ becomes:

A. $\int_1^2 \frac{1-u^2}{u} du$

B. $\int_2^4 \frac{1-u^2}{u} du$

C. $\int_1^2 \frac{1-u^2}{2u} du$

D. $\int_1^2 \frac{1-u^2}{4u} du$

E. $\int_2^4 \frac{1-u^2}{2u} du$

SECTION B: SHORT ANSWER/ANALYSIS**Question 6 (7 marks)**

A wine glass is formed by rotating, around the y -axis, the graph defined by function

$$f : [0, 2] \rightarrow \mathbb{R}, f(x) = \frac{1}{10}(2 + 5x^3). \text{ All measurements are in cm.}$$

- a) Sketch the graph of $f(x)$ clearly labelling coordinates of endpoints.

2 marks

- b) State a definite integral that would find the volume of the glass formed, when full, after it is rotated around the y -axis.

2 marks

- c) Evaluate this volume, in cubic centimetres.

1 mark


The curve $f : [0, a] \rightarrow \mathbb{R}, f(x) = \frac{1}{10}(2 + 5x^3)$ is rotated about the x -axis now and the volume of the solid obtained in this way is equal by $\frac{22\pi}{175}$ cubic centimetres.

- d) Find the value of a .

2 marks

Question 7 (6 marks)

Let f be a function defined by $f(x) = x + 2 \cos x$, $x \in [0, 2\pi]$. The diagram below shows a region S bound by the graph of f and the line $y = x$.



A and C are the points of intersection of the line $y = x$ and the graph of f , and B is the minimum point of f .

- a) If A , B and C have x -coordinates $\frac{a\pi}{2}$, $\frac{b\pi}{2}$ and $\frac{c\pi}{2}$, where $a, b, c \in \mathbb{Z}^+$, find the values of a , b and c .

3 marks

- b) Write down a definite integral which would find the area of region S .

2 marks

- c) Hence find the area of the region S .

1 mark

END OF PAPER TWO