

Year 2017

VCE

Specialist Mathematics

Trial Examination 2

Solutions



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9018 5376
FAX: (03) 9817 4334
kilbaha@gmail.com
<http://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from **Copyright Agency Limited**. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

Reproduction and communication for educational purposes The Australian Copyright Act 1968 (the Act) allows a maximum of one chapter or 10% of the pages of this work, to be reproduced and/or communicated by any educational institution for its educational purposes provided that educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact
CAL, Level 15, 233 Castlereagh Street, Sydney, NSW, 2000

Tel: (02) 9394 7600

Fax: (02) 9394 7601

Email: info@copyright.com.au

Web: <http://www.copyright.com.au>

- While every care has been taken, no guarantee is given that these answers are free from error. Please contact us if you believe you have found an error.

SECTION 1

ANSWERS

1	A	B	C	D	E
2	A	B	C	D	E
3	A	B	C	D	E
4	A	B	C	D	E
5	A	B	C	D	E
6	A	B	C	D	E
7	A	B	C	D	E
8	A	B	C	D	E
9	A	B	C	D	E
10	A	B	C	D	E
11	A	B	C	D	E
12	A	B	C	D	E
13	A	B	C	D	E
14	A	B	C	D	E
15	A	B	C	D	E
16	A	B	C	D	E
17	A	B	C	D	E
18	A	B	C	D	E
19	A	B	C	D	E
20	A	B	C	D	E

SECTION A



Question 1

Answer E

$$f(x) = \frac{x^3 + ax^2 - ax + x - a^2}{x^2 - a} = x + a + \frac{x}{x^2 - a}$$

vertical asymptotes at $x = \sqrt{a}$ and $x = -\sqrt{a}$

and an oblique asymptote at $y = x + a$




Question 2

Answer B

The domain of $f(x) = \frac{2}{\pi} \tan^{-1}\left(\frac{2x+a}{a}\right) + a$

is R and the range is

$$\frac{2}{\pi} \times \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) + a = (-1 + a, 1 + a)$$



Question 3

Answer C


Since the coefficients are real, by the conjugate root theorem, the roots occur in conjugate pairs, the roots are $z = a$, $z = a \pm ai$ and $z = a \pm i$, there are 5 roots, the minimum degree of the polynomial is 5.

Question 4 **Answer E**

$$\frac{x^2}{x^3 - a^3} = \frac{x^2}{(x-a)(x^2 + ax + a^2)}$$

since we have the non-linear factor,
the partial fractions are given by

$$\frac{A}{x-a} + \frac{Bx+C}{x^2 + ax + a^2}$$



Question 5 **Answer B**

$$\underline{r}(t) = a \sec(t) \underline{i} + b \tan^2(t) \underline{j}$$

The parametric equations are
 $x = a \sec(t)$ and $y = b \tan^2(t)$.


$$\sec(t) = \frac{x}{a}, \quad \tan^2(t) = \frac{y}{b}$$

$$1 + \tan^2(t) = \sec^2(t)$$

$$1 + \frac{y}{b} = \frac{x^2}{a^2}$$

$$y = b \left(\frac{x^2}{a^2} - 1 \right)$$

The particle moves on part of a parabola.



Question 6 **Answer E**

Given $\underline{a} = -\underline{i} + y \underline{j} - 3 \underline{k}$ and $\underline{b} = 2 \underline{i} - 4 \underline{j} + 6 \underline{k}$

- A. Is true, when $y = 1$ (or any value) the vectors \underline{a} and \underline{b} are coplanar.
- B. Is true, when $y = 2$, then $-2 \underline{a} = \underline{b}$ so the vectors \underline{a} and \underline{b} are parallel.
 $\underline{a} \cdot \underline{b} = -2 - 4y - 18 = -4y - 20$
- C. Is true, if $y > -5$, $\underline{a} \cdot \underline{b} < 0$ therefore the angle between the vectors \underline{a} and \underline{b} is obtuse.
- D. Is true, if $y < -5$, $\underline{a} \cdot \underline{b} > 0$ therefore the angle between the vectors \underline{a} and \underline{b} is acute.
- E. $|\underline{a}| = \sqrt{1 + y^2 + 9} = \sqrt{10 + y^2}$ and $|\underline{b}| = \sqrt{4 + 16 + 36} = \sqrt{56}$ when $y = 8$,
 $|\underline{a}| = \sqrt{10 + 8^2} = \sqrt{74}$, the vectors \underline{a} and \underline{b} are not equal in length, **E.** is false

Question 7**Answer C**

$$z = \sqrt{\sqrt{2} + 2} + \sqrt{2 - \sqrt{2}}i = 2 \operatorname{cis}\left(\frac{\pi}{8}\right)$$

$$\bar{z} = 2 \operatorname{cis}\left(-\frac{\pi}{8}\right)$$

$$\frac{1}{\bar{z}} = \frac{1}{2} \operatorname{cis}\left(\frac{\pi}{8}\right)$$

$$\operatorname{Arg}\left(\frac{1}{\bar{z}^{10}}\right) = 10 \times \frac{\pi}{8} - 2\pi = -\frac{3\pi}{4}$$

Question 8**Answer C**

One of the roots of $z^4 - a^4i = 0$ is

$$z = \frac{a}{2} \left(\sqrt{\sqrt{2} + 2} + \sqrt{2 - \sqrt{2}}i \right) = a \operatorname{cis}\left(\frac{\pi}{8}\right)$$

From question 7. In total, there are 4 roots, they all lie on a circle of radius a , and are

equally spaced around the circle by $\frac{\pi}{2}$.

The roots do not occur in conjugate pairs, Option C. is correct.

Question 9**Answer E**

Since \underline{u} is a unit vector, $|\underline{u}| = 1$, now $\underline{u} \cdot \underline{v} = \sqrt{3} = |\underline{u}||\underline{v}|\cos(\theta) = 1 \times 2 \cos(\theta)$

$\cos(\theta) = \frac{\sqrt{3}}{2} \Rightarrow \theta = 30^\circ$, Amanda is correct the angle between the vectors \underline{u} and \underline{v} is 30° .

The scalar resolute of \underline{u} in the direction of \underline{v} is equal to $\underline{u} \cdot \hat{\underline{v}} = \frac{\underline{u} \cdot \underline{v}}{|\underline{v}|} = \frac{\sqrt{3}}{2}$.

Brianna is correct.

The scalar resolute of \underline{v} in the direction of \underline{u} is equal to $\underline{v} \cdot \hat{\underline{u}} = \frac{\underline{v} \cdot \underline{u}}{|\underline{u}|} = \frac{\sqrt{3}}{1} = \sqrt{3}$.

Colin is correct.

$$|\underline{u} + \underline{v}|^2 = (\underline{u} + \underline{v}) \cdot (\underline{u} + \underline{v}) = \underline{u} \cdot \underline{u} + 2\underline{u} \cdot \underline{v} + \underline{v} \cdot \underline{v} = |\underline{u}|^2 + 2\underline{u} \cdot \underline{v} + |\underline{v}|^2 = 1 + 2\sqrt{3} + 4$$

$$|\underline{u} + \underline{v}| = \sqrt{5 + 2\sqrt{3}}. \text{ Dianne is correct.}$$

$$|\underline{u} - \underline{v}|^2 = (\underline{u} - \underline{v}) \cdot (\underline{u} - \underline{v}) = \underline{u} \cdot \underline{u} - 2\underline{u} \cdot \underline{v} + \underline{v} \cdot \underline{v} = |\underline{u}|^2 - 2\underline{u} \cdot \underline{v} + |\underline{v}|^2 = 1 - 2\sqrt{3} + 4$$

$$|\underline{u} - \underline{v}| = \sqrt{5 - 2\sqrt{3}}. \text{ Edward is correct. So all are correct.}$$

Question 10 **Answer D**

$$\frac{dy}{dx} = f(x) = \sin^3(2x) \quad y_0 = 2 \quad x_0 = 0 \quad h = \frac{\pi}{8},$$

using Euler's Method


$$y_1 = y_0 + hf(x_0) = 2 + \frac{\pi}{8} \sin^3(0) = 2 \quad \text{and} \quad x_1 = \frac{\pi}{8}$$

$$y_2 = y_1 + hf(x_1)$$

$$= 2 + \frac{\pi}{8} \sin^3\left(\frac{\pi}{4}\right) = 2 + \frac{\pi}{8} \times \frac{\sqrt{2}}{4} \quad \text{and} \quad x_2 = \frac{\pi}{4}$$

$$y_3 = y_2 + hf(x_2)$$

$$= 2 + \frac{\pi}{8} \times \frac{\sqrt{2}}{4} + \frac{\pi}{8} \sin^3\left(\frac{\pi}{2}\right) = 2 + \frac{\pi}{8} \left(\frac{\sqrt{2}}{4} + 1\right)$$



Question 11 **Answer A**


$$\dot{r}(t) = t \cos(t) \underline{i} + t \sin(t) \underline{j}$$

$$r(t) = \int t \cos(t) dt \underline{i} + \int t \sin(t) dt \underline{j}$$

$$= (\cos(t) + t \sin(t)) \underline{i} + (\sin(t) - t \cos(t)) \underline{j} + \underline{c}$$

$$r(0) = \underline{0} = \underline{i} + \underline{c} \Rightarrow \underline{c} = -\underline{i}$$

$$r(t) = (\cos(t) + t \sin(t) - 1) \underline{i} + (\sin(t) - t \cos(t)) \underline{j}$$



Question 12 **Answer A**

$$p(t) = (2t - 2) \underline{i} + (t^2 - 6t + 9) \underline{j}$$

$$q(t) = (3t - 4) \underline{i} + (t^2 - 4t + 5) \underline{j}$$


$$\underline{i}: (1) \quad 2t - 2 = 3t - 4 \Rightarrow t = 2$$

$$\underline{j}: (2) \quad t^2 - 6t + 9 = t^2 - 4t + 5 \Rightarrow t = 2$$

$$p(2) = q(2) = 2 \underline{i} + \underline{j}$$


both particles move on parabolic paths

and P and Q are in the same position when $t = 2$.



Question 13

Answer D



resolving up parallel to plane around the m_1 kg mass (1) $T - m_1g \sin(\theta) = m_1a$

resolving downwards around the m_2 kg mass (2) $m_2g - T = m_2a$

adding to eliminate the tension in the string, to find the acceleration a , of the system

$$(1)+(2) \quad m_2g - m_1g \sin(\theta) = m_1a + m_2a \Rightarrow a = \frac{g(m_2 - m_1 \sin(\theta))}{m_1 + m_2}$$

$a > 0$ when $m_2 > m_1 \sin(\theta)$ and $a = 0$ when $m_2 = m_1 \sin(\theta)$

B. and **C.** are correct.

$$(1) \times m_2 \quad Tm_2 - m_1m_2g \sin(\theta) = m_1m_2a$$

$$(2) \times m_1 \quad m_2m_1g - Tm_1 = m_1m_2a \text{ subtracting to eliminate the acceleration, gives}$$

the tension in the string is equal to $\frac{m_1m_2g(1 + \sin(\theta))}{m_1 + m_2}$ newtons or $\frac{m_1m_2(1 + \sin(\theta))}{m_1 + m_2}$ kg-wt.

A. is correct.

When $m_2 = 2m_1$ and $\theta = 30^\circ$, $T = \frac{m_1 \times 2m_1g \left(1 + \frac{1}{2}\right)}{m_1 + 2m_1} = m_1g$ newtons. **D.** is incorrect

When $m_2 = 2m_1$ and $\theta = 30^\circ$, $a = \frac{g \left(2m_1 - \frac{m_1}{2}\right)}{m_1 + 2m_1} = \frac{g}{2} \text{ ms}^{-2}$ **E.** is correct

SECTION B

Question 1

a. solving $x(t) = 8 = t - \sin\left(\frac{\pi t}{2}\right)$

$$7 - \sin\left(\frac{7\pi}{2}\right) = 7 - (-1) = 8 \quad \text{A1}$$

b. $y(0) = 1$, $y(7) = 3$

initial height when released was 1 metre above the ground, and hits the wall 3 metres above the ground. A1

c. $\frac{dx}{dt} = \dot{x} = 1 - \frac{\pi}{2} \cos\left(\frac{\pi t}{2}\right)$, $\frac{dy}{dt} = \dot{y} = \pi \sin\left(\frac{\pi t}{2}\right)$ M1

The drone is flying horizontally, when $\dot{y} = 0$ but $\dot{x} \neq 0$, so when

$$\sin\left(\frac{\pi t}{2}\right) = 0 \Rightarrow \frac{\pi t}{2} = \pi, 2\pi, 3\pi$$

$$t = 2, 4, 6 \text{ seconds.} \quad \text{A1}$$


$$x(2) = 2, y(2) = 5, x(4) = 4, y(4) = 1, x(6) = 6, y(6) = 5$$

flying horizontally when $t = 2$ at $(2, 5)$, $t = 4$ at $(4, 1)$, $t = 6$ at $(6, 5)$ A1

d.

t	0	1	2	3	4	5	6	7
x	0	0	2	4	4	4	6	8
y	1	3	5	3	1	3	5	3

G2



e. $v(t) = \left(1 - \frac{\pi}{2} \cos\left(\frac{\pi t}{2}\right)\right) \underline{i} + \pi \sin\left(\frac{\pi t}{2}\right) \underline{j}$ velocity vector, A1

when it hits the wall $v(7) = \left(1 - \frac{\pi}{2} \cos\left(\frac{7\pi}{2}\right)\right) \underline{i} + \pi \sin\left(\frac{7\pi}{2}\right) \underline{j} = \underline{i} - \pi \underline{j}$

speed when it hits the wall $|v(7)| = \sqrt{1 + \pi^2}$ m/s A1

f. Angle at which it hits the wall $\tan(\theta) = \frac{1}{\pi}$

$\theta = \tan^{-1}\left(\frac{1}{\pi}\right) \approx 17.7^\circ$ A1

g.i. $s = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

$s = \int_0^7 \sqrt{\left(1 - \frac{\pi}{2} \cos\left(\frac{\pi t}{2}\right)\right)^2 + \pi^2 \sin^2\left(\frac{\pi t}{2}\right)} dt$ A1

ii. $s = 18.33$ metres. A1

Define $x1(t) = t - \sin\left(\frac{\pi t}{2}\right)$	Done
Define $y1(t) = 3 - 2 \cdot \cos\left(\frac{\pi t}{2}\right)$	Done
Define $r(t) = [x1(t) \ y1(t)]$	Done
$r(2)$	[2 5]
$r(4)$	[4 1]
$r(6)$	[6 5]
$r(7)$	[8 3]
Define $v(t) = \frac{d}{dt}(r(t))$	Done
$v(t)$	$\begin{bmatrix} \pi \cos\left(\frac{\pi t}{2}\right) \\ 1 - \frac{\pi}{2} \quad \pi \sin\left(\frac{\pi t}{2}\right) \end{bmatrix}$
$v(7)$	[1 -π]
$\frac{\tan^{-1}\left(\frac{1}{\pi}\right) \cdot 180}{\pi}$	17.7
$\int_0^7 \text{norm}(v(t)) dt$	18.33

Question 2

a. $f(x) = 10 \sec\left(\frac{\pi x}{n}\right) - 10$

$$f(10) = 10 \Rightarrow 10 = 10 \sec\left(\frac{10\pi}{n}\right) - 10$$

$$\sec\left(\frac{10\pi}{n}\right) = 2$$

$$\cos\left(\frac{10\pi}{n}\right) = \frac{1}{2}$$

M1

$$\frac{10\pi}{n} = \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3}$$

$$n = 30$$

b. $g(x)$ is the inverse function, swap x and y

$$g: x = 10 \sec\left(\frac{\pi y}{30}\right) - 10$$

$$\sec\left(\frac{\pi y}{30}\right) = \frac{x+10}{10}$$

$$\cos\left(\frac{\pi y}{30}\right) = \frac{10}{x+10}$$

M1

$$\frac{\pi y}{30} = \cos^{-1}\left(\frac{10}{x+10}\right)$$

$$y = g(x) = f^{-1}(x) = \frac{30}{\pi} \cos^{-1}\left(\frac{10}{x+10}\right) \Rightarrow k = 30, a = 10$$

c.i. $A = \int_0^{10} \left(\frac{30}{\pi} \cos^{-1}\left(\frac{10}{x+10}\right) - 10 \sec\left(\frac{\pi x}{30}\right) + 10 \right) dx$

or $A = 2 \int_0^{10} \left(\frac{30}{\pi} \cos^{-1}\left(\frac{10}{x+10}\right) - x \right) dx$

A1

or $A = 2 \int_0^{10} \left(x - 10 \sec\left(\frac{\pi x}{30}\right) + 10 \right) dx$

ii. $A = \frac{300}{\pi} (\pi - 2 \log_e(2 + \sqrt{3})) \approx 48.480 \text{ cm}^2$

A1

d. $y = \sec(kx) \Rightarrow \frac{dy}{dx} = k \tan(kx) \sec(kx)$
 $f(x) = 10 \sec\left(\frac{\pi x}{30}\right) - 10 \Rightarrow f'(x) = 10 \times \frac{\pi}{30} \tan\left(\frac{\pi x}{30}\right) \sec\left(\frac{\pi x}{30}\right)$
 $f'(x) = \frac{\pi}{3} \tan\left(\frac{\pi x}{30}\right) \sec\left(\frac{\pi x}{30}\right)$ A1

e. $y = \frac{30}{\pi} \cos^{-1}\left(\frac{10}{x+10}\right) = \frac{30}{\pi} \cos^{-1}(u)$, $u = \frac{10}{x+10} = 10(x+10)^{-1}$
 $\frac{dy}{du} = \frac{-30}{\pi \sqrt{1-u^2}}$ $\frac{du}{dx} = -10(x+10)^{-2}$
 $g'(x) = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{-30}{\pi \sqrt{1-\left(\frac{10}{x+10}\right)^2}} \times \frac{-10}{(x+10)^2}$ M1
 $= \frac{300}{\pi(x+10)^2} \times \frac{1}{\sqrt{\frac{(x+10)^2 - 10^2}{(x+10)^2}}}$
 $= \frac{300}{\pi|x+10|\sqrt{x(x+20)}}$ A1

f.i. $s = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ the arc length
 $s = \int_0^{10} \sqrt{1 + \frac{\pi^2}{9} \tan^2\left(\frac{\pi x}{30}\right) \sec^2\left(\frac{\pi x}{30}\right)} dx + \int_0^{10} \sqrt{1 + \frac{90000}{\pi^2(x+10)^2 x(x+20)}} dx$
 or $s = 2 \int_0^{10} \sqrt{1 + \frac{\pi^2}{9} \tan^2\left(\frac{\pi x}{30}\right) \sec^2\left(\frac{\pi x}{30}\right)} dx$ A1
 or $s = 2 \int_0^{10} \sqrt{1 + \frac{90000}{\pi^2(x+10)^2 x(x+20)}} dx$

ii. $s = 30.674$ cm A1

Define $f1(x) = 10 \cdot \sec\left(\frac{\pi \cdot x}{30}\right) - 10$	Done
Define $f2(x) = \frac{30}{\pi} \cdot \cos^{-1}\left(\frac{10}{x+10}\right)$	Done
$f1(10)$	10
$f2(10)$	10
$2 \cdot \int_0^{10} (x - f1(x)) dx$	48.480
$2 \cdot \int_0^{10} (f2(x) - x) dx$	48.480
$\int_0^{10} (f2(x) - f1(x)) dx$	48.480
$\frac{d}{dx}(f1(x))$	$\frac{\pi \cdot \sin\left(\frac{\pi \cdot x}{30}\right)}{3 \cdot \left(\cos\left(\frac{\pi \cdot x}{30}\right)\right)^2}$
$\Delta \frac{d}{dx}(f2(x))$	$\frac{300}{\pi \cdot \sqrt{x \cdot (x+20)} \cdot x+10 }$
$2 \cdot \int_0^{10} \sqrt{1 + \left(\frac{d}{dx}(f1(x))\right)^2} dx$	30.674
$\Delta 2 \cdot \int_0^{10} \sqrt{1 + \left(\frac{d}{dx}(f2(x))\right)^2} dx$	30.674
$\Delta \int_0^{10} \sqrt{1 + \left(\frac{d}{dx}(f1(x))\right)^2} dx + \int_0^{10} \sqrt{1 + \left(\frac{d}{dx}(f2(x))\right)^2} dx$	30.674

Question 3

a. $U(\sqrt{2}-1, \sqrt{2}), V(-\sqrt{2}-1, \sqrt{2}), C(-1, 0)$
 $\overline{CU} = \overline{OU} - \overline{OC} = \sqrt{2}\underline{i} + \sqrt{2}\underline{j}, |\overline{CU}| = 2$
 $\overline{CV} = \overline{OV} - \overline{OC} = -\sqrt{2}\underline{i} + \sqrt{2}\underline{j}, |\overline{CV}| = 2$ A1
 $\overline{CU} \cdot \overline{CV} = -2 + 2 = 0$
 angle between \overline{CU} and \overline{CV} is 90° (or $\frac{\pi}{2}$) A1


b. $u = \sqrt{2}-1 + \sqrt{2}i$
 $|u| = \sqrt{(\sqrt{2}-1)^2 + (\sqrt{2})^2} = \sqrt{2-2\sqrt{2}+1+2} = \sqrt{5-2\sqrt{2}}$
 $b=5, a=2$ A1
 $\text{Arg}(u) = \tan^{-1}\left(\frac{\sqrt{2}}{\sqrt{2}-1}\right) = \tan^{-1}\left(\frac{\sqrt{2}}{\sqrt{2}-1} \times \frac{\sqrt{2}+1}{\sqrt{2}+1}\right) = \tan^{-1}(2+\sqrt{2})$ A1

c. $S = \{z : |z+1| = 2, z \in C\}, z = x + yi$
 $|(x+1) + yi| = 2$
 $\sqrt{(x+1)^2 + y^2} = 2$
 $(x+1)^2 + y^2 = 4$
 circle centre $(-1, 0)$ radius 2 A1

d. $R = \{z : \text{Arg}(z+1) = \frac{3\pi}{4}, z \in C\}, z = x + yi$
 $\text{Arg}(x+1 + yi) = \frac{3\pi}{4}$
 $\tan\left(\frac{3\pi}{4}\right) = -1 = \frac{y}{x+1}$
 ray $y = -(x+1)$ for $x < -1$ A1

e. $T = \{z : |z| = |z+1-i|, z \in C\}, z = x + yi$
 $|x + yi| = |(x+1) + (y-1)i|$
 $\sqrt{x^2 + y^2} = \sqrt{(x+1)^2 + (y-1)^2}$
 $x^2 + y^2 = x^2 + 2x + 1 + y^2 - 2y + 1$
 line $y = x + 1$ A1

- f.** solving $(x+1)^2 + y^2 = 4$ and $y = -(x+1)$ with $x < -1$
 $2y^2 = 4$
 $y^2 = 2$
 $y = \sqrt{2}$ $x = -\sqrt{2} - 1$
 $(-\sqrt{2} - 1, \sqrt{2})$ A1
- g.** solving $(x+1)^2 + y^2 = 4$ and $y = x+1$
 $2y^2 = 4$
 $y^2 = 2$
 $y = \pm\sqrt{2}$ $x = \sqrt{2} - 1$, $-\sqrt{2} - 1$
 $(\sqrt{2} - 1, \sqrt{2})$, $(-\sqrt{2} - 1, -\sqrt{2})$ A1
- h.** open circle at the point $(-1,0)$ as the point is not included, for R . G3



$u := \sqrt{2} - 1 + \sqrt{2} \cdot i$	$\sqrt{2} - 1 + \sqrt{2} \cdot i$
$ u $	$\sqrt{5 - 2 \cdot \sqrt{2}}$
$\text{angle}(u)$	$\tan^{-1}(\sqrt{2} + 2)$
$z := x + y \cdot i$	$x + y \cdot i$
$s := z + 1 = 2$	$\sqrt{x^2 + 2 \cdot x + y^2 + 1} = 2$
$(\sqrt{x^2 + 2 \cdot x + y^2 + 1} = 2)^2$	$x^2 + 2 \cdot x + y^2 + 1 = 4$
$\text{completeSquare}(x^2 + 2 \cdot x + y^2 + 1 = 4, \{x, y\})$	$(x + 1)^2 + y^2 = 4$
$r := \text{angle}(z + 1) - \frac{3 \cdot \pi}{4}$	$\tan^{-1}\left(\frac{x + 1}{y}\right) + \frac{\pi \cdot \text{sign}(y)}{2} - \frac{3 \cdot \pi}{4}$
$\text{solve}(r = 0, y)$	$y = -(x + 1) \text{ and } x + 1 < 0$
$t := z - z + 1 - i $	$\sqrt{x^2 + y^2} - \sqrt{x^2 + 2 \cdot x + y^2 - 2 \cdot y + 2}$
$\text{solve}(t = 0, y)$	$y = x + 1$
$\text{solve}((x + 1)^2 + y^2 = 4 \text{ and } y = -(x + 1), \{x, y\}) x < -1$	$x = -(\sqrt{2} + 1) \text{ and } y = \sqrt{2}$
$\text{solve}((x + 1)^2 + y^2 = 4 \text{ and } y = x + 1, \{x, y\})$	$x = -(\sqrt{2} + 1) \text{ and } y = -\sqrt{2} \text{ or } x = \sqrt{2} - 1 \text{ and } y = \sqrt{2}$

Question 4

a. $V = \pi \int_a^b x^2 dy \quad y = \frac{1}{2}(x^2 - 3) \Rightarrow x^2 = 2y + 3$

$$V = \pi \int_0^9 (2y + 3) dy = \pi [y^2 + 3y]_0^9 = \pi(81 + 27 - 0)$$

$$V = 108\pi \text{ cm}^3 \tag{A1}$$

b. $V(h) = \pi \int_0^h (2y + 3) dy = \pi [y^2 + 3y]_0^h$

$$V(h) = \pi(h^2 + 3h) \tag{A1}$$

c. $\frac{dV}{dt} = -3\sqrt{h} \quad , \quad \frac{dV}{dh} = \pi(2h + 3) \tag{A1}$

$$\frac{dt}{dh} = \frac{dt}{dV} \frac{dV}{dh} = \frac{\pi(2h + 3)}{-3\sqrt{h}}$$

$$t = -\frac{\pi}{3} \int_9^0 \left(\frac{2h + 3}{\sqrt{h}} \right) dh = \frac{\pi}{3} \int_0^9 \left(2h^{\frac{1}{2}} + 3h^{-\frac{1}{2}} \right) dh \tag{M1}$$

$$t = \frac{\pi}{3} \left[\frac{4}{3} h^{\frac{3}{2}} + 6h^{\frac{1}{2}} \right]_0^9 = \frac{\pi}{3} \left(\frac{4}{3} \times \sqrt{9^3} + 6\sqrt{9} - 0 \right)$$

$$t = 18\pi \text{ sec} \tag{A1}$$

d. let the circle have centre at $(0, a)$, so $x^2 + (y - a)^2 = 16$
 at the point of contact, gradients are equal, using implicit differentiation
 $2x + 2(y - a) \frac{dy}{dx} = 0$, but $y = \frac{1}{2}(x^2 - 3) \Rightarrow \frac{dy}{dx} = x$ M1
 $2x + 2(y - a)x = 0 \Rightarrow 2x[1 + y - a] = 0 \Rightarrow y - a = -1$
 substitute into $x^2 + (y - a)^2 = 16 \Rightarrow x^2 + 1 = 16 \Rightarrow x = \pm\sqrt{15}$
 substitute into $y = \frac{1}{2}(x^2 - 3) \Rightarrow y = \frac{1}{2}(15 - 3) = 6 \Rightarrow a = 7$ A1
 point of contact $(\pm\sqrt{15}, 6)$ A1

e. since the radius of the ice block is 4, the bottom of the ice block is at $y = 3$
 $V = V(6) - \pi \int_3^6 (16 - (y - 7)^2) dy$ A1
 $V = V(6) - \left[16y - \frac{1}{3}(y - 7)^3 \right]_3^6$
 $V = \pi(36 + 18) - \pi \left[(16 \times 6) - \frac{1}{3} \times (-1)^3 - (16 \times 3) + \frac{1}{3} \times (-4)^3 \right]$
 $V = 27\pi \text{ cm}^3$ A1

$\pi \int_0^9 (2 \cdot y + 3) dy$	108 · π
$\pi \int_0^h (2 \cdot y + 3) dy$	h · (h+3) · π
Define $v(h) = \pi \int_0^h (2 \cdot y + 3) dy$	Done
$\pi \int_0^9 \frac{2 \cdot h + 3}{3 \cdot \sqrt{h}} dh$	18 · π
$v(6) - \pi \int_3^6 (16 - (y - 7)^2) dy$	27 · π

Question 5


a. $m = 4.5 \text{ kg}$ $R = kv^2$ $k = 0.225$

$$m\ddot{x} = -(mg + kv^2)$$

$$4.5\ddot{x} = -(4.5 \times 9.8 + 0.225v^2)$$

$$\ddot{x} = -\left(9.8 + \frac{0.225v^2}{4.5}\right) = -\left(9.8 + \frac{v^2}{20}\right) = -\left(\frac{9.8 \times 20 + v^2}{20}\right)$$

$$\ddot{x} = -\frac{(196 + v^2)}{20}$$



M1

b.i. Use $\ddot{x} = \frac{dv}{dt} = -\frac{(196 + v^2)}{20}$, $v(0) = 3.5$

inverting $\frac{dt}{dv} = -\frac{20}{(196 + v^2)}$

$$t = \int \frac{-20}{196 + v^2} dv$$

A1

$$t = -\frac{20}{\sqrt{196}} \tan^{-1}\left(\frac{v}{\sqrt{196}}\right) + c$$

$$t = -\frac{10}{7} \tan^{-1}\left(\frac{v}{14}\right) + c$$

A1

to find c use $v = 3.5$ when $t = 0$

$$0 = -\frac{10}{7} \tan^{-1}\left(\frac{3.5}{14}\right) + c \Rightarrow c = \frac{10}{7} \tan^{-1}\left(\frac{1}{4}\right)$$

$$t = \frac{10}{7} \left(\tan^{-1}\left(\frac{1}{4}\right) - \tan^{-1}\left(\frac{v}{14}\right) \right)$$

$$\frac{7t}{10} = \tan^{-1}\left(\frac{1}{4}\right) - \tan^{-1}\left(\frac{v}{14}\right)$$

A1

$$\tan^{-1}\left(\frac{v}{14}\right) = \tan^{-1}\left(\frac{1}{4}\right) - \frac{7t}{10}$$

$$v = 14 \tan\left(\tan^{-1}\left(\frac{1}{4}\right) - \frac{7t}{10}\right)$$

ii. When $v = 0$ $t = \frac{10}{7} \tan^{-1}\left(\frac{1}{4}\right) \approx 0.350$ seconds

A1

- c.i.** use $\ddot{x} = v \frac{dv}{dx} = -\frac{(196+v^2)}{20}$
- $$\frac{dv}{dx} = -\frac{(196+v^2)}{20v} \quad \text{M1}$$
- inverting $\frac{dx}{dv} = \frac{-20v}{196+v^2}$
- $$D = \int_{3.5}^0 \frac{-20v}{196+v^2} dv \quad \text{A1}$$
- alternatively $v = \frac{dx}{dt} = 14 \tan\left(\tan^{-1}\left(\frac{2}{7}\right) - \frac{7t}{10}\right)$ M1
- $$D = 14 \int_0^{0.350} \tan\left(\tan^{-1}\left(\frac{1}{4}\right) - \frac{7t}{10}\right) dt \quad \text{A1}$$
- ii.** $D = 0.606$ metres A1

$\text{deSolve}\left(v' = -\frac{(196+v^2)}{20} \text{ and } v(0) = \frac{7}{2}, t, v\right)$	$\frac{\tan^{-1}\left(\frac{v}{14}\right) - \tan^{-1}\left(\frac{1}{4}\right)}{14} = -\frac{t}{20}$
$\text{solve}\left(\frac{\tan^{-1}\left(\frac{v}{14}\right) - \tan^{-1}\left(\frac{1}{4}\right)}{14} = -\frac{t}{20}, v\right)$	$v = -14 \cdot \tan\left(\frac{7 \cdot t}{10} - \tan^{-1}\left(\frac{1}{4}\right)\right) \text{ and } 7 \cdot t - 10 \cdot \tan^{-1}\left(\frac{1}{4}\right) \geq -5 \cdot \pi \text{ and } 7 \cdot t - 10 \cdot \tan^{-1}\left(\frac{1}{4}\right) \leq 5 \cdot \pi$
$\int_{3.5}^0 \frac{-20 \cdot v}{196+v^2} dv$	0.606246
$14 \cdot \int_0^{0.35} \tan\left(\tan^{-1}\left(\frac{1}{4}\right) - \frac{7 \cdot t}{10}\right) dt$	0.606246
$d: 0.6062462136117$	0.606246

Question 6

a.i. $A \stackrel{d}{=} N(30, 5^2)$, $\bar{A} \stackrel{d}{=} N\left(30, \frac{5^2}{30}\right)$

$\Pr(\bar{A} < 28) = 0.0142$ A1

ii. $n = 30$, $\bar{x} = 30$ $s = 5$ 95% $z = 1.96$

$$\bar{x} \pm z \times \frac{s}{\sqrt{n}} = 30 \pm 1.96 \times \frac{5}{\sqrt{30}}$$

$(28.21, 31.79)$ A1

"Title"	"z Interval"
"CLower"	28.2108
"CUpper"	31.7892
"x̄"	30.
"ME"	1.78919
"n"	30.
"σ"	5.

b.i. $A \stackrel{d}{=} N(30, 5^2)$, $B \stackrel{d}{=} N(25, 4^2)$

$T = A - B$

$E(T) = E(A) - E(B) = 30 - 25 = 5$ M1

$\text{Var}(T) = \text{Var}(A) + \text{Var}(B) = 5^2 + 4^2 = 41$

$\Pr(T < 0) = 0.2174$ A1

ii. $\Pr\left(T < \frac{0 - (5 - m)}{\sqrt{41}}\right) = 0.9 \Rightarrow \frac{m - 5}{\sqrt{41}} = 1.282$ M1

$m = 13.21$ minutes A1

c.i. $H_0: \mu = 30$
 $H_1: \mu < 30$ one sided to test his mean time has decreased A1

ii. $\bar{x} = 28, \mu = 30, \sigma = 5, n = 30$

$$p = \Pr(\bar{X} < 28) = \Pr\left(Z < \frac{28-30}{\frac{5}{\sqrt{30}}}\right) = \Pr(Z < -2.1909)$$

$p = 0.0142$ A1

iii. since $p < 0.05$ there is evidence to support the alternative hypothesis H_1 ,
 yes it is a quicker route to get to school. A1

$\frac{28-30}{\frac{5}{\sqrt{30}}}$	-2.1909														
normCdf(-∞, -2.1909, 0, 1)	0.0142														
zTest 30, 5, 28, 30, -1: stat.results	<table border="1"> <tr> <td>"Title"</td> <td>"z Test"</td> </tr> <tr> <td>"Alternate Hyp"</td> <td>"$\mu < \mu_0$"</td> </tr> <tr> <td>"z"</td> <td>-2.1909</td> </tr> <tr> <td>"PVal"</td> <td>0.0142</td> </tr> <tr> <td>"\bar{x}"</td> <td>28.0000</td> </tr> <tr> <td>"n"</td> <td>30.0000</td> </tr> <tr> <td>"σ"</td> <td>5.0000</td> </tr> </table>	"Title"	"z Test"	"Alternate Hyp"	" $\mu < \mu_0$ "	"z"	-2.1909	"PVal"	0.0142	" \bar{x} "	28.0000	"n"	30.0000	" σ "	5.0000
"Title"	"z Test"														
"Alternate Hyp"	" $\mu < \mu_0$ "														
"z"	-2.1909														
"PVal"	0.0142														
" \bar{x} "	28.0000														
"n"	30.0000														
" σ "	5.0000														
invNorm(0.05, 0, 1)	-1.6449														

END OF SECTION B SUGGESTED ANSWERS