



**SUZANNE CORY**  
High School

Name : \_\_\_\_\_ Teacher : \_\_\_\_\_

## 2015 MATHEMATICAL METHODS (CAS) UNIT 4

### SAC 5 – Analysis Task

### Chapters 14, 15, 16, 17 & 18 - Probability

Reading time: 10 minutes

Writing time: 85 minutes

### QUESTION AND ANSWER BOOKLET

#### Structure of Booklet

<i>Section</i>	<i>Number of Questions</i>	<i>Number of questions to be answered</i>	<i>Number of Marks</i>
1	4	4	60

- Students are permitted to bring into the test room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, one bound reference (which will be collected for the duration of the SAC), one approved CAS calculator (memory DOES NOT need to be cleared) and, if desired, one scientific calculator. For approved computer-based CAS, their full functionality may be used.
- Students are NOT permitted to bring into the test room: blank sheets of paper and/or white out liquid/tape.

#### Materials supplied

- Question and answer booklet.

#### Instructions

- Write your **name** and **teacher** in the space provided above on this page.
- All written responses must be in English.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the test room.**

## Mathematical Methods (CAS) Formulas

### Mensuration

area of a trapezium:	$\frac{1}{2}(a+b)h$	volume of a pyramid:	$\frac{1}{3}Ah$
curved surface area of a cylinder:	$2\pi rh$	volume of a sphere:	$\frac{4}{3}\pi r^3$
volume of a cylinder:	$\pi r^2 h$	area of a triangle:	$\frac{1}{2}bc \sin A$
volume of a cone:	$\frac{1}{3}\pi r^2 h$		

### Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x  + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$	

product rule:  $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$       quotient rule:  $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

chain rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$       approximation:  $f(x+h) \approx f(x) + hf'(x)$

### Probability

$\Pr(A) = 1 - \Pr(A')$        $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$

$\Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)}$       transition matrices:  $S_n = T^n \times S_0$

mean:  $\mu = E(X)$       variance:  $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

Probability distribution		Mean	Variance
discrete	$\Pr(X=x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$

END OF FORMULA SHEET

### Instructions for Section 1

Answer all questions in the spaces provided.

In all questions where a numerical answer is required an **exact value** must be given unless otherwise specified.

In questions where more than 1 mark is available, appropriate working must be shown.

#### Question 1 (20 marks)

Canine Bio-Engineering is a bio-tech company that has bred a genetically modified Dalmation dog (*canis familiaris minima maculosa*) for spot-intolerant people who want a Dalmation dog but don't want it to have too many spots. The number of spots,  $X$ , on the modified Dalmation dog is a discrete random variable with probability distribution as shown in the following table:

$x$	2	3	4	5	6
$\Pr(X=x)$	0.3	0.4	$k$	0.05	0.02

- a. Show that  $k = 0.23$ . 1 mark

---

---

- b. For the number of spots on the modified Dalmation dog:

- i. find the mean 2 marks

---

---

---

- ii. find the median 1 mark

---

- iii. show that the variance is 0.9019 1 mark

---

---

---

- iv. find  $\text{Var}(3X - 2)$  2 marks

---

---

---

---

- v. find  $\Pr(\mu - 2\sigma \leq X \leq \mu + 2\sigma)$  **and** interpret your result. 3 marks

---

---

---

---

---

---

---

The large population of genetically modified dogs are kept in a secure open range enclosure.

- c. If two dogs are randomly chosen from the enclosure, find the probability that
- i. each dog has three spots, correct to 2 decimal places 1 mark

---

---

- ii. the total number of spots on both dogs is equal to seven 2 marks

---

---

---

---

---

---

---

- iii. one of the dogs has two spots, given that the total number of spots on both dogs is equal to seven, correct to 4 decimal places. 2 marks

---

---

---

---

---

---

---

---

---

---

**d.** Eight modified Dalmation dogs are randomly chosen from the enclosure.

**i.** Find, correct to 4 decimal places, the probability that a dog chosen at random from the eight dogs has two spots. 1 mark

---

---

**ii.** Find, correct to 4 decimal places, the probability that at least four of the dogs have two spots. 1 mark

---

---

**e.** A random group of genetically modified Dalmation dogs manage to escape from the enclosure. Find the smallest number of dogs in the group if the probability that at least two of the dogs in the group, having less than four spots, is greater than 0.99. 3 marks

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**THIS PAGE IS BLANK**

**Question 2** (10 marks)

It is found that when an **unmarried** couple go and see a movie together, there is a 58% chance that they will see a romantic comedy (RomCom) if the previous movie they saw together was a RomCom, and a 46% chance that they will see an action movie if the previous movie they saw together was an action movie.

- a. Create a transition matrix for the probabilities of an unmarried couple seeing either a RomCom or an action movie together, given the previous movie seen together. 1 mark

$$T = \begin{bmatrix} & \\ & \end{bmatrix}$$

It is found that when a **married** couple go and see a movie together there is a 92% chance that they will see a RomCom if the previous movie they saw together was an action movie, and a 5% chance that they will see an action movie if the previous movie they saw together was a RomCom. A transition matrix for the probabilities of a married couple seeing either a RomCom or an action movie together given the previous movie seen together is

$$T = \begin{bmatrix} 0.95 & 0.92 \\ 0.05 & 0.08 \end{bmatrix}$$

- b. John and Susan are a **married** couple who decide to go and see a movie together. It is equally likely that they will see either a RomCom or an action movie the first time that they go to the movies together. Find the probability that the second time they go to the movies together, John and Susan will see a RomCom 2 marks

---

---

---

---

---

---

---

---

c. As it happens, the first movie that John and Susan saw in 2015 was an action movie. Find:

i. the probability that the next three movies John and Susan see together are RomComs. 1 mark

---

---

ii. the probability that the fifth movie they see together in 2015 will be a RomCom, correct to 4 decimal places. 2 marks

---

---

---

---

---

---

iii. the steady state probability that they will see an action movie together (an exact value is required) 2 marks

---

---

---

---

---

---

On a particular day it was found that 84 married couples saw an action movie together and 212 married couples saw a RomCom together.

d. Find, correct to the nearest whole number, the number of married couples expected to see each type of movie in the long term. 2 marks

---

---

---

---

---

---

---

---



**Question 3** (14 marks)

A social researcher carried out a lifestyle survey on a group of people. A mathematical model was then developed from the result. The time, in hours, spent each week on sport and recreation by people surveyed is a continuous random variable,  $T$ , with probability density function

$$f(t) = \begin{cases} kt^2 e^{-\frac{t}{2}} & \text{for } 0 \leq t \leq 8 \\ -\frac{rt}{13e^4} + \frac{116}{13e^4} & \text{for } 8 < t \leq 14.5 \\ 0 & \text{elsewhere} \end{cases}$$

- a.** Given that  $f(t)$  is continuous at  $t = 8$ , show that  $k = \frac{1}{16}$  and  $r = 8$ . 2 marks

---



---



---



---



---



---




---



---

- b.** Sketch the probability density function  $f(t)$  on the axes below. Label any intercept(s) and turning point(s) using coordinates. 3 marks



- c.** Find the median time spent on sport and recreation by this group, in hours, correct to 2 decimal places. 2 marks

---



---



---

**d.** Find the modal time spent on sport and recreation by this group. 1 mark

---

---

**e.** A person is chosen at random from the group. Find the probability that the person:

**i.** spends at least 5 hours on sport and recreation, correct to four decimal places 1 mark

---

---

**ii.** spends less than 8 hours on sport and recreation, correct to four decimal places 1 mark

---

---

**iii.** spent less than 8 hours on sport and recreation, given that they spent at least 5 hours on sport and recreation, correct to 2 decimal places. 2 marks

---

---

---

---

---

The social researcher decides to interview 6 people from the group to see whether or not the people spend at least 5 hours on sport and recreation each week.

**f.** Find the probability, correct to two decimal places, that three people spend at least 5 hours on sport and recreation each week. 2 marks

---

---

---

**Question 4** (16 marks)

Records kept by Massachusetts Institute of Technology suggest that the distribution of the total fees paid by graduating students is normal, with mean \$60000 and standard deviation \$5000.

**a.** What proportion of the university's graduates pay:

**i.** less than \$55000, correct to 4 decimal places? 1 mark

---

---

**ii.** more than \$50000 but less than \$74000, correct to 4 decimal places? 1 mark

---

---

**b.** The university's advertising manager wishes to claim that '90% of our graduating students pay less than  $c$  in total fees. What should  $c$  be, correct to the nearest dollar? 1 mark

---

---

Jeremy is practising shots from the free-throw line on the university basketball court. If Jeremy makes a shot, the probability he will make the next shot is 0.6. If he misses a shot, the probability he will make the next shot is 0.3. Jeremy misses his first shot.

**c.** What is the probability Jeremy will make his next three shots? 1 mark

---

---

**d.** What is the probability Jeremy will make exactly two of his next three shots? 3 marks

---

---

---

---

---

---

---

---

---

---

- e. Given that Jeremy made at least one of the next three shots, what is the probability he made all three shots, correct to 2 decimal places? 2 marks

---

---

---

---

---

The ATAR scores of applicants for a particular university course are normally distributed. The university decided to accept the top 15% of applicants. Jeremy was accepted into the university with an ATAR score of 86. Only 10% of applicants had a better score than him. His friend Jody was offered a scholarship. Her ATAR score was 95 and only 1% of applicants had a better score than hers.

- f. What was the mean and standard deviation of the ATAR scores of applicants? Give your answers correct to two decimal places. 4 marks

---

---

---

---

---

---

---

---

Jody wants to join one of the university's sports clubs. The probability she will be selected for soccer and not rugby is  $p$ . The probability she will be selected for rugby is  $p^3$  and the probability she will be selected for neither sport is  $p^2$ . Being selected for soccer is independent of being selected for rugby.

- g. What is the probability Jody will be selected for both sports? Give your answer correct to two decimal places. 3 marks

---

---

---

---

---

---

---

---

**END OF QUESTION AND ANSWER BOOKLET**