



Scotch Student ID #				
Circle the relevant digits	0	0	0	0
	1	1	1	1
	2	2	2	2
	3	3	3	3
	4	4	4	4
	5	5	5	5
	6	6	6	6
	7	7	7	7
	8	8	8	8
	9	9	9	9

Scotch College

Teacher's Name

MATHEMATICAL METHODS

Unit 3-SAC 1b – Application Task: Test

Thursday 2nd June 2022

Reading Time	none
Writing Time	45 minutes

Task Sections	Marks	Your Marks
Extended Response Questions	30	
Total Marks	30	

Declaration

I declare that any work I have submitted for this VCE assessment is wholly my own, unless properly referenced or authorised for use by my teacher. I have had no assistance from any person in my home nor have I been assisted by, or given assistance to, a boy in my class or cohort unless specifically permitted to do so by my teacher. I have not used the internet or other sources to assist me in my responses unless specifically permitted by my teacher. I acknowledge my work may be reproduced, communicated, compared and archived for the purposes of detecting plagiarism and collusion.

Signature: _____

General Instructions

- Answer all questions in the spaces provided.
- In all questions where a numerical answer is required, an exact value must be given unless otherwise specified.
- In questions where more than one mark is available, appropriate working must be shown.
- Unless otherwise indicated, the diagrams in this task are not drawn to scale.

Allowed Materials

- Calculators are not allowed
- Notes and/or references are not allowed.

At the end of the task


- Ensure you cease writing upon request.

Electronic Devices

Students are **not** allowed to have a mobile phone, smart watch and/or any other unauthorised electronic device in the SAC, unless it is **TURNE**D OFF and is placed on the front teacher desk.

Question 1 (3 marks)

Sketch the graph of $y = \log_e(2x - 1)$. Label all axis intercepts with their co-ordinates and all asymptotes with their equations.



Question 2 (3 marks)

Solve each of the equations below.

a. $2 \times 3^x = 5$

1 mark

b. $\log_5(2) - \log_5(x) = 3$

2 marks

Question 3 (4 marks)

a. The graph given by the rule $y = a \times 10^x - 2$ passes through the point $(\log_{10}(3), 4)$.

Find the value of a .

2 marks

b. The graph given by the rule $y = 3 \times \log_2(x - b)$ passes through the point $\left(\frac{3}{2}, -3\right)$.

Find the value of b .

2 marks

Question 4 (5 marks)

Consider the polynomial $P(x) = 2x^3 + 3x^2 - 12x - 6$.

a. Find $P'(x)$.

1 mark

b. Hence, find the co-ordinates of all stationary points of the graph of $y = e^{2x^3 + 3x^2 - 12x - 6}$.

4 marks

Question 5 (6 marks)

Consider the function $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = \log_e(x^2)$.

- a.** Find the rule of the derivative, $f'(x)$. 1 mark

- b.** Show that the tangent to the graph of $y = f(x)$ at the point where $x = a$ is given by


$$y = \frac{2x}{a} + 2(\log_e(a) - 1). \quad \text{2 marks}$$

- c.** Find the value of a such that the tangent to the graph passes through the origin. 1 mark

- d.** Using the value of a you found in **part c**, find the equation of the normal to the graph of $y = f(x)$ at the point $x = a$. 2 marks

Question 6 (9 marks)

- a.** Sketch the graphs of $y_1 = x^2$, $x \geq 0$ and $y_2 = x^{\frac{1}{2}}$, $x \geq 0$, on the same axes below, labelling any axis intercepts, endpoints and points of intersection with their co-ordinates. 3 marks



- b.** For each of the graphs in **part a** find the co-ordinates of the point where the gradient of the tangent to the curve is one. 3 marks

- c. Find the value of k so that the graphs with equations $y = x^2 + k$ and $y = (x - k)^{\frac{1}{2}}$ have a common tangent at a single point of intersection.

1 mark

- d. Let $f : [0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^3 + b$, and

$$g : [b, \infty) \rightarrow \mathbb{R}, g(x) = (x - b)^{\frac{1}{3}}.$$

Find the value of b such that the graphs of $y = f(x)$ and $y = g(x)$ have a common tangent at a single point of intersection.

2 marks

END OF SAC 1b

Mathematical Methods formulas

Mensuration

area of a trapezium	$\frac{1}{2}(a+b)h$	volume of a pyramid	$\frac{1}{3}Ah$
curved surface area of a cylinder	$2\pi rh$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$		

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$		
$\frac{d}{dx}((ax+b)^n) = an(ax+b)^{n-1}$	$\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c, n \neq -1$		
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$		
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e(x) + c, x > 0$		
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$		
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$		
$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$			
product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$		

Probability

$\Pr(A) = 1 - \Pr(A')$		$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$	
$\Pr(A B) = \frac{\Pr(A \cap B)}{\Pr(B)}$			
mean	$\mu = E(X)$	variance	$\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

Probability distribution		Mean	Variance
discrete	$\Pr(X = x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$

Sample proportions

$\hat{p} = \frac{X}{n}$		mean	$E(\hat{P}) = p$
standard deviation	$\text{sd}(\hat{P}) = \sqrt{\frac{p(1-p)}{n}}$	approximate confidence interval	$\left(\hat{p} - z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right)$